Chemical Science

EDGE ARTICLE

Check for updates

Cite this: Chem. Sci., 2024, 15, 19037

O All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 5th September 2024 Accepted 23rd October 2024

DOI: 10.1039/d4sc06005a

rsc.li/chemical-science

Introduction

During the last decades, C-H activation has surfaced as a powerful and sustainable tool in modern organic synthesis.¹⁻⁶ In this context, ruthenium has emerged as a privileged metal for this type of transformation.^{7,8} Particularly, the direct aryl addition to C-C double or triple bonds via hydroarylation turned out to be an attractive and highly atom-economic approach for the direct functionalization of aromatic C(sp²)-H bonds. Different approaches were developed to realize the transition metal-catalyzed anti-Markovnikov hydroarylation of unactivated olefins. Prominent transition metal-catalysts for this transformation⁹ are primarily based on ruthenium¹⁰⁻¹⁵ – with pioneering contributions by Lewis¹⁶ and Murai¹⁷ - rhodium¹⁸⁻²³ or iridium.²⁴⁻²⁹ However, these processes are of limited utility for late-stage functionalization, as they require elevated reaction temperatures. For the same reason, the degree of C-H bond functionalization can be difficult to control, as overfunctionalization is observed in some cases. To overcome this challenge, Yoshikai³⁰⁻³² reported on the cobalt-catalyzed hydroarylation of olefins at room temperature. Although those reactions were performed at reduced temperature and therefore prevented the formation of the bis-alkylated product, they required a substoichiometric amount of Grignard reagents which translates into reduced functional

^aWöhler Research Institute for Sustainable Chemistrv (WISCh). Georg-August-Universität, Tammannstraße 2, 37077 Göttingen, Germany. E-mail: Lutz.Ackermann@chemie.uni-goettingen.de

Visible light-induced ruthenium(II)-catalyzed hydroarylation of unactivated olefins[†]

Sven Trienes, 🗓 ‡^{ab} Stéphane Golling, ‡^a Matthew H. Gieuw, ^a Marco Di Matteo^a and Lutz Ackermann D*ab

Hydroarylation reactions have emerged as a valuable tool for the direct functionalization of C-H bonds with ideal atom economy. However, common catalytic variants for these transformations largely require harsh reaction conditions, which often translate into reduced selectivites. In contrast, we herein report on a photo-induced hydroarylation of unactivated olefins at room temperature employing a readily available ruthenium(II) catalyst. Our findings include high position- and regio-selectivity and remarkable tolerance of a wide range of functional groups, which further enabled the late-stage diversification.

> group tolerance, and hence potentially limits viable latestage transformations of complex molecules (Scheme 1A).

> To circumvent the harsh reaction conditions represented either by high reaction temperatures or Grignard reagents, we envisioned that a photochemical approach could realize the desired transformation under significantly milder conditions. Additionally, visible light irradiation appears as sustainable and easily tunable driver for chemical reactions, comparable to electrochemistry,³³⁻³⁶ that has proven to be highly efficient in organic chemistry.³⁷⁻⁴³ Among a variety of different photochemical approaches, dual-catalytic systems with a transition metal-catalyst and an exogeneous photosensitizer, often



Scheme 1 (A) Previous work: metal-catalyzed anti-Markovnikov hydroarylation of olefins; (B) this work: photo-induced rutheniumcatalyzed hydroarylation of non-activated olefins at room temperature.

View Article Online View Journal | View Issue

ROYAL SOCIETY

OF CHEMISTRY

^bDZHK (German Centre for Cardiovascular Research), Potsdamer Straße 58, 10875 Berlin, Germany

[†] Electronic supplementary information (ESI) available. See DOI: https://doi.org/10.1039/d4sc06005a

[‡] These authors contributed equally to this work.

based on ruthenium or iridium, showed remarkable efficiency. Alternatively, in some cases the transition metalcatalyst itself can show interesting photophysical properties, thereby superseding the addition of a separate photocatalyst. In the context of photochemical C-H functionalization, both with and without an exogenous photocatalyst, transformations utilizing transition metals,44,45 such as palladium,⁴⁶⁻⁵¹ rhodium,^{52,53} copper⁵⁴⁻⁵⁶ or iron⁵⁷ have been exploited. Another metal with remarkable synergistic effects of merging photochemistry and C-H activation is ruthenium.58 Within our program on photoinduced, rutheniumcatalyzed C-H functionalizations,59-63 we wondered whether a photo-induced ruthenium-catalyzed hydroarylation of nonactivated olefins would indeed be viable at room temperature (Scheme 1B).

Results and discussion

We commenced our studies by probing reaction conditions for the envisioned photo-enabled, room-temperature rutheniumcatalyzed hydroarylation of olefins with ketimine 1a as model substrate. [Ru(OAc)₂(p-cymene)] turned out to be the catalyst of choice for this transformation, affording the corresponding alkylated ketone 2a in 85% yield after hydrolysis (Table 1, entry 1). Other ruthenium(II) catalysts, such as $[RuCl_2(p-cymene)]_2$ or $[Ru(O_2CMes)_2(p-cymene)]$, allowed the formation of the desired product 2a, albeit with lower efficacy (Table 1, entries 2 and 3). Several other arene-containing ruthenium-complexes were also employed giving comparable efficacy.64 The biscationic ruthenium-aqua complex65 $[Ru(H_2O)(t-BuCN)_5][BF_4]_2$ gave low yields, also in the presence of KOAc (Table 1, entry 4). Likewise, the arene-free, biscationic complex⁶⁶ [Ru(MeCN)₆][BF₄]₂ failed to furnish the desired product 2a in the absence of light, highlighting the unique features of the metallaphotocatalysis (Table 1, entry 5). $RuCl_3 \cdot 3H_2O$ and $Ru_3(CO)_{12}$ failed to give the desired product (Table 1, entries 6, 7). Additional carboxylate sources⁶⁷ did not improve the catalytic performance (Table 1, entries 8, 9). Finally, a similar result was obtained when the p-methoxyphenyl (PMP) moiety was substituted by a 3,4,5-trimethoxyphenyl group (TMP) (Table 1, entry 10). Control experiments were conducted, leading to a complete inhibition of reactivity in the dark or without the ruthenium catalyst, mirroring their essential role in the catalysis (Table 1, entry 11).

Substrate scope evaluation

With the optimized conditions in hand, we explored the versatility of the direct hydroarylation of 1-octene with various ketimines **1** (Scheme 2A). Initially, the synthetic utility was successfully demonstrated by upscaling our standard reaction on 1.2 mmol scale yielding the alkylated compound **2a** in 90% yield. Subsequently, a wide range of substituted ketimines **1** were subjected to our reaction conditions to evaluate the functional group compatibility. Noteworthy, numerous functionalities were well tolerated, and the mono-alkylated product was selectively obtained in all cases. Different alkyl substituents in

Table 1 Optimized conditions and deviation effects^a



^{*a*} Reaction conditions: **1a** (0.20 mmol), 1-octene (0.60 mmol), [Ru(OAc)₂(*p*-cymene)] (10 mol%), 1,4-dioxane (1.0 mL), room temperature (30–35 °C). Subsequent hydrolysis with 1 M HCl (3.0 mL). Detailed experimental procedures are shown in the ESI. ^{*b*} Yields of isolated product. Yields in parentheses determined by ¹H NMR using an internal standard. ^{*c*} 24 h, 30 °C, in the dark. ^{*d*} Using [RuCl₂(*p*cymene)]₂ as catalyst. PMP = *p*-methoxyphenyl.

the para-position (2b-2e) or a phenyl-group (2f) afforded the desired compounds in excellent yields. Additionally, the use of various electron donating groups such as methoxy (2g), thioether (2h), dimethylamine (2i), 1,3-dioxole (2j) or mesylate (2k) led to the targeted products in high yields. It is noteworthy that only one isomer was observed in the case of 1,3-dioxole derivative (2j). Other interesting functionalities, such as methyl ester (21), *p*-fluoro (2m), *p*-chloro (2n), *p*-trifluoromethyl (20) or even *p*-mesyl (2**p**) were well tolerated. Subsequently, the use of ketimines bearing a heterocycle in para-position were also smoothly transformed into the corresponding hydroarylation products with yields up to 95%, demonstrating the excellent functional group tolerance of this approach (2q-2s). Finally, substituents in the meta-position of the aromatic ring were also tolerated, and only one constitutional isomer was obtained (2t, 2u) in 59% and 74% yield, respectively.

Then, the effect of substituents on the alkene was explored (Scheme 2B). A longer alkyl chain did not impact the reactivity as ketone 2v was obtained in 95% yield. The use of an alkene bearing a phenyl ring led to an excellent yield of 98% (2w). A sensitive bromide derivative (2x) and a free alcohol (2y) were also compatible and afforded the corresponding compounds in good yields. Different silyl groups (2z-ab) proved to be suitable as well. Similarly, a boronic acid masking group (MIDA) was well tolerated (2ac). Finally, terminal alkenes incorporated in important biorelevant compounds were successfully transformed into the desired hydroarylation products. Therefore, the reaction was performed in the presence of ibuprofen (2ad) and



Scheme 2 Substrate scope for various ketimines 1 with 1-octene. Reaction conditions: 1 (0.20 mmol), 1-octene (0.60 mmol), $[Ru(OAc)_2(p-cymene)]$ (10 mol%), 1,4-dioxane (1.0 mL), room temperature (30–35 °C). Detailed experimental procedures are shown in the ESI.† (a) Subsequent hydrolysis with 1 M HCl (3.0 mL). (b) 96 h reaction time. (c) 1.6 equiv. of 1-octene.

cholesterol (2ae) derivatives giving the targeted products in acceptable to excellent yields.

Additionally, the ruthenium-catalyzed hydroarylation proved viable for different directing groups (Scheme 2C). The use of a bis(*p*-methoxyphenyl)imine led to the corresponding monoalkylated product **3a** in a satisfactory yield of 59%. A good yield of 58% was observed for bis(diethylamino phenyl)imine **3b**. Moreover, a cyclic imine was employed to furnish the corresponding dihydroisoquinoline **3c** in moderate yield. Furthermore, heteroarenes also enabled the photo-induced hydroarylation. Therefore, 2-phenylpyridine gave access to the alkylated product **3d** in 80% yield. The mild conditions of the photo-induced C–H activation translated into improved monoselectivities as compared to thermal mode of action.^{10,64} A reduced efficacy was observed for a pyrimidyl coordinating group (**3e**) and an aldimine (**3f**).

Mechanistic investigations

To gain insights into the working mode of the metallaphotocatalysis, mechanistic studies were conducted. Initially, the role of the blue light irradiation was investigated in detail with an on-off experiment (Scheme 3A). Here, the essential role of the light was confirmed. Next, the reaction was monitored *via* ¹H NMR spectroscopy, with free *p*-cymene being observed.⁵⁹ The addition of a substoichiometric amount of *p*-cymene to the reaction mixture inhibited catalysis.⁶⁴ Additionally, a quantum yield⁶⁴ of 1.2% was in good agreement. Finally, UV-spectroscopy allowed us to identify that both the ruthenium complex and ruthenacycle **Ru-1** show a strong absorbance of light in the range between 380–460 nm. Analysis of the reaction mixture before and after irradiation with blue light also show the formation of new species (Scheme 3C).⁶⁴

To gain further insights into the potential mode of action, an intermolecular competition experiment was performed between electron-rich ketimine **1b** and electron-deficient ketimine **1o** (Scheme 4A). After 24 h, the trifluoromethyl-substituted compound **2o** was obtained in 30% yield against 5% for the methyl-substituted product **2b**. Consequently, the C-H activation may occur preferentially on electron-poor arenes with weaker C-H bonds indicating a carboxylate-assisted



Scheme 3 Key mechanistic findings on the role of the blue light. Detailed experimental procedures are shown in ESI.† (a) Relative yields referring to the initial amount of phenylpyridine (0.2 mmol) or $[Ru(OAc)_2(p-cymene)]$ (0.02 mmol). RM = reaction mixture.



Scheme 4 Key mechanistic findings.

cycloruthenation. This tendency was also observed during the evaluation of the substrate scope of the ketimine 1, as slightly higher yields were obtained in the case of electron-poor ketimines (Scheme 2). To pursue our studies, the independently synthesized ruthenacycle Ru-1 was employed as catalyst under our standard conditions. Notably, this complex was only effective in the presence of a catalytic amount of potassium acetate (20 mol%), further substantiating that a carboxylate-assisted ruthenation is involved in the catalytic cycle (Scheme 4B). Interestingly, product formation was not observed in the dark with complex Ru-1 as catalyst, being suggestive of lightindependent cyclometallation, along with a light-induced activation - arguably through decoordination of the *p*-cymene ligand.⁶⁴ Next, deuterium-labelling experiments were carried out. When ketimine $1a \cdot d_5$ was treated with $[Ru(OAc)_2(p \cdot cym \cdot d_5)]$ ene)] under blue light irradiation, a significant H/D exchange was observed (Scheme 4C). This result points to the reversibility of the cleavage of the ortho C-H bond. The hydroarylation reaction between 1a-d₅ and dimethyl(phenyl)(vinyl)silane was also examined. Deuterium incorporation was only observed at the β -position of the hydroarylation product **2ac-d**₅ while no deuterium was introduced at the α -position (Scheme 4C). This observation is in contrast to rhodium(1)-catalyzed alkylations of aromatic amides reported by Chatani whose mechanistic proposal was considered to undergo a carbene mechanism which was confirmed by deuterium labelling experiments and DFT calculations.68 The carbene mechanism was therefore considered unlikely for the photo-induced hydroarylation.

(0)



Scheme 5 Proposed catalytic cycle.

Based on our experiments and literature precedents,⁵⁹⁻⁶¹ a plausible mechanism for the photo-enabled hydroarylation of olefins involves an initial C–H ruthenation of ketimine **1** to generate ruthenacycle **A** (Scheme 5). After light-induced *p*-cymene dissociation, the catalytic active species **B** is formed. Subsequently, the coordination of the olefin *via* intermediate **C**, along with migratory insertion into the Ru–C bond forms intermediate **D**. Then, acetic acid enables proto-demetallation to generate the species **E**. Finally, ligand exchange with another equivalent of the starting material releases the desired product **2a'**, while regenerating catalyst **B** through cyclometalation.

Conclusions

In summary, we reported on versatile C–H alkylations enabled by photo-induced ruthenium(π)-catalyzed hydroarylation of non-activated alkenes. In sharp contrast to established catalytic systems, the carboxylate-assisted ruthenium(π) catalysis proved efficient under exceedingly mild reaction conditions, namely at ambient temperature, enabled by visible light irradiation. The ruthenaphotocatalysis was easily conducted on gram-scale and its robustness was mirrored by a broad functional group tolerance.

Data availability

The data supporting this article have been uploaded as part of the ESI.†

Author contributions

Conceptualization, L. A.; funding acquisition, L. A.; investigation S. T., S. G., M. H. G., M. D. M., methodology, S. T., S. G.; resources, L. A., supervision, L. A.; writing – original draft, S. T., S. G., L. A., writing – reviewing & editing, all authors.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

The authors gratefully acknowledge generous support by the DZHK, the DFG (Gottfried Wilhelm Leibniz award to L.A.) and the ERC Advanced Grant no. 101021358 (L.A.).

Notes and references

- T. Rogge, N. Kaplaneris, N. Chatani, J. Kim, S. Chang, B. Punji, L. L. Schafer, D. G. Musaev, J. Wencel-Delord, C. A. Roberts, R. Sarpong, Z. E. Wilson, M. A. Brimble, M. J. Johansson and L. Ackermann, C-H activation, *Nat. Rev. Methods Primers*, 2021, 1, 43.
- 2 S. Rej, Y. Ano and N. Chatani, Bidentate Directing Groups: An Efficient Tool in C–H Bond Functionalization Chemistry for the Expedient Construction of C–C Bonds, *Chem. Rev.*, 2020, **120**, 1788–1887.
- 3 P. Gandeepan, T. Müller, D. Zell, G. Cera, S. Warratz and L. Ackermann, 3d Transition Metals for C-H Activation, *Chem. Rev.*, 2019, **119**, 2192–2452.
- 4 C. Sambiagio, D. Schönbauer, R. Blieck, T. Dao-Huy, G. Pototschnig, P. Schaaf, T. Wiesinger, M. F. Zia, J. Wencel-Delord, T. Besset, B. U. W. Maes and M. Schnürch, A comprehensive overview of directing groups applied in metal-catalysed C–H functionalisation chemistry, *Chem. Soc. Rev.*, 2018, 47, 6603–6743.
- 5 J. A. Leitch and C. G. Frost, Regioselective Transition-Metal-Catalyzed C–H Functionalization of Anilines, *Synthesis*, 2018, **50**, 2693–2706.
- 6 J. Wencel-Delord and F. Glorius, C–H bond activation enables the rapid construction and late-stage diversification of functional molecules, *Nat. Chem.*, 2013, 5, 369–375.
- 7 J. A. Leitch and C. G. Frost, Ruthenium-catalysed σ -activation for remote *meta*-selective C–H functionalisation, *Chem. Soc. Rev.*, 2017, **46**, 7145–7153.
- 8 P. B. Arockiam, C. Bruneau and P. H. Dixneuf, Ruthenium(II)-Catalyzed C-H Bond Activation and Functionalization, *Chem. Rev.*, 2012, **112**, 5879–5918.
- 9 Z. Dong, Z. Ren, S. J. Thompson, Y. Xu and G. Dong, Transition-Metal-Catalyzed C-H Alkylation Using Alkenes, *Chem. Rev.*, 2017, **117**, 9333-9403.
- 10 M. Schinkel, I. Marek and L. Ackermann, Carboxylate-Assisted Ruthenium(II)-Catalyzed Hydroarylations of Unactivated Alkenes through C-H Cleavage, *Angew. Chem.*, *Int. Ed.*, 2013, **52**, 3977–3980.

- 11 G. Rouquet and N. Chatani, Ruthenium-catalyzed *ortho*-C-H bond alkylation of aromatic amides with α,β-unsaturated ketones *via* bidentate-chelation assistance, *Chem. Sci.*, 2013, 4, 2201–2208.
- 12 L. Ackermann, S. I. Kozhushkov and D. S. Yufit, Ruthenium-Catalyzed Hydroarylation of Methylenecyclopropanes through C-H Bond Cleavage: Scope and Mechanism, *Chem.-Eur. J.*, 2012, **18**, 12068–12077.
- 13 M.-O. Simon, G. Ung and S. Darses, Tandem Catalysis: Alcohol Oxidation and C–C Bond Formation *via* C–H Bond Activation, *Adv. Synth. Catal.*, 2011, **353**, 1045–1048.
- 14 S. I. Kozhushkov, D. S. Yufit and L. Ackermann, Ruthenium-Catalyzed Hydroarylations of Methylenecyclopropanes: Mild C-H Bond Functionalizations with Conservation of Cyclopropane Rings, *Org. Lett.*, 2008, **10**, 3409–3412.
- 15 R. Martinez, R. Chevalier, S. Darses and J.-P. Genet, A Versatile Ruthenium Catalyst for C–C Bond Formation by C–H Bond Activation, *Angew. Chem., Int. Ed.*, 2006, **45**, 8232–8235.
- 16 L. N. Lewis and J. F. Smith, Catalytic carbon-carbon bond formation via ortho-metalated complexes, J. Am. Chem. Soc., 1986, 108, 2728–2735.
- 17 S. Murai, F. Kakiuchi, S. Sekine, Y. Tanaka, A. Kamatani, M. Sonoda and N. Chatani, Efficient catalytic addition of aromatic carbon-hydrogen bonds to olefins, *Nature*, 1993, 366, 529–531.
- 18 K. Shibata, T. Yamaguchi and N. Chatani, Rhodium-Catalyzed Alkylation of C-H Bonds in Aromatic Amides with Styrenes via Bidentate-Chelation Assistance, *Org. Lett.*, 2015, **17**, 3584-3587.
- 19 J. Kwak, Y. Ohk, Y. Jung and S. Chang, Rollover Cyclometalation Pathway in Rhodium Catalysis: Dramatic NHC Effects in the C–H Bond Functionalization, *J. Am. Chem. Soc.*, 2012, **134**, 17778–17788.
- 20 C.-H. Jun, C. W. Moon, J.-B. Hong, S.-G. Lim, K.-Y. Chung and Y.-H. Kim, Chelation-Assisted RhI-Catalyzed *ortho*-Alkylation of Aromatic Ketimines or Ketones with Olefins, *Chem.-Eur. J.*, 2002, **8**, 485–492.
- 21 C.-H. Jun, J.-B. Hong, Y.-H. Kim and K.-Y. Chung, The Catalytic Alkylation of Aromatic Imines by Wilkinson's Complex: The Domino Reaction of Hydroacylation and *ortho*-Alkylation, *Angew. Chem., Int. Ed.*, 2000, **39**, 3440–3442.
- 22 Y.-G. Lim, J.-B. Kang and Y. H. Kim, Regioselective alkylation of 2-phenylpyridines with terminal alkenes *via* C–H bond activation by a rhodium catalyst, *J. Chem. Soc., Perkin Trans.* 1, 1996, 2201–2206.
- 23 Y.-G. Lim, Y. H. Kim and J.-B. Kang, Rhodium-catalysed regioselective alkylation of the phenyl ring of 2-phenylpyridines with olefins, *J. Chem. Soc., Chem. Commun.*, 1994, 2267–2268.
- 24 J. Kim, S.-W. Park, M.-H. Baik and S. Chang, Complete Switch of Selectivity in the C–H Alkenylation and Hydroarylation Catalyzed by Iridium: The Role of Directing Groups, *J. Am. Chem. Soc.*, 2015, **137**, 13448–13451.
- 25 T. Shibata and T. Shizuno, Iridium-Catalyzed Enantioselective C-H Alkylation of Ferrocenes with Alkenes

Using Chiral Diene Ligands, Angew. Chem., Int. Ed., 2014, 53, 5410-5413.

- 26 S. Pan, N. Ryu and T. Shibata, Iridium(I)-Catalyzed Direct C–H Bond Alkylation of the C-7 Position of Indolines with Alkenes, *Adv. Synth. Catal.*, 2014, **356**, 929–933.
- 27 S. Pan, N. Ryu and T. Shibata, Ir(I)-Catalyzed C–H Bond Alkylation of C2-Position of Indole with Alkenes: Selective Synthesis of Linear or Branched 2-Alkylindoles, *J. Am. Chem. Soc.*, 2012, **134**, 17474–17477.
- 28 K. Tsuchikama, M. Kasagawa, Y.-K. Hashimoto, K. Endo and T. Shibata, Cationic iridium–BINAP complex-catalyzed addition of aryl ketones to alkynes and alkenes via directed C–H bond cleavage, *J. Organomet. Chem.*, 2008, 693, 3939–3942.
- 29 R. Dorta and A. Togni, Addition of the *ortho*-C–H bonds of phenol across an olefin catalysed by a chiral iridium(I) diphosphine complex, *Chem. Commun.*, 2003, 760–761.
- 30 W. Xu and N. Yoshikai, Highly Linear Selective Cobalt-Catalyzed Addition of Aryl Imines to Styrenes: Reversing Intrinsic Regioselectivity by Ligand Elaboration, *Angew. Chem., Int. Ed.*, 2014, **53**, 14166–14170.
- 31 K. Gao and N. Yoshikai, Regioselectivity-Switchable Hydroarylation of Styrenes, *J. Am. Chem. Soc.*, 2011, **133**, 400–402.
- 32 K. Gao and N. Yoshikai, Cobalt–Phenanthroline Catalysts for the *ortho* Alkylation of Aromatic Imines under Mild Reaction Conditions, *Angew. Chem., Int. Ed.*, 2011, **50**, 6888–6892.
- 33 Y.-K. Xing, Z.-H. Wang, P. Fang, C. Ma and T.-S. Mei, Divergent synthesis of aryl amines and dihydroquinazolinones via electrochemistry-enabled rhodium-catalyzed C-H functionalization, *Sci. China Chem.*, 2023, **66**, 2863–2870.
- 34 Y. Zhang, Z. Cai, S. Warratz, C. Ma and L. Ackermann, Recent advances in electrooxidative radical transformations of alkynes, *Sci. China Chem.*, 2023, **66**, 703–724.
- 35 M. Yan, Y. Kawamata and P. S. Baran, Synthetic Organic Electrochemical Methods Since 2000: On the Verge of a Renaissance, *Chem. Rev.*, 2017, **117**, 13230–13319.
- 36 Y. Wang, S. Dana, H. Long, Y. Xu, Y. Li, N. Kaplaneris and
 L. Ackermann, Electrochemical Late-Stage
 Functionalization, *Chem. Rev.*, 2023, **123**, 11269–11335.
- 37 P. Bellotti, H.-M. Huang, T. Faber and F. Glorius, Photocatalytic Late-Stage C-H Functionalization, *Chem. Rev.*, 2023, **123**, 4237–4352.
- 38 J. Großkopf, T. Kratz, T. Rigotti and T. Bach, Enantioselective Photochemical Reactions Enabled by Triplet Energy Transfer, *Chem. Rev.*, 2022, **122**, 1626–1653.
- 39 C.-S. Wang, P. H. Dixneuf and J.-F. Soulé, Photoredox Catalysis for Building C–C Bonds from C(sp²)–H Bonds, *Chem. Rev.*, 2018, **118**, 7532–7585.
- 40 L. Marzo, S. K. Pagire, O. Reiser and B. König, Visible-Light Photocatalysis: Does It Make a Difference in Organic Synthesis?, *Angew. Chem., Int. Ed.*, 2018, 57, 10034–10072.
- 41 J. Twilton, C. Le, P. Zhang, M. H. Shaw, R. W. Evans and D. W. C. MacMillan, The merger of transition metal and photocatalysis, *Nat. Rev. Chem*, 2017, **1**, 0052.

- 42 N. A. Romero and D. A. Nicewicz, Organic Photoredox Catalysis, *Chem. Rev.*, 2016, **116**, 10075–10166.
- 43 J. M. R. Narayanam and C. R. J. Stephenson, Visible light photoredox catalysis: applications in organic synthesis, *Chem. Soc. Rev.*, 2011, **40**, 102–113.
- 44 K. P. S. Cheung, S. Sarkar and V. Gevorgyan, Visible Light-Induced Transition Metal Catalysis, *Chem. Rev.*, 2022, **122**, 1543–1625.
- 45 R. Kancherla, K. Muralirajan, A. Sagadevan and M. Rueping, Visible Light-Induced Excited-State Transition-Metal Catalysis, *Trends Chem.*, 2019, **1**, 510–523.
- 46 Y.-C. Hsu, V. C. C. Wang, K.-C. Au-Yeung, C.-Y. Tsai, C.-C. Chang, B.-C. Lin, Y.-T. Chan, C.-P. Hsu, G. P. A. Yap, T. Jurca and T.-G. Ong, One-Pot Tandem Photoredox and Cross-Coupling Catalysis with a Single Palladium Carbodicarbene Complex, *Angew. Chem., Int. Ed.*, 2018, 57, 4622–4626.
- 47 W.-J. Zhou, G.-M. Cao, G. Shen, X.-Y. Zhu, Y.-Y. Gui, J.-H. Ye, L. Sun, L.-L. Liao, J. Li and D.-G. Yu, Visible-Light-Driven Palladium-Catalyzed Radical Alkylation of C-H Bonds with Unactivated Alkyl Bromides, *Angew. Chem., Int. Ed.*, 2017, 56, 15683–15687.
- 48 D. Kalyani, K. B. McMurtrey, S. R. Neufeldt and M. S. Sanford, Room-Temperature C-H Arylation: Merger of Pd-Catalyzed C-H Functionalization and Visible-Light Photocatalysis, J. Am. Chem. Soc., 2011, 133, 18566–18569.
- 49 A. Saha, S. Guin, W. Ali, T. Bhattacharya, S. Sasmal, N. Goswami, G. Prakash, S. K. Sinha, H. B. Chandrashekar, S. Panda, S. S. Anjana and D. Maiti, Photoinduced Regioselective Olefination of Arenes at Proximal and Distal Sites, *J. Am. Chem. Soc.*, 2022, 144, 1929–1940.
- 50 K. Pak Shing Cheung, J. Fang, K. Mukherjee, A. Mihranyan and V. Gevorgyan, Asymmetric intermolecular allylic C–H amination of alkenes with aliphatic amines, *Science*, 2022, **378**, 1207–1213.
- 51 S. Sarkar, K. P. S. Cheung and V. Gevorgyan, Recent Advances in Visible Light Induced Palladium Catalysis, *Angew. Chem., Int. Ed.*, 2024, **63**, e202311972.
- 52 J. Tanaka, Y. Nagashima, A. J. Araujo Dias and K. Tanaka, Photo-Induced *ortho*-C-H Borylation of Arenes through In Situ Generation of Rhodium(II) Ate Complexes, *J. Am. Chem. Soc.*, 2021, 143, 11325–11331.
- 53 J. Thongpaen, R. Manguin, V. Dorcet, T. Vives, C. Duhayon, M. Mauduit and O. Baslé, Visible Light Induced Rhodium(I)-Catalyzed C–H Borylation, *Angew. Chem., Int. Ed.*, 2019, 58, 15244–15248.
- 54 S. Trienes, J. Xu and L. Ackermann, Photoinduced C-H arylation of 1,3-azoles *via* copper/photoredox dual catalysis, *Chem. Sci.*, 2024, **15**, 7293–7299.
- 55 C. Li, B. Chen, X. Ma, X. Mo and G. Zhang, Light-Promoted Copper-Catalyzed Enantioselective Alkylation of Azoles, *Angew. Chem., Int. Ed.*, 2021, **60**, 2130–2134.

- 56 F. Yang, J. Koeller and L. Ackermann, Photoinduced Copper-Catalyzed C-H Arylation at Room Temperature, *Angew. Chem., Int. Ed.*, 2016, **55**, 4759–4762.
- 57 A. M. Messinis, T. von Münchow, M. Surke and L. Ackermann, Room temperature photo-promoted ironcatalysed arene C–H alkenylation without Grignard reagents, *Nat. Catal.*, 2024, 7, 273–284.
- 58 X. Wang, G. An and G. Li, Visible Light-Induced Ru-Catalyzed Directing-Group-Assisted C-H Activation of Arenes, *ChemCatChem*, 2024, **16**, e202301523.
- 59 Y. Wang, S. Chen, X. Chen, A. Zangarelli and L. Ackermann, Photo-Induced Ruthenium-Catalyzed Double Remote C(sp²)-H/C(sp³)-H Functionalizations by Radical Relay, *Angew. Chem., Int. Ed.*, 2022, **61**, e202205562.
- 60 J. Struwe, K. Korvorapun, A. Zangarelli and L. Ackermann, Photo-Induced Ruthenium-Catalyzed C–H Benzylations and Allylations at Room Temperature, *Chem.–Eur. J.*, 2021, 27, 16237–16241.
- 61 K. Korvorapun, J. Struwe, R. Kuniyil, A. Zangarelli, A. Casnati, M. Waeterschoot and L. Ackermann, Photo-Induced Ruthenium-Catalyzed C-H Arylations at Ambient Temperature, *Angew. Chem., Int. Ed.*, 2020, **59**, 18103–18109.
- 62 P. Gandeepan, J. Koeller, K. Korvorapun, J. Mohr and L. Ackermann, Visible-Light-Enabled Ruthenium-Catalyzed *meta*-C–H Alkylation at Room Temperature, *Angew. Chem., Int. Ed.*, 2019, **58**, 9820–9825.
- 63 T. Michiyuki, I. Maksso and L. Ackermann, Photo-Induced Ruthenium-Catalyzed C–H Arylation Polymerization at Ambient Temperature, *Angew. Chem., Int. Ed.*, 2024, **63**, e202400845.
- 64 For detailed information, see the ESI.†
- 65 G. McArthur, J. H. Docherty, M. D. Hareram, M. Simonetti, I. J. Vitorica-Yrezabal, J. J. Douglas and I. Larrosa, An airand moisture-stable ruthenium precatalyst for diverse reactivity, *Nat. Chem.*, 2024, **16**, 1141–1150.
- 66 C. C. Underwood, B. S. Stadelman, M. L. Sleeper and J. L. Brumaghim, Synthesis and electrochemical characterization of $[Ru(NCCH_3)_6]^{2+}$, tris(acetonitrile) tris(pyrazolyl)borate, and tris(acetonitrile) tris(pyrazolyl) methane ruthenium(II) complexes, *Inorg. Chim. Acta*, 2013, **405**, 470–476.
- 67 L. Ackermann, Carboxylate-Assisted Transition-Metal-Catalyzed C–H Bond Functionalizations: Mechanism and Scope, *Chem. Rev.*, 2011, **111**, 1315–1345.
- 68 T. Yamaguchi, S. Natsui, K. Shibata, K. Yamazaki, S. Rej, Y. Ano and N. Chatani, Rhodium-Catalyzed Alkylation of C–H Bonds in Aromatic Amides with Non-activated 1-Alkenes: The Possible Generation of Carbene Intermediates from Alkenes, *Chem.–Eur. J.*, 2019, **25**, 6915– 6919.