

Nanoscale Advances

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: H. Tiwari, S. Singh, R. Kumar, A. Mandal, A. Pathak, N. K. Varma, L. Kumar and V. Gautam, *Nanoscale Adv.*, 2025, DOI: 10.1039/D5NA00596E.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Title

Novel Advancements in Nanomaterials-Based Contrast Agents Across Multimodal Imaging and Theranostic Applications

Authors

Harshita Tiwari¹, Swati Singh¹, Rajiv Kumar¹, Abhijit Mandal², Abhishek Pathak³, Navin Kumar Verma^{4,5,6,7*}, Lalit Kumar⁸, Vibhav Gautam^{1*}

Affiliations

¹*Centre of Experimental Medicine and Surgery, Institute of Medical Sciences, Banaras Hindu University, Varanasi 221005, Uttar Pradesh, India*

²*Department of Radiotherapy and Radiation Medicine, Institute of Medical Sciences, Banaras Hindu University, Varanasi 221005, Uttar Pradesh, India*

³*Department of Neurology, Institute of Medical Sciences, Banaras Hindu University, Varanasi 221005, Uttar Pradesh, India*

⁴*Lee Kong Chian School of Medicine, Nanyang Technological University Singapore, Clinical Sciences Building, 11 Mandalay Road, Singapore 308232, Singapore*

⁵*Singapore Eye Research Institute, The Academia, 20 College Road, Singapore 169856, Singapore*

⁶*National Skin Centre, 1 Mandalay Road, Singapore 308205, Singapore*

⁷*Skin Research Institute of Singapore, Clinical Sciences Building, 11 Mandalay Road, Singapore 308232, Singapore*

⁸*Department of Urology, Institute of Medical Sciences, Banaras Hindu University, Varanasi 221005, Uttar Pradesh, India*

***Corresponding Authors**

Dr. Vibhav Gautam

Address

Centre of Experimental Medicine and Surgery
Institute of Medical Sciences
Banaras Hindu University, Varanasi-221005,
Uttar Pradesh, India
Email ID: vibhav.gautam4@bhu.ac.in
Contact No. +918860182113

&

Dr. Navin Kumar Verma

View Article Online
DOI: 10.1039/D5NA00596E

Address

Lee Kong Chian School of Medicine
Nanyang Technological University, Singapore
Clinical Sciences Building, 11 Mandalay Road,
Singapore 308232, Singapore
Email ID: nkverma@ntu.edu.sg

Abstract

Nanomaterials offer significant potential for non-invasive multimodal imaging due to their multifunctionality and tunable nanoscale features. Advances in their design and conjugation with organic and inorganic materials have enhanced their production and utility. Functionalizing nanoparticles (NPs) with imaging agents enables high-contrast imaging with spatial precision. Plasmonic NPs, lanthanide NPs, semiconductor-based quantum dots (QDs), and biogenic NPs have been employed as contrast agents for sensitive and specific imaging. Diseases such as cancer, neurological, gastrointestinal, and cardiovascular conditions demand early diagnosis for effective therapy. Therefore, functionalized NPs are employed to enhance molecular imaging by penetrating cells and targeting biomolecules, improving imaging modalities like positron emission tomography (PET), X-ray computed tomography (CT), near-infrared fluorescence (NIRF), magnetic resonance imaging (MRI), and photoacoustic imaging (PAI). This review highlights novel NP applications for image-guided surgery and treatment, emphasizing their role in combining imaging techniques for precision diagnostics. Challenges such as clinical translation and toxicity are discussed, underscoring the need for further research. NP-based contrast agents have emerged as an effective tool for bridging the gap between traditional diagnostics and personalized treatments, enabling real-time therapeutic monitoring and early stage theranostics.


Keywords: Nanoparticles; Multimodal Imaging; Contrast Agents; Molecular Diagnostics; Theranostics

1. Introduction

Non-invasive imaging technologies offer sophisticated visualization aimed at early-stage disease diagnosis without entering and damaging the local body tissues due to which they have drawn significant attention in both biological and medical imaging ¹. Unlike invasive methods, non-invasive imaging avoids direct contact with organs or tissues, thus reducing complications, improving repeatability, and enabling longitudinal studies. Contrast agents enhance the performance of imaging modalities by increasing the specificity and sensitivity which is essential for tracking the activity of disease and their response to certain therapeutics ². The application of diverse range of imaging tools has broadly expanded from anatomical imaging to including functional and molecular imaging, thereby enhancing the accuracy of diagnosis and effectiveness of therapeutics. The use of molecular probes has not only resulted in the generation of high-resolution images of anatomical structures, but also aided in determining



tissue permeability and organ perfusion ³. Conventional contrast agents, such as iodinated compounds and gadolinium-based agents, have long been used to enhance anatomical imaging by altering the physical properties of tissues ^{4 5}. While conventionally synthesized contrast agents have enabled high-contrast imaging at the macro level, achieving micro-level imaging required for visualizing dynamic behaviors and specific cellular functions still remains challenging. In recent years, stimuli-responsive nanoprobe capable of activating their imaging signals only in specific tumor microenvironments, such as acidic pH or hypoxic regions, have gained significant interest for improving signal-to-noise ratios and minimizing off-target effects ^{6 7}. Magnetic particle imaging has also emerged as an excellent modality exploiting the superparamagnetic properties of iron oxide NPs offering radiation-free and highly sensitive detection for quantitative theranostic applications ⁸. Other than this, hybrid nanoplateforms integrating organic, inorganic, and biological components have also played a significant role in offering multifunctionality for simultaneous multimodal imaging and combinatorial therapies ⁹. Genetically engineered and endogenously produced contrast agents have surpassed conventional ones, emerging as essential tools for probing specific molecules in pathophysiological conditions. Genetically engineered contrast agents are biologically encoded reporter proteins that are genetically modified to emit optoacoustic signals or produce chromophores that emit such signals to offer more benefits than exogenous contrast agents for prolonged duration ¹⁰. Diverse range of contrast agents such as ferritin ¹¹, luciferase ¹², fluorescent proteins ¹⁰, aquaporins ¹³, have been used across different imaging modalities for longitudinal imaging. On the other hand, endogenously produced contrast agents are designed to be synthesized inside the body by cells, often through the introduction of specific reporter genes which means that, unlike conventional exogenous agents, there is no need to repeatedly administer chemical contrast media, which can cause toxicity, immune reactions, or off-target effects. For example, reporter enzymes such as β -galactosidase, tyrosinase, or synthetic enzymes can catalyze the production or accumulation of paramagnetic or diamagnetic metabolites that alter local MRI signal intensities ^{14 15}. This provides a way to visualize gene expression or metabolic activity directly within target tissues. The use of gas vesicles (GVs) as air-filled protein nanostructures naturally produced by certain microbes like cyanobacteria or archaea is yet another good example of endogenous contrast agents. Researchers have designed novel genetically encoded acoustic reporters by transferring the genetic machinery for GV production into mammalian cell ¹⁶. These GV scatter ultrasound waves, generating strong acoustic contrast precisely at the sites where engineered cells produce them. Notably, GV are synthesized by the body cells only, there is no need for repeated administration of synthetic



microbubbles or external NP emulsions, which typically have short circulation times and may pose safety concerns^{17 18}.

Nano-sized imaging and fluorescent materials are rapidly emerging as practical revolutionary tools for improving disease diagnosis across a broad spectrum of *in vivo* imaging techniques¹⁹. These agents provide a sensitive approach for non-invasive diagnostics because of their unique surface attributes which can be fabricated with a diverse range of targeting agents capable of high-contrast imaging. Latest advancement in the versatility of nanoprobe has enabled the functioning of novel imaging tools surpassing the intrinsic drawbacks of conventional diagnostics in the field of clinical diagnostics and bio-imaging at molecular level²⁰. The electromagnetic and optical properties of a diverse range of NPs including plasmonic NPs, lanthanide NPs, semiconductor-nanocrystal based QDs and biogenic NPs have confronted a plethora of biomedical imaging-based applications (**Table 1**). The co-delivery of therapeutic and diagnostic agents into a single NP have enabled the development and application of novel theranostic tools in biomedical sector. NP-encapsulated theranostic agents hold great promise in real-time monitoring of the intracellular accumulation, targeted delivery to specific tissue/organs and controlled release of the drug, resulting in increased efficacy and decreased side-effects. For example, in a recent study a glutathione-responsive nano prodrug was designed where drug release and fluorescence signal are tightly coupled, allowing simultaneous real-time visualization of intracellular uptake and therapeutic action²¹. NPs can serve as excellent contrast agents for imaging purpose because of their small size, surface properties and ultrasensitive detection. Despite rapid progress, challenges such as long-term safety profile, bioaccumulation, and standardized large-scale synthesis still need to be addressed before wide clinical translation of NP-based contrast agents.


This article aims to provide a detailed overview of the evolving role of nanomaterials, such as plasmonic NPs, lanthanide NPs, QDs, and biogenic NPs in biomedical imaging and theranostics. Furthermore, the review emphasizes the potential of the nanomaterials for early detection of complex diseases, including neurological, cardiovascular, gastrointestinal disorders, and cancer. Specifically, we highlight the application of NPs as multimodal contrast agents in different imaging techniques like PET, X-Ray CT, NIRF, MRI and PAI to enhance the diagnostic accuracy. We emphasize ongoing research efforts to harness the potential of the nanomaterials for early detection of complex diseases. Moreover, we have also reviewed recent advancements in nanomaterial-based combinatorial image-guided therapies, promises and

perspectives on the ongoing development of early theranostics for improved patient-centred outcomes.


2. Types of Nanomaterials-Based Bioimaging Agents

2.1 Diagnostic Potential of Plasmonic NPs

Plasmonic NPs are metal-based nanoformulations, typically composed of gold, silver, or copper, that exhibit localized surface plasmon resonance (LSPR) phenomenon where conduction electrons collectively oscillate in response to light resulting in intense optical properties such as enhanced absorption and scattering, which are widely exploited in imaging and biosensing applications ²². Plasmonic NPs have revolutionized biomedical and clinical imaging by enhancing the resolution, specificity, and sensitivity of diagnostic techniques. When these NPs interact with light, their conduction electrons undergo collective oscillations, leading to significant absorption and scattering at specific resonance frequencies thereby resulting in the production of strong electromagnetic fields. The enhanced electromagnetic fields generated by LSPR can enhance the signals in imaging modalities leading to improved sensitivity and resolution ²³. These NPs have emerged as potent candidates in the field of nanotechnological sciences due to their physical and chemical properties, such as small size, high surface-area-to-volume ratio, ease of functionalization, and tunable properties. This tunability facilitates the development of multimodal imaging agents that can operate across different imaging platforms, providing comprehensive diagnostic details ²⁴. Metallic NPs are categorized as clusters or colloidal particles depending on the crystal lattice and particle size ranging from 1-100 nm ²⁵. The size of NPs is mainly governed by its synthesis technique, which can be adjusted by modifying the temperature, pH, ratio of the reactants and concentration of the solvent ²⁶. Apart from this, the type and quantity of ligands used during synthesis also play a crucial role in controlling nucleation and growth dynamics ²⁷. The unique optoelectronic and light scattering properties of these NPs make them suitable to be exploited for diagnostic and therapeutic resulting in controlled release of theranostic agent. The application of metallic NPs in different imaging modalities including PET, CT scan, ultrasound (US), surface enhanced Raman spectroscopy (SERS), PAI and MRI has led to the advancement in the real-time diagnosis and treatment of various diseases. In this context, gold NPs (AuNPs), silver NPs (AgNPs), and copper NPs (CuNPs) offer unique advantages for specific imaging modalities. In a recent research, colloidal AuNPs in conjugation with arginine-glycine-aspartic acid (RGD) peptides have been used to perform optical coherence tomography (OCT) and photoacoustic



microscopy (PAM) for the observation of choroidal neovascularization²⁸. Furthermore, thiol-capped AuNPs synthesized from Turkevich method have also displayed high contrast in spectral photon counting CT (SPCCT), providing a proof-of concept as an ideal candidate for contrast imaging of cancer²⁹. Moreover, gold nanospheres have also been used as PAI contrast agents for high contrast imaging of breast cancer³⁰. The application of AgNPs as contrast imaging agent and biosensors is also of considerable interest to many researchers. The plasmonic properties of silver is better than that of gold as it experiences lower optical losses and delivers enhanced performance in plasmonic applications³¹. There is significant demand of AgNPs in light-based nanotechnologies that involve production and regulation of light through surface plasmon resonance (SPR). Due to their optical properties influenced by extinction coefficient, size and wavelength, AgNPs are widely utilized in imaging, photocatalysis, and biosensing applications, enabling trace detection of certain elements and facilitating the study of biological interactions and *in vivo* monitoring techniques³². Recent research has reported the fabrication of silver-iron oxide NPs (AgIONPs) that binds specifically and effectively to thrombus due to which it has been exploited as PAI and NIRF bimodal contrast imaging agent and photothermal therapeutic agent for thrombosis³³. Likewise, in a computational simulation based finding it was observed that polyvinyl alcohol coated silver triangular nanoprisms were used for PAI and photothermal therapy of breast cancer in mouse models³⁴. Apart from the utilization of monometallic NPs, infusion of an additional metal often leads to enhanced performance because the introduction of intermetallic polar bonds and structural irregularities increases the number of active sites. Additionally bimetallic NPs tend to form intricate structures comprising of core-shell, hollow or porous structures that further leads to improved SPR effects³⁵. In a research, doxorubicin-functionalized bimetallic gold-core palladium-shell nanocomplex (Au@PdNDs.PEG/DOX) was designed for demonstrating theranostic effect against breast cancer enabling multimodal plasmon-based intracellular imaging alongside potent cytotoxic activity³⁶. In another study, Fe–Au core–satellite NPs synthesized via pulsed laser ablation have been investigated for their multifunctional potential, serving both as bimodal MRI and CT contrast agents, and as effective sensitizers for photothermal therapy of cancer³⁷. Likewise, polyethylene glycol (PEG)ylated AgNPs and AuNPs synthesized by one pot chemical reduction method have been used for CT imaging agent and radiosensitizing agent for the treatment of oral carcinomas³⁸. The use of bimetallic NPs in cancer therapy is still in its early stages and will require extensive future research to unlock their full potential, particularly in advancing novel formulations and expanding applications into areas like gene therapy and immunotherapy. Conclusively,



plasmonic NPs offer a unique platform for their exploitation as versatile tools due to their ability to bind with several ligands, therapeutic agents and radioisotopes, opening up the prospect of therapeutic administration and multimodal imaging. However, despite their promising applications, challenges such as toxicity, aggregation, and size-related limitations must be addressed to ensure the safe and effective use of these NPs in clinical settings. The dearth of certain noble metallic NPs may be attributed to their high price and limited availability. Despite the latest developments in the field of nano-imaging, the effect of long-term exposure of metallic NPs to patients is still unknown. Detailed research and comprehensive experimental data are required to figure out the ecological safety and biological efficacy of metallic NPs. The future of metallic NPs in clinical imaging holds great promise, particularly in the realms of theranostics, where diagnostic imaging and therapy are integrated into a single treatment regimen.

2.2 Optimization of Lanthanide NPs for Multimodal Luminescence Imaging

Lanthanides, a group of 15 rare-earth elements from atomic numbers 57 to 71, are characterized by partially filled 4f orbitals that impart them with unique optical and paramagnetic properties due to which they have emerged as versatile platforms for multimodal imaging applications. Their paramagnetic nature and the presence of unpaired 4f electrons enhance contrast in imaging which increase the relaxation of surrounding protons, improving MRI sensitivity³⁹. Moreover, these materials display exceptional photonic characteristics, including upconversion and downshifting which refer to the emission of ultraviolet or visible light and near-infrared (NIR) light, respectively, upon NIR excitation⁴⁰. This optical versatility in combination with their inherent X-ray attenuation properties, makes lanthanide NPs exceptionally suitable for integrated multimodal imaging. They are highly suitable for bioimaging due to their sharp emission peaks and long luminescence lifetimes, which allow for time-gated detection and significantly reduce background noise⁴¹. Their emission arises from electronic f–f transitions that have low absorption coefficients due to which an external chromophore is typically used to harvest excitation energy and transfer it non-radiatively to the lanthanide ion, triggering luminescence⁴². The emitted light spans a broad spectral region, ranging from ultraviolet (e.g., Gd³⁺), through visible wavelengths (e.g., Tm³⁺: blue, Tb³⁺: green, Dy³⁺: yellow, Sm³⁺: orange, Eu³⁺: red), to the near-infrared region (e.g., Pr³⁺, Nd³⁺, Ho³⁺, Er³⁺, Yb³⁺), enabling their exploitation across different imaging modalities⁴³. For example, PEGylated terbium nanorods that were designed in a recent investigation displayed high X-ray attenuation and strong green luminescence for MRI and X-ray CT imaging in mouse model⁴⁴. In another research, sodium



lanthanide tungstate-based NPs $\text{NaDy}(\text{WO}_4)_2$ and $\text{NaHo}(\text{WO}_4)_2$ were employed as bimodal contrast agents for *in vivo* CT and high-field MRI exhibiting significant biocompatibility and tumor accumulation via enhanced permeability and retention effect ⁴⁵. Recently, researchers also developed biocompatible lanthanide vanadate core-shell-shell NPs $\text{DyVO}_4@YVO_4@Nd$ -doped GdVO_4 as a multimodal system offering dual T_1 – T_2 MRI contrast and strong NIR luminescence, with optimized relaxivity and decreased quenching through layered structural design ⁴⁶. However, incorporating multiple lanthanide ions can result in undesirable energy transfer, resulting in luminescence quenching. For example, recent work on Yb^{3+} – Tm^{3+} co-doped systems shows that back energy transfer from Tm^{3+} activators to Yb^{3+} sensitizers significantly weaken the upconversion emission, and that spatially segregating these ions in core-shell-shell structure effectively reduces this quenching ⁴⁷. While many studies have displayed significant biocompatibility of lanthanide nanomaterials in animal models, comprehensive evaluations of their long-term safety profile, including aspects like water solubility, cytotoxicity, and excretion pathways, still need to be addressed ⁴⁸. The functionalization of lanthanide NPs often involves complex synthesis processes, which can affect scalability and reproducibility, posing challenges for clinical translation ⁴⁹. The existing challenges can be addressed by optimizing the structure of NP to control energy transfer pathways, precise selection of dopant concentrations to prevent quenching, and improving surface modification techniques for better biocompatibility and targeted delivery. Additionally, advancing synthesis methods for the formation of stable lanthanide NPs and integrating them with contrast agents can further enhance their effectiveness in multimodal imaging and disease diagnosis.

2.3 Emerging Applications of Semiconductor-based QDs as Contrast Agents

Semiconductor nanocrystals, commonly referred to as QDs, are nanoscale imaging probes known for their excellent optical, structural, electrical, and magnetic properties, due to which they have shown encouraging advancements in the field of non-invasive clinical imaging. The pertinent favourable properties of QDs include size-tunable fluorescence, enhanced signal brightness, photobleaching resistance, adjustable light emission, and synchronized excitation of several fluorescence colours ⁵⁰. While conventional QDs are typically composed of cadmium or lead based metals, recent advances have shifted focus toward less toxic alternatives such as copper indium sulfide (CuInS_2) ⁵¹, indium arsenide (InAs) ⁵², silver sulfide (Ag_2S) ⁵³, indium phosphide (InP) ⁵⁴, silicon nanocrystals ⁵⁵, and biogenic carbon QDs to improve biocompatibility and clinical relevance. A single light source may concurrently excite multiple



colours of QDs with slight spectrum overlapping offering substantial benefits for ultrasensitive detection of target molecule. The optical properties of QDs are largely affected by any alteration in core size, shell coating, surface chemistry and composition. The structure of the QDs has a core which is semiconducting in nature with fluorescent and optical properties. The core is coated with shell which protects it from nonradiative recombination thereby resisting photobleaching and improving the brightness and stability of QDs which is essential for high contrast and enhanced bio-imaging⁵⁶. Additionally, surface coating strategies are especially critical when using heavy-metal-based cores to reduce toxicity and prevent leaching, although these materials are gradually being replaced by non-heavy-metal-based QDs due to regulatory and safety concerns^{57 20}. Any variation in core composition and size can result in customized emission profile with a precise maximum anywhere from ultraviolet (UV) to NIR electromagnetic spectral region. The ability of QDs to be tailored for specific imaging modalities has led to their growing use in diagnostics, particularly in fluorescence imaging, multiplexed imaging, and *in vivo* tracking. The narrow emission spectra of QDs allow for precise wavelength discrimination, which is crucial in multi-colour imaging and multiplexed assays. The uptake of modified QD conjugates with multifunctional capabilities offers a significant time and cost advantage over single-colour assays which results in the identification of complex cellular proteins in patient samples. QD-labeled cells hold great promise for intricate detection and real-time monitoring of phenotypic and functional abnormalities in diseased condition offering systemized patient-centred diagnosis and treatment. For instance, recently carboxyl-modified QDs were used for image-guided high resolution detection of bone fracture by the aid of NIR-IIb fluorescence imaging⁵⁸. Lead/cadmium sulfide QDs (PbS@CdS QDs) were also used for fluorescence-guided surgical removal of tumor in a recent investigation⁵⁹. Likewise, zinc-doped silver telluride QDs (Zn: Ag₂Te QDs) were used for non-invasive imaging of cerebral vasculature of mice after brain injury⁶⁰. Moreover, a bimodal QD-based nanoprobe has also been used for fluorometric MRI of mesenchymal stem cells for the detection of adipogenic differentiation in a recent research⁶¹. QDs have also been used to broaden the horizon of multiphoton fluorescence for multiplex imaging of subcortical structures of brain⁶². Recent research has expanded beyond traditional Cd-based QDs to exploring safer semiconductor nanocrystals. In a study, researchers have designed eco-friendly, non-toxic Cd-free glyco-CuInS₂ QDs with dual visible/NIR emission and significant tumor penetration, making them effective and affordable fluorescent bio-probes for *in vivo* cancer imaging⁶³. Recent investigations have also developed silicon QDs for *in vivo* fluorescence imaging of osteosarcoma, leveraging their photoluminescence and favourable safety profile⁶⁴.



Another study has revealed that the fabrication of carbon QDs from agro waste biomass served as excellent contrast agents for fluorescence imaging because of their significant biocompatibility⁶⁵. Moreover, to address the toxicity of Cd-based QDs and weak photoluminescence of other Cd-free alternatives, a group of researchers have designed bright and biocompatible *in vivo* tumor-targeting Cd-free SiO₂@InP QDs@SiO₂ NPs by compactly embedding InP/ZnS QDs in silica, demonstrating their potential as superior fluorescent nanoprobes for bioimaging⁶⁶. Currently, QDs are being rigorously exploited for multimodal imaging because of their charge transfer property for the emission of NIR fluorescence. In addition, efforts are being made to use QDs for MRI as well. QDs can be incorporated with magnetic materials (such as iron oxide), resulting in magnetic QDs that provide dual-modal imaging (optical and magnetic). Efforts are being made to conjugate QDs semiconductor nanocrystals which exhibit comparable imaging performance without the heavy-metal-related toxicity concerns. By tagging QDs with radioisotopes, they can be used for non-invasive, whole-body imaging, which is particularly valuable for cancer diagnostics, monitoring, and treatment planning. In theranostic applications, QDs are being investigated for their ability to combine both diagnostic imaging and therapeutic functions. Functionalized QDs can deliver therapeutic agents, such as drugs or genes, to specific tissues while simultaneously enabling real-time imaging to monitor the treatment's efficacy. However, it would be challenging to fabricate specific QDs for different medical conditions surpassing the current non-specific imaging agents. The chemical stability of the core structure can be improved by refining the shell and surface coating techniques, resolving the toxicity issues that QDs continue to encounter. These enhanced NIR QDs can eventually replace existing conventional imaging modalities allowing their advancement from preclinical to clinical conditions, opening up new avenues for disease diagnosis and therapy. While QDs have remarkable potential in clinical imaging, challenges remain. One of the main concerns with QDs, especially those based on cadmium (CdSe, CdTe), is their potential toxicity. The heavy metals used in some QDs can be cytotoxic and pose a risk of accumulation in tissues. To address this, researchers have focused on developing less toxic alternatives, such as core-shell QDs, where the toxic core is encapsulated by a biocompatible shell. The use of QDs in clinical settings therefore requires careful evaluation of their safety, biocompatibility, and long-term effects. Regulatory agencies such as the FDA have stringent requirements for the clinical use of nanomaterials, and more research is needed to establish clear guidelines. Large-scale production of QDs with consistent quality for clinical applications remains another challenge. Efforts are underway to design QDs with safer core materials, such as silica, graphene oxide, and carbon-based QDs, which are less



toxic than traditional cadmium-based QDs. Enhancing the targeting capabilities of QDs through better functionalization with biomolecules will improve their specificity and reduce off-target effects. The integration of QDs with therapeutic agents could create multifunctional nanoplatforms that can be used for diagnosis, therapy, and monitoring, revolutionizing personalized medicine. Despite the above-mentioned challenges, ongoing research continues to address limitations, making QDs a promising tool for early disease diagnosis, real-time imaging, and personalized medicine. As the field progresses, the development of safer, more efficient QDs will likely pave the way for their widespread use in clinical settings.

2.4 Advancement of Biogenic NPs in Multimodal Imaging

In recent years, researchers and scientific community are becoming more interested in the quest for alternative large-scale technologies that are both economical and eco-friendly. One such technology is green synthesis, which is now being employed in applications that are both medically and ecologically acceptable. Biogenic NPs based imaging contrast agents have seen remarkable advancements in the field of multimodal imaging, offering improved specificity and versatility in visualizing biological processes. These agents leverage natural or biological materials to provide contrast in multiple imaging modalities, allowing for a more comprehensive understanding of pathophysiology of disease. Their ability to integrate with different imaging techniques has opened new avenues for more accurate diagnosis, real-time treatment monitoring, and research in various biomedical sectors. Biogenic nanomaterials have shown encouraging developments in the field of molecular imaging surpassing the limitations of conventional imaging modalities resulting in reduced toxicity and enhanced efficacy. Broadly, these nanomaterials can be categorized as either intrinsically biogenic, produced directly by living organisms or biofabricated via green synthesis routes using biological extracts as reducing and stabilizing agents ⁶⁷. Both classes are increasingly integrated into advanced imaging and theranostic platforms. For instance, AgNPs synthesized from the ethanolic leaf extract of *Zinnia elegans* were used as potent theranostic agents for non-invasive NIR-mediated imaging and cancer therapeutics ⁶⁸. Cysteamine-folic acid coated AgNPs synthesized from the leaf extract of *Coffea arabica* was used as a potent contrast agent for CT imaging of cancer cells ⁶⁹. Dextran-fabricated cerium oxide (Dex-CeO₂) NPs have been used as CT contrast agent for the imaging of gastrointestinal (GI) tract and inflammatory bowel disease (IBD) ⁷⁰. Beyond metal-based examples, biological entities themselves, such as oncolytic bacteria have been exploited for image-guided disease therapy paving the way for considerable development in the field of nanotheranostics ⁷¹. For example, *Staphylococcus*



aureus cells used for the synthesis of silver selenide QDs (Ag₂Se QDs) with catalase have been used as photoacoustic agent in bioimaging⁷². Additionally, bacterial magnetosomes have been engineered for MRI-guided tumor targeting and hyperthermia due to their high magnetization and biocompatibility⁷³. Moreover, naturally secreted exosomes have gained traction as endogenous nanoscale vesicles were employed as MRI/CT contrast agents that have demonstrated image guided photothermal therapy against cancer⁷⁴. In addition to this, plant virus NPs have also been exploited to be used as multimodal imaging contrast agents with therapeutic potential⁷⁵. Complementary to these, lipid and polymer-based nanomaterials have also been used as novel contrast agents to assemble complex imaging modalities. For instance, lipid-coated iron oxide (Fe₃O₄) NPs have been used as MRI contrast agent in a latest investigation⁷⁶. Terpolymer-lipid based manganese dioxide (MnO₂) NPs have also been used as MRI agent for enhanced tumor detection⁷⁷. Furthermore, a DNA-based lipid nanodevice was endogenously designed for high contrast imaging of miRNA in tumor cells⁷⁸. In addition to this AS1411 aptamer and RGD fabricated chitosan-based poly(lactic-co-glycolic acid) (PLGA) NPs were used for real-time imaging and co-delivery of docetaxel paving the way for novel theranostics⁷⁹. Despite being ecofriendly and sustainable, biogenic nanomaterials frequently require large scale culture and tedious synthesis methods. Even though the precise mechanisms underlying biogenic synthesis are still unreliable, ongoing research aims to shed light on this process and has revealed novel capabilities of green synthesis routes with enormous potential for plethora of biomedical applications.

3. Nanomaterials-Driven Diagnostic Advancements in Disease Detection

3.1 Nanomaterial-Based Strategies for Brain Disease Diagnosis and Therapy

The increasing prevalence of neurological diseases has emerged as a leading cause of disability and mortality, imposing a significant need to address them in near future. A large number of patients are being diagnosed with neuropathic diseases, but complete recovery remains rare due to the irreversible loss of neurons in affected tissues. Neurological diseases encompass a wide range of conditions, including brain tumors, traumatic injuries, vascular disorders, and neurodegenerative diseases. These diseases can cause irreversible damage to the central or peripheral nervous system, underscoring the critical importance of precise diagnosis and clinical evaluation. However, present diagnostic tools are relatively elementary with limited resolution and face multiple challenges in accurately locating the site of injury which significantly hampers the ability to provide precise diagnosis and therapy. Due to their unique




physicochemical attributes, substantial efforts have been dedicated to exploit nanomaterials in advancing research and clinical translation of neural disease theranostics. Nanomaterials can cross the blood–brain barrier (BBB) to accumulate selectively at targeted site, and enable multimodal imaging or controlled drug release. For instance, in a latest work, amyloid- β ($A\beta$) specific Gd^{3+} NPs ($NP@SiO_2@F-SLOH$) was used as NIR imaging agent of MRI for the successful real-time monitoring of $A\beta$ level which is a prognostic biomarker of Alzheimer's disease ⁸⁰. TAT-Polyp-QL which is a colour-convertible nanoprobe was designed for fluorescent diagnosis of Parkinson's disease with enhanced specificity and sensitivity ⁸¹. Furthermore, recently rapamycin loaded NP was used as a potent theranostic agent for improved MRI and NIR fluorescence imaging of acute ischaemic stroke facilitating precise multimodal imaging with least background interference, thereby significantly enhancing drug tracking and diagnostic precision ⁸². Beyond these examples, novel magnetic-based nanotheranostics are being explored for non-invasive brain tumor imaging and magnetic hyperthermia ⁸³. Moreover, smart nanocarriers functionalized with targeting ligands or exosomes are also emerging as strong candidates for precise delivery of RNA therapeutics and neuroprotective agents across the BBB ^{84 85}. Recent advancements have also emphasized on the development of stimuli-responsive nanomaterials to enhance brain-targeting efficiency and reduce systemic toxicity. For example, cell membrane-coated NPs, such as macrophage or neutrophil membrane camouflaged nanocarriers, have demonstrated prolonged circulation, immune evasion, and enhanced BBB penetration for precise neuroinflammatory disease imaging ⁸⁶. Moreover, efforts are underway to integrate clustered regularly interspaced short palindromic repeats/CRISPR-associated protein 9 (CRISPR/Cas9) systems with nanoscale delivery carriers such as lipid NPs and exosomes to enable gene editing directly within brain tissues with simultaneous molecular imaging for precise visualization and monitoring of therapeutic gene modulation ⁸⁷. In parallel, the development of multifunctional nanorobots and magnetic field-guided NPs offer significant control over navigation and accumulation at deep brain sites ⁸⁸. These next-generation strategies, combined with advanced imaging modalities, hold promise for early-stage, non-invasive diagnosis and personalized treatment of complex brain disorders. However, their clinical translation will require extensive *in vivo* safety validation and long-term monitoring to ensure biocompatibility and therapeutic efficacy. Conclusively, nanomaterials have the ability to cross the blood brain barrier thereby allowing significant diagnosis and therapy of neurological disorders. While the research on nanomaterials for brain disease imaging remains in its early stages, it encounters significant challenges, including biosafety and clinical translation. However, with the ongoing



interdisciplinary advancements in nanotechnology and multimodal imaging, there is strong optimism that these nanomaterials will progress from the laboratory to clinical research offering encouraging patient centred treatment options for neurological diseases.

3.2 Next-Generation Imaging Agents in Nano-Oncology


Cancer remains a significant worldwide health challenge, characterized by uncontrolled cell proliferation and metastasis, which disrupts and damage normal healthy cells. Over the past several decades, there has been significant advantage in the scientific knowledge of the molecular basis of cancer due to the rapid breakthroughs in molecular biology research. However, only a few experimental approaches have been able to withstand rigorous testing and become clinically viable. Molecular imaging is an area that has emerged from the great drive towards developing non-invasive imaging techniques to observe molecular alterations in *in vitro* and *in vivo* conditions⁸⁹. For instance, H-ferritin-nanocaged gadolinium NPs (Gd-HFn) have been designed for MRI-based ultrasensitive detection of tumor biomarkers⁹⁰. Carcinoembryonic antigen-conjugated fluorescent silica NPs (CEA-FSNs) were developed for immunofluorescence imaging of colorectal cancer serving as potent imaging agent for early diagnosis of colorectal cancer⁹¹. Glucose and casein coated ultra-small superparamagnetic iron oxide (USPIO) NPs were recently developed for their conjugation with a Cy7.5-K-8AOC-bombesin peptide for the construction of USPIO(Cy7.5)-BBN NPs. These NPs showed promising results when tested *in vivo* and *ex vivo* NIRF imaging because of their high specificity towards gastrin-releasing peptide receptors (GRPs) of pancreatic cancer⁹². Zwitterionic charge-convertible NIR cyclodextrin derivative was also fabricated with pheophorbide-conjugated ferrocene for ultra specific imaging and therapeutics of rectal cancer thus serving as a potent theranostic agent⁹³. Additionally, a methylene blue-integrated fibroblast activation protein inhibitor (FAPI) nanoprobe was also constructed for enhanced PET/CT fluorescence imaging of tumor which demonstrates its potential to serve as powerful multimodal imaging agent⁹⁴. In recent years, researchers have also focused on developing smart and stimuli-responsive nanoprobe to further improve tumor specificity and minimize off-target effects. For instance, pH-responsive polymeric nanocarriers embedded with indocyanine green (ICG) have shown promising results for real-time NIRF and PAI of acidic tumor microenvironments⁹⁵. Moreover, enzyme-activated probes, such as matrix metalloproteinase (MMP)-sensitive iron oxide nanoclusters, have been developed to enhance MRI contrast precisely at tumor sites exhibiting high MMP expression⁹⁶. Another noteworthy advancement is the integration of DNA nanotechnology with imaging, enabling programmable



nanodevices that can selectively hybridize with tumor-related microRNAs and trigger a detectable fluorescence or MRI signal ⁹⁷. Researchers are also developing hybrid nanostructures combining gold nanorods (GNRs) and QDs to achieve synergistic benefits of PAI and NIRF for intraoperative tumor margin delineation ⁹⁸. NPs-based theranostic and multimodal imaging have been extensively researched as they present several opportunities to overcome the drawbacks of conventional imaging modalities. However, there are several challenges in actual clinical translations of NPs due to their intricate nanostructures including generation of immune response, safety profile, and significant differences between imaging from actual cancer patients and the current *in vivo* animal model. Therefore, prior to its clinical application, unusual adverse effects on animal models post NP administration should be examined. The sustainable development of nano-oncology in upcoming future will continue to drive next generation of cancer detection by the aid of multimodal imaging.

3.3 Multimodal Imaging Approaches in Cardiovascular Nanomedicine

The clinical practice of cardiology exploits the versatility of a wide range of NPs to aid in accurate diagnosis, risk assessment and real-time therapeutic visualization. Most of the imaging modalities provide significant details of anatomical cardiovascular structures, however, fundamental molecular processes are yet to be reflected accurately. Cardiovascular diseases present different challenges as compared to cancer biology when it comes to multimodal imaging including small and dynamic structures of cardiovascular system, short time interval between different cardiovascular processes, blood flow and presence of pathological constituents. Recent investigations have examined the application of certain NP platforms for theranostics of cardiovascular disorders. For example, a nanoplateform (CNA35-GNR/PFP@NPs) comprising of collagen targeted and lipid NP encapsulated perfluoropentane (PFP) and GNRs was constructed for the multimodal high-resolution US, CT and PAI of myocardial fibrosis ⁹⁹. A ratiometric semiconducting polymeric NP was designed for successful PAI of pneumonia-procured atherosclerotic plaque that resulted in strong photoacoustic signaling for the accurate and specific detection of superoxide anions ¹⁰⁰. Additionally, a novel phase change material and superparamagnetic iron oxide loaded macrophage membrane-modified biomimetic nanoprobe was used as an imaging agent for multimodal imaging of autoimmune myocarditis ¹⁰¹. Moreover, silver iron oxide NPs have also been exploited for multimodal imaging and photothermal therapy (PTT) of thrombosis ³³. In addition to this, a perfluoro-crown ether payload (¹⁹F-HDL) based high-density lipoprotein-derived nanotracer was used for hot-spot multimodal imaging of myeloid cell egression in



ischaemic heart disease¹⁰². Recent progress has also focused on the development of enzyme-responsive gold nanoclusters coated with fibrin-targeting peptides that have demonstrated enhanced sensitivity in detecting early-stage thrombi via combined photoacoustic and MRI¹⁰³. Researchers have also developed ROS-sensitive polymeric micelles loaded with NIR dyes, enabling real-time imaging of oxidative stress within atherosclerotic plaques¹⁰⁴. In another notable advancement, multifunctional exosome-mimetic nanovesicles decorated with iron oxide and fluorophores have been employed for T1-weighted MRI of inflamed endothelial regions, providing improved specificity for pathological angiogenesis¹⁰⁵. Moreover, targeted lipid-polymer hybrid NPs carrying ultrasound contrast agents and PET isotopes have been tested for non-invasive assessment of myocardial ischaemia, offering both functional and molecular information in a single diagnostic session¹⁰⁶. It is clearly evident that the bioavailability of imaging agents for the detection of cardiovascular diseases has significantly enhanced by the application of NP-based multimodal imaging agents. The potential to design accessible nanotracers that provide site-specific molecular imaging has been made feasible due to the specific microenvironment of cardiovascular diseases. However, in-depth research is still required to elucidate the underlying mechanism, associated pharmacokinetics and intraorganellar uptake. Additionally, comprehensive safety assessment must be conducted prior to clinical trials to minimize the level of potential side effects.

3.4 Nanomaterials-Based Advanced Imaging Approaches for Detection of Gastrointestinal (GI) Diseases

Nanomaterials have been extensively exploited for the diagnosis of GI disorders because of their ability to be conjugated with multiple imaging modalities for ultrasensitive and specific detection. Conventional techniques like digestive endoscopy and ultrasonography are mostly used to diagnose GI disorders. However, these techniques often result in discomfort, respiratory problems, allergy, and microbial infections in patients during or post investigation. The involuntary peristalsis in GI tract and densely interconnected abdominal tissues makes it challenging for the conventional techniques to diagnose GI disorders. Therefore, most of the latest research work is inclining towards the application of NP-based diagnosis in associated with wide range of multimodal imaging and contrast agents for specific biodistribution and enhanced imaging of GI disorders. For instance, dextran coated bismuth oxide NPs (Bi₂O₃-Dex NPs) were used for improved CT imaging of inflammatory bowel disease (IBD)¹⁰⁷. Similarly, a bismuth-pectin based microgel network (Bi-GLUE) was designed for real time imaging of GI tract using X-RAY and MRI due to its significant mucoadhesive property¹⁰⁸. Apart from



this, in a novel work, gadolinium oxide carbonate and mesoporous silica coated GNRs (AuNR-SiO₂-Gd) was used for high contrast imaging using CT, MRI and PAI for the diagnosis of pancreatic ductal adenocarcinoma¹⁰⁹. The microemulsion method was used for the construction of bimodal-polymer based NP which displayed significant high resolution imaging after being integrated with NIR fluorescence imaging and PAI of GI tract¹¹⁰. An innovative one-pot solvothermal method was used for the synthesis of PEGylated BaGdF₅ NPs for high contrast MRI/CT bimodal *in vivo* imaging of GI tract¹¹¹. Recently, macrophage-membrane coated bismuth oxyiodide nanodots (BiOI@M) have been shown to provide robust X-ray/CT contrast in the GI tract with excellent biocompatibility and mucoadhesive behavior for improved lesion localization¹¹². Furthermore, PEGylated AuNPs have been demonstrated to offer size-independent, high-contrast CT imaging in murine models of ulcerative colitis¹¹³. Another promising direction involves spectral CT imaging using bismuth or rhenium-sulfide NPs, which enable dual-energy differentiation of GI lesions and reduce artefacts, as evidenced in recent rodent studies¹¹⁴. These multifunctional nanoprobe offer higher sensitivity, rapid clearance, and clinical potential. In summary, ongoing developments of NP-based imaging modalities offer potential diagnostic approach for the early detection of GI diseases. The conjugation of nanoprobe with multiple contrast agents have not only improved the specificity but have also provided a platform for the selective accumulation of theranostic agents at the targeted site. However, multiple challenges including precise targeting, guaranteed safety and cost effectiveness are yet to be confronted for successful clinical implementation of NP-based imaging agents in gastroenterology.

4. Applications of NP-Based Contrast Agents in Advanced Bioimaging Modalities


4.1 Role of Nanomaterials in PET scanning for Molecular Imaging

PET is a highly sensitive, non-invasive radionuclide imaging modality that offers great diagnostic advantage in preclinical and clinical fields due to its ability to quantitatively visualize physiological and molecular processes. PET imaging is generally required because of the development of radiotracers that can exhibit targeted action against specific biomarkers of certain disease. In this context, nanomaterials play a pivotal role by serving as vehicles for PET isotopes and enabling improved pharmacokinetics, enhanced signal retention, and precise molecular targeting¹¹⁵. Various types of biomolecules including proteins, antibodies or lipids can serve as potent radiotracers opening the avenues for PET bioimaging by the aid of nanoprobe. For instance, radiolabeled polymeric NPs are heavily utilized for evaluating their



biodistribution through PET scanning. In an interesting work, ^{89}Zr PLGA-NH₂ NPs labeled monocytetes were used for radioactive *in vivo* cellular tracking of breast tumors as well as *Staphylococcus aureus* bacterial infection using PET imaging ¹¹⁶. Moreover, anti-CD64 antibody conjugated gold and methotrexate encapsulating PLGA NPs were used for the diagnosis and therapy of rheumatoid arthritis ¹¹⁷. Ga-labeled amphiphilic polymeric NPs have been used to diagnose lymph node metastasis through PET imaging ¹¹⁸. Apart from this, lipid NPs are being exploited in improved PET based diagnosis of certain diseases. For example, ^{68/67}Ga-radiolabeled sphingolipid nanoemulsions serve as a potent nanoprobe for PET imaging of breast and lung carcinoma ¹¹⁹. A group of researchers have recently developed a novel nanocomplex (DOX@TLNPs) composed of tannic acid, 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-[methoxy(polyethylene glycol)-2000 (DSPE-PEG2k), and doxorubicin, which demonstrated significant tumor targeting, sustained drug release as confirmed by PET imaging using ^{89}Zr -labeled particles, and displayed effective anticancer activity with minimal toxicity ¹²⁰. The potential of radioactive imaging agents can be significantly highlighted through the aforementioned examples and advancements. Radiolabeled positron emitters offer a platform for precise molecular imaging and substantial image-guided therapy planning of various diseases. However, there are a lot of issues that are the main roadblocks in NPs based PET imaging. For example, before the selection of a suitable NP synthesis procedure, the half-life and biological targets should be kept in mind. Secondly, the appropriate labeling strategy for ideal fabrication of NP is also a matter of concern for many researchers. Additionally, there is a need to design novel biofabricated radiolabelled NPs to minimize the toxicity during PET imaging. The stability of the radiolabel, potential toxicity, and *in vivo* fate of the NP formulation are some of the additional concerns. Last but not the least, lack of evaluation of the radiolabeled NPs accumulation in certain organs including liver and kidneys often complicates image interpretation and safety profiles. Future research should focus on optimizing biocompatible and biodegradable nanomaterials. that offer site-specific targeting, high radiolabel stability, and minimal off-target accumulation. Moreover, systematic evaluation of pharmacodynamics and organ-specific distribution is necessary for their clinical translation. Nevertheless, further investigations in this intriguing field are expected to be planned by exploiting polymeric or lipid NPs that can offer selective delivery of therapeutic agents at target sites, allowing for enhanced *in vivo* tracking using PET technology.


4.2 Optimization of Nanomaterials-Based X-Ray CT Contrast Agents for Clinical Applications



X-Ray CT has emerged as one of the most prevalent non-invasive clinical diagnostic tools due to its low cost, optimal performance, rapid acquisition time and ability to provide high-resolution anatomical details. CT contrast agents can offer precise molecular imaging by corresponding diagnostic signals with biological processes. The specificity and sensitivity of CT imaging is directly proportional to its X-ray attenuation coefficient. However, conventional iodinated contrast agents often confer several challenges including short circulation time, nephrotoxicity, and lack of specificity. Therefore, designing nanomaterial-based CT contrast agents with improved pharmacokinetics, lower toxicity, and enhanced imaging performance is a pressing need in biomedical multimodal imaging¹²¹. A wide range of NPs, particularly metal NPs, have been constructed to serve as a potent CT contrast agent for specific bioimaging. For example, tantalum nanodots were synthesized for efficient CT imaging along with significant biocompatibility, good water solubility and renal clearance¹²². In addition, in an innovative research work, hydroxyapatite based nanoformulations, AuNPs and copper doped graphene oxide were separately evaluated for X-ray CT scanning. The results demonstrated that hydroxyapatite microspheres and thermally treated hydroxyapatite can serve as better contrast agents as compared to other nanoformulations¹²³. Moreover, polymer-fabricated lead oxide NPs have also been designed to be used as efficient CT contrast agents to facilitate high contrast tumor imaging¹²⁴. Despite these advances, many NP-based CT agents are still in their experimental stage due to long-term safety concerns, non-biodegradability, and potential organ accumulation. NP-based agents must display good biocompatibility, colloidal stability, suitable hydrodynamic size (<100 nm for renal clearance)¹²⁵, and less toxicity related concerns for successful clinical translation. Therefore, addressing these requirements by involving surface modification, biodegradable polymer coatings, and ligand targeted conjugation is required to enhance site-specific imaging while reducing off-target effects. Future research should focus on developing hybrid nanoplatforms that combine CT imaging with therapeutic capabilities (theranostics), evaluating long-term *in vivo* biodistribution, and complying with regulatory safety standards to to expand their applications in biomedical field.

4.3 Fluorescent Nanoprobes in NIRF Imaging

NIRF imaging has drawn substantial attention in the field of multimodal imaging due to its ultra sensitivity, contrast enhancement property, and versatility resulting in the application of a wide range of fluorescent nanoprobes for capturing images of biological components. The images obtained through fluorescence are studied by the observation of several factors including transfer of resonance energy, emission spectra, intensity and lifetime of fluorescent



agents. These factors provide insights into dynamic molecular interactions, localization, and metabolic processes within the biological system, allowing for a comprehensive understanding of cellular functions. Conventional imaging techniques often struggle with limited sensitivity, low specificity and poor spatial resolution. NIRF imaging has been reported in overcoming these limitations by enabling real-time, high-resolution imaging with reduced background autofluorescence. NIR fluorescent nanoprobes must exhibit certain critical features such as high quantum yield, excellent photostability, biocompatibility, minimal toxicity, and tunable emission within the NIR window (650–1700 nm) to function effectively in clinical bioimaging. In particular, the NIR-II window (1000–1700 nm) is now being actively investigated for superior in-depth imaging and signal-to-noise ratio ¹²⁶.

In order to accomplish specific targeted bioimaging, the surface of the fluorophores or NPs are modified with proteins or ligands to make them capable of recognising prognostic biomarkers of particular disease. In a novel approach, bislactosyl-fabricated BODIPY-TPE fluorescent probe NPs (BTL-Leus) was designed for the identification of leucine aminopeptidase (LAP) through fluorescence imaging in hepatoma cells ¹²⁷. Indocyanine green loaded calcium-based carbon NPs (Ca-CNPs@ICG) have been used for NIRF imaging of tumor thereby aiding in enhanced photodynamic therapy (PDT) ¹²⁸. Furthermore, in an innovative research erbium-based lanthanide NPs were designed for fluorescence imaging mediated surgery of orthotopic glioma ¹²⁹. Fluorescent imaging is rapidly emerging as an advanced imaging technique due to substantial development in microscopy, spectroscopy and material sciences. Despite these advances, certain limitations still persist that needs to be addressed in the coming future. The selection of fluorescent materials is often complicated due to the need to balance properties such as brightness, toxicity, and *in vivo* stability. Many conventional fluorophores suffer from rapid photobleaching or short fluorescence lifetimes, limiting their use in long-term tracking. Moreover, non-specific accumulation and rapid clearance pose challenges for achieving sustained imaging at target sites. To address these issues, current research focuses on engineering novel NIR fluorophores that aim to deliver improved decay lifetimes, higher contrast ratios, and the potential for multimodal fluorescent imaging, broadening their utility in both diagnostics and theranostics.

4.4 Advancements in Nanomaterials-Based High Contrast MRI

MRI is one of the most dynamic imaging modalities known for its non-invasive nature, high spatial resolution, excellent soft tissue contrast and ability of three-dimensional imaging. MRI



contrast agents, particularly those based on nanomaterials, have significantly enhanced the imaging of tissues by altering their magnetic properties, thereby improving diagnostic precision. The application of magnetic contrast agents has enabled the modification of intrinsic properties of tissues making the visualization flexible thereby enhancing the specificity and sensitivity of MRI (**Figure 1**). Wide range of nanomaterials such as superparamagnetic iron oxide, paramagnetic manganese oxide, and lanthanide-based compounds are being engineered for this purpose. In an earlier study, carbon-coated iron oxide NPs ($\text{Fe}_3\text{O}_4@\text{C}$ NPs) were used as MRI contrast agents for *in vivo* liver and kidney imaging¹³⁰. Similarly, a manganese oxide based MRI contrast agent has also been exploited for MRI of tumor in a novel work¹³¹. PEGylated manganese zinc ferrite NPs have also been designed to serve as MRI contrast agent for high contrast MRI¹³². Moreover, iron oxide NPs with citric acid coating have been used for determining the effect of pH and the results revealed that acidic coating enhanced the biocompatibility and stability making it suitable to be used as high contrast MRI agent¹³³. Many MRI nanoprobes are being functionalized with targeting ligands or therapeutic molecules to further improve specificity and multimodal utility. This dual-functional approach opens new possibilities in image-guided therapy. However, challenges such as long-term biocompatibility, biodistribution, clearance, and potential off-target accumulation still require deeper investigation. The understanding of magnetic effects is crucial for the development of novel MRI contrast agents. In addition to this, extensive *in vivo* research is also required to comprehend their mode of action in various organs for the purpose of MRI. Future efforts should focus on the development of biocompatible and stimuli-responsive MRI contrast agents with optimized relaxivity. Moreover, integrating artificial intelligence (AI)-assisted image analysis and predictive modeling could further enhance the clinical utility and precision of NP-based MRI platforms in personalized medicine.

4.5 Role of Nanoparticles in PAI for Enhanced Diagnosis

PAI is an emerging non-invasive full body bioimaging technique that works on the principle of optical imaging and ultrasound and offers high resolution and high contrast imaging due to deep tissue penetrability. PAI is widely used for the investigation and biodistribution of therapeutic or diagnostic agents thereby offering precise quantification of these agents inside the body (**Figure 2**). This unique hybrid approach enables high-resolution, high-contrast bioimaging at depths beyond the reach of purely optical methods, making it highly promising for clinical diagnostics and therapy monitoring. However, the effectiveness of PAI heavily depends on the availability of exogenous contrast agents that can strongly absorb NIR light and




exhibit minimal photobleaching with significant biocompatibility, and preferably offer multifunctional therapeutic capabilities. To address this, diverse range of NPs have been used for the purpose of fabrication of PAI contrast agents because of their intrinsic optical properties. For instance, acrylate-substituted thiadiazoloquinoxaline-diketopyrrolopyrrole polymeric NPs (PATQ-DPP) were developed as novel photoacoustic contrast agents for imaging and PTT of nasopharyngeal cancer ¹³⁴. Likewise, PLGA decorated methylene blue NPs have also been designed to serve as PAI and phototherapeutic agents ¹³⁵. Naphthalene diimide conjugated polycyclic molecule was nano precipitated with CoFe₂O₄ for the synthesis of NDI-S@CoFe₂O₄ that served as potent PAI agent for multimodal photodynamic and sonodynamic therapy ². Macrophage membrane-coated photoacoustic nanoprobes were developed for PAI of neuroinflammation in murine model ¹³⁶. Moreover, hyaluronic-acid-modified polydopamine NPs (PDA@HA) were constructed to serve as PAI contrast agent for the detection and real time therapeutic monitoring of endometriosis lesions ¹³⁷. While various kinds of functional nanomaterials have been investigated for enhancing high contrast PAI, there is still a need for comprehensive research on the *in vivo* safety profile of these nanomaterials. In addition to this, critical requirement for any PAI nanocomplex is also required to ensure strong and stable optical absorption with low toxicity, biodegradability, and clearance from the body to avoid long-term accumulation. Therefore, NPs should be subjected to rigorous toxicity assessment before proceeding for clinical application. Future research should focus not only on designing novel nanoformulation with optimized photoacoustic performance but also on comprehensive pharmacokinetic and regulatory assessments to overcome the challenges before clinical translation. The development of highly efficient and sensitive photoacoustic systems is a key area of research for spatial, dynamic, real-time, high contrast 3D multimodal bioimaging.

5. Role of Multifunctional Nanoparticles for Image-Guided Theranostics

Disease therapy can be facilitated by multifaceted NPs with specialized multi-modal imaging abilities, which can offer a novel approach for precise diagnosis and image-guided real-time treatment. The image guided approach can aid in conventional therapy by providing low dose of therapeutic agents and its controlled release with minimal side effects and specific targeting. In recent years, novel multifunctional NP formulations have advanced the field of image-guided theranostics by integrating multimodal imaging with targeted therapies to achieve diagnostic precision and treatment efficacy while minimizing systemic toxicity. For example, novel pyropheophorbide a-bisaminoquinoline lipid NPs (PPBC LNPs) have demonstrated combined PAI and fluorescence imaging alongside potent photothermal and photodynamic




therapy, facilitating real-time treatment monitoring and effective tumor eradication in bladder cancer models ¹³⁸. Similarly, enzyme-activatable theranostic nanomedicines designed to respond to tumor-associated enzymes such as MMP-14 and cathepsin have displayed enhanced MRI contrast and site-specific release of chemotherapeutics coupled with photothermal ablation, thereby improving treatment selectivity in case of glioblastoma ⁹⁶. Upconversion NPs (UCNPs) co-doped with lanthanides like gadolinium and iron oxide have been engineered for trimodal photoluminescence, CT, and MRI, offering high-resolution imaging and enabling precise monitoring of photoactivated tumor therapy deep within tissues ¹³⁹. Magnetic particle imaging has also recently emerged as an innovative tool for guiding magnetic fluid hyperthermia, where magnetic NPs serve dual functions as imaging tracers and localized heat sources, allowing for non-invasive, real-time mapping of treatment zones and temperature control with high spatial resolution ¹⁴⁰. Researchers have also designed pH-responsive nanogels encapsulating gadolinium or manganese-based MRI agents, which display enhanced relaxivity in the acidic tumor microenvironment and provide superior imaging contrast with minimal systemic toxicity due to their improved biodegradability and clearance profiles ¹⁴¹. To overcome long-term bioaccumulation concerns of noble metal NPs, ultrasmall gold-in-nano architectures have been developed, where sub-5 nm gold nanoclusters are confined within biodegradable silica capsules, ensuring effective photothermal tumor ablation under near-infrared irradiation while enabling complete renal clearance post-treatment ¹⁴². Additionally, hybrid theranostic nanoplateforms integrating gene editing systems like (CRISPR/Cas9) with MRI-visible nanocarriers have opened new avenues for real-time monitoring of gene therapy efficacy and off-target effects ^{143 144}. NPs are loaded with therapeutic moieties such as antigens, cytokines, immune checkpoint inhibitors, photosensitizers, chemotherapeutic drugs, polymers or fluorophores and can be subsequently monitored for efficient imaging and theranostics. For example, an iRGD peptide-based nanocomplex for its conjugation with dual immune checkpoint inhibitors for successful crossing of blood brain barrier and offering efficient immunotherapy of glioblastoma by blocking CXCL12/CXCR4 and PD-1/PD-L1 pathways ¹⁴⁵. Another fascinating research has focused on the conjugation of Ag₂S QDs with Pluronic F-127 (an amphiphilic polymer) followed by its functionalization with cancer cell membrane-based adjuvant for the construction of a potent nanocomplex Ag₂S@P@M-A capable of sonodynamic immunotherapy of colon cancer ¹⁴⁶. Chitosan coated paclitaxel was conjugated with MoS₂ having photothermal and optical properties for combined image-guided chemotherapy and PTT of breast cancer by the aid of MoS₂@PTX-CS-K237 NP ¹⁴⁷. The application of NPs as multimodal contrast agents for the purpose of image guided therapy has



also broadened its horizon in the field of précised surgical treatment. For instance, PLGA-cyclic RGD, perfluorohexane, and indocyanine green conjugated multifunctional NPs (PLGA-cRGD-PFH-ICG NPs) were designed for bimodal PET and NIRF imaging and promoting dissolution of activated platelets for the treatment of coronary microthrombosis¹⁴⁸. Moreover, a polystyrene nanoprobe has also been recently designed for precise fluorescence imaging guided resection of metastatic lesions in a latest work¹⁴⁹. An *in situ* spraying method was used for fluorescence guided surgery of aminopeptidase overexpressing metastatic cancer in a novel research work¹⁵⁰. Furthermore, in a recent work, bi-doped iron selenide NPs were fabricated for multimodal CT and MRI guided combined PTT and chemodynamic therapy of colorectal cancer¹⁵¹. The effectiveness of nanomaterials-based image guided therapy can be evidenced by the aid of aforementioned examples and research areas. The trend towards combining multiple therapeutic modalities such as chemotherapy, immunotherapy, phototherapy, and chemodynamic therapy within a single NP system, complemented by dual or triple imaging modes, holds immense potential to revolutionize personalized treatment regimens. However, a number of challenges including half-life of NPs in blood, pre-planning of surgical process, precise targeting and safety profile of NPs still need to be resolved for high contrast multimodal imaging and theranostics of certain diseases with minimum side effects.

6. Concluding Remarks and Future Prospects

Nanomaterials-based multimodal contrast agents have drawn significant attention due to their several advantages including homogeneity, high dispersibility, stability, tunable size and functional modification ability. The versatility of wide range of NPs makes them an ideal candidate to be integrated with existing conventional imaging modalities to enhance their spatial resolution and improving the diagnostic accuracy. Herein, we have provided recent data about the novel nanomaterials-based contrast and imaging agents that have been exploited for enhancing the efficacy of different multimodal imaging techniques along with their applications in certain common diseases. The designing of high contrast NP based multimodal imaging agents is an active area of research, however, it is still challenging for many researchers to construct multifunctional low-cost contrast agents on a large scale. This is a challenging field and has a lot of room to expand before advancing the application of NP in clinical imaging. Apart from the challenges in scaling-up-processes, the characterization of NPs has also not been given significant attention by nano-engineers which has resulted in limited success in their *in vitro* and *in vivo* imaging applications. Substantial characterization of NPs is important for their delivery in adequate doses according to the sensitivity of any



particular imaging modality, route of administration, pharmacokinetics, dispersibility and biocompatibility. The safety profile and toxicity of novel NP-based imaging moieties are yet other factors that need to be addressed for minimizing the adverse effects of imaging or theranostics. The presence of heavy transition metals restricts the application of these contrast agents in clinical bioimaging and diagnostic tests due to underlying toxicity issues. To address these complex issues, scientists and researchers from different biological backgrounds employ different assays to confirm or disprove toxicity which often results in different conclusions. Toxicological nature of any nanomaterial can be measured by ROS production, determination of lethal dose (LD₅₀) value, haemolytic assay and *in vivo* biodistribution profile. However, the application of commercial kits for *in vitro* toxicity assays is not completely valid for testing the safety profile of NPs and it is not necessary that *in vitro* results align with *in vivo* analysis. Therefore, a standard is required for testing the toxicity of all fabricated NPs and determining their potential to be conjugated with imaging or therapeutic agents. The fabrication, durability and efficacy of NP-based contrast agents highly depend on certain pharmacokinetic attributes including route of administration. Although the intravenous injection is the most preferred route of administration for the delivery of NP for bio-imaging, normal body vasculature has the tendency to limit the biodistribution resulting in delayed vascular equilibration. Additionally, certain NPs become prone to reticuloendothelial system (RES) uptake and immune clearance which makes it challenging for them to reach at the target site. Although, the clinical application of NPs is restricted due to their limited fabrication and lack of patient trials, NPs based imaging and diagnostics have paved the way to delve deeper into the pathophysiology of any disease or medical condition. It is believed that NP-based contrast agents would present a multitude of advantages including early-stage theranostics that will ultimately progress to clinical trials. Comprehensive research and innovative advancements are still required to bridge the gap between experimental results and clinical applications.

Acknowledgements

HT would like to thank the Junior Research Fellowship from Department of Biotechnology, Government of India. SS would like to thank University Grants Commission, New Delhi, India for Junior Research Fellowship. VG acknowledges funding support to his lab from Institution of Eminence Seed and Bridge Grant, Banaras Hindu University, Varanasi, India. The authors also acknowledge the initial support provided by the members of the Proteomics Laboratory during the preparation of this manuscript.



Author Information

Corresponding Authors

Dr. Vibhav Gautam - ¹Centre of Experimental Medicine and Surgery, Institute of Medical Sciences, Banaras Hindu University, Varanasi-221005, India; **ORCID:** 0000-0001-7956-9555; Email: vibhav.gautam4@bhu.ac.in

Dr. Navin Kumar Verma – ⁴Lee Kong Chian School of Medicine, Nanyang Technological University Singapore, Clinical Sciences Building, 11 Mandalay Road, Singapore 308232, Singapore

⁵Singapore Eye Research Institute, The Academia, 20 College Road, Singapore 169856, Singapore

⁶National Skin Centre, 1 Mandalay Road, Singapore 308205, Singapore

⁷Skin Research Institute of Singapore, Clinical Sciences Building, 11 Mandalay Road, Singapore 308232, Singapore, Email: nkverma@ntu.edu.sg

Authors

Harshita Tiwari - Centre of Experimental Medicine and Surgery, Institute of Medical Sciences, Banaras Hindu University, Varanasi-221005, India; **Email:** harshitatiwari@bhu.ac.in

Swati Singh - Centre of Experimental Medicine and Surgery, Institute of Medical Sciences, Banaras Hindu University, Varanasi-221005, India; **Email:** swati.7672@bhu.ac.in

Rajiv Kumar - Centre of Experimental Medicine and Surgery, Institute of Medical Sciences, Banaras Hindu University, Varanasi-221005, India; **Email:** rajiv.kumar@bhu.ac.in

Abhijit Mandal - Department of Radiotherapy and Radiation Medicine, Institute of Medical Sciences, Banaras Hindu University, Varanasi 221005, Uttar Pradesh, India; Email:

Abhishek Pathak - Department of Neurology, Institute of Medical Sciences, Banaras Hindu University, Varanasi 221005, Uttar Pradesh, India; Email: abhishek.neuro@bhu.ac.in

Lalit Kumar - Department of Urology, Institute of Medical Sciences, Banaras Hindu University, Varanasi 221005, Uttar Pradesh, India; Email: lalitkumarims@bhu.ac.in



Conflict of interest

The corresponding author on behalf of all the authors declares no potential conflict of interest.

Funding details

VG would like to thank Institution of Eminence Seed and Bridge Grant, Banaras Hindu University, Varanasi, India for providing funding support to his laboratory.

REFERENCES


- McKay A, Pantoja C, Hall R, Matthews S, Spalding P, Banerjee R. Patient understanding and experience of non-invasive imaging diagnostic techniques and the liver patient pathway. *Journal of Patient-Reported Outcomes*. 2021;5:1-14.
- Xie C-Y, Pang C-L, Chan B, Wong EY-Y, Dou Q, Vardhanabhuti V. Machine learning and Radiomics applications in esophageal cancers using non-invasive imaging Methods—A critical review of literature. *Cancers*. 2021;13(10):2469.
- Lu S, Dai Z, Cui Y, Kong D-M. Recent development of advanced fluorescent molecular probes for organelle-targeted cell imaging. *Biosensors*. 2023;13(3):360.
- Iacobellis F, Di Serafino M, Russo C, et al. Safe and informed use of gadolinium-based contrast agent in body magnetic resonance imaging: where we were and where we are. *Journal of Clinical Medicine*. 2024;13(8):2193.
- Sengar A, Vijayanandan A. Comprehensive review on iodinated X-ray contrast media: complete fate, occurrence, and formation of disinfection byproducts. *Science of the total environment*. 2021;769:144846.
- Chen S, Chen M, Yang J, et al. Design and engineering of hypoxia and acidic pH dual-stimuli-responsive intelligent fluorescent nanoprobe for precise tumor imaging. *Small*. 2021;17(28):2100243.
- Ren X, Han S, Li Y, et al. Tumor Microenvironment-Activatable Phototheranostic: Leveraging Nitric Oxide and Weak Acidity as Dual Biomarkers for Ratiometric Fluorescence, Photoacoustic Imaging, and Photothermal Therapy. *Analytical Chemistry*. 2024;96(21):8689-8695.
- Rezaei B, Tay ZW, Mostufa S, et al. Magnetic nanoparticles for magnetic particle imaging (MPI): design and applications. *Nanoscale*. 2024;16(25):11802-11824.
- Yanar F, Carugo D, Zhang X. Hybrid nanoplatforms comprising organic nanocompartments encapsulating inorganic nanoparticles for enhanced drug delivery and bioimaging applications. *Molecules*. 2023;28(15):5694.
- Wu Y, Zeng F, Zhao Y, Wu S. Emerging contrast agents for multispectral optoacoustic imaging and their biomedical applications. *Chemical Society Reviews*. 2021;50(14):7924-7940.
- Song N, Zhang J, Zhai J, Hong J, Yuan C, Liang M. Ferritin: a multifunctional nanoplatform for biological detection, imaging diagnosis, and drug delivery. *Accounts of chemical research*. 2021;54(17):3313-3325.
- Liu S, Su Y, Lin MZ, Ronald JA. Brightening up biology: advances in luciferase systems for in vivo imaging. *ACS chemical biology*. 2021;16(12):2707-2718.
- Chowdhury R, Wan J, Gardier R, et al. Molecular imaging with aquaporin-based reporter genes: quantitative considerations from Monte Carlo diffusion simulations. *ACS synthetic biology*. 2023;12(10):3041-3049.



14. Yue P, Nagendraraj T, Wang G, Jin Z, Angelovski G. The role of responsive MRI probes in the past and the future of molecular imaging. *Chemical Science*. 2024;15(48):20122-20154.
15. Zeng Z, Tan Y, Luo T, et al. Tyrosinase-Activated MRI and PET Probes for Selective Melanoma Imaging. *ACS sensors*. 2025;10(4):3023-3032.
16. Hurt RC, Buss MT, Duan M, et al. Genomically mined acoustic reporter genes for real-time in vivo monitoring of tumors and tumor-homing bacteria. *Nature Biotechnology*. 2023;41(7):919-931.
17. Lin L, Du Y, Wang Y, et al. Genetically engineered gas vesicle proteins with proliferative potential for synergistic targeted tumor therapy. *RSC advances*. 2025;15(1):157-166.
18. Hahmann J, Ishaqat A, Lammers T, Herrmann A. Sonogenetics for monitoring and modulating biomolecular function by ultrasound. *Angewandte Chemie International Edition*. 2024;63(13):e202317112.
19. Kumarasamy RV, Natarajan PM, Umapathy VR, Roy JR, Mironescu M, Palanisamy CP. Clinical applications and therapeutic potentials of advanced nanoparticles: A comprehensive review on completed human clinical trials. *Frontiers in Nanotechnology*. 2024;6:1479993.
20. Li S, Wei J, Yao Q, Song X, Xie J, Yang H. Emerging ultrasmall luminescent nanoprobes for in vivo bioimaging. *Chemical Society Reviews*. 2023;52(5):1672-1696.
21. Nisar S, Starosta E, Elayyan M, Regmi A, Sui B. Photoinduced electron transfer-based glutathione-sensing theranostic nanoprodrug with self-tracking and real-time drug release monitoring for cancer treatment. *ACS Applied Materials Interfaces*. 2024;16(6):6859-6867.
22. Hang Y, Wang A, Wu N. Plasmonic silver and gold nanoparticles: shape-and structure-modulated plasmonic functionality for point-of-care sensing, bio-imaging and medical therapy. *Chemical Society Reviews*. 2024;53(6):2932-2971.
23. Zhou Y, Lu Y, Liu Y, Hu X, Chen H. Current strategies of plasmonic nanoparticles assisted surface-enhanced Raman scattering toward biosensor studies. *Biosensors Bioelectronics*. 2023;228:115231.
24. Mcoyi MP, Mpofo KT, Sekhwama M, Mthunzi-Kufa P. Developments in localized surface plasmon resonance. 2024;Plasmonics:1-40.
25. Fritea L, Banica F, Costea TO, et al. Metal nanoparticles and carbon-based nanomaterials for improved performances of electrochemical (Bio) sensors with biomedical applications. *Materials*. 2021;14(21):6319.
26. Elishav O, Blumer O, Vanderlick TK, Hirshberg B. The effect of ligands on the size distribution of copper nanoclusters: Insights from molecular dynamics simulations. *The Journal of Chemical Physics*. 2024;160(16)
27. Choi J, Kim BH. Ligands of Nanoparticles and Their Influence on the Morphologies of Nanoparticle-Based Films. *Nanomaterials*. 2024;14(20):1685.
28. Nguyen VP, Qian W, Li Y, et al. Chain-like gold nanoparticle clusters for multimodal photoacoustic microscopy and optical coherence tomography enhanced molecular imaging. *Nature Communications*. 2021;12(1):34.
29. Ibrahim YO, Maalej N, Pirzada BM, et al. Gold nanoparticles spectral CT imaging and limit of detectability in a new materials contrast-detail phantom. *Physica Medica*. 2024;120:103326.
30. Sun I-C, Dumani DS, Emelianov SY. Applications of the Photocatalytic and Photoacoustic Properties of Gold Nanorods in Contrast-Enhanced Ultrasound and Photoacoustic Imaging. *ACS nano*. 2024;18(4):3575-3582.
31. Sotiriou GA. Biomedical applications of multifunctional plasmonic nanoparticles. *Wiley Interdisciplinary Reviews: Nanomedicine Nanobiotechnology*. 2013;5(1):19-30.



32. Singh KR, Natarajan A, Pandey SS. Bioinspired multifunctional silver nanoparticles for optical sensing applications: a sustainable approach. *ACS Applied Bio Materials*. 2023;6(11):4549-4571.
33. Vazquez-Prada KX, Moonshi SS, Wu Y, et al. A spiky silver-iron oxide nanoparticle for highly efficient targeted photothermal therapy and multimodal imaging of thrombosis. *Small*. 2023;19(11):2205744.
34. Mondal S, Montaña-Priede JL, Park S, et al. Computational analysis of drug free silver triangular nanoprism theranostic probe plasmonic behavior for in-situ tumor imaging and photothermal therapy. *Journal of Advanced Research*. 2022;41:23-38.
35. Niu G, Gao F, Wang Y, Zhang J, Zhao L, Jiang Y. Bimetallic nanomaterials: a promising nanoplatform for multimodal cancer therapy. *Molecules*. 2022;27(24):8712.
36. Oladipo AO, Unuofin JO, Iku SI, Nkambule TT, Mamba BB, Msagati TA. Bimetallic Au@ Pd nanodendrite system incorporating multimodal intracellular imaging for improved doxorubicin antitumor efficiency. *International Journal of Pharmaceutics*. 2021;602:120661.
37. Griaznova OY, Belyaev IB, Sogomonyan AS, et al. Laser synthesized core-satellite Fe-Au nanoparticles for multimodal in vivo imaging and in vitro photothermal therapy. *Pharmaceutics*. 2022;14(5):994.
38. Baijal G, Somashekar R, Iyer S, Nayak V. Comparative study of one pot synthesis of PEGylated gold and silver nanoparticles for imaging and radiosensitization of oral cancers. *Radiation Physics Chemistry*. 2022;194:109990.
39. Rahmani AA, Jia Q, Bahti HH, Fauzia RP, Wyantuti S. Recent advances in lanthanide-based nanoparticle contrast agents for magnetic resonance imaging: Synthesis, characterization, and applications. *OpenNano*. 2024:100226.
40. Liu N, Homann C, Morfin S, et al. Core-multi-shell design: unlocking multimodal capabilities in lanthanide-based nanoparticles as upconverting, T 2-weighted MRI and CT probes. *Nanoscale*. 2023;15(48):19546-19556.
41. Zhu X, Wang X, Zhang H, Zhang F. Luminescence lifetime imaging based on lanthanide nanoparticles. *Angewandte Chemie International Edition*. 2022;61(42):e202209378.
42. Monteiro JH. Recent advances in luminescence imaging of biological systems using lanthanide (III) luminescent complexes. *Molecules*. 2020;25(9):2089.
43. Alexander C, Guo Z, Glover PB, Faulkner S, Pikramenou Z. Luminescent Lanthanides in Biorelated Applications: From Molecules to Nanoparticles and Diagnostic Probes to Therapeutics. *Chemical Reviews*. 2025;125(4):2269-2370.
44. Caro C, Paez-Muñoz JM, Beltran AM, Pernia Leal M, García-Martín ML. PEGylated terbium-based nanorods as multimodal bioimaging contrast agents. *ACS Applied Nano Materials*. 2021;4(4):4199-4207.
45. Gómez-González E, Caro C, Núñez NO, et al. Sodium lanthanide tungstate-based nanoparticles as bimodal contrast agents for in vivo high-field MRI and CT imaging. *Journal of Materials Chemistry B*. 2024;12(43):11123-11133.
46. Gómez-González E, Núñez NO, Caro C, García-Martín ML, Becerro AI, Ocaña M. Lanthanide vanadate-based nanoparticles as multimodal T1-T2 MRI contrast agent and NIR luminescent imaging probe. *Journal of Alloys Compounds*. 2024;1003:175647.
47. Huang D, Li F, Ågren H, Chen G. Inhibiting concentration quenching in Yb³⁺-Tm³⁺ upconversion nanoparticles by suppressing back energy transfer. *Nature Communications*. 2025;16(1):1-8.
48. Zhang Q, O'Brien S, Grimm J. Biomedical applications of lanthanide nanomaterials, for imaging, sensing and therapy. *Nanotheranostics*. 2022;6(2):184.



- 1044 49. Du K, Feng J, Gao X, Zhang H. Nanocomposites based on lanthanide-doped
1045 upconversion nanoparticles: diverse designs and applications. *Light: Science Applications*.
1046 2022;11(1):222.
- 1047 50. Chen G-H, Chen P-H, Lin C-T, Jang T-W, Yang P, Chen H-S. Enhanced photostability
1048 of core/shell quantum dots under intense blue light irradiation through positive photoaging
1049 mechanism. *ACS Applied Materials Interfaces*. 2023;15(45):52795-52805.
- 1050 51. Lee HK, Kim T, Jang YA, Jeong Y, Lee SW, Park CH. Near-Infrared Emissive
1051 CuInS₂/ZnS Quantum Dot-Embedded Polymer Scaffolds for Photon Upconversion Imaging.
1052 *Advanced Materials*. 2025:2502333.
- 1053 52. Jalali HB, De Trizio L, Manna L, Di Stasio F. Indium arsenide quantum dots: an
1054 alternative to lead-based infrared emitting nanomaterials. *Chemical Society Reviews*.
1055 2022;51(24):9861-9881.
- 1056 53. Ding C, Huang Y, Shen Z, Chen X. Synthesis and bioapplications of Ag₂S quantum
1057 dots with near-infrared fluorescence. *Advanced Materials*. 2021;33(32):2007768.
- 1058 54. Almeida G, Ubbink RF, Stam M, du Fossé I, Houtepen AJ. InP colloidal quantum dots
1059 for visible and near-infrared photonics. *Nature Reviews Materials*. 2023;8(11):742-758.
- 1060 55. Li Z, Mahajan A, Andaraarachchi HP, Lee Y, Kortshagen UR. Water-soluble
1061 luminescent silicon nanocrystals by plasma-induced acrylic acid grafting and PEGylation. *ACS*
1062 *applied bio materials*. 2021;5(1):105-112.
- 1063 56. K. Tiwari P, Sahu M, Kumar G, Ashourian M. Pivotal role of quantum dots in the
1064 advancement of healthcare research. *Computational Intelligence Neuroscience*.
1065 2021;2021(1):2096208.
- 1066 57. Sobhanan J, Rival JV, Anas A, Shibu ES, Takano Y, Biju V. Luminescent quantum
1067 dots: Synthesis, optical properties, bioimaging and toxicity. *Advanced Drug Delivery Reviews*.
1068 2023;197:114830.
- 1069 58. Zhang P, Wang Y, Liu X, et al. Carboxyl-Modified Quantum Dots for NIR-IIb Bone
1070 Marrow Imaging. *ACS Applied Materials Interfaces*. 2024;16(7):8509-8517.
- 1071 59. Li H, Du Z, Zhu L, et al. Ultrabright NIR-IIb Fluorescence Quantum Dots for Targeted
1072 Imaging-Guided Surgery. *ACS Applied Materials Interfaces*. 2024;
- 1073 60. Zhou Y, Huang B, Chen S-H, Liu S-L, Zhang M, Cui R. Ultra-bright near-infrared-IIb
1074 emitting Zn-doped Ag₂Te quantum dots for noninvasive monitoring of traumatic brain injury.
1075 *Nano Research*. 2023;16(2):2719-2727.
- 1076 61. Yang D, Wei X, Piao Z, et al. Constructing bimodal nanoprobe based on Gd:
1077 AgInS₂/ZnS quantum dots for fluorometric/magnetic resonance imaging in mesenchymal stem
1078 cells. *Journal of Materials Science Technology*. 2023;148:116-122.
- 1079 62. Tong S, Zhong J, Chen X, et al. In vivo deep-brain 3-and 4-photon fluorescence
1080 imaging of subcortical structures labeled by quantum dots excited at the 2200 nm window. *ACS*
1081 *nano*. 2023;17(4):3686-3695.
- 1082 63. Guan X, Zhang L, Lai S, et al. Green synthesis of glyco-CuInS₂ QDs with visible/NIR
1083 dual emission for 3D multicellular tumor spheroid and in vivo imaging. *Journal of*
1084 *Nanobiotechnology*. 2023;21(1):118.
- 1085 64. Chen G, Wang L, He P, et al. Biodistributions and Imaging of Poly (ethylene glycol)-
1086 Conjugated Silicon Quantum Dot Nanoparticles in Osteosarcoma Models via Intravenous and
1087 Intratumoral Injections. *ACS Applied Bio Materials*. 2023;6(11):4856-4866.
- 1088 65. Manjubaashini N, Bargavi P, Balakumar S. Carbon quantum dots derived from agro
1089 waste biomass for pioneering bioanalysis and in vivo bioimaging. *Journal of Photochemistry*
1090 *Photobiology A: Chemistry*. 2024;454:115702.
- 1091 66. Ham K-M, Kim M, Bock S, et al. Highly bright silica-coated InP/ZnS quantum dot-
1092 embedded silica nanoparticles as biocompatible nanoprobe. *International Journal of*
1093 *Molecular Sciences*. 2022;23(18):10977.



67. El-Seedi HR, El-Shabasy RM, Khalifa SA, et al. Metal nanoparticles fabricated by green chemistry using natural extracts: Biosynthesis, mechanisms, and applications. *RSC advances*. 2019;9(42):24539-24559.
68. Haque S, Norbert CC, Acharyya R, Mukherjee S, Kathirvel M, Patra CR. Biosynthesized silver nanoparticles for cancer therapy and in vivo bioimaging. *Cancers*. 2021;13(23):6114.
69. Lian W, Gan M. Fabrication of folic acid–cysteamine-modified silver nanoparticles as promising contrast agent for computed tomography imaging. *Biotechnology Bioprocess Engineering*. 2024;29(1):185-196.
70. Naha PC, Hsu JC, Kim J, et al. Dextran-coated cerium oxide nanoparticles: a computed tomography contrast agent for imaging the gastrointestinal tract and inflammatory bowel disease. *ACS nano*. 2020;14(8):10187-10197.
71. Zeng X, Chen Q, Chen T. Nanomaterial-assisted oncolytic bacteria in solid tumor diagnosis and therapeutics. *Bioengineering Translational Medicine*. 2024:e10672.
72. Chen SH, Liu H, Huang B, et al. Biosynthesis of NIR-II Ag₂Se Quantum Dots with Bacterial Catalase for Photoacoustic Imaging and Alleviating-Hypoxia Photothermal Therapy. *Small*. 2024:2310795.
73. Ren G, Zhou X, Long R, et al. Biomedical applications of magnetosomes: state of the art and perspectives. *Bioactive Materials*. 2023;28:27-49.
74. Yang M, Wang X, Pu F, et al. Engineered exosomes-based photothermal therapy with MRI/CT imaging guidance enhances anticancer efficacy through deep tumor nucleus penetration. *Pharmaceutics*. 2021;13(10):1593.
75. Shahgolzari M, Venkataraman S, Osano A, Akpa PA, Hefferon K. Plant virus nanoparticles combat cancer. *Vaccines*. 2023;11(8):1278.
76. Zhang L, Tong S, Zhang Q, Bao G. Lipid-encapsulated Fe₃O₄ nanoparticles for multimodal magnetic resonance/fluorescence imaging. *ACS Applied Nano Materials*. 2020;3(7):6785-6797.
77. Yen T-YC, Abbasi AZ, He C, et al. Biocompatible and bioactivable terpolymer-lipid-MnO₂ Nanoparticle-based MRI contrast agent for improving tumor detection and delineation. *Materials Today Bio*. 2024;25:100954.
78. Li C, Jia H, Zhao W, et al. Endogenously Activated and Self-Reinforced DNA Lipid Nanodevice for Spatial-Specific and High-Contrast Imaging of MicroRNA in Cells and Animals. *Advanced Functional Materials*. 2024:2405780.
79. Chauhan M, Shekhar S, Yadav B, et al. AS1411 aptamer/RGD dual functionalized theranostic chitosan-PLGA nanoparticles for brain cancer treatment and imaging. *Biomaterials Advances*. 2024;160:213833.
80. Wang C, Wang X, Chan HN, et al. Amyloid- β oligomer-targeted gadolinium-based NIR/MR dual-modal theranostic nanoprobe for Alzheimer's disease. *Advanced Functional Materials*. 2020;30(16):1909529.
81. Yang J, Wang L, Su Y, et al. Color-convertible fluorescent nanoprobe for Parkinson's disease diagnosis. *Chemical Engineering Journal*. 2022;429:132368.
82. Cheng Y, Cheng A, Jia Y, et al. pH-Responsive multifunctional theranostic rapamycin-loaded nanoparticles for imaging and treatment of acute ischemic stroke. *ACS Applied Materials Interfaces*. 2021;13(48):56909-56922.
83. Xu S, Zhang G, Zhang J, Liu W, Wang Y, Fu X. Advances in brain tumor therapy based on the magnetic nanoparticles. *International Journal of Nanomedicine*. 2023:7803-7823.
84. Liu X, Xia T, Fang Y, et al. Overcoming the blood–brain barrier by using a multistage exosome delivery system to inhibit central nervous system lymphoma. *Nanomedicine: Nanotechnology, Biology Medicine*. 2022;41:102523.



- 1143 85. Ahmad F, Varghese R, Panda S, et al. Smart nanoformulations for brain cancer
1144 theranostics: Challenges and promises. *Cancers*. 2022;14(21):5389.
- 1145 86. Li J, Wei Y, Zhang C, et al. Cell-membrane-coated nanoparticles for targeted drug
1146 delivery to the brain for the treatment of neurological diseases. *Pharmaceutics*.
1147 2023;15(2):621.
- 1148 87. Seijas A, Cora D, Novo M, Al-Soufi W, Sánchez L, Arana ÁJ. CRISPR/Cas9 Delivery
1149 Systems to Enhance Gene Editing Efficiency. *International Journal of Molecular Sciences*.
1150 2025;26(9):4420.
- 1151 88. Xu M, Qin Z, Chen Z, et al. Nanorobots mediated drug delivery for brain cancer active
1152 targeting and controllable therapeutics. *Discover Nano*. 2024;19(1):183.
- 1153 89. Siddique S, Chow JCL. Recent Advances in Functionalized Nanoparticles in Cancer
1154 Theranostics. *Nanomaterials*. 2022;12(16):2826.
- 1155 90. Zhang J, Yuan C, Kong L, et al. H-ferritin-nanocaged gadolinium nanoparticles for
1156 ultra-sensitive MR molecular imaging. *Theranostics*. 2024;14(5):1956.
- 1157 91. Chung S-J, Hadrick K, Nafiujjaman M, et al. Targeted Biodegradable Near-Infrared
1158 Fluorescent Nanoparticles for Colorectal Cancer Imaging. *ACS Applied Bio Materials*. 2024;
- 1159 92. Xu H, Yu P, Bandari RP, et al. Bimodal MRI/fluorescence nanoparticle imaging
1160 contrast agent targeting prostate cancer. *Nanomaterials*. 2024;14(14):1177.
- 1161 93. Nguyen D-T, Baek M-J, Lee SM, et al. Photobleaching-mediated charge-convertible
1162 cyclodextrin nanoparticles achieve deep tumour penetration for rectal cancer theranostics.
1163 *Nature Nanotechnology*. 2024:1-12.
- 1164 94. Zhao X, Zhang G, Chen J, et al. A rationally designed nuclei-targeting FAPI 04-based
1165 molecular probe with enhanced tumor uptake for PET/CT and fluorescence imaging. *European
1166 Journal of Nuclear Medicine Molecular Imaging*. 2024;51(6):1593-1604.
- 1167 95. Wu C, Sun Q, Liu X, Sun X, Chen Z, Shan H. Indocyanine Green-Loaded Liposomes-
1168 Assisted Photoacoustic Computed Tomography for Evaluating In Vivo Tumor Penetration of
1169 Liposomal Nanocarriers. *Micromachines*. 2023;15(1):90.
- 1170 96. Shokri Varniab Z, Chang E, Wang J, et al. Dual-enzyme activated theranostic
1171 nanoparticles for image-guided glioblastoma therapy. *Scientific Reports*. 2025;15(1):13540.
- 1172 97. Yue S, Zhan J, Xu X, Xu J, Bi S, Zhu J-J. A “dual-key-and-lock” DNA nanodevice
1173 enables spatially controlled multimodal imaging and combined cancer therapy. *Chemical
1174 Science*. 2024;15(29):11528-11539.
- 1175 98. Jokerst JV, Cole AJ, Van de Sompel D, Gambhir SS. Gold nanorods for ovarian cancer
1176 detection with photoacoustic imaging and resection guidance via Raman imaging in living
1177 mice. *ACS nano*. 2012;6(11):10366-10377.
- 1178 99. Li F, Chen L, Zhong S, et al. Collagen-Targeting Self-Assembled Nanoprobes for
1179 Multimodal Molecular Imaging and Quantification of Myocardial Fibrosis in a Rat Model of
1180 Myocardial Infarction. *ACS nano*. 2024;18(6):4886-4902.
- 1181 100. Ma Y, Xu L, Yin B, et al. Ratiometric semiconducting polymer nanoparticle for reliable
1182 photoacoustic imaging of pneumonia-induced vulnerable atherosclerotic plaque in vivo. *Nano
1183 letters*. 2021;21(10):4484-4493.
- 1184 101. Yin D, Zheng M, Zhang Q, Li M, Xiang P, Tian J. Macrophage membrane-modified
1185 targeted phase-change nanoparticles for multimodal imaging of experimental autoimmune
1186 myocarditis. *Nanoscale*. 2024;16(13):6680-6695.
- 1187 102. Senders ML, Meerwaldt AE, van Leent MM, et al. Probing myeloid cell dynamics in
1188 ischaemic heart disease by nanotracer hot-spot imaging. *Nature nanotechnology*.
1189 2020;15(5):398-405.
- 1190 103. Setia A, Mehata AK, Priya V, et al. Nanoparticles for Thrombus Diagnosis and
1191 Therapy: Emerging Trends in Thrombus-theranostics. *Nanotheranostics*. 2024;8(2):127.



- 1192 104. Liu J, Jia B, Li Z, Li W. Reactive oxygen species-responsive polymer drug delivery
1193 systems. *Frontiers in Bioengineering Biotechnology*. 2023;11:1115603. View Article Online
DOI:10.1039/D3NA00596E
- 1194 105. Zhang H, Mao Y, Nie Z, et al. Iron Oxide nanoparticles Engineered macrophage-
1195 derived exosomes for targeted pathological angiogenesis therapy. *ACS nano*.
1196 2024;18(10):7644-7655.
- 1197 106. Liu C, Fan Z, He D, et al. Designer functional nanomedicine for myocardial repair by
1198 regulating the inflammatory microenvironment. *Pharmaceutics*. 2022;14(4):758.
- 1199 107. Shu G, Zhang C, Wen Y, Pan J, Zhang X, Sun S-K. Bismuth drug-inspired ultra-small
1200 dextran coated bismuth oxide nanoparticles for targeted computed tomography imaging of
1201 inflammatory bowel disease. *Biomaterials*. 2024:122658.
- 1202 108. Tian M, Mu X, Fan D, et al. A transformable mucoadhesive microgel network for
1203 noninvasive multimodal imaging and radioprotection of a large area of the gastrointestinal
1204 tract. *Advanced Materials*. 2023;35(40):2303436.
- 1205 109. Zhao Y, Ye F, Brismar TB, et al. Multimodal imaging of pancreatic ductal
1206 adenocarcinoma using multifunctional nanoparticles as contrast agents. *ACS applied materials
1207 interfaces*. 2020;12(48):53665-53681.
- 1208 110. Ma L, Huang S, He S, Wang Z, Cheng Z. Polydopamine-coated downconversion
1209 nanoparticle as an efficient dual-modal near-infrared-II fluorescence and photoacoustic
1210 contrast agent for non-invasive visualization of gastrointestinal tract in vivo. *Biosensors
1211 Bioelectronics*. 2020;151:112000.
- 1212 111. Feng-Zhi C, Jian-Hua L, Yang L, et al. Synthesis of PEGylated BaGdF5 Nanoparticles
1213 as Efficient CT/MRI Dual-modal Contrast Agents for Gastrointestinal Tract Imaging. *Chinese
1214 Journal of Analytical Chemistry*. 2020;48(8):1004-1011.
- 1215 112. Jin M, Ling Y, Jin L, Wang X, Jiang X, Li Q. Cell membrane-functionalized bismuth
1216 oxyiodide nanodots as potential contrast agent for gastrointestinal tract imaging. *Colloids
1217 Surfaces B: Biointerfaces*. 2025;245:114235.
- 1218 113. Rosario-Berrios DN, Pang A, Liu LP, et al. The effect of the size of gold nanoparticle
1219 contrast agents on CT imaging of the gastrointestinal tract and inflammatory bowel disease.
1220 *Bioconjugate Chemistry*. 2025;
- 1221 114. Yue N-n, Xu H-m, Xu J, et al. Application of nanoparticles in the diagnosis of
1222 gastrointestinal diseases: a complete future perspective. *International Journal of
1223 Nanomedicine*. 2023;4143-4170.
- 1224 115. Najdian A, Beiki D, Abbasi M, et al. Exploring innovative strides in radiolabeled
1225 nanoparticle progress for multimodality cancer imaging and theranostic applications. *Cancer
1226 Imaging*. 2024;24(1):127.
- 1227 116. Krekorian M, Cortenbach KR, Boswinkel M, et al. In vivo PET imaging of monocytes
1228 labeled with [89Zr] Zr-PLGA-NH2 nanoparticles in tumor and Staphylococcus aureus
1229 infection models. *Cancers*. 2021;13(20):5069.
- 1230 117. Gandhi S, Shende P. Anti-CD64 Antibody-Conjugated PLGA Nanoparticles
1231 Containing Methotrexate and Gold for Theranostics Application in Rheumatoid Arthritis.
1232 *AAPS PharmSciTech*. 2024;25(1):22.
- 1233 118. Chen Q, Fu X, Cai H, et al. 68Ga-labeled amphiphilic polymer nanoparticles for PET
1234 imaging of sentinel lymph node metastasis. *Regenerative Biomaterials*. 2023;10:rbad029.
- 1235 119. Díez-Villares S, Pellico J, Gómez-Lado N, et al. Biodistribution of 68/67Ga-
1236 radiolabeled sphingolipid nanoemulsions by pet and spect imaging. *International journal of
1237 nanomedicine*. 2021:5923-5935.
- 1238 120. Fan Y, Miao W, Yan R, Li C, Wang X. Tannylated lipid nanoparticles for prolonged
1239 circulation and PET imaging-guided cancer therapy. *Biomaterials Advances*. 2025/10/01/
1240 2025;175:214325.



- 1241 121. Owens TC, Anton N, Attia MF. CT and X-ray contrast agents: current clinical
1242 challenges and the future of contrast. *Acta Biomaterialia*. 2023;171:19-36.
- 1243 122. Dong Y, Shu G, Wei Y, Pan J, Li D, Sun S-K. Gram-Scale Synthesis of Renal-Clearable
1244 Tantalum Nanodots with High Water Solubility for Computed Tomography Imaging In Vivo.
1245 *ACS nano*. 2024;
- 1246 123. Pugliese Pereira B, Antoine C, de Barros AODS, et al. Evaluating Hydroxyapatite, Gold
1247 Nanoparticles, and Graphene-Copper as Bimodal agents for X-ray and computed tomography.
1248 *Bioengineering*. 2023;10(2):238.
- 1249 124. Al Saidi AKA, Baek A, Liu S, et al. Lead Oxide Nanoparticles as X-ray Contrast Agents
1250 for In Vitro and In Vivo Imaging. *ACS Applied Nano Materials*. 2023;6(21):20129-20138.
- 1251 125. Mellor RD, Uchegbu IF. Ultrasmall-in-nano: why size matters. *Nanomaterials*.
1252 2022;12(14):2476.
- 1253 126. Zheng F, Huang X, Ding J, et al. NIR-I dye-based probe: a new window for bimodal
1254 tumor theranostics. *Frontiers in Chemistry*. 2022;10:859948.
- 1255 127. Zhou W, Liu Y-c, Ma Q-y, et al. AIE-active lysosome-targeted fluorescent organic
1256 nanoparticles for leucine aminopeptidase-activatable fluorescent imaging and precision
1257 photodynamic therapy potential. *Dyes Pigments*. 2024;221:111781.
- 1258 128. Pang E, Li X, Zhao S, et al. Calcium-enriched carbon nanoparticles loaded with
1259 indocyanine green for near-infrared fluorescence imaging-guided synergistic calcium
1260 overload, photothermal therapy, and glutathione-depletion-enhanced photodynamic therapy.
1261 *Journal of Materials Chemistry B*. 2024;12(7):1846-1853.
- 1262 129. Ren F, Liu H, Zhang H, et al. Engineering NIR-IIb fluorescence of Er-based lanthanide
1263 nanoparticles for through-skull targeted imaging and imaging-guided surgery of orthotopic
1264 glioma. *Nano Today*. 2020;34:100905.
- 1265 130. Yue H, Zhao D, Tegafaw T, et al. Core-Shell Fe₃O₄@ C Nanoparticles as Highly
1266 Effective T2 Magnetic Resonance Imaging Contrast Agents: In Vitro and In Vivo Studies.
1267 *Nanomaterials*. 2024;14(2):177.
- 1268 131. Yang G, Xia J, Dai X, et al. A Targeted Multi-Crystalline Manganese Oxide as a
1269 Tumor-Selective Nano-Sized MRI Contrast Agent for Early and Accurate Diagnosis of
1270 Tumors. *International Journal of Nanomedicine*. 2024:527-540.
- 1271 132. Cheraghali S, Dini G, Caligiuri I, Back M, Rizzolio F. PEG-coated MnZn ferrite
1272 nanoparticles with hierarchical structure as MRI contrast agent. *Nanomaterials*.
1273 2023;13(3):452.
- 1274 133. Alzoubi F, Noqta OA, AlZoubi T, et al. Exploring the impact of pH on the properties
1275 of citric acid-coated iron oxide nanoparticles as high-performance T2 contrast agent for MRI
1276 applications. *Results in Engineering*. 2023;18:101206.
- 1277 134. Li Z, Lin Y, Qiu T, et al. Noninvasive Photothermal Therapy of Nasopharyngeal Cancer
1278 Guided by High Efficiency Optical-Absorption Nanomaterial Enhanced by NIR-II
1279 Photoacoustic Imaging. *International Journal of Nanomedicine*. 2024:7817-7830.
- 1280 135. Xu X, Wu H, Yang Y, et al. PLGA-coated methylene blue nanoparticles for
1281 photoacoustic imaging and photodynamic/photothermal cascaded precisely synergistic therapy
1282 of tumor. *RSC advances*. 2022;12(3):1543-1549.
- 1283 136. Pan Y, Chen J, Zhang Y, et al. Second Near-Infrared Macrophage-Biomimetic
1284 Nanoprobes for Photoacoustic Imaging of Neuroinflammation. *Molecular Pharmaceutics*.
1285 2024;21(4):1804-1816.
- 1286 137. Lv Q, Zhang Y, Yang R, et al. Photoacoustic imaging endometriosis lesions with
1287 nanoparticulate polydopamine as a contrast agent. *Advanced Healthcare Materials*.
1288 2024;13(2):2302175.



- 1289 138. Tang M, Mahri S, Shiao Y-P, et al. Multifunctional and Scalable Nanoparticles for
 1290 Bimodal Image-Guided Phototherapy in Bladder Cancer Treatment. *Nano-Micro Letters*.
 1291 2025;17(1):222. View Article Online
DOI: 10.1039/D5NA00596E
- 1292 139. Min Y, Li J, Liu F, Padmanabhan P, Yeow EK, Xing B. Recent advance of biological
 1293 molecular imaging based on lanthanide-doped upconversion-luminescent nanomaterials.
 1294 *Nanomaterials*. 2014;4(1):129-154.
- 1295 140. Lu Y, Rivera-Rodriguez A, Tay ZW, et al. Combining magnetic particle imaging and
 1296 magnetic fluid hyperthermia for localized and image-guided treatment. *International Journal*
 1297 *of Hyperthermia*. 2020;37(3):141-154.
- 1298 141. Ma G, Zhang X, Zhao K, et al. Polydopamine nanostructure-enhanced water interaction
 1299 with pH-responsive manganese sulfide nanoclusters for tumor magnetic resonance contrast
 1300 enhancement and synergistic ferroptosis–photothermal therapy. *ACS nano*. 2024;18(4):3369-
 1301 3381.
- 1302 142. Cassano D, Santi M, D’Autilia F, Mapanao AK, Luin S, Voliani V. Photothermal effect
 1303 by NIR-responsive excretable ultrasmall-in-nano architectures. *Materials Horizons*.
 1304 2019;6(3):531-537.
- 1305 143. Cho HY, Yoo M, Pongkulapa T, et al. Magnetic Nanoparticle-Assisted Non-Viral
 1306 CRISPR-Cas9 for Enhanced Genome Editing to Treat Rett Syndrome. *Advanced Science*.
 1307 2024;11(24):2306432.
- 1308 144. Zhu H, Zhang L, Tong S, Lee CM, Deshmukh H, Bao G. Spatial control of in vivo
 1309 CRISPR–Cas9 genome editing via nanomagnets. *Nature biomedical engineering*.
 1310 2019;3(2):126-136.
- 1311 145. Wei R, Li J, Lin W, et al. Nanoparticle-mediated blockade of CXCL12/CXCR4
 1312 signaling enhances glioblastoma immunotherapy: Monitoring early responses with MRI
 1313 radiomics. *Acta Biomaterialia*. 2024;177:414-430.
- 1314 146. Zhang Y, Zhang Y, Li Y, et al. Harnessing Ag2S quantum dots with immune adjuvant
 1315 for NIR-II fluorescence imaging-guided sonodynamic immunotherapy of colon cancer.
 1316 *Chemical Engineering Journal*. 2023;474:145685.
- 1317 147. Wang P, Peng Z, Zhang Y, et al. A chitosan-camouflaged nanomedicine triggered by
 1318 hierarchically stimuli to release drug for multimodal imaging-guided chemotherapy of breast
 1319 cancer. *Carbohydrate Polymers*. 2024;335:122073.
- 1320 148. Bai S, Liao J, Zhang B, et al. Multimodal and multifunctional nanoparticles with
 1321 platelet targeting ability and phase transition efficiency for the molecular imaging and
 1322 thrombolysis of coronary microthrombi. *Biomaterials Science*. 2020;8(18):5047-5060.
- 1323 149. Xu H, Yuan L, Shi Q, Tian Y, Hu F. Ultrabright NIR-II Nanoprobe for Image-Guided
 1324 Accurate Resection of Tiny Metastatic Lesions. *Nano Letters*. 2024;24(4):1367-1375.
- 1325 150. Li H, Yao Q, Sun W, et al. Aminopeptidase N activatable fluorescent probe for tracking
 1326 metastatic cancer and image-guided surgery via in situ spraying. *Journal of the American*
 1327 *Chemical Society*. 2020;142(13):6381-6389.
- 1328 151. Liu Z, Hong C, Pan C, et al. Synergistically Enhanced Chemodynamic Therapy Induced
 1329 by NIR-II Photothermal Therapy with MRI/CT Imaging Guidance for Colorectal Cancer. *ACS*
 1330 *Materials Letters*. 2024;6(5):1593-1605.
- 1331 152. Chen Y, Wu H, Zhou H, et al. PEGylated indium nanoparticles: a metallic contrast
 1332 agent for multiwavelength photoacoustic imaging and second near-infrared photothermal
 1333 therapy. *ACS Applied Materials Interfaces*. 2021;13(39):46343-46352.
- 1334 153. Choi G, Kim H-K, Baek AR, et al. Multifunctional imaging of amyloid-beta peptides
 1335 with a new gadolinium-based contrast agent in Alzheimer’s disease. *Journal of Industrial*
 1336 *Engineering Chemistry*. 2020;83:214-223.



- 1337 154. Verry C, Dufort S, Lemasson B, et al. Targeting brain metastases with ultrasmall
1338 theranostic nanoparticles, a first-in-human trial from an MRI perspective. *Science Advances*.
1339 2020;6(29):eay5279.
- 1340 155. Deblock L, Descamps B, Goemaere I, et al. Dual-Modality Hafnium Oxide
1341 Nanocrystals for in Vivo Computed Tomography and Fluorescence Imaging of Sentinel
1342 Lymph Nodes. *Chemistry of Materials*. 2023;35(21):8883-8896.
- 1343 156. Zhang J, Ning Y, Zhu H, et al. Fast detection of liver fibrosis with collagen-binding
1344 single-nanometer iron oxide nanoparticles via T 1-weighted MRI. *Proceedings of the National
1345 Academy of Sciences*. 2023;120(18):e2220036120.
- 1346 157. Waddington DE, Boele T, Maschmeyer R, Kuncic Z, Rosen MS. High-sensitivity in
1347 vivo contrast for ultra-low field magnetic resonance imaging using superparamagnetic iron
1348 oxide nanoparticles. *Science Advances*. 2020;6(29):eabb0998.
- 1349 158. Zhang T, Li J, Lu J, et al. Enhanced tumor-targeting ability of transferrin-functionalized
1350 magnetic nanoparticles by in vivo AMF stimulation. *Biomaterials*. 2025;315:122925.
- 1351 159. Jazrawi A, Pantiora E, Abdsaleh S, et al. Magnetic-guided axillary ultrasound (MagUS)
1352 sentinel lymph node biopsy and mapping in patients with early breast cancer. a phase 2, single-
1353 arm prospective clinical trial. *Cancers*. 2021;13(17):4285.
- 1354 160. Elhalawani H, Awan MJ, Ding Y, et al. Data from a terminated study on iron oxide
1355 nanoparticle magnetic resonance imaging for head and neck tumors. *Scientific Data*.
1356 2020;7(1):63.
- 1357 161. Mohajeri M, Salehi P, Heidari B, et al. PEGylated Pemetrexed and PolyNIPAM
1358 Decorated Gold Nanoparticles: A Biocompatible and Highly Stable CT Contrast Agent for
1359 Cancer Imaging. *ACS Applied Bio Materials*. 2024;7(9):5977-5991.
- 1360 162. Oliveira GNdSA, Silva JYR, do Nascimento Santos DKD, et al. Design of a magnetic
1361 nanocarrier containing phyllacanthone as delivery of anticancer phytochemical:
1362 Characterization and theranostic in vitro applications. *Journal of Alloys Compounds*.
1363 2024;177860.
- 1364 163. Jameel MS, Aziz AA, Dheyab MA, Mehrdel B, Khaniabadi PM, Khaniabadi BM.
1365 Green sonochemical synthesis platinum nanoparticles as a novel contrast agent for computed
1366 tomography. *Materials Today Communications*. 2021;27:102480.
- 1367 164. Guo C, Sun J, Dong J, et al. A natural anthocyanin-based multifunctional theranostic
1368 agent for dual-modal imaging and photothermal anti-tumor therapy. *Journal of Materials
1369 Chemistry B*. 2021;9(36):7447-7460.
- 1370 165. Xue N, Zhou C, Chu Z, Chen L, Jia N. Barley leaves mediated biosynthesis of Au
1371 nanomaterials as a potential contrast agent for computed tomography imaging. *Science China
1372 Technological Sciences*. 2021;64(2):433-440.
- 1373 166. Molaei MJ. Turmeric-derived gadolinium-doped carbon quantum dots for
1374 multifunctional fluorescence imaging and MRI contrast agent. *Journal of Luminescence*.
1375 2023;257:119692.
- 1376 167. Amiri A, Fazaeli Y, Zare H, et al. Radiolabeled florescent-magnetic graphene oxide
1377 nanosheets: Probing the biodistribution of a potential PET-MRI hybrid imaging agent for
1378 detection of fibrosarcoma tumor. *Annals of Nuclear Medicine*. 2024;38(5):350-359.
- 1379 168. Alhasan A, Sani SA, Tajuddin HA, et al. Synthesis of I@ MPA-Mn: ZnSe as an
1380 efficient contrast agent for CT/fluorescence bi-modal imaging application. *Radiation Physics
1381 Chemistry*. 2023;209:110947.
- 1382 169. Saladino GM, Kilic NI, Brodin B, et al. Carbon quantum dots conjugated rhodium
1383 nanoparticles as hybrid multimodal contrast agents. *Nanomaterials*. 2021;11(9):2165.
- 1384 170. Deng H, Konopka CJ, Prabhu S, et al. Dextran-mimetic quantum dots for multimodal
1385 macrophage imaging in vivo, ex vivo, and in situ. *ACS nano*. 2022;16(2):1999-2012.



Table 1: Overview of Different Types of NPs with their Examples, Stage of Trial, Applications in Imaging Modalities and Theranostic Properties

S.No	Type of NPs	Examples of Nanoformulations	Stage of trial	Imaging modalities	Theranostic Properties	References
1	Metallic NPs	Fe-AuNPs	<i>In vivo</i>	MRI/CT	Theranostic agent for laser induced photothermal treatment of tumor.	37
		PEGylated Indium NPs	<i>In vivo</i>	PAI	High contrast imaging and photothermal therapy of breast tumor.	152
		Chalcone conjugated Gd chelate (Gd-DO3A-Chal)	<i>In vivo</i>	MRI/Fluorescence optical imaging	Use of amyloid beta targeting contrast agents for the detection of Alzheimer's disease	153
		Gd-based (Activation and Guidance of Irradiation by X-ray) AGuIX NPs	Phase I clinical trial	MRI	Brain tumor imaging	154
		IRDye 800CW coupled hafnium oxide (HfO ₂) nanocrystals	<i>In vivo</i>	X-Ray CT/NIRF imaging	Detection of sentinel lymph nodes	155
2	Magnetic NPs	Single-nanometer iron oxide (SNIO) fabricated collagen-binding peptide (CBP) NPs	<i>In vivo</i>	MRI	Non-invasive detection of liver fibrosis	156
		Superparamagnetic iron oxide NPs (SPIONs)	<i>In vivo</i>	MRI	High contrast imaging of liver and kidney at ultra-low magnetic field (ULF)	157
		Transferrin fabricated magnetic NPs (SPIO-Tf)	<i>In vivo</i>	MRI	Magnetothermally stimulated theranostic NPs	158
		Superparamagnetic iron oxide NPs (SPIONs)	Phase II clinical trial	MRI-lymphography MRI-LG/magnetic-guided axillary ultrasound (MagUS)	Minimally invasive sentinel lymph node biopsy in breast cancer patients reducing the risk of diagnostic surgery.	159
		Ultra-small superparamagnetic iron oxide (USPIO) NPs	Clinical trial	Dynamic contrast enhanced (DCE)-weighted MRI	Detection of head and neck squamous cell carcinomas (HNSCCs)	160





3	Biogenic NPs	PEGylated Pemetrexed and PolyNIPAM decorated AuNPs	<i>In vivo</i>	CT contrast agent	Biocompatible and stable contrast agent for the imaging of breast tumor.	161
		β -cyclodextrin coated and phyllacanthone-loaded magnetic iron oxide NPs ($\text{Fe}_3\text{O}_4@ \beta\text{CD-PHY}$)	<i>In vivo</i>	MRI/PAI	Theranostic application in cancer by the induction of magnetic hyperthermia.	162
		<i>Prosopis farcta</i> derived PtNPs	<i>In vitro</i>	CT	Biocompatible green contrast agents for enhanced bioimaging.	163
		Gd chelated anthocyanin-based NP (ANP-Gd)	<i>In vivo</i>	PAI/MRI	Multifunctional theranostic agent for image guided photothermal therapy of tumor.	164
		Barley leaves derived AuNPs (BL-Au NPs)	<i>In vivo</i>	CT	Biocompatible NPs with good colloidal stability and high X-Ray attenuation capacity.	165
4	Quantum Dots	Turmeric-derived Gd-doped carbon QDs	<i>In vitro</i>	MRI/Fluorescence imaging	Multifunctional biological contrast agents with significant cell penetration ability.	166
		Cysteine modified magnetic graphene oxide nanosheets conjugated cadmium telluride QDs ($\text{GO}@ \text{Fe}_3\text{O}_4\text{-cys-CdTe}$ QDs)	<i>In vivo</i>	PET/MRI	Radiolabelled imaging agent for the diagnosis of fibrosarcoma tumor.	167
		Iohexol conjugated mercaptopropionic acid capped Mn:ZnSe QDs (I@MPA-Mn:ZnSe QDs)	<i>In vitro</i>	CT/Fluorescence imaging	High luminescence photostable cell imaging with good X-ray attenuation ability.	168
		Carbon QDs conjugated rhodium NPs	<i>In vitro</i>	X-ray Fluorescence Computed Tomography (XFCT)	Highly photostable bioimaging agents with low photobleaching property.	169
		Dextran mimetic QDs	<i>In vivo</i>	PET/CT	Single cell imaging capacity with high resolution.	170

Figure legends:

View Article Online
DOI: 10.1039/D5NA00596E

Figure 1: Diagrammatic representation of dual role of magnetic NPs as brain MRI contrast and therapeutic agents. Magnetic NPs can serve as a potent MRI contrast agent for high contrast imaging with spatial resolution for specific detection of brain tumors. In addition to this, magnetic NPs can also serve as potent therapeutic agent for the treatment of brain tumor by inducing magnetic hyperthermia. Magnetic hyperthermia results in nuclear ablation and HSP70 and HSP90 blockade leading to reactive oxygen species (ROS) mediated oxidative DNA damage.

Figure 2: Pictorial illustration of NPs-based PAI guided photothermal therapy of breast cancer. The administration of NPs results in high contrast PAI of breast cancer to provide real time monitoring of breast cancer therapy. The NPs get activated by the NIR light and results in light to heat conversion resulting in generation of ROS and temperature induced oxidative stress for photothermal ablation of tumor.






Figure 1: Diagrammatic representation of dual role of magnetic NPs as brain MRI contrast and therapeutic agents. Magnetic NPs can serve as a potent MRI contrast agent for high contrast imaging with spatial resolution for specific detection of brain tumors. On the other hand, magnetic NPs can also serve as potent therapeutic agent for the treatment of brain tumor by inducing magnetic hyperthermia. Magnetic hyperthermia results in nuclear ablation and HSP70 and HSP90 blockade leading to reactive oxygen species (ROS) mediated oxidative DNA damage.




Figure 2: Pictorial illustration of NPs-based PAI guided photothermal therapy of breast cancer. The administration of NPs results in high contrast PAI of breast cancer to provide real time monitoring of breast cancer therapy. The NPs get activated by the NIR light and results in light to heat conversion resulting in generation of ROS and temperature induced oxidative stress for photothermal ablation of tumor



Data Availability Statement

[View Article Online](#)
DOI: 10.1039/D5NA00596E

The data supporting the findings of this study are available within the manuscript and its supplementary materials. No additional datasets were generated or analyzed for this review article.

