

RSC Chemical Biology

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: S. C. Hatfield, A. N. Mattingley, K. K. Sujeta, L. D. Humphrey, T. Crook, H. Aranda, C. H. Freckleton, J. V. Clayson, C. Renstrom and J. L. Price, *RSC Chem. Biol.*, 2025, DOI: 10.1039/D5CB00326A.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

ARTICLE

Folded-State Compatibility and Unfolded-State Constraint Govern Staple-Based Stabilization: Guidelines from a Coiled-Coil ModelSamantha C. Hatfield,^a Alexa N. Mattingley,^a Kayla K. Sujeta,^a Logan D. Humphrey,^a Taylor Crook,^a Hiram Aranda,^a Christian H. Freckleton,^a Joseph V. Clayson,^a Chase Renstrom,^a Joshua L. Price^{*a}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Peptide stapling has emerged as a powerful strategy for stabilizing protein conformation, improving proteolytic resistance, and enhancing biomolecular recognition. Yet design principles for selecting staple sites remain elusive, so advances in stapling have depended largely on trial and error. Here we establish quantitative guidelines for staple placement by exploiting the well-defined geometry of an α -helical coiled coil to compare alternative staple sites in a controlled way. Using both experimental measurements and molecular simulations, we find that (1) staples that link residue pairs that normally form interhelical salt bridges yield greater stabilization than those linking non-salt-bridged pairs; (2) N-terminal staples are more stabilizing than C-terminal staples, where an existing interhelical disulfide constraint reduces their impact; and (3) mismatches between the staple length and site spacing can cause destabilization by forcing the structure into a compressed, non-native geometry. Together, these results show that staple-based stabilization depends on two underlying factors: *unfolded-state constraint* (the entropic advantage gained when the staple limits how far apart the linked residues can separate in the unfolded ensemble) and *folded-state compatibility* (how well the staple's maximum accessible span matches the native separation of those residues in the folded structure). These principles provide a predictive framework for rational stapled peptide design, offering a path beyond empirical screening toward principle-guided development of stabilized peptide therapeutics.

Introduction

Peptide-based therapeutics continue to expand as a drug class, offering high target specificity and the ability to engage extended protein-protein interaction surfaces that are inaccessible to small molecules,¹⁻⁷ yet their clinical potential is often limited by rapid clearance, conformational instability, and proteolysis.⁸ Macrocyclization or stapling⁹⁻¹² provides a powerful strategy for mitigating these liabilities by covalently linking two residues (referred to here as the *anchor positions*), thereby constraining a peptide in a conformation that resembles its folded or bound state, effectively prepaying part of the entropic cost of folding or binding. Here we use the term "stapling" for any covalent crosslinking or macrocyclization reaction that connects side-chain and/or backbone groups in a peptide or protein, whether the linked residues reside within a single secondary structural element (e.g., a helix) or across a larger folded architecture. Though early applications of stapling focused primarily on α -helical peptides,¹³⁻¹⁵ subsequent work has applied the same concept broadly to non-helical peptides and folded proteins.¹⁶⁻²¹

Diverse chemistries, including lactams^{22, 23} and disulfide linkages,²⁴ along with thiol alkylation/arylation,²⁵⁻³¹ olefin metathesis,^{13-15, 32-34} and azide-alkyne cycloaddition,³⁵⁻⁴¹ have expanded the accessible range of staples, with additional approaches continuing to emerge.⁴²⁻⁵¹ Stapling can improve both conformational stability and proteolytic resistance,⁵²⁻⁵⁴ and in favorable cases also enhances cell-permeability and target binding affinity.^{15, 25, 27, 44, 52, 55-59} Despite these successes, the geometric and energetic determinants that make a given site/linker pair stabilizing remain incompletely understood.

Existing design strategies for stapling (most extensively developed for intrahelical stapling of α -helical peptides) typically rely on empirical site selection and iterative screening of multiple linker lengths.^{13-15, 22-24, 41-43, 48, 51} Collaborative efforts have yielded qualitative "rules of thumb" for intrahelical stapling,⁶⁰ but there remains little mechanistic guidance on how staple geometry maps onto stabilization energy. Comparable uncertainties arise in the development of PROTACs⁶¹⁻⁶⁴ and antibody-drug conjugates,⁶⁵⁻⁶⁷ where some linker choices significantly limit efficacy. Though computational strategies have been applied to optimize linker design in PROTACs,⁶⁸⁻⁷² comparable efforts to derive predictive guidelines for peptide stapling are limited, with little attention to why certain staples succeed at particular sites or whether such outcomes could be anticipated. Our previous work showed that staple-induced increases in conformational stability correlate with enhanced proteolytic resistance,⁵²⁻⁵⁴ suggesting that a quantitative

^a Address here.^b Address here.^c Address here.

† Footnotes relating to the title and/or authors should appear here.

Supplementary Information available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x






Figure 1. A) Sequence and structure of coiled coil **A/B** shown as a ribbon diagram (PDB: 1kd9), with e-positions in subunit **A** and g-positions in subunit **B** highlighted in orange and blue, respectively. Inset shows close-up view of *nsb e/g'* pair 27e/29g' vs. *sb e/g'* pair 27e/22g'. (B) Sequences of stapled variants **s6e/8g'-z4x**, **s13e/15g'-z4x**, **s20e/22g'-z4x**, **s27e/29g'-z4x**, and **s27e/29g'-x4x** (in which the staple is incorporated at an *nsb e/g'* site) and stapled variants **s6e/1g'-x4x**, **s13e/8g'-x4x**, **s20e/15g'-x4x**, **s27e/22g'-z4x**, **s27e/22g'-x4x**, and **s34e/29g'-x4x** (in which the staple is incorporated at an *sb e/g'* site). **N** represents Asn-linked PEG azide **z4**; **X** represents propargylglycine **x**; **N-X** represents staple **z4x** and **X-X** represents staple **x4x**. (C) The structures of staples **z4x** and **x4x** and components **z4**, **x**, and bis-azido PEG **4**.

understanding the structural factors that govern staple-based stabilization would meaningfully accelerate the rational design of stapled peptide and proteins with improved proteolytic stability.

To interrogate the geometric factors that govern staple-based stabilization, we used the disulfide-linked α -helical coiled-coil heterodimer **A/B**⁷³ as a spatially well-defined model system. Others have previously applied interhelical stapling to α -helical coiled coils⁷⁴⁻⁷⁶ or helix bundles⁷⁷⁻⁸⁰, but with limited attention to the energetic consequences of specific site-staple combinations. **A/B** folds cooperatively and exhibits a clean two-state thermal unfolding transition, allowing its conformational stability (ΔG) to be quantified unambiguously. The regular heptad repeat in **A/B** creates predictable inter-residue spacings,⁸¹⁻⁸⁴ effectively providing a molecular ruler for varying the positions of the anchor residues and quantifying how site-staple geometry influences stabilization. Its covalent C-terminal disulfide tether fixes its oligomeric state even when unfolded, enabling differences in folding free energy ($\Delta\Delta G$) to be attributed directly to staple installation. By combining experiment with replica-exchange molecular dynamics simulations, we identify the structural and thermodynamic determinants of staple-based stabilization in **A/B**—principally folded-state compatibility and unfolded-state constraint—providing mechanistic principles that extend beyond this archetypal coiled-coil scaffold and support the rational design

of stapled peptides and proteins. Whereas our prior work on this system (discussed briefly below) examined a small number of representative sites, the present study expands the sequence space systematically, allowing us to isolate the geometric and thermodynamic factors that govern staple-based stabilization in a controlled framework.

Results and Discussion

Stapling at *e/g'* positions: when the staple is just right

In coiled coil **A/B**,⁷³ e residues pair with g' residues from the preceding heptad of the opposite subunit: for example, Glu27e from the fourth heptad of subunit **A** forms a salt bridge with Lys22g' from the third heptad of subunit **B**, rather than with Lys29g' (Figure 1; apostrophes denote residues in subunit **B**). Accordingly, we refer to positions like 27e/22g' as salt-bridged *e/g'* pairs (*sb*) and positions like 27e/29g' as non-salt-bridged *e/g'* pairs (*nsb*). We previously prepared four variants of **A/B** in which an Asn-linked four-unit PEG azide (**z4**) and propargylglycine (**x**) occupy the e and g' positions of *nsb* pairs 6e/8g', 13e/15g', 20e/22g', or 27e/29g', respectively: we call these variants **6e/8g'-z4x**, **13e/15g'-z4x**, **20e/22g'-z4x**, and **27e/29g'-z4x**.^{53, 54} We then stapled **z4** and **x** via copper-catalyzed azide-alkyne cycloaddition (CuAAC), yielding four stapled variants in which the **z4x** triazole staple connects the



Table 1. Observed vs. simulated melting temperatures (T_m) for stapled coiled-coil variants and their non-stapled counterparts; the observed vs. simulated impact of stapling on coiled-coil conformational stability ($\Delta\Delta G$); and simulated C β -C β distances for staple-site residues of each variant at the indicated temperatures, sorted by staple location.^a

| Site | Variant | T_m (°C) | | Impact of stapling $\Delta\Delta G$ (kcal/mol) | | C β -C β Distances | | |
|-------------|----------------------|------------|-------------|--|--------------|--------------------------------|--------------|----------------|
| | | observed | simulated | observed | simulated | 310 K | 450 K | Δ 450 K |
| nsb e/g' | 6e/8g'-z4x | 39.5 ± 0.2 | 81.8 ± 0.1 | | | 10.5 ± 0.1 | 59.5 ± 0.2 | |
| | s6e/8g'-z4x | 69.1 ± 0.2 | 101.8 ± 0.3 | -2.53 ± 0.04 | -2.00 ± 0.06 | 10.09 ± 0.01 | 14.40 ± 0.01 | -45.1 ± 0.2 |
| | 13e/15g'-z4x | 42.1 ± 0.1 | 83.6 ± 0.1 | | | 10.18 ± 0.04 | 50.0 ± 0.3 | |
| | s13e/15g'-z4x | 57.7 ± 0.1 | 92.8 ± 0.2 | -1.33 ± 0.02 | -1.10 ± 0.04 | 10.32 ± 0.01 | 14.65 ± 0.01 | -35.3 ± 0.3 |
| | 20e/22g'-z4x | 41.8 ± 0.2 | 82.9 ± 0.1 | | | 10.45 ± 0.10 | 38.8 ± 0.1 | |
| | s20e/22g'-z4x | 54.3 ± 0.1 | 89.0 ± 0.1 | -1.09 ± 0.02 | -0.84 ± 0.02 | 10.41 ± 0.01 | 14.07 ± 0.02 | -24.8 ± 0.1 |
| | 27e/29g'-z4x | 41.1 ± 0.2 | 85.4 ± 0.1 | | | 11.6 ± 0.2 | 23.22 ± 0.04 | |
| | s27e/29g'-z4x | 48.2 ± 0.1 | 87.5 ± 0.1 | -0.65 ± 0.02 | -0.27 ± 0.02 | 10.58 ± 0.01 | 13.55 ± 0.02 | -9.68 ± 0.04 |
| sb e/g' | 27e/29g'-xx | 39.8 ± 0.2 | 87.4 ± 0.1 | | | 11.5 ± 0.1 | 22.07 ± 0.03 | |
| | s27e/29g'-x4x | 52.9 ± 0.2 | 90.5 ± 0.1 | -1.08 ± 0.03 | -0.48 ± 0.02 | 10.55 ± 0.02 | 13.62 ± 0.02 | -8.45 ± 0.04 |
| | 6e/1g'-xx | 48.7 ± 0.1 | 87.4 ± 0.1 | | | 10.66 ± 0.05 | 62.5 ± 0.4 | |
| | s6e/1g'-x4x | 77.5 ± 0.1 | 103.6 ± 0.1 | -2.96 ± 0.04 | -2.03 ± 0.03 | 9.74 ± 0.02 | 13.61 ± 0.02 | -48.9 ± 0.4 |
| | 13e/8g'-xx | 41.8 ± 0.2 | 82.9 ± 0.1 | | | 10.01 ± 0.04 | 53.9 ± 0.3 | |
| | s13e/8g'-x4x | 75.6 ± 0.1 | 101.7 ± 0.4 | -3.13 ± 0.05 | -1.76 ± 0.08 | 9.52 ± 0.01 | 15.08 ± 0.02 | -38.8 ± 0.3 |
| | 20e/15g'-xx | 43.2 ± 0.1 | 83.8 ± 0.1 | | | 9.45 ± 0.01 | 44.3 ± 0.2 | |
| | s20e/15g'-x4x | 74.4 ± 0.1 | 93.5 ± 0.2 | -3.13 ± 0.04 | -0.80 ± 0.03 | 9.71 ± 0.01 | 14.85 ± 0.02 | -29.4 ± 0.2 |
| f/b' | 27e/22g'-z4x | 43.2 ± 0.1 | 84.6 ± 0.1 | | | 9.46 ± 0.02 | 32.26 ± 0.07 | |
| | s27e/22g'-z4x | 63.6 ± 0.1 | 91.1 ± 0.2 | -2.01 ± 0.03 | -0.79 ± 0.03 | 9.62 ± 0.02 | 14.33 ± 0.01 | -17.93 ± 0.07 |
| | 27e/22g'-xx | 42.5 ± 0.2 | 85.7 ± 0.1 | | | 9.58 ± 0.02 | 31.3 ± 0.1 | |
| | s27e/22g'-x4x | 63.7 ± 0.1 | 92.0 ± 0.1 | -2.07 ± 0.01 | -0.72 ± 0.02 | 9.45 ± 0.01 | 14.61 ± 0.01 | -16.7 ± 0.1 |
| | 34e/29g'-xx | 51.8 ± 0.1 | 89.3 ± 0.1 | | | 10.02 ± 0.08 | 14.98 ± 0.02 | |
| | s34e/29g'-x4x | 61.9 ± 0.1 | 92.8 ± 0.1 | -1.07 ± 0.01 | -0.42 ± 0.02 | 9.06 ± 0.03 | 11.20 ± 0.02 | -3.78 ± 0.03 |
| f/b' | 7f/10b'-xx | 48.9 ± 0.2 | 92.8 ± 0.1 | | | 16.11 ± 0.02 | 49.2 ± 0.3 | |
| | s7f/10b'-x4x | 58.0 ± 0.2 | 98.5 ± 0.4 | -0.64 ± 0.01 | -0.40 ± 0.05 | 11.8 ± 0.4 | 14.73 ± 0.02 | -34.5 ± 0.3 |
| | 14f/17b'-xx | 43.4 ± 0.2 | 88.3 ± 0.1 | | | 16.34 ± 0.01 | 43.0 ± 0.2 | |
| | s14f/17b'-x4x | --- | 75.6 ± 0.9 | --- | 0.66 ± 0.06 | 14.8 ± 0.4 | 14.65 ± 0.02 | -28.4 ± 0.2 |
| | 21f/24b'-xx | 41.9 ± 0.2 | 86.6 ± 0.1 | | | 16.08 ± 0.04 | 32.8 ± 0.1 | |
| | s21f/24b'-x4x | --- | 77.8 ± 0.7 | --- | 0.56 ± 0.06 | 13.8 ± 0.2 | 14.14 ± 0.02 | -18.6 ± 0.1 |
| f/b' | 28f/31b'-xx | 45.7 ± 0.1 | 87.2 ± 0.1 | | | 16.03 ± 0.05 | 17.57 ± 0.02 | |
| | s28f/31b'-x4x | 37.8 ± 0.3 | 82.7 ± 0.2 | 0.62 ± 0.03 | 0.43 ± 0.03 | 11.1 ± 0.2 | 12.79 ± 0.02 | -4.78 ± 0.03 |

^a $\Delta\Delta G$ values (observed and simulated) for each stapled variant are given \pm standard error relative to its non-stapled counterpart at the T_m of the non-stapled counterpart (observed or simulated, as appropriate). Observed T_m and $\Delta\Delta G$ values for stapled variants **s6e/8g'-z4x**, **s13e/15g'-z4x**, **s20e/22g'-z4x**, **s27e/29g'-z4x**, **s27e/29g'-x4x** and their non-stapled counterparts are from reference 53, whereas simulated T_m and $\Delta\Delta G$ values for these variants are from reference 93. All other data in the table are newly reported here. In all cases, observed T_m and $\Delta\Delta G$ and values were obtained from variable temperature CD experiments at 15 μ M protein concentration in 20 mM sodium phosphate buffer (pH 7) + 4.0 M GdnHCl, whereas simulated T_m , $\Delta\Delta G$, and C β -C β distances were obtained from implicit-solvent T-REMD simulations of each variant (see Supporting Information for details). Note that simulated T_m values are higher in magnitude than the corresponding observed T_m values. As we have explained previously in reference 93, this difference likely reflects the fact that our experiments were conducted in high concentrations of denaturant, whereas our simulations were conducted with implicit solvent and without denaturant.

nsb pairs: **s6e/8g'-z4x**, **s13e/15g'-z4x**, **s20e/22g'-z4x**, and **s27e/29g'-z4x**. We used variable-temperature CD to quantify the stability of each stapled variant relative to its non-stapled counterpart, expressing the effect of the **z4x** staple as a change in folding free energy ($\Delta\Delta G$). We found that the **z4x** staple is most stabilizing (-2.53 ± 0.04 kcal/mol) at the most distal nsb

site **6e/8g'**, with progressively smaller effects at sites that are more proximal to the C-terminal Cys33-Cys33' disulfide: -1.33 ± 0.02 kcal/mol at **13e/15g'**; -1.09 ± 0.02 kcal/mol at **20e/22g'**; and -0.65 ± 0.02 kcal/mol at **27e/29g'** (Table 1). This trend echoes our earlier observations that staples that bridge residues farther apart in sequence can confer greater



stabilization than staples at nearby sites.^{53, 54, 85} Related ideas have emerged from theoretical and experimental studies of disulfide-bonded proteins,⁸⁶⁻⁹² which proposed that crosslinks that span longer loops are more stabilizing because they more strongly restrict conformational freedom in the unfolded ensemble. We test this proposal quantitatively using the molecular dynamics simulations described below.

Subsequent implicit-solvent parallel temperature replica exchange molecular dynamics (T-REMD) simulations⁹³ on stapled variants **s6e/8g'-z4x**, **s13e/15g'-z4x**, **s20e/22g'-z4x**, and **s27e/29g'-z4x** and their non-stapled counterparts sampled 16 parallel temperatures from 310 to 450 K, yielding simulated $\Delta\Delta G$ values that follow a similar trend. Although the implicit-solvent model systematically underestimates the absolute $\Delta\Delta G$ values, it reliably captures their relative order. Analysis of the trajectories from each variant provides a compelling rationale for the observations described above. In all trajectories, the Cys33-Cys33' disulfide constrains the C-termini of subunits **A** and **B** to remain close, regardless of stapling. However, in the most distal variant **s6e/8g'-z4x**, the staple also holds their N-termini together, even at elevated temperatures. As a result, the unfolded ensemble of **s6e/8g'-z4x** is more compact and has less conformational freedom than its non-stapled counterpart, leading to a higher probability of refolding, and a greater stability at elevated temperature. In contrast, the staple in the most proximal variant **s27e/29g'-z4x** lies much farther from the N-termini of subunits **A** and **B**, which remain nearly as free to separate as when no staple is present. Consequently, the **z4x** staple in variant **s27e/29g'-z4x** affords less protection from thermal unfolding than in **s6e/8g'-z4x**.

Having established that the **z4x** staple's effects arise from its geometric restraint of the unfolded ensemble (i.e., unfolded-state constraint), we next asked whether a more synthetically convenient staple would show comparable behavior. Our earlier stapling efforts relied on the **z4x** staple,^{53, 54} which requires access to Fmoc-protected derivatives of propargylglycine **x** (commercially available) and of Asn-linked PEG-azide **z4** (synthesized in our lab). As a more synthetically convenient alternative, we envisioned connecting two **x** residues with a single bis-azido PEG via CuAAC: for example, four-unit bis-azido PEG **4** connects two **x** residues to form the **x4x** staple (Figure 1). Previous experiments confirmed that the **x4x** staple is a reasonable surrogate for the **z4x** staple: it is moderately more stabilizing at *nsb* 27e/29g' (-1.08 ± 0.03 kcal/mol vs. -0.65 ± 0.02 kcal/mol), and similarly stabilizing at *sb* 27e/22g' (-2.07 ± 0.03 vs. -2.01 ± 0.02 kcal/mol), giving an average difference of $\Delta\Delta\Delta G = -0.25 \pm 0.03$ kcal/mol for **x4x** vs. **z4x**.⁵³ Previously reported T-REMD simulations of **s27e/29g'-x4x** and **s27e/29g'-z4x**,⁹³ together with new simulations of **s27e/22g'-x4x** and **s27e/22g'-z4x** (Table 1; see Supporting Information), are consistent with these observations: the **x4x** staple is slightly more stabilizing at *nsb* 27e/29g' (-0.48 ± 0.02 kcal/mol vs. -0.27 ± 0.02 kcal/mol), and similarly stabilizing at *sb* 27e/22g' (-0.72 ± 0.02 vs. -0.79 ± 0.03 kcal/mol), giving an average simulated difference of $\Delta\Delta\Delta G = -0.07 \pm 0.02$ kcal/mol for **x4x** vs. **z4x**. Therefore, we consider the **z4x** and **x4x** staples

interchangeable and use the more synthetically convenient **x4x** staple in the new variants described below. DOI: 10.1039/D5CB00326A

We next asked whether staple-based stabilization at salt-bridged (*sb*) e/g' pairs would show a similar dependence on distance from the C-terminal Cys33-Cys33' disulfide as observed previously for non-salt-bridged (*nsb*) e/g' pairs. We explored this possibility by preparing non-stapled variants **6e/1g'-xx**, **13e/8g'-xx**, **20e/15g'-xx**, **27e/22g'-xx**, or **34e/29g'-xx**, in which we incorporated propargylglycine **x** residues at the corresponding *sb* e/g' pairs. We next connected the paired **x** residues in each construct with bis-azido PEG **4** via CuAAC, yielding stapled variants **s6e/1g'-x4x**, **s13e/8g'-x4x**, **s20e/15g'-x4x**, **s27e/22g'-x4x**, or **s34e/29g'-x4x**. Variable temperature CD experiments (Table 1) reveal that **x4x**-based stabilization at *sb* e/g' pairs increases with distance from the C-terminal Cys33-Cys33' disulfide, rising from proximal 34e/29g' (-1.07 ± 0.01 kcal/mol) to more distal 27e/22g' (-2.07 ± 0.03 kcal/mol) and then plateauing beyond 20e/15g' (-3.13 ± 0.04 kcal/mol for 20e/15g'; -3.13 ± 0.05 kcal/mol for 13e/8g'; -2.96 ± 0.04 kcal/mol for 6e/1g'), an observation examined in more detail below. In contrast, at *nsb* e/g' pairs, **z4x** stapling provides stabilization that increases approximately linearly from proximal 27e/29g' to distal 6e/8g', without the leveling off seen at *sb* sites.

T-REMD simulations on these **x4x**-stapled variants and their non-stapled counterparts qualitatively mirror the experimental proximal-to-distal stabilization trend described above for *sb* e/g' sites, but with an important difference: in the simulations, the plateau in staple-based stabilization occurs later: stabilization is weakest at proximal 34e/29g' (-0.42 ± 0.02 kcal/mol), increases modestly at 27e/22g' and 20e/15g' (-0.72 ± 0.02 kcal/mol; -0.80 ± 0.03 kcal/mol), and reaches the largest predicted values at 13e/8g' and 6e/1g' (-1.76 ± 0.08 kcal/mol; -2.03 ± 0.03 kcal/mol). Comparison of simulated and experimental T_m and ΔG values suggests that the simulations underestimate the stability of **s20e/15g'-x4x** relative to the other variants. Because the implicit-solvent model we used does not fully capture site-specific solvation effects, modest discrepancies of this kind are not unexpected. In any case, the purpose of our simulations was to reveal qualitative mechanistic trends rather than to reproduce absolute $\Delta\Delta G$ values. One possible explanation for this discrepancy is that **x4x**-based stabilization at 20e/15g' arises in part from a structural feature not represented in the implicit-solvent model. For example, if stapling at 20e/15g' protects a critical hydrogen bond or other non-covalent interaction from solvent exposure, the resulting stabilization could exceed that predicted by preorganization alone; however, our current data do not provide a definitive explanation.

Both experiments and simulations show that stapling confers greater stabilization at *sb* than at *nsb* e/g' sites that are similarly distal to the C-terminal disulfide, though they differ in the magnitude of this effect (-1.42 ± 0.02 kcal/mol in experiments vs. -0.28 ± 0.03 kcal/mol in simulations). One possible explanation is that the **z4x** and **x4x** staples may be better matched to the folded-state geometry of the *sb* than *nsb* e/g' sites, an idea we refer to here as *folded-state compatibility*.






Figure 2. Ribbon diagrams of median-energy frames extracted from the 310 K and 450 K trajectories from T-REMD simulations of (A) non-stapled variant **6e/8g'-z4x** and its stapled counterpart **s6e/8g'-z4x** (B). Side chains are shown as sticks, with *nsb* site **6e/8g'** highlighted in orange. Cβ-Cβ distances for the *nsb* site **6e/8g'** within these frames are shown with dashed double-headed arrow; Cβ-Cβ distances for the *nsb* site **6e/8g'** are plotted vs. simulation time for each trajectory (gray lines). Orange lines represent average Cβ-Cβ distance within a moving window of the 50 points that surround each corresponding point in the gray line. The solid blue line shows the average Cβ-Cβ distance over the trajectory from $t = 40$ ns to $t = 2$ μs. Dotted blue lines show this trajectory average Cβ-Cβ distance \pm one standard deviation.

Specifically, if these staples comfortably span the side-chain β -carbon distance of the *e/g'* anchor positions (the Cβ-Cβ distance, used here as measure of folded-state anchor spacing) at *sb* sites but approach their maximum span at *nsb* sites, then the resulting geometric strain could attenuate stabilization at *nsb* positions. We explored this possibility by examining (1) the intrinsic folded-state anchor spacing of the prospective *e/g'* staple site in each non-stapled variant; (2) the folded-state anchor spacing at the same site in the corresponding stapled variant; and (3) several independent estimates of the maximum span accessible to the **z4x** and **x4x** staples.

To assess the folded-state compatibility of the **z4x** and **x4x** staples with *nsb* vs. *sb* *e/g'* sites, we compared the Cβ-Cβ distance for the prospective *e/g'* staple site in the folded conformation of each non-stapled variant with the corresponding distance at the same site its stapled counterpart, using the 310 K T-REMD trajectories described above. In the non-stapled variants, these 310 K Cβ-Cβ distances reflect the intrinsic folded-state anchor spacing of each site: *nsb* *e/g'* positions span 10.1–11.6 Å (average 10.86 ± 0.05 Å), whereas *sb* *e/g'* positions span 9.4–10.7 Å (average 9.86 ± 0.02 Å). This difference is consistent with the possibility that the **z4x** and **x4x** staples are well matched to the shorter native spacing of *sb* *e/g'* sites, whereas the slightly longer intrinsic spacing of *nsb* sites might approach or exceed the staples' maximum accessible span. However, in the stapled variants, the 310 K Cβ-Cβ distances remain essentially unchanged—10.1–10.6 Å for *nsb* sites (average 10.39 ± 0.01 Å) and 9.1–9.7 Å for *sb* sites (average

9.52 ± 0.01 Å)—and each adopts a well-folded coiled coil conformation like that of its non-stapled counterpart. Because neither staple distorts native coiled-coil geometry or forces the anchor residues from their intrinsic folded-state positions, we conclude that **z4x** and **x4x** show comparable folded-state compatibility at both *nsb* and *sb* *e/g'* sites.

We sought to validate this conclusion by examining three independent estimates of the staples' maximum accessible span and comparing them to the intrinsic folded-state anchor spacings described above for *nsb* vs. *sb* *e/g'* sites. First, DFT-optimized extended conformations of isolated **z4x** and **x4x** staple models yield Cβ-Cβ distances of 19.3 and 20.4 Å, respectively, values that likely overestimate the true maximum span, because such fully extended conformations are unlikely when the staple is anchored within a coiled coil. Second, MD simulations of unconstrained **z4x** and **x4x** staple models yield average Cβ-Cβ distances of 10.96 ± 0.04 Å and 10.66 ± 0.04 Å, respectively. Third, the **z4x** and **x4x** staples reach substantially larger spans in the 450 K T-REMD trajectories of the stapled variants (Figure 2), where the unfolded peptide chain pulls on the staple from both ends: 13.5–14.7 Å for **z4x** (average 13.83 ± 0.01 Å) and 11.2–15.1 Å for **x4x** (average 14.20 ± 0.01 Å). These unfolded-state Cβ-Cβ distances likely provide the most realistic estimate of maximum accessible staple span. In all cases, the accessible staple lengths exceed the intrinsic folded-state spacing of both *nsb* and *sb* *e/g'* sites, indicating that better folded-state compatibility is not the source of the enhanced






Figure 3. Plots of observed and simulated staple-based stabilization $\Delta\Delta G$ vs. unfolded $\Delta C\beta-C\beta$ at *nsb e/g'* sites (A,B); *sb e/g'* sites (C,D); and all *e/g'* sites (E,F) together. Dotted lines represent fits of the data to a linear equation, with the indicated intercept, slope, and R^2 values.

stabilization observed at *sb* sites, and that that the **z4x** and **x4x** staples can readily span both classes of anchor positions.

Having ruled out folded-state compatibility as the primary source of the different stabilization observed at *nsb* vs. *sb e/g'* sites, we next examined how stapling alters the separation of the anchor residues in the unfolded ensemble. The reduction in unfolded-state (450 K) anchor spacing (i.e., $C\beta-C\beta$ distance) in stapled variants relative to their non-stapled counterparts provides a quantitative measure of *unfolded-state constraint* that we call $\Delta C\beta-C\beta$ (Figure 2). Unfolded-state anchor spacings remain similar across the **z4x**- or **x4x**-stapled variants (Table 1), but their non-stapled counterparts show much greater variation, increasing sharply with distance from the C-terminal disulfide. For example, at non-stapled *nsb e/g'* pairs, unfolded $C\beta-C\beta$ distances are 23.22 ± 0.04 Å (**z4x**) or 22.07 ± 0.03 Å (**x4x**) at proximal 27e/29g', and 38.8 ± 0.1 Å, 50.0 ± 0.3 Å, and 59.5 ± 0.2 Å at 20e/22g', 13e/15g', and distal 6e/8g', respectively. The covalent constraint of the staple collapses these unfolded anchor spacings to the shorter values observed in the stapled variants (Table 1; Figure 2 illustrates this for **s6e/8g-z4x** vs. **6e/8g-z4x**), giving $\Delta C\beta-C\beta$ values that become progressively more negative at more distal sites. For *nsb e/g'* pairs, $\Delta C\beta-C\beta$ is largest in magnitude at distal 6e/8g' (-45.1 ± 0.2 Å) and decreases at 13e/15g' (-35.3 ± 0.3 Å), 20e/22g' (-24.8 ± 0.1 Å), and proximal 27e/29g' (-9.68 ± 0.04 Å for **z4x**; -8.45 ± 0.04 for

x4x), correlating strongly with the observed and simulated magnitudes of staple-based stabilization ($\Delta\Delta G$) at these sites (Figure 3A,B).

We observed similar trends at *sb e/g'* pairs: $\Delta C\beta-C\beta$ is again largest in magnitude at distal 6e/1g' (-48.9 ± 0.4 Å) and decreases at 13e/8g' (-38.9 ± 0.3 Å), 20e/15g' (-29.4 ± 0.2 Å), 27e/22g' (-16.7 ± 0.1 for **x4x**; -17.93 ± 0.07 for **z4x**), and proximal 34e/29g' (-3.78 ± 0.03 Å), correlating strongly with both the observed and simulated $\Delta\Delta G$ values at these sites (Figure 3C,D). Across sites located at comparable distances from the C-terminal disulfide, unfolded $\Delta C\beta-C\beta$ tends to be larger in magnitude for *sb* vs. *nsb e/g'* pairs (Table 1; compare *sb* 6e/1g' with *nsb* 6e/8g'), consistent with the notion that stapling overcomes more conformational entropy at *sb e/g'* sites and therefore might confer greater stabilization. If unfolded-state constraint ($\Delta C\beta-C\beta$) were the sole determinant of $\Delta\Delta G$, then plotting $\Delta\Delta G$ vs. $\Delta C\beta-C\beta$ across all *e/g'* sites should yield a single linear relationship encompassing both *nsb* and *sb* data points. This expectation is borne out in the simulations, where simulated $\Delta\Delta G$ values scale linearly with $\Delta C\beta-C\beta$ across all *nsb* and *sb e/g'* sites combined (Figure 3F, $R^2 = 0.89$); in addition, separate fits of simulated $\Delta\Delta G$ vs. $\Delta C\beta-C\beta$ for the *nsb* and *sb* subsets give similar slopes, intercepts, and R^2 values (Figures 3B,D). In contrast, the experimental $\Delta\Delta G$ values correlate more weakly with $\Delta C\beta-C\beta$ when both *nsb* and *sb e/g'* sites are



considered together (Figure 3E, $R^2 = 0.53$). Separate fits of experimental $\Delta\Delta G$ vs. $\Delta C\beta-C\beta$ for *nsb* and *sb* sites yield similar slopes and R^2 values, but markedly different intercepts (Figures 3A,C), indicating that although $\Delta C\beta-C\beta$ influences stabilization similarly within each class, stapling at *sb* sites is intrinsically more stabilizing than at *nsb* sites, for reasons not explained by unfolded-state constraint alone.

Taken together, these results suggest that unfolded-state constraint, as captured by $\Delta C\beta-C\beta$, provides a useful quantitative descriptor of staple-based stabilization within either the *nsb* or *sb* *e/g'* subset, but does not fully account for the systematic enhancement observed at *sb* *e/g'* positions. Our implicit-solvent T-REMD simulations reproduce the $\Delta\Delta G$ vs. $\Delta C\beta-C\beta$ scaling expected if unfolded-state constraint were the dominant contributor and reveal no systematic difference between *nsb* and *sb* sites, implying that the additional stabilization observed experimentally at *sb* sites arises from effects not captured by the implicit-solvent model. One possibility is that solvent-mediated interactions, differential solvent exposure, or other environment-dependent contributions differ between *sb* and *nsb* *e/g'* sites in ways that are not reflected in $\Delta C\beta-C\beta$ alone. Despite this limitation, plots of observed vs. simulated T_m and $\Delta\Delta G$ values show good overall agreement, indicating that the simulations correctly identify the most and least stabilizing *e/g'* sites, even though they underestimate the additional *sb*-specific stabilization observed experimentally (see Supporting Information).

Sequence separation as a broadly useful surrogate for unfolded-state constraint

The strong correlation between $\Delta\Delta G$ and $\Delta C\beta-C\beta$ at *e/g'* sites suggests that unfolded-state constraint is a key contributor to staple-based stabilization whenever the staple is long enough to accommodate the site's intrinsic folded-state spacing. Because $\Delta C\beta-C\beta$ requires simulation to compute, we wondered whether a simpler, experimentally accessible metric might capture the same underlying effect. Across the *e/g'* positions in **A/B**, the magnitude of $\Delta C\beta-C\beta$ tracks closely with the *sequence separation* between the two anchor residues. This relationship is expected: residues that are widely separated in sequence can access larger average separations in the unfolded ensemble, so constraining them with a staple should yield a larger entropic benefit. To test whether sequence separation serves as a useful predictor beyond the **A/B** coiled coil variants explored so far, we examined previously reported $\Delta\Delta G$ values for stapled variants in several additional protein scaffolds that differ substantially from **A/B**: the WW domain, the SH3 domain, and the HER2 helix bundle affibody. Though these systems differ from each other in topology (**A/B** is a disulfide-bound coiled coil; HER2 affibody is a monomeric three-helix bundle; SH3 and WW domain are both β -sheet rich monomers), a consistent trend emerges cross staple sites. When $\Delta\Delta G$ values from all four scaffolds are plotted against sequence separation, the data fall along a common trajectory: staples between anchor positions that are farther apart in sequence produce greater stabilization, reflecting greater imposed unfolded-state constraint. Consistent with the $\Delta C\beta-C\beta$ analysis above, the **A/B** data




Figure 4. Plot of observed staple-based stabilization ($\Delta\Delta G$) within coiled-coil **A/B**, and previously reported WW, SH3, and HER2 affibody scaffolds vs. the sequence separation of the staple anchor positions shows a linear relationship ($R^2 = 0.63$)

presented here follow this relationship closely, as do the cross-scaffold data, with only modest scaffold-dependent scatter (Figure 4).

These results suggest that sequence separation provides a broadly applicable simulation-free surrogate for the unfolded-state constraint contribution to staple-based stabilization, so long as the staple is geometrically compatible with the folded-state spacing of the prospective anchor positions. Practically, sequence separation offers a rapid and intuitive first-pass screen for selecting candidate staple sites: residues that are close in space in the folded state but far apart in the sequence are expected to yield the greatest entropic benefit upon stapling. Having established the general connection between sequence separation, unfolded-state constraint, and stabilization, we next turned to positions where this framework breaks down: the *f/b'* sites, where intrinsic folded-state spacing exceeds the span of the **z4x** and **x4x** staples. These positions illustrate the contrasting regime where folded-state geometric incompatibility rather than unfolded state entropy dominates the behavior of stapled variants.

Stapling at *f/b'* positions: when the staple is too short

Stapling at *f/b'* positions presents a qualitatively different geometric challenge from stapling at *e/g'* sites and serves as a useful boundary case for assessing the biophysical consequences when staple length becomes limiting. In coiled coil **A/B**, *f* residues in subunit **A** and *b'* residues in subunit **B** point away from the helix-helix interface (Figure 5), placing them substantially farther apart ($C\beta-C\beta$ distance of 15.4 Å for 7*f* and 10*b'* in the **A/B** crystal structure) than the *nsb* or *sb* *e/g'* pairs described above (Table). However, the $C\beta-C\beta$ distance for 7*f*/10*b'* is only slightly beyond the ~14 Å maximum accessible span inferred above for the **z4x** or **x4x** staples (Figure 2, Table 1), and it was unclear what kind of energetic penalty might be






Figure 5. Sequence and structure of coiled coil **A/B** shown as a ribbon diagram (PDB: 1kd9), with f-positions in subunit **A** and b-positions in subunit **B** highlighted in orange and blue, respectively. Inset shows close-up view of f/b' pair 7f/10b'. Also shown are sequences of stapled variants **s7f/10b'-x4x**, **s14f/17b'-x4x**, **s21f/24b'-x4x**, and **s28f/31b'-x4x**. **X** represents propargyl glycine **x** and **X-X** represents staple **x4x**.

incurred as a result. Based on simple distance comparisons alone, such sites might therefore appear marginally compatible with the **z4x** or **x4x** staples. Consistent with this ambiguity, we previously found that stapling at 7f/10b' with **z4x** was stabilizing (-0.61 ± 0.03 kcal/mol),⁵³ but substantially less so than at nearby *nsb* 6e/8g' (-2.53 ± 0.04 kcal/mol), even though the two sites have nearly the same sequence separation. This result suggested that f/b' positions lie near a geometric threshold where small mismatches between staple span and intrinsic folded-state anchor spacing could lead to qualitatively different energetic outcomes. To determine whether this behavior is specific to 7f/10b' or reflects a general failure mode for f/b' sites, and to assess whether such cases could be identified prospectively, we systematically examined the additional f/b' positions in **A/B** using both experiment and T-REMD simulations.

We prepared and characterized stapled variants **s7f/10b'-x4x**, **s21f/24b'-x4x**, and **s28f/31b'-x4x**, together with their non-stapled counterparts, but were unable to prepare stapled **s14f/17b'-x4x** despite repeated efforts. Variable temperature CD experiments reveal that **x4x** stapling is only weakly stabilizing at distal 7f/10b' (-0.23 ± 0.02 kcal/mol); moderately destabilizing at proximal 28f/31b' (0.62 ± 0.03 kcal/mol); and strongly destabilizing at 21f/24b' where **s21f/24b'-x4x** was too unstable to obtain reliable T_m or ΔG values). The synthetic intractability of **s14f/17b'-x4x** is consistent with an expectation of comparable or greater destabilization. Collectively, these results indicate that the intrinsic folded-state geometry of f/b' positions places the anchor residues beyond the effective reach of the **x4x** (and by analogy **z4x**) staples, and that this folded-state geometric incompatibility leads to destabilization rather than stabilization.

f/b' variants

View Article Online

DOI: 10.1039/D5CB00326A

s7f/10b'-x4x

Ac-EVKQLE**X**EVEELESELWHLLENEVARLEKENAEC**X**EA-NH₂

Ac-KV**X**QLKAKVEELKSKLWHLKLNKVARLKKKNAEC**X**KA-NH₂

s14f/17b'-x4x

Ac-EVKQLEAEVEELE**X**ELWHLLENEVARLEKENAEC**X**EA-NH₂

Ac-KVKQLKAKV**X**ELKSKLWHLKLNKVARLKKKNAEC**X**KA-NH₂

s21f/24b'-x4x

Ac-EVKQLEAEVEELESELWHL**X**EVARLEKENAEC**X**EA-NH₂

Ac-KVKQLKAKVEELKSKL**X**HLLKLNKVARLKKKNAEC**X**KA-NH₂

s28f/31b'-x4x


Ac-EVKQLEAEVEELESELWHLLENEVARLE**X**ENAEC**X**EA-NH₂

Ac-KVKQLKAKVEELKSKLWHLKLNKV**X**RLKKKNAEC**X**KA-NH₂


T-REMD simulations corroborate these experimental trends. In the simulations, **x4x** stapling is slightly stabilizing at 7f/10b' (-0.40 ± 0.05 kcal/mol); strongly destabilizing at 14f/17b' (0.66 ± 0.06 kcal/mol) and 21f/24b' (0.56 ± 0.06 kcal/mol); and modestly destabilizing at 28f/31b' (0.43 ± 0.03 kcal/mol; Table 1). Notably, the simulations predict the greatest destabilization at 14f/17b': the same site where we were unable to install the staple experimentally. The geometric origin of this trend becomes clear when comparing the intrinsic folded-state anchor spacing of the non-stapled variants with the maximum effective span of the **x4x** staple. In the unfolded (450 K) conformations of stapled variants **s7f/10b'-x4x**, **s14f/17b'-x4x**, **s21f/24b'-x4x**, and **s28f/31b'-x4x**, the f/b' C β -C β distances converge to 12.79–14.73 Å (average 14.08 ± 0.01 Å) consistent with the maximum effective reach inferred above for **x4x** at e/g' sites. In contrast, in the folded (310 K) conformations of the corresponding non-stapled variants, native f/b' C β -C β distances (16.03–16.34 Å) exceed the ~ 14 Å reach of the **x4x** staple, indicating that the staple is incompatible with the intrinsic folded-state spacing at these positions. Accordingly, stapling at f/b' sites necessarily forces the coiled coil into a compressed non-native geometry that shortens the natural f/b' anchor spacing. Indeed, 310 K T-REMD simulations of the stapled variants show that the f/b' spacing collapses to 11.1–14.8 Å, confirming that the staple imposes a non-native geometry on the folded state. Thus, folded-state incompatibility rather than unfolded-state constraint dominates the behavior of f/b' sites and explains the destabilization seen experimentally and computationally.



nsb e/g' pairs



sb e/g' pairs



f/b' pairs




Figure 6. Ribbon diagrams of median-energy frames from 310 K T-REMD trajectories of stapled variants **s6e/8g'-z4x**, **s13e/15g'-z4x**, **s20e/22g'-z4x**, **s27e/29g'-z4x**, and **s27e/29g'-x4x** (nsb e/g' sites); **s6e/1g'-x4x**, **s13e/8g'-x4x**, **s20e/15g'-x4x**, **s27e/22g'-z4x**, **s27e/22g'-x4x**, and **s34e/29g'-x4x** (sb e/g' sites); **s7f/10b'-x4x**, **s14f/17b'-x4x**, **s21f/24b'-x4x**, and **s28f/31b'-x4x** (f/b' sites), with side chains shown as sticks. Atoms in each stapled variant are colored by the difference in root-mean-square fluctuation (Δ RMSF) for that atom relative to the corresponding atom in its non-stapled counterpart, according to the heat-map scale to the right, where red, green, and blue represent increased (≥ 5.0 Å), unchanged (0.0 Å), and decreased (≤ -5.0 Å) RMSF, respectively. Because the components of the **z4x** staple are present in both stapled and non-stapled variants, we can directly compare of RMSF values for the staple atoms before and after stapling, so the atoms of **z4x** are colored. In contrast, this is not possible for the **x4x** staple, where the atoms of bis-azido PEG 4 are not present in the non-stapled variants; therefore, the atoms of **x4x** are colored gray.

We next examined how this staple-enforced contraction propagates structural distortion through the coiled coil by computing atom-wise root-mean-square fluctuation (RMSF) differences between stapled and non-stapled variants in the 310 K T-REMD simulations (Figure 6). Stapling at distal **7f/10b'** or proximal **28f/31b'** produces modest locally confined RMSF increases, limited primarily to the immediately adjacent

residues: N-terminal to the staple for **7f/10b'** or C-terminal for **28f/31b'**. In contrast, stapling at the centrally located **14f/17b'** and **21f/24b'** positions generates substantial RMSF increases on both sides of the staple. For example, **s21f/24b'-x4x** displays elevated RMSF at both termini, and **s14f/17b'-x4x** exhibits widespread increases throughout the structure, which deviates substantially from ideal coiled-coil geometry (Figure 5). These



differing consequences of folded-state geometric incompatibility at peripheral vs. central f/b' sites suggest that near the termini, the coiled coil can absorb deformation locally, but that in the central region enforced shortening of the intrinsic f/b' anchor spacing propagates into nearby a and d positions, disrupting the hydrophobic core and strongly destabilizing the fold. In contrast, when the enforced and ideal C β -C β distances coincide—as with the folded-state compatible **z4x** or **x4x** staples at the *nsb* or *sb* e/g' sites—the staple does not introduce substantial RMSF changes. Together, these results indicate that the observed destabilization at f/b' pairs represents a predictable failure mode for **x4x** and **z4x** stapling, arising specifically from geometric incompatibility of the maximum accessible staple span with the intrinsic folded-state anchor spacing. Importantly, this incompatibility can be identified using simulation-derived metrics, providing a prospective means to avoid staples that are intrinsically too short for a given site.

Re-examination of previously published data reveals a similar sensitivity to folded-state geometric compatibility. In the **A/B** scaffold, we previously found that a shorter **z2x** staple was destabilizing at the *nsb* site 27e/29g' (0.31 ± 0.01 kcal/mol), a location where its longer counterpart **z4x** was stabilizing (-0.65 ± 0.02 kcal/mol). In contrast, **z2x** was strongly stabilizing at the nearby *sb* 27e/22g' site (-2.04 ± 0.03 kcal/mol), matching the effect of **z4x**, and illustrating that the shorter staple is compatible with the tighter intrinsic *sb* spacing, but not with the wider *nsb* spacing (Table 1). We observed similar behavior in the WW domain: a short two-ethylene-oxide staple linking positions 16 and 19 destabilized the fold by 0.17 ± 0.02 kcal/mol when installed at anchor positions 16 and 19, whereas a five-unit staple at the same positions was stabilizing (-0.29 ± 0.02 kcal/mol). Together with the f/b' results above, these observations support the general principle that staple length must be matched to the intrinsic folded-state anchor spacing at a prospective staple site to avoid introducing geometric strain.

Conclusions

Here we combined experiments and molecular simulations to identify the structural determinants that govern staple-based stabilization in a coiled-coil model system. Three guidelines emerge for staple placement within the **A/B** scaffold:

1. **Salt-bridged vs. non-salt-bridged sites:** Staples at salt-bridged (*sb*) e/g' pairs are more stabilizing than those at non-salt-bridged (*nsb*) pairs positioned at comparable distances from the C-terminal disulfide.
2. **Positional context:** Staples placed farther from the C-terminal disulfide provide greater stabilization than proximal staples, where the pre-existing covalent tether limits their entropic contribution.
3. **Geometric compatibility:** Stabilization requires the staple to match the intrinsic folded-state C β -C β spacing of the prospective site. When the staple is too short it forces the linked residues into a compressed non-native geometry, as observed for **x4x** at f/b' positions.

Although these guidelines were derived from a single coiled-coil scaffold, the two mechanistic principles underlying them appear more broadly applicable to stapled proteins and peptides in general.

1. **Reducing unfolded-state separation enhances stabilization:** Staples are most stabilizing when they substantially restrict separation of the linked residues in the unfolded ensemble. The larger the reduction in unfolded-state distance upon stapling (i.e., the greater the unfolded-state constraint), the larger the entropic benefit. In practice, this means that (i) staples that link residues close in sequence are typically less stabilizing than those that link residues farther apart, and (ii) staples placed near an existing covalent connection (e.g., a disulfide or another staple) have a diminished effect.
2. **Compatibility with folded-state geometry enhances stabilization:** Staples should reinforce, rather than distort, the native folded-state spacing between anchor residues. Across the **A/B** coiled coil and the WW domain, short staples (e.g., **x4x** at f/b' sites and **z2x** at *nsb* e/g' sites in **A/B**, and the two-unit staple at positions 16 and 19 in WW) destabilize the fold when the intrinsic anchor spacing exceeds the staple's accessible span. These observations emphasize a general principle: stabilization requires folded-state compatibility between the intrinsic folded-state spacing of the anchor residues and the finite geometric reach of the staple.

The strong agreement between our experimental measurements and T-REMD simulations suggests straightforward strategies for applying these guidelines prospectively. A useful workflow is: (1) simulate the non-stapled protein; (2) inspect C β -C β distances in both folded and unfolded ensembles, to identify sites that are close in the folded state yet widely separated in the unfolded state; and (3) match these sites to staples of an appropriate span, using the ~ 14 Å effective reach of **x4x** as a reference point and extending the analysis to other linkers by analogous simulations. This simulation-guided approach offers a rational predictive framework for selecting both staple sites and staple lengths, reducing dependence on empirical screening. More broadly, the mechanistic principles of unfolded-state constraint and folded-state compatibility provide a general foundation for designing stabilizing linkers in diverse protein folds, potential applications in enhancing conformational stability, protease resistance, and biomolecular recognition in peptide- and protein-based therapeutics.

Author contributions

S.H. performed stapling, purification, and circular dichroism experiments. S.H. and J.L.P. performed molecular dynamic simulations. All other authors assisted with synthesis of peptides. The manuscript was written and revised by S.H. and J.L.P. All authors have given approval to the final version of the manuscript.



Conflicts of interest

There are no conflicts to declare.

Data availability

The supporting data has been provided as part of the Supplementary information, which contains supporting figures, detailed and complete experimental procedures, compound purification and characterization, variable-temperature circular dichroism data, and molecular dynamic simulation data. Video files showing the last 200 ns of each of the 16 temperature-centric trajectories of each variant are available, as are initial PDB files for beginning T-REMD simulations on each variant (note that these PDB files are the starting non-minimized non-equilibrated models built as described in the Supplementary information)

Acknowledgements

We acknowledge support from NIH NIGMS 2R15GM116055–02.

Notes and references

- (1) Lau, J. L.; Dunn, M. K. Therapeutic peptides: Historical perspectives, current development trends, and future directions. *Bioorgan Med Chem* **2018**, *26*, 2700–2707.
- (2) Muttenthaler, M.; King, G. F.; Adams, D. J.; Alewood, P. F. Trends in peptide drug discovery. *Nat. Rev. Drug Discov.* **2021**, *20*, 309–325.
- (3) Wang, L.; Wang, N.; Zhang, W.; Cheng, X.; Yan, Z.; Shao, G.; Wang, X.; Wang, R.; Fu, C. Therapeutic peptides: current applications and future directions. *Signal Transduct. Target. Ther.* **2022**, *7*, 48.
- (4) Jain, S.; Gupta, S.; Patiyal, S.; Raghava, G. P. S. THPdb2: compilation of FDA approved therapeutic peptides and proteins. *Drug Discov. Today* **2024**, *29*, 104047.
- (5) Arkin, M. R.; Wells, J. A. Small-molecule inhibitors of protein–protein interactions: progressing towards the dream. *Nat. Rev. Drug Discov.* **2004**, *3*, 301–317.
- (6) J. Wilson, A. Inhibition of protein–protein interactions using designed molecules. *Chem. Soc. Rev.* **2009**, *38*, 3289–3300.
- (7) Conte, L. L.; Chothia, C.; Janin, J. The atomic structure of protein–protein recognition sites. *J. Mol. Biol.* **1999**, *285*, 2177–2198.
- (8) Diao, L.; Meibohm, B. Pharmacokinetics and Pharmacokinetic–Pharmacodynamic Correlations of Therapeutic Peptides. *Clin. Pharmacokinet.* **2013**, *52*, 855–868.
- (9) Cram, D. J. Preorganization - from Solvents to Spherands. *Angew. Chem. Int. Ed.* **1986**, *25*, 1039–1057.
- (10) Lehn, J. M. Supramolecular Chemistry - Scope and Perspectives Molecular, Supermolecules, and Molecular Devices. *Angew. Chem. Int. Ed.* **1988**, *27*, 89–112.
- (11) Pedersen, C. J. The Discovery of Crown Ethers (Nobel Lecture). *Angew. Chem. Int. Ed.* **1988**, *27*, 1021–1027.
- (12) Hruby, V. J. Conformational Restrictions of Biologically-Active Peptides Via Amino-Acid Side-Chain Groups. *Life Sci* **1982**, *31*, 189–199.
- (13) Blackwell, H. E.; Grubbs, R. H. Highly efficient synthesis of covalently cross-linked peptide helices by ring-closing metathesis. *Angew. Chem. Int. Ed.* **1998**, *37*, 3281–3284.
- (14) Schafmeister, C. E.; Po, J.; Verdine, G. L. An all-hydrocarbon cross-linking system for enhancing the helicity and metabolic stability of peptides. *J. Am. Chem. Soc.* **2000**, *122*, 5891–5892.
- (15) Walensky, L. D.; Kung, A. L.; Escher, I.; Malia, T. J.; Barbuto, S.; Wright, R. D.; Wagner, G.; Verdine, G. L.; Korsmeyer, S. J. Activation of apoptosis in vivo by a hydrocarbon-stapled BH3 helix. *Science* **2004**, *305*, 1466–1470.
- (16) Mahesh, S.; Adebomi, V.; Muneeswaran, Z. P.; Raj, M. Bioinspired Nitroalkylation for Selective Protein Modification and Peptide Stapling. *Angew. Chem. Int. Ed.* **2020**, *59*, 2793–2801.
- (17) Morewood, R.; Nitsche, C. A biocompatible stapling reaction for generation of constrained peptides. *Chem. Sci.* **2021**, *12*, 669–674.
- (18) Iannuzzelli, J. A.; Bacik, J. P.; Moore, E. J.; Shen, Z. F.; Irving, E. M.; Vargas, D. A.; Khare, S. D.; Ando, N.; Fasan, R. Tuning Enzyme Thermostability via Computationally Guided Covalent Stapling and Structural Basis of Enhanced Stabilization. *Biochemistry* **2022**, *61*, 1041–1054.
- (19) Galan, S. R. G.; Raj, R.; Mamalis, D.; Jones, L. H.; Mohammed, S.; Davis, B. G. The Minimum Protein Staple? - Towards 'bio'-Baldwin's rules inter-phosphosite linking in the MEK1 activation loop. *Chem. Sci.* **2024**, *15*, 1306–1317.
- (20) Wang, N.; Xu, Z. Y.; Shen, H. X.; Cong, W.; Zhang, S.; Hu, H. G.; Li, X. Tyrosinase-Catalyzed Peptide Stapling Using Para-Amino Phenylalanine and Tyrosine Anchoring Residues. *Angew. Chem. Int. Ed.* **2025**, *64*,
- (21) Shaikh, D. S.; Rajput, S.; Kalia, D. Rapid and High-Yielding Disulfide Conjugation at Any Desired Site in Proteins. *J. Am. Chem. Soc.* **2026**,
- (22) Chorev, M.; Roubini, E.; Mckee, R. L.; Gibbons, S. W.; Goldman, M. E.; Caulfield, M. P.; Rosenblatt, M. Cyclic Parathyroid-Hormone Related Protein Antagonists - Lysine-13 to Aspartic-Acid 17 [I to (I + 4)] Side-Chain to Side-Chain Lactamization. *Biochemistry* **1991**, *30*, 5968–5974.
- (23) Osapay, G.; Taylor, J. W. Multicyclic Polypeptide Model Compounds. 2. Synthesis and Conformational Properties of a Highly Alpha-Helical Uncosapeptide Constrained by 3 Side-Chain to Side-Chain Lactam Bridges. *J. Am. Chem. Soc.* **1992**, *114*, 6966–6973.
- (24) Jackson, D. Y.; King, D. S.; Chmielewski, J.; Singh, S.; Schultz, P. G. General-Approach to the Synthesis of Short Alpha-Helical Peptides. *J. Am. Chem. Soc.* **1991**, *113*, 9391–9392.
- (25) Brunel, F. M.; Dawson, P. E. Synthesis of constrained helical peptides by thioether ligation: application to analogs of gp41. *Chem. Commun.* **2005**, 2552–2554.
- (26) Zhang, F. Z.; Sadovski, O.; Xin, S. J.; Woolley, G. A. Stabilization of folded peptide and protein structures via distance matching with a long, rigid cross-linker. *J Am Chem Soc* **2007**, *129*, 14154–14155.
- (27) Wang, Y. X.; Chou, D. H. C. A Thiol-Ene Coupling Approach to Native Peptide Stapling and Macrocyclization. *Angew. Chem. Int. Ed.* **2015**, *54*, 10931–10934.
- (28) Peraro, L.; Siegert, T. R.; Kritzer, J. A. Conformational Restriction of Peptides Using Dithiol Bis-Alkylation. *Methods Enzymol.* **2016**, *580*, 303–332.
- (29) Moore, E. J.; Zorine, D.; Hansen, W. A.; Khare, S. D.; Fasan, R. Enzyme stabilization via computationally guided protein stapling. *Proc. Natl. Acad. Sci. U.S.A.* **2017**, *114*, 12472–12477.
- (30) Rojas, A. J.; Zhang, C.; Vinogradova, E. V.; Buchwald, N. H.; Reilly, J.; Pentelute, B. L.; Buchwald, S. L. Divergent unprotected peptide macrocyclisation by palladium-mediated cysteine arylation. *Chem. Sci.* **2017**, *8*, 4257–4263.



- (31) Dangi, A.; Angera, I. J.; Del Valle, J. R. Intrastrand Peptide Staples That Promote β -Sheet Folding, Self-Assembly, and Amyloid Seeding. *J. Am. Chem. Soc.* **2025**, *147*, 29930–29938.
- (32) Chapman, R. N.; Dimartino, G.; Arora, P. S. A highly stable short alpha-helix constrained by a main-chain hydrogen-bond surrogate. *J. Am. Chem. Soc.* **2004**, *126*, 12252–12253.
- (33) Patgiri, A.; Jochim, A. L.; Arora, P. S. A Hydrogen Bond Surrogate Approach for Stabilization of Short Peptide Sequences in alpha-Helical Conformation. *Acc. Chem. Res.* **2008**, *41*, 1289–1300.
- (34) Verdine, G. L.; Hilinski, G. J. Stapled Peptides for Intracellular Drug Targets. *Methods Enzymol.* **2012**, *503*, 3–33.
- (35) Roice, M.; Johannsen, I.; Meldal, M. High capacity poly(ethylene glycol) based amino polymers for peptide and organic synthesis. *Qsar. Comb. Sci.* **2004**, *23*, 662–673.
- (36) Cantel, S.; Isaad, A. L. C.; Scrima, M.; Levy, J. J.; DiMarchi, R. D.; Rovero, P.; Halperin, J. A.; D'Ursi, A. M.; Papini, A. M.; Chorev, M. Synthesis and conformational analysis of a cyclic peptide obtained via i to i+4 intramolecular side-chain to side-chain azide - Alkyne 1,3-dipolar cycloaddition. *J. Org. Chem.* **2008**, *73*, 5663–5674.
- (37) Lau, Y. H.; Wu, Y. T.; de Andrade, P.; Galloway, W. R. J. D.; Spring, D. R. A two-component 'double-click' approach to peptide stapling. *Nat Protoc* **2015**, *10*, 585–594.
- (38) Haney, C. M.; Werner, H. M.; McKay, J. J.; Horne, W. S. Thermodynamic origin of alpha-helix stabilization by side-chain cross-links in a small protein. *Org. Biomol. Chem.* **2016**, *14*, 5768–5773.
- (39) Tran, P. T.; Larsen, C. O.; Rondbjerg, T.; De Foresta, M.; Kunze, M. B. A.; Marek, A.; Loper, J. H.; Boyhus, L. E.; Knuhtsen, A.; Lindorff-Larsen, K.; et al. Diversity-Oriented Peptide Stapling: A Third Generation Copper-Catalysed Azide-Alkyne Cycloaddition Stapling and Functionalisation Strategy. *Chem. Eur. J.* **2017**, *23*, 3490–3495.
- (40) Scrima, M.; Le Chevalier-Isaad, A.; Rovero, P.; Papini, A. M.; Chorev, M.; D'Ursi, A. M. Cu-I-Catalyzed Azide-Alkyne Intramolecular i-to-(i+4) Side-Chain-to-Side-Chain Cyclization Promotes the Formation of Helix-Like Secondary Structures. *Eur. J. Org. Chem.* **2010**, *2010*, 446–457.
- (41) Kawamoto, S. A.; Coleska, A.; Ran, X.; Yi, H.; Yang, C. Y.; Wang, S. M. Design of Triazole-Stapled BCL9 alpha-Helical Peptides to Target the beta-Catenin/B-Cell CLL/lymphoma 9 (BCL9) Protein-Protein Interaction. *J. Med. Chem.* **2012**, *55*, 1137–1146.
- (42) Kumita, J. R.; Smart, O. S.; Woolley, G. A. Photo-control of helix content in a short peptide. *Proc. Natl. Acad. Sci. U.S.A.* **2000**, *97*, 3803–3808.
- (43) Leduc, A. M.; Trent, J. O.; Wittliff, J. L.; Bramlett, K. S.; Briggs, S. L.; Chirgadze, N. Y.; Wang, Y.; Burris, T. P.; Spatola, A. F. Helix-stabilized cyclic peptides as selective inhibitors of steroid receptor-coactivator interactions. *Proc. Natl. Acad. Sci. U.S.A.* **2003**, *100*, 11273–11278.
- (44) Brunel, F. M.; Zwick, M. B.; Cardoso, R. M. F.; Nelson, J. D.; Wilson, I. A.; Burton, D. R.; Dawson, P. E. Structure-function analysis of the epitope for 4E10, a broadly neutralizing human immunodeficiency virus type 1 antibody. *J. Virol.* **2006**, *80*, 1680–1687.
- (45) Haney, C. M.; Loch, M. T.; Horne, W. S. Promoting peptide alpha-helix formation with dynamic covalent oxime side-chain cross-links. *Chem. Commun.* **2011**, *47*, 10915–10917.
- (46) Ullrich, S.; George, J.; Coram, A. E.; Morewood, R.; Nitsche, C. Biocompatible and Selective Generation of Bicyclic Peptides. *Angew. Chem. Int. Ed.* **2022**, *61*, e202208400.
- (47) Voss, S.; Rademann, J.; Nitsche, C. Peptide–Bismuth Bicycles: In Situ Access to Stable Constrained Peptides with Superior Bioactivity. *Angew. Chem. Int. Ed.* **2022**, *61*, e202113857.
- (48) Morewood, R.; Nitsche, C. Bioinspired peptide stapling generates stable enzyme inhibitors. *Chem. Commun.* **2022**, *58*, 10817–10820.
- (49) Keyes, E. D.; Mifflin, M. C.; Austin, M. J.; Alvey, B. J.; Lovely, L. H.; Smith, A.; Rose, T. E.; Buck-Koehntop, B. A.; Motwani, J.; Roberts, A. G. Chemoselective, Oxidation-Induced Macrocyclization of Tyrosine-Containing Peptides. *J. Am. Chem. Soc.* **2023**, *145*, 10071–10081.
- (50) Coram, A. E.; Morewood, R.; Voss, S.; Price, J. L.; Nitsche, C. Exploring biocompatible chemistry to create stapled and photoswitchable variants of the antimicrobial peptide aurein 1.2. *J. Pept. Sci.* **2024**, *30*, e3551.
- (51) Fang, P. Y.; Pang, W. K.; Xuan, S. H.; Chan, W. L.; Leung, K. C. F. Recent advances in peptide macrocyclization strategies. *Chem. Soc. Rev.* **2024**, *53*, 11725–11771.
- (52) Bird, G. H.; Madani, N.; Perry, A. F.; Princiotto, A. M.; Supko, J. G.; He, X. Y.; Gavathiotis, E.; Sodroski, J. G.; Walensky, L. D. Hydrocarbon double-stapling remedies the proteolytic instability of a lengthy peptide therapeutic. *P Natl Acad Sci USA* **2010**, *107*, 14093–14098.
- (53) Xiao, Q.; Jones, Z. B.; Hatfield, S. C.; Ashton, D. S.; Dalley, N. A.; Dyer, C. D.; Evangelista, J. L.; Price, J. L. Structural guidelines for stabilization of alpha-helical coiled coils via PEG stapling. *RSC Chem Biol* **2022**, *3*, 1096–1104.
- (54) Xiao, Q.; Ashton, D. S.; Jones, Z. B.; Thompson, K. P.; Price, J. L. Long-range PEG Stapling: Macrocyclization for Increased Protein Conformational Stability and Resistance to Proteolysis. *RSC Chem Biol* **2020**, *1*, 273–280.
- (55) Stewart, M. L.; Fire, E.; Keating, A. E.; Walensky, L. D. The MCL-1 BH3 helix is an exclusive MCL-1 inhibitor and apoptosis sensitizer. *Nature Chemical Biology* **2010**, *6*, 595–601.
- (56) Baek, S.; Kutchukian, P. S.; Verdine, G. L.; Huber, R.; Holak, T. A.; Lee, K. W.; Popowicz, G. M. Structure of the Stapled p53 Peptide Bound to Mdm2. *J Am Chem Soc* **2012**, *134*, 103–106.
- (57) Chang, Y. S.; Graves, B.; Guerlavais, V.; Tovar, C.; Packman, K.; To, K. H.; Olson, K. A.; Kesavan, K.; Gangurde, P.; Mukherjee, A.; et al. Stapled alpha-helical peptide drug development: A potent dual inhibitor of MDM2 and MDMX for p53-dependent cancer therapy. *P Natl Acad Sci USA* **2013**, *110*, E3445–E3454.
- (58) Huhn, A. J.; Guerra, R. M.; Harvey, E. P.; Bird, G. H.; Walensky, L. D. Selective Covalent Targeting of Anti-Apoptotic BFL-1 by Cysteine-Reactive Stapled Peptide Inhibitors. *Cell Chem Biol* **2016**, *23*, 1123–1134.
- (59) Meng, G. P.; Pu, J.; Li, Y.; Han, A. X.; Tian, Y. L.; Xu, W.; Zhang, T. H.; Li, X.; Lu, L.; Wang, C.; et al. Design and Biological Evaluation of m-Xylene Thioether-Stapled Short Helical Peptides Targeting the HIV-1 gp41 Hexameric Coiled-Coil Fusion Complex. *J Med Chem* **2019**, *62*, 8773–8783.
- (60) Chandramohan, A.; Josien, H.; Yuen, T. Y.; Duggal, R.; Spiegelberg, D.; Yan, L.; Juang, Y. C. A.; Ge, L.; Aronica, P. G.; Kaan, H. Y. K.; et al. Design-rules for stapled peptides with in vivo activity and their application to Mdm2/X antagonists. *Nat. Commun.* **2024**, *15*, 489.
- (61) Zorba, A.; Nguyen, C.; Xu, Y. R.; Starr, J.; Borzilleri, K.; Smith, J.; Zhu, H. Y.; Farley, K. A.; Ding, W. D.; Schiemer, J.; et al. Delineating the role of cooperativity in the design of potent PROTACs for BTK. *Proc. Natl. Acad. Sci. U.S.A.* **2018**, *115*, E7285–E7292.
- (62) Wurzb, R. P.; Dellamaggiore, K.; Dou, H.; Javier, N.; Lo, M. C.; McCarter, J. D.; Mohl, D.; Sastri, C.; Lipford, J. R.; Ceet, V. J. A "Click



Chemistry Platform" for the Rapid Synthesis of Bispecific Molecules for Inducing Protein Degradation. *J. Med. Chem.* **2018**, *61*, 453–461.

(63) Sun, Y.; Xue, Y. L.; Sun, P. K.; Mu, S. Y.; Liu, H. B.; Sun, Y.; Wang, L.; Wang, J. K.; Wu, T. X.; Yin, W. B.; et al. Discovery of the First Potent, Selective, and In Vivo Efficacious Polo-like Kinase 4 Proteolysis Targeting Chimera Degradator for the Treatment of TRIM37-Amplified Breast Cancer. *J. Med. Chem.* **2023**, *66*, 8200–8221.

(64) Dong, Y. W.; Ma, T. T.; Xu, T.; Feng, Z. Y.; Li, Y. G.; Song, L. L.; Yao, X. J.; Ashby, C. J. r.; Hao, G. F. Characteristic roadmap of linker governs the rational design of PROTACs. *Acta Pharm Sin B* **2024**, *14*, 4266–4295.

(65) Su, D.; Kozak, K. R.; Sadowsky, J.; Yu, S. F.; Fourie-O'Donohue, A.; Nelson, C.; Vandlen, R.; Ohri, R.; Liu, L. N.; Ng, C.; et al. Modulating Antibody-Drug Conjugate Payload Metabolism by Conjugation Site and Linker Modification. *Bioconjugate Chem.* **2018**, *29*, 1155–1167.

(66) Samantasinghar, A.; Sunildutt, N. P.; Ahmed, F.; Soomro, A. M.; Salih, A. R. C.; Parihar, P.; Memon, F. H.; Kim, K. H.; Kang, I. S.; Choi, K. H. A comprehensive review of key factors affecting the efficacy of antibody drug conjugate. *Biomed Pharmacother* **2023**, *161*, 114408.

(67) Su, D. A.; Zhang, D. L. Linker Design Impacts Antibody-Drug Conjugate Pharmacokinetics and Efficacy via Modulating the Stability and Payload Release Efficiency. *Front Pharmacol* **2021**, *12*, 687926.

(68) Igashov, I.; Stärk, H.; Vignac, C.; Schneuing, A.; Satorras, V. G.; Frossard, P.; Welling, M.; Bronstein, M.; Correia, B. Equivariant 3D-conditional diffusion model for molecular linker design. *Nat. Mach. Intell.* **2024**, *6*, 417–427.

(69) Neeser, R. M.; Akdel, M.; Kovtun, D.; Naef, L. Reinforcement Learning-Driven Linker Design via Fast Attention-based Point Cloud Alignment. *arXiv* **2023**, 2303.08166 [cs.LG].

(70) Guo, J.; Knuth, F.; Margreitter, C.; Janet, J. P.; Papadopoulos, K.; Engkvist, O.; Patronov, A. Link-INVENT: generative linker design with reinforcement learning. *Digital Discovery* **2023**, *2*, 392–408.

(71) Imrie, F.; Bradley, A. R.; van der Schaar, M.; Deane, C. M. Deep Generative Models for 3D Linker Design. *J. Chem. Inf. Model.* **2020**, *60*, 1983–1995.

(72) Kudo, G.; Hirao, T.; Harada, R.; Shigeta, Y.; Hirokawa, T.; Yoshino, R. Construction of PROTAC-Mediated Ternary Complex Structure Distribution Profiles Using Extensive Conformational Search. *J. Chem. Inf. Model.* **2025**, *65*, 6939–6948.

(73) Keating, A. E.; Malashkevich, V. N.; Tidor, B.; Kim, P. S. Side-chain repacking calculations for predicting structures and stabilities of heterodimeric coiled coils. *Proc. Natl. Acad. Sci. U.S.A.* **2001**, *98*, 14825–14830.

(74) Wuo, M. G.; Mahon, A. B.; Arora, P. S. An Effective Strategy for Stabilizing Minimal Coiled Coil Mimetics. *J. Am. Chem. Soc.* **2015**, *137*, 11618–11621.

(75) Wang, C.; Li, X.; Yu, F.; Lu, L.; Jiang, X. F.; Xu, X. Y.; Wang, H. X.; Lai, W. Q.; Zhang, T. H.; Zhang, Z. Q.; et al. Site-specific Isopeptide Bridge Tethering of Chimeric gp41 N-terminal Heptad Repeat Helical Trimers for the Treatment of HIV-1 Infection. *Sci. Rep.* **2016**, *6*, 32161.

(76) Wuo, M. G.; Hong, S. H.; Singh, A.; Arora, P. S. Synthetic Control of Tertiary Helical Structures in Short Peptides. *J. Am. Chem. Soc.* **2018**, *140*, 16284–16290.

(77) Ekblad, T.; Tolmachev, V.; Orlova, A.; Lendel, C.; Abrahmsen, L.; Karlstrom, A. E. Synthesis and Chemoselective Intramolecular Crosslinking of a HER2-Binding Affibody. *Biopolymers* **2009**, *92*, 116–123.

(78) Lindgren, J.; Karlstrom, A. E. Intramolecular Thioether Crosslinking of Therapeutic Proteins to Increase Proteolytic Stability. *Chembiochem* **2014**, *15*, 2132–2138.

(79) Nilsson, A.; Lindgren, J.; Karlstrom, A. E. Intramolecular Thioether Crosslinking to Increase the Proteolytic Stability of Affibody Molecules. *Chembiochem* **2017**, *18*, 2056–2062.

(80) Pelay-Gimeno, M.; Bange, T.; Hennig, S.; Grossmann, T. N. In Situ Cyclization of Native Proteins: Structure-Based Design of a Bicyclic Enzyme. *Angew. Chem. Int. Ed.* **2018**, *57*, 11164–11170.

(81) Crick, F. H. C. The Packing of Alpha-Helices - Simple Coiled-Coils. *Acta Crystallogr* **1953**, *6*, 689–697.

(82) Burkhard, P.; Stetefeld, J.; Strelkov, S. V. Coiled coils: a highly versatile protein folding motif. *Trends Cell Biol.* **2001**, *11*, 82–88.

(83) Woolfson, D. N. The design of coiled-coil structures and assemblies. *Adv. Protein Chem.* **2005**, *70*, 79–112.

(84) Lupas, A. N.; Gruber, M. The structure of alpha-helical coiled coils. *Adv. Protein Chem.* **2005**, *70*, 37–78.

(85) Xiao, Q.; Becar, N. A.; Brown, N. P.; Smith, M. S.; Stern, K. L.; Draper, S. R. E.; Thompson, K. P.; Price, J. L. Stapling of two PEGylated side chains increases the conformational stability of the WW domain via an entropic effect. *Org. Biomol. Chem.* **2018**, *16*, 8933–8939.

(86) Flory, P. J. Theory of Elastic Mechanisms in Fibrous Proteins. *J. Am. Chem. Soc.* **1956**, *78*, 5222–5234.

(87) Pace, C. N.; Grimsley, G. R.; Thomson, J. A.; Barnett, B. J. Conformational Stability and Activity of Ribonuclease-T1 with Zero, One, and 2 Intact Disulfide Bonds. *J. Biol. Chem.* **1988**, *263*, 11820–11825.

(88) Betz, S. F. Disulfide Bonds and the Stability of Globular Proteins. *Protein Sci.* **1993**, *2*, 1551–1558.

(89) Dani, V. S.; Ramakrishnan, C.; Varadarajan, R. MODIP revisited: re-evaluation and refinement of an automated procedure for modeling of disulfide bonds in proteins. *Protein Eng.* **2003**, *16*, 187–193.

(90) Bastolla, U.; Demetrius, L. Stability constraints and protein evolution: the role of chain length, composition and disulfide bonds. *Protein Eng. Des. Sel.* **2005**, *18*, 405–415.

(91) Dombkowski, A. A.; Sultana, K. Z.; Craig, D. B. Protein disulfide engineering. *FEBS Lett.* **2014**, *588*, 206–212.

(92) Lister, J. G. R.; Loewen, M. E.; Loewen, M. C.; St-Jacques, A. D. Rational design of disulfide bonds to increase thermostability of catechol 1,2 dioxygenase. *Biotechnol. Bioeng.* **2024**, *121*, 3389–3401.

(93) Price, J. L. Parallel Temperature Replica-Exchange Molecular Dynamics Simulations Capture the Observed Impact of Stapling on Coiled-Coil Conformational Stability. *J. Phys. Chem. B* **2025**, *129*, 866–875.



The supporting data has been provided as part of the Supplementary information, which contains supporting figures, detailed and complete experimental procedures, compound purification and characterization, variable-temperature circular dichroism data, and molecular dynamic simulation data. Video files showing the last 200 ns of each of the 16 temperature-centric trajectories of each variant are available, as are initial PDB files for beginning T-REMD simulations on each variant (note that these PDB files are the starting non-minimized non-equilibrated models built as described in the Supplementary information)

Video files:

(filenames follow the names of each variant in Figures 1 and 4, but with forward slashes and apostrophes omitted):

6e1g-xx.mp4
s6e1g-x4x.mp4
13e8g-xx.mp4
s13e8g-x4x.mp4
20e15g-xx.mp4
s20e15g-x4x.mp4
27e22g-xx.mp4
s27e22g-x4x.mp4
27e22g-z4x.mp4
s27e22g-z4x.mp4
34e29g-xx.mp4
s34e29g-x4x.mp4
7f10b-xx.mp4
s7f10b-x4x.mp4
14f17b-xx.mp4
s14f17b-x4x.mp4
21f24b-xx.mp4
s21f24b-x4x.mp4
28f31b-xx.mp4
s28f31b-x4x.mp4

Initial PDB files:

6e1g-xx.pdb
s6e1g-x4x.pdb
13e8g-xx.pdb
s13e8g-x4x.pdb
20e15g-xx.pdb
s20e15g-x4x.pdb
27e22g-xx.pdb
s27e22g-x4x.pdb
27e22g-z4x.pdb
s27e22g-z4x.pdb
34e29g-xx.pdb
s34e29g-x4x.pdb
7f10b-xx.pdb
s7f10b-x4x.pdb
14f17b-xx.pdb
s14f17b-x4x.pdb
21f24b-xx.pdb
s21f24b-x4x.pdb
28f31b-xx.pdb
s28f31b-x4x.pdb



Initial parameter/topology files for T-REMD simulations:

6e1g-xx.prmtop
s6e1g-x4x.prmtop
13e8g-xx.prmtop
s13e8g-x4x.prmtop
20e15g-xx.prmtop
s20e15g-x4x.prmtop
27e22g-xx.prmtop
s27e22g-x4x.prmtop
27e22g-z4x.prmtop
s27e22g-z4x.prmtop
34e29g-xx.prmtop
s34e29g-x4x.prmtop
7f10b-xx.prmtop
s7f10b-x4x.prmtop
14f17b-xx.prmtop
s14f17b-x4x.prmtop
21f24b-xx.prmtop
s21f24b-x4x.prmtop
28f31b-xx.prmtop
s28f31b-x4x.prmtop

