

# RSC Chemical Biology

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: O. Perzanowska, J. Kowalska and J. Jemielity, *RSC Chem. Biol.*, 2025, DOI: 10.1039/D6CB00033A.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

# Enhancing mRNA Stability and Translational Potential Through Tailored Modifications at the 3' End

New Article Online  
DOI: 10.1039/D6CB00033A

Olga Perzanowska,<sup>a,b</sup> Joanna Kowalska<sup>a\*</sup>, Jacek Jemielity<sup>b\*</sup>

<sup>a</sup> Division of Biophysics, Faculty of Physics, University of Warsaw, Pasteura 5, 02-093, Warsaw, Poland

<sup>b</sup> Centre of New Technologies, University of Warsaw, Banacha 2c, 02-097, Warsaw, Poland

**E-mail:** j.jemielity@cent.uw.edu.pl; Joanna.Kowalska@fuw.edu.pl

## Abstract

Poly(A) tails regulate mRNA turnover and translation through multiple RNA–protein interactions, and deadenylation is the initiating step of the major cytoplasmic decay pathways. Here, we report a simple post-transcriptional strategy to protect the 3' end of synthetic mRNA by enzymatically ligating short, chemically modified 5'-phosphorylated dinucleotides. Using T4 RNA ligase 1, we attached 2'-O-methyl and/or phosphorothioate-containing A- or G-dinucleotides to the 3' end of model oligoadenylates and to an IVT Gaussia luciferase (GLuc) mRNA bearing a ~150-nt poly(A) tail. All ligated products showed strong resistance to human CNOT7-mediated deadenylation *in vitro*, whereas the unmodified control mRNA underwent poly(A) removal. In rabbit reticulocyte lysate, 3'-modified GLuc mRNAs translated comparably to the control, indicating minimal interference with the translational machinery. In mammalian cells (A549, JAWSII and HEK293), protein output depended on the dinucleotide structure and cell type; two adenosine donors—pApA<sub>m</sub> and the D1 phosphorothioate stereoisomer pApsA<sub>m</sub>—consistently performed best, increasing cumulative GLuc production up to 163% in HEK293 and up to 79% in A549. These results establish dinucleotide ligation as a minimal and modular 3'-end engineering approach that enhances resistance to deadenylation and can improve the translational performance of therapeutic mRNA candidates.

## Introduction

The use of *in vitro* transcribed (IVT) mRNAs as potential drugs has been considered since the 1990s.<sup>1</sup> Due to great technological advances, for the last decade mRNA has become an increasingly promising tool in vaccine development<sup>2–4</sup> and protein replacement therapies.<sup>5,6</sup> The recent SARS-CoV-2 pandemic showcased the potential of mRNA-based vaccines in a spectacular fashion.<sup>7,8</sup> The main advantages of mRNA virus vaccines over more conventional methods (such as killed/attenuated virus or DNA-based vaccines) are several: relatively low-cost and fast manufacturing, no risk of infection or insertional mutagenesis and, most importantly for this study, IVT mRNAs can be easily modified in various ways to improve their stability and translational potential *in vivo*.<sup>9–12</sup> Recent work has also shown that engineering the poly(A) tail architecture itself (e.g., segmented poly(A) designs) can improve robustness of IVT mRNA production and increase translatability.<sup>13,14</sup> In addition to virus vaccines, mRNA cancer vaccines are being extensively studied as a non-toxic, highly specific alternative to conventional cancer treatments.<sup>15–17</sup> Still, the core problem of mRNA-based drugs remains – mRNA is a highly unstable molecule due to omnipresent ribonucleases (RNases) that degrade RNA.<sup>18</sup> Although significant achievements have been made in the realm of IVT mRNA proliferation, both at the point of delivery<sup>19</sup> and further cellular metabolism, we still face a number of obstacles, the main of which is the efficiency of *in vivo* protein production after therapeutic mRNA administration.

Regardless of the decay pathway, mRNA degradation in mammalian cells is initiated by deadenylation, which involves the removal of the poly(A) tail located at the 3' end of most



mature eukaryotic mRNAs.<sup>20</sup> Deadenylation is predominantly executed by the CCR4–NOT complex and is increasingly recognized as a post-transcriptional hub that integrates mRNA turnover with translational control through multiple, regulated interactions with RNA-binding proteins and ribosomes.<sup>21–23</sup> The poly(A) tail acts as one of the main regulators of mRNA metabolism<sup>24</sup> by protecting the coding regions of mRNA from degradation<sup>25</sup> and playing an important role in the translation initiation process through protein–RNA interactions.<sup>26,27</sup> Beyond tail length, mounting evidence indicates that poly(A) tails can encode regulatory information through their composition and dynamics, including the presence and positioning of non-adenosine residues, providing a conceptual basis for minimal 3'-end interventions.<sup>28</sup> Both of these features, combined with the non-coding nature of poly(A) tail, make it a perfect target for introducing advantageous modifications to increase stability and translational potential of IVT mRNAs without hindering the binding of the ribosomal machinery. In recent years, several in-depth studies tackling poly(A) modifications for mRNA labeling and increasing mRNA stability have been reported.<sup>13,14,29–32</sup> Transcriptome-wide poly(A) tail profiling has further revealed that non-A residues within and at the termini of poly(A) tails are widespread and dynamic across biological contexts, supporting the notion that modest compositional changes at the 3' end can substantially affect RNA fate.<sup>33</sup> In line with this, poly(A) tail engineering—ranging from chemical modifications to architectural redesign—has emerged as a practical route to improve IVT mRNA performance. This greatly inspired our studies, in addition to our previous work concerning structure–activity relationship between modified poly(A) tail analogs and poly(A) binding protein (PABP).<sup>34</sup> Among other functions, PABP promotes translation initiation by playing an intermediary role between the poly(A) tail and the rest of the translational machinery. Thus, we previously considered investigating the ligand specificity of PABP as an important step in further development of therapeutic mRNAs. In our research, we concluded that several of the investigated nucleotide modifications (2'-O-methyladenosine, 2'-O-fluoroadenosine and interchain phosphorothioates) had the ability to fully inhibit the poly(A) degradation process carried out by CNOT7 deadenylase. Additionally, by strategically placing these modifications within the poly(A) chain, such as at the 3' end, any significant decrease in binding affinity for PABP could be avoided. Importantly, the specificity and rates of CCR4–NOT-mediated deadenylation can be tuned by adaptor proteins and signaling inputs, implying that the impact of 3'-end engineering may vary across cellular contexts.<sup>35</sup>

In this study, we tested a novel chemo-enzymatic approach to the modification of mRNA 3' end. We synthesized two sets of 5'-monophosphorylated dinucleotides, containing adenine (A) or guanine (G) nucleobases. In each set, there were three differently modified compounds, all containing 2'-O-methyl combined either with a regular phosphate (pAp<sub>A</sub>m/pGpG<sub>m</sub>) or a phosphorothioate linkage (two pAp<sub>S</sub>Am stereoisomers or two pGp<sub>S</sub>Gm stereoisomers). First, we incorporated each of the dinucleotide compounds into the 3' end of a short 12-mer oligoadenylate analog (A<sub>12</sub> OA) via enzymatic ligation. 14-mer oligonucleotide products were then tested for susceptibility to specific degradation. The same dinucleotide compounds were then ligated to the 3' end of mRNA encoding *Gaussia Luciferase* reporter protein. Following the ligation, all modified mRNAs were again tested for susceptibility to deadenylase activity. Protein expression experiments with the modified mRNAs were conducted both in the rabbit reticulocyte lysate (RRL) system and three different cell lines. This line of study led to the identification of mRNAs that yield significantly increased protein output compared to unmodified mRNAs.

## Results and discussion


### Short oligoadenylate analogs modified with dinucleotides at the 3' end are fully resistant to deadenylation by recombinant CNOT7



Most 3' end chemical modifications proposed to date require either co-transcriptional incorporation<sup>29</sup> or the preparation of relatively long and complex compounds<sup>31</sup> for the ligation approach. Poly(A) tail engineering is increasingly explored as a practical platform lever to improve IVT mRNA performance, including architectural redesign of tails while leaving the coding region unchanged.<sup>13</sup> We aimed to achieve both increased enzymatic stability and translational potential of mRNA with small, easy-to-obtain compounds that could be incorporated to the 3' end of any RNA chain, from short oligonucleotides to fully functional mRNAs. Drawing from our previous results,<sup>34</sup> we selected two nucleotide modifications: 2'-O-methylation and 5',3'-phosphorothioate to introduce into 5'-monophosphorylated 5',3'-dinucleotides that could act as donors in an enzymatic ligation reaction. Our goal was to achieve a high level of enzymatic protection of the 3' end, with an emphasis on protection against deadenylation, without hindering other important biological processes. The 3' end of the RNA chain can be modified and, therefore, stabilized via enzymatic ligation carried out with T4 RNA Ligase 1. Due to the substrate specificity of T4 RNA Ligase 1, it is possible to attach chemically modified oligonucleotides of up to several dozen subunits in length to the 3' end of any RNA precursor.<sup>31</sup> The only limitation is the availability of the phosphate group at the 5' end of the oligonucleotide donor and a free 3'-OH group at the 3' end of the RNA substrate/acceptor.


As all known eukaryotic deadenylases exhibit high substrate preference<sup>31,36</sup> or indirect dependence<sup>37,38</sup> on polyadenylate chains, we decided to design compounds containing either adenine or guanine nucleobases, in addition to the aforementioned chemical modifications. Incorporating G subunits at the 3' end of RNA chain could provide additional protection against deadenylation by further interfering with recognition and binding by deadenylases.<sup>39</sup> Notably, non-adenosine residues at poly(A) tails are now recognized as widespread and dynamic, supporting the notion that modest 3'-end compositional changes can have disproportionate consequences for RNA fate.<sup>28,33</sup> At the same time, altering the nucleobases of the terminal subunits of the poly(A) tail could impede cellular processes necessary for high mRNA translational output. In addition to achieving improved stability, we intended to identify compounds capable of increasing protein production, and so we considered it important to test different chemical modifications and nucleobase variations.

All 5'-monophosphorylated dinucleotides (pApA<sub>m</sub>, pAp<sub>S</sub>A<sub>m</sub> stereoisomers D1 and D2, pGpG<sub>m</sub>, pGp<sub>S</sub>G<sub>m</sub> stereoisomers D1 and D2; Figure 1) were obtained by solid-phase synthesis on solid support loaded with either 2'-O-Me-A or 2'-O-Me-G phosphoramidites. All compounds were purified via semi-preparative reversed-phase high-performance liquid chromatography (RP-HPLC) and characterized by high-resolution mass spectrometry (HRMS; for full characterization see Supplementary Information). Based on previous reports linking phosphorus stereochemistry in dinucleotides with chromatographic mobility, we established that the stereoisomer D1 has *R<sub>P</sub>* configuration, while stereoisomer D2 – *S<sub>P</sub>*.<sup>40,41</sup>



**Figure 1.** 5'-Monophosphorylated dinucleotides containing chemical modifications studied in this work. D1 and D2 marks refer to the elution order of two P-diastereomers during RP-HPLC separation (D1 is faster,  $R_F$  and  $S_P$ , respectively).

The ligation conditions and the enzymatic deadenylation protocol were optimized on a short oligoadenylate analog ( $A_{12}$ ) before attempting to introduce the modifications to the poly(A) tails of functional mRNA (Figure 2A). For the ligation reaction with T4 RNA Ligase 1, a 5-fold molar excess of dinucleotide donor over oligonucleotide acceptor was enough to accomplish full conversion of the  $A_{12}$  substrate to the 14-mer product overnight at room temperature (Figure S1A).<sup>43</sup> We noticed that in the case of pGp<sub>s</sub>G<sub>m</sub> D2 dinucleotide donor, a small amount of doubly-ligated product was formed (Figure S1B), which arises likely from the fact that the 2'-O-methylation of the ribose hinders, but does not completely block, the formation of 3',5'-phosphodiester bond and elongation of the chain.<sup>42</sup> The main ligation products were in each case purified by RP-HPLC purification and freeze-dried.



**Figure 2.** Modification of 3'-end of short RNA with chemically modified dinucleotides. (A) Schematic representation of 3'-end modification of oligonucleotides via enzymatic ligation followed by stability experiments and PAGE analysis. (B) PAGE analysis of ligation products after degradation experiments with CNOT7 deadenylyase. Samples were separated on 18% polyacrylamide urea gel. Gels were stained with SYBR Gold before scanning.

To test the 14-mer ligation products for their susceptibility to enzymatic degradation, they were subjected to recombinant human CNOT7 deadenylyase. CNOT7 is one of the catalytic subunits of the protein complex CCR4-NOT, which is mainly responsible for the removal of poly(A) tails from cytoplasmic eukaryotic mRNA.<sup>43,44</sup> Beyond its catalytic role, CCR4-NOT functions as a regulatory hub coupling deadenylation with broader mRNP remodeling and translational control, making 3'-end protection strategies particularly attractive in therapeutic mRNA design. Modified oligonucleotides were incubated with CNOT7 for 30 minutes at 37°C.<sup>21-23</sup> Starting at 0 minutes, aliquots were taken from the reaction mixture every 5 minutes, and the reaction progress was quenched by dilution with a Loading Dye/EDTA mixture. Samples were analyzed using high-resolution urea PAGE (Figure 2B). Unlike the unmodified  $A_{12}$  chain, which underwent gradual shortening as a result of enzymatic digestion, all modified 14-mers




remained stable in the presence of CNOT7. These results confirmed our assumptions that incorporating selected, chemically modified dinucleotide structures at the 3' end of polyadenylate chain can protect it from the CNOT7 enzymatic activity and potentially other 3'-5' exonucleases.

### mRNAs modified with dinucleotides at the 3' end are not susceptible to CNOT7 in vitro

Given inevitable differences in reaction conditions when working with full mRNA moieties rather than short oligonucleotides, we made necessary changes to our ligation protocol. The main concern was the lower concentration of mRNA in the solution, or rather the concentration of 3'-OH groups available as acceptors in the reaction. This required a substantial increase in the amount of dinucleotides in the ligation mixture, from a 5-fold molar excess to a 100-fold excess. Also, 15% of PEG800 was added to the reaction mixture to increase molecular crowding and accelerate the ligation process.<sup>45</sup> The incubation time was shortened from overnight to just 1 hour to avoid exposing vulnerable mRNAs to unfavorable conditions.

To facilitate later biological studies, we chose mRNA encoding *Gaussia* Luciferase (GLuc) as a reporter protein for the 3' end modification experiments.<sup>46</sup> Following *in vitro* transcription, GLuc mRNA, capped at the 5' end with cap-1 analog (m<sup>7</sup>GpppA<sub>m</sub>G) and containing a ~150-long poly(A) tail at the 3' end, was purified using RP-HPLC to increase the quality of the samples and to remove double-stranded RNA contaminants.<sup>47</sup> Each ligation reaction was prepared according to the improved protocol (for the full ligation protocol see Experimental Section), including an unmodified control mRNA that was incubated without any 5'-monophosphorylated dinucleotides to ensure an identical preparation process for all mRNA samples (Table 1, Figure 3A). After 1 h of ligation, samples were purified using Monarch RNA cleanup spin columns. Similar to previously described 14-mer oligonucleotides, in order to test the effectiveness of the ligation procedure and examine the mRNAs for resistance to deadenylation, all mRNA samples were subjected to enzymatic degradation by CNOT7 deadenylase (Figure 3B) for 30 min. As per previous results, ligated mRNAs remained stable, whereas the unmodified control sample was shortened by removal of the poly(A) tail.



**Figure 3.** Modification and evaluation of *Gaussia* Luciferase mRNA. (A) 3' end of Gluc mRNA is modified with different dinucleotides via ligation. (B) PAGE analysis of modified mRNAs, along with unmodified control GLuc mRNA, after 30 minutes of incubation with CNOT7. All modified mRNAs remained stable and full-length after incubation with CNOT7, whereas unmodified GLuc was shortened through removal of its poly(A) tail. Samples were




denatured for 5 minutes at 75 °C prior to analysis on 7% polyacrylamide gel with urea in TBE buffer. After separation, the gel was stained with SYBR Gold before scanning. (C) Table of abbreviations. Article Online  
DOI: 10.1039/D6CB00033A


## Translation efficiency in cells is increased by 3'-end dinucleotide modifications of mRNA

Knowing that all ligated mRNAs were highly resistant to deadenylation, we hypothesized that at least some would lead to increased *Gaussia* Luciferase protein production in cells, at least in part by impeding poly(A) shortening and thereby potentially prolonging functional mRNA lifetime.<sup>21–23</sup> At the same time, we wanted to account for differences in translation mechanisms, mRNA decay rates, and transfection efficiencies in different cell lines. To this end, we chose three cell lines of significantly different morphologies and origins. A549 epithelial lung cancer cells are a popular model for testing anti-cancer therapies.<sup>48</sup> JAWSII and other dendritic cells have been long used as models for cellular mediators in anti-cancer vaccinations with capability to induce an immune response against tumors.<sup>49,50</sup> Lastly, HEK 293 kidney cells are highly versatile, easy to transfect, and characterized by an innate high level of translation and protein production.<sup>51</sup>

Prior to cell line experiments, all modified mRNAs were also used in translation efficiency assays in the rabbit reticulocyte lysate (RRL) system. All exhibited translation levels similar to or slightly higher than unmodified GLuc mRNA (Figure S2). Given the nature of the RRL system, in which the cellular machinery responsible for mRNA degradation is reduced to a minimum, these results were satisfactory. Since the mRNA modifications we introduced were intended to increase mRNA stability, this effect would not be detectable in the RRL system. Importantly, based on the result of the experiment, we could assume that none of the 3' end modifications interfered significantly with the translational machinery, at least in the RRL system.

All ligated mRNAs were next transfected into A549, JAWSII and HEK293 cell lines, along with unmodified control mRNA. *Gaussia* Luciferase expression was monitored at several time points (16, 40, 64, 88 h after transfection) by collecting cell culture medium and measuring the activity level of secreted *Gaussia* Luciferase protein. Total protein expression levels of all modified mRNAs were normalized to the unmodified GLuc mRNA expression.





**Figure 4.** Translational properties of IVT mRNAs modified at the 3' end with various stabilizing dinucleotides. (A) GLuc activity in cell culture medium of A549, HEK293 and JAWSII cells measured after 16, 40, 64 and 88 h from transfection with 3'-modified GLuc mRNAs and control unmodified mRNA. Data presented as mean value  $\pm$  SD of 2 (A549) or 3 (HEK293 and JAWSII) independent biological replicates, each consisting of 3 independent transfections. (B) Total protein production of *Gaussia* Luciferase over four days, calculated based on the same results. All translation levels of modified mRNAs were normalized to control GLuc mRNA and presented as mean values  $\pm$  SD. (C) Structures of A1 (pApAm) and A2 (pApsAm D1) dinucleotides that exhibited the best translation enhancement properties in cell lines.

As expected, protein expression varied greatly depending on both mRNA type and cell line (Figure 4). In adenocarcinomic cell line A549, only mRNA **A2** exhibited translational potential higher than the control (79% increase in protein production). The rest were expressed similarly to control mRNA, indicating that either these modifications did not stabilize mRNA in this cell line or any translational enhancement brought about by increased stability was undercut by modifications of the 3' end negatively affecting the translation process (Figure 4A, B). In JAWSII cell line, we also identified one 3' end modification that resulted in enhanced translation (mRNA **A1**, 70%). Interestingly, both D1 (mRNA **A2**) and D2 (mRNA **A3**) stereoisomers of pAp<sub>s</sub>Am decreased the translation levels, in case of D2 by 40%. In the case of mRNAs ligated with guanosine dinucleotides (mRNA **G1**, **G2**, **G3**), their translational output remained similar to the control. In HEK 293 cells, differences in protein expression were most pronounced. Both mRNA **A1** and **A2** led to significantly increased protein expression (by 105% and 163% respectively), while modification in mRNA **A3** once again proved to be the least beneficial in terms of improving the translational potential. mRNA **G1** was also translated very efficiently (118% increase in protein production) and both **G2** and **G3** led to improved expression (by 65% and 49% respectively). The pronounced cell-type dependence is consistent with mechanistic evidence that CCR4–NOT-mediated deadenylation rates can be tuned by adaptor recruitment and signaling inputs, which may modulate how strongly a given 3'-end modification shifts the balance between translation and decay.<sup>35</sup> It is also possible that other general or cell-



specific degradation mechanisms are responsible for the removal of the 3'-terminal dinucleotides from mRNA. Overall, two dinucleotides provided the best results in translational output enhancement: pApA<sub>m</sub> (mRNA **A1**) and pAp<sub>s</sub>A<sub>m</sub> D1 (mRNA **A2**) (Figure 4C). pApA<sub>m</sub> increased translation in both JAWSII and HEK293 cell lines, whereas pAp<sub>s</sub>A<sub>m</sub> D1 provided translation enhancement in A549 and HEK293 cells. pGpG<sub>m</sub> modification also proved beneficial and could be of further interest considering its potentially even greater stability against deadenylases than in case of adenosine dinucleotides.

### 3'-terminal dinucleotide modification increases cytoplasmic stability of mRNA

We next aimed to investigate whether the 3'-terminal modification of mRNA affects its cytoplasmic stability. However, standard methods, such as transfection followed by qPCR, cannot accurately address this question as they are confounded by transfection kinetics and endosome-trapped mRNA. As such, we have microinjected mRNAs encoding a fluorescent reporter (eGFP) directly into cells' cytoplasm and monitored their translational activity over time by confocal microscopy. Single A549 cells were injected with two mRNA variants: unmodified eGFP and modified with A2 (eGFP-pAp<sub>s</sub>A<sub>m</sub> D1). The data on eGFP protein concentration over time were then analyzed using mathematical modeling to estimate protein and mRNA half-lives ( $t_{dp}$  and  $t_{dm}$ , respectively). To verify the assumption of a constant rate of mRNA degradation rate, and thus the possibility of using the eGFP reporter in the study, we used the simplest, first-order kinetic formulas:

$$\frac{d[P]}{dt} = k_p[mRNA] - k_{dp}[P], \text{ Eq. 1}$$

$$\frac{d[mRNA]}{dt} = -k_{dm}[mRNA], \text{ Eq. 2}$$

where [mRNA] and [P] are concentrations of mRNA and protein, respectively;  $k_p = t_{dp}^{-1}$  corresponds to the rate of eGFP production,  $k_{dp} = t_{dp}^{-1}$  is the rate of eGFP degradation, and  $k_{dm} = t_{dm}^{-1}$  is the rate of degradation of mRNA.

Based on the data collected from three independent replicates (Fig. S3), the estimated half-life was roughly twofold longer for modified eGFP-pAp<sub>s</sub>A<sub>m</sub> D1 ( $t_{dm} \sim 66\text{--}71$  h), when compared to its unmodified counterpart ( $t_{dm} \sim 27\text{--}28$  h). This corresponds well with the overall increased protein output of GLuc-pAp<sub>s</sub>A<sub>m</sub> D1 (**A2**) mRNA in A549 cells compared to the unmodified control mRNA and indicates that this type 3'-end modification may significantly increase the cytoplasmic half-life of a mRNA.

## Conclusions

In summary, we developed a simple and efficient post-transcriptional strategy for engineering the 3' end of synthetic mRNA using enzymatic ligation of short, chemically modified 5'-phosphorylated dinucleotides. We designed and synthesized six dinucleotide donors (A- and G-based), each bearing a 2'-O-methyl modification at the second nucleotide and, in selected cases, a stereodefined phosphorothioate linkage. Using T4 RNA ligase 1, these donors were appended to the 3' end of a short oligoadenylate model (A<sub>12</sub>) and to an IVT Gaussia luciferase (GLuc) mRNA carrying a ~150-nt poly(A) tail. In both formats, 3'-end dinucleotide installation conferred strong resistance to CNOT7-mediated deadenylation *in vitro*, whereas the unmodified mRNA control underwent poly(A) removal.

In mammalian cells, the impact on protein output depended on the dinucleotide structure and the cellular context. Among the tested constructs, two adenosine-based donors (**A1** and **A2**) yielded the most consistent increases in reporter production across cell lines. A2 modification



also increased mRNA half-life after microinjection into A549 cells. Taken together, these results demonstrate that minimal 3'-end modifications can effectively protect the poly(A) tail from enzymatic shortening in vitro and, in favorable contexts, increase protein output in living cells without the need for altering the coding sequence. Future work should extend this approach to additional mRNA cargos and delivery settings, including in vivo models, and directly quantify mRNA abundance/functional lifetime in cells to further clarify the relationship between the susceptibility to deadenylation and protein output.

Article Online  
DOI: 10.1039/D6CB00033A

## Experimental section

### Synthesis of modified dinucleotides

Solid-phase synthesis was performed using AKTA Oligopilot synthesizer. Reagents and solvents were sourced commercially. Five phosphoramidite equivalents were used for each coupling step. Oligonucleotides were synthesized on a 5  $\mu\text{mol}$  scale.

Cleavage from the solid support with simultaneous deprotection was performed using AMA (1:1 concentrated aqueous ammonia and concentrated aqueous methylamine) at 50  $^{\circ}\text{C}$  for 1 h, followed by filtration (to remove solid support particles) and freeze-drying. Final deprotection of the 2'-OH group was performed using triethylamine trihydrofluoride in DMSO at 60  $^{\circ}\text{C}$  for 4 h. After precipitation in n-butanol with sodium acetate (final concentration 75  $\mu\text{M}$ ), analogs were purified using an Agilent Tech. Series 1200 HPLC with a Clarity Oligo-RP C-18 column (150  $\times$  10 mm, flow rate 5 mL/min) in triethylammonium acetate (TEAAc) buffer with a linear acetonitrile gradient, coupled with UV-Vis detection at 260 nm. The structure and homogeneity of the synthesized dinucleotides were confirmed using high-resolution mass spectrometry with negative electrospray ionization.

**pApA<sub>m</sub>**: 2'OMe-A(Bz) RNA CPG 1000/110 support (142 mg, 5  $\mu\text{mol}$ ; LinkTech, Biosearch Technologies) was suspended in dry acetonitrile. rA (Bz) CE-phosphoramidite (74 mg, 75  $\mu\text{mol}$ ; LinkTech, Biosearch Technologies) and Bis(2-cyanoethyl)-N,N-diisopropylphosphoramidite (21 mg, 75  $\mu\text{mol}$ , Sigma-Aldrich) were dissolved in dry acetonitrile to a final concentration of 0.2 M. The synthesis product was cleaved, deprotected, and purified according to the general procedure.

**pApsA<sub>m</sub>**: Synthesis was performed using the same procedure as for pApA<sub>m</sub>, but with additional sulfurization step with 3-ethoxy-1,2,4-dithiazoline-5-one (0.05 M solution in acetonitrile) following the first coupling step.

**pGpG<sub>m</sub>**: 2'OMe-G (dmf) RNA CPG 1000/110 support (167 mg, 5  $\mu\text{mol}$ ; LinkTech, Biosearch Technologies) was suspended in dry acetonitrile. Ac-G-CE Phosphoramidite (70 mg, 75  $\mu\text{mol}$ ; GlenResearch) and Bis(2-cyanoethyl)-N,N-diisopropylphosphoramidite (21 mg, 75  $\mu\text{mol}$ , Sigma-Aldrich) were dissolved in dry acetonitrile to a final concentration of 0.2 M. The synthesis product was cleaved, deprotected, and purified according to the general procedure.

**pGpsG<sub>m</sub>**: Synthesis was performed using the same procedure as for pGpG<sub>m</sub>, but with additional sulfurization step with 3-ethoxy-1,2,4-dithiazoline-5-one (0.05 M solution in acetonitrile) following the first coupling step.

### mRNA synthesis and purification

mRNA encoding *Gaussia* luciferase was obtained using pJET\_T7\_GLuc\_31A plasmid, which was previously digested with AarI restriction enzyme (ThermoFisher Scientific). The plasmid was obtained by cloning the T7 promoter sequence and coding sequence of *Gaussia* luciferase



into pJET\_luc\_128A.<sup>52</sup> *In vitro* transcription reaction (120  $\mu$ l) was incubated at 37 °C for 3 h in transcription buffer (40 mM Tris-HCl pH 7.9, 20 mM MgCl<sub>2</sub>, 1 mM DTT, and 2 mM spermidine) with 0.125 mg/ml T7 RNA polymerase, 1 U/ $\mu$ l RiboLock RNase Inhibitor, 4 mM ATP/UTP/CTP, 2 mM GTP, 4 mM m<sup>7</sup>GpppA<sub>m</sub>pG (cap 1 analog), 0.002U/ $\mu$ l inorganic pyrophosphatase and 40 ng/ $\mu$ l of the template. After 3h of incubation, 1 U/ $\mu$ l DNase was added to the reaction mixture and the incubation was continued for another 30 min at 37 °C. Reaction was then diluted 2x and crude mRNA was purified with Monarch RNA Cleanup Columns T2050 (New England BioLabs). Quality of the transcript was evaluated on native 1% 1 $\times$  Tris-Borate-EDTA (TBE) agarose gel and the concentration of the sample was determined spectrophotometrically.

The same *in vitro* transcription protocol was used to obtain the eGFP mRNA. Template DNA for eGFP was prepared by AarI digestion (Thermo) of the pJET1.2\_T7\_Egfp\_3'utr-beta-globin\_A128.

The poly(A) tail of mRNA was elongated to approximately 150 adenosines using Poly(A) Polymerase Tailing Kit (LuciGen) and following the standard protocol suggested by the manufacturer. The reaction (~400  $\mu$ l) was incubated for 30 min at 37 °C in buffer (50 mM Tris-HCl, pH 8.0, 0.25 M NaCl, 10 mM MgCl<sub>2</sub>) with 1 mM ATP, 1 U/ $\mu$ l RiboLock RNase Inhibitor, 0.8 U/ $\mu$ l Poly(A) Polymerase and 0.6  $\mu$ g/ $\mu$ l of mRNA. The reaction was stopped by adding EDTA to the final concentration of 11 mM and purified again using the same cleanup columns as previously mentioned. The quality of the mRNA was evaluated on an agarose gel as described previously. Final purification was performed on an Agilent Technologies Series 1200 HPLC using RNASep™ Prep – RNA Purification Column (ADS Biotec) at 55 °C. A linear gradient of buffer B (0.1M triethylammonium acetate, pH 7.0, and 50% of acetonitrile) from 20% to 25% in buffer A (0.1M triethylammonium acetate, pH 7.0) over 20 minutes at 0.9 ml/min flow was used. mRNA was later precipitated from eluted fractions using 1.2 volumes of isopropanol with 0.3 M NaOAc (pH 5.2) and 1  $\mu$ l/ml of glycogen, followed by a washing step with 80% ethanol. mRNA pellets were dried under reduced pressure and dissolved in RNase-free water. The concentration of the final sample was determined spectrophotometrically.

## Ligation protocols

**A<sub>12</sub> ligation:** 5'-phosphorylated modified dinucleotides were ligated to the 3' end of the 12-mer oligoadenylate analog using T4 RNA Ligase 1 by employing the following protocol: ligation reaction mixture (100  $\mu$ l) containing 10  $\mu$ l of 10x concentrated T4 Ligase buffer, 10  $\mu$ l of DMSO, 1 mM ATP, 3 mM DTT, 1.2 U/ $\mu$ l RiboLock RNase Inhibitor, 1 U/ $\mu$ l T4 RNA Ligase 1, 1  $\mu$ M A<sub>12</sub>, 5  $\mu$ M of chosen dinucleotide, was incubated overnight (16 h) at 24 °C. The reaction was stopped by a 10x dilution with water. Before purification, chloroform/phenol extraction was carried out to remove protein from the solution, followed by precipitation in isopropanol. Purification was performed on an Agilent Technologies Series 1200 HPLC using Kinetex 2.6  $\mu$ m C18 column (Phenomenex). A linear gradient of buffer B (0.1M triethylammonium acetate, pH 7.0, and 50% of acetonitrile) from 0% to 30% in buffer A (0.1M triethylammonium acetate, pH 7.0) over 20 minutes and 30% to 50% until 30 minutes, at 1 ml/min flow was used. After purification, samples were freeze-dried and dissolved in water. The concentration of the final sample was determined spectrophotometrically.

**mRNA ligation:** 5'-phosphorylated modified dinucleotides were ligated to the 3' end of the *Gaussia* Luciferase mRNA using T4 RNA Ligase 1 by employing the following protocol: ligation reaction mixture (5  $\mu$ l) containing 0.5  $\mu$ l of 10x concentrated T4 Ligase buffer, 0.5  $\mu$ l of DMSO, 1 mM ATP, 1.2 U/ $\mu$ l RiboLock RNase Inhibitor, 2U/ $\mu$ l T4 RNA Ligase 1, 15% PEG800, 1.5  $\mu$ g mRNA, 81  $\mu$ M of chosen dinucleotide, was incubated for 1 h at 24 °C. Reaction was stopped



by 10x dilution with water, and the mRNA products were purified using Monarch RNA Cleanup Columns T2030 (New England BioLabs). The quality of the modified mRNAs was evaluated on native 1% 1× Tris-Borate-EDTA (TBE) agarose gel, and the concentration of the sample was determined spectrophotometrically. Isolated yield after ligation and purification was usually ~70%.

### CNOT7 degradation assay

For 14-mer oligonucleotide products, CNOT7 digestion experiments were conducted as described previously<sup>34</sup>.

For modified mRNA products, poly(A) tail degradation (deadenylation) reaction (5 µl) was carried out in a reaction buffer (20 mM Tris, 20 mM DTT, 10 mM MgCl<sub>2</sub>, 100 mM KCl, pH 8.0) with 60 ng of chosen mRNA and 0.2 µg/µl of CNOT7<sup>53</sup> at 37 °C for 30 min. The reaction was stopped with EDTA (12.5 mM) and a 10x dilution with water. Reaction products were separated electrophoretically on 7% PAA polyacrylamide gel with urea in TBE buffer. The gel was later stained with SYBR Gold prior to scanning.

### Translation efficiency experiments in Rabbit Reticulocyte Lysate system


Translation efficiency experiments were carried out in the RRL system (Promega). 4 µl of RRL was diluted with 5 µl of buffer (40 µM amino acid mixture Δcysteine, 40 µM amino acid mixture Δmethionine, 2 mM MgCl<sub>2</sub>, 380 mM potassium acetate) and incubated for 1 h at 30 °C. Next, 1 µl of chosen modified *Gaussia* Luciferase mRNA dilution (7.8 pMol – 500 pMol) was added to 9 µl of RRL/buffer mixture and incubated for another 1 h at 30 °C. The reaction was quenched by freezing in liquid nitrogen. To detect luminescence, 50 µl of 10 ng/ml of h-coelenterazine (NanoLight) in PBS buffer was added to 10 µl of thawed translation mixture. Luminescence was measured using a Synergy H1 (BioTek) microplate reader. Linear regression was fitted to the points corresponding to the dependence of luminescence on mRNA concentration. The slope coefficients (a) of the linear equations obtained from the linear fit corresponded to the translation efficiency characteristic of a given mRNA molecule. All data was normalized to the average translation efficiency of unmodified GLuc mRNA.

### Protein expression studies

HEK-293 cells (ATCC CRL-1573) and A549 cells (ATCC CCL-185) were grown in DMEM (Gibco) supplemented with 10% FBS, GlutaMax and 1% penicillin/streptomycin (Gibco) at 5% CO<sub>2</sub> and 37 °C. The murine immature dendritic cell line JAWS II (ATCC CRL-11904) was grown in RPMI 1640 (Gibco) supplemented with 10% FBS, sodium pyruvate (Gibco), 1% penicillin/streptomycin, and 5 ng/ml GM-CSF (PeproTech) at 5% CO<sub>2</sub> and 37 °C. In all experiments, the passage number for the cells ranged from 5 to 25. On the day of transfection, 10<sup>4</sup> cells were seeded in 100 µl of medium per well, without antibiotics, in a 96-well plate. The transfection mixture for each well contained 0.3 µl Lipofectamine MessengerMAX Transfection Reagent (Invitrogen) and 5 ng of mRNA in 10 µl of OptiMEM (Gibco). Transfection lasted 16 h. *Gaussia* Luciferase expression was determined at multiple time points by fully removing the medium from each well and replacing it with fresh medium once every 24 h for 4 days. Luminescence was detected by adding 50 µl of 10 ng/ml of h-coelenterazine in PBS to 10 µl of cell culture medium. Luminescence was measured using a Synergy H1 (BioTek) microplate reader. Total protein expression for each mRNA over 4 days was reported as a mean value ± SD normalized to unmodified *Gaussia* Luciferase mRNA.

### Kinetics of reporter protein and mRNA decay in single cells

A549 Cells were microinjected with either the eGFP or eGFP-pAp<sub>s</sub>A<sub>m</sub> D1 mRNA. Next, the concentration of eGFP was measured at subsequent time points. The measurements were



performed using the NIKON A1 confocal microscopy with the LSM Upgrade KIT from PicoQuant consisting of three pulse laser lines and two single photon avalanche diode (SPAD) detectors. The images of single cells were acquired and stored as the .ptu files; one file per cell per time point. The registered files were analysed using the smICA software<sup>54</sup>. First, each PTU file was filtered to subtract the signal from each fluorescence channel. Next, using the calibration data, we applied the smICA software to estimate the concentrations of fluorescently-labelled dextrans and eGFP at each pixel in the imaged cells. The values for all pixels were averaged, and the mean value was stored. The procedure was repeated at every time point. Finally, the time evolution of normalised mean values of the eGFP concentration was fitted with the kinetic model given by the formula (Eq. 3), derived directly from Eq. 1 and Eq. 2:

$$\frac{C}{C_0} = [RNA]_0 \frac{k_p e^{-k_{dm}(t-t_0)} - e^{-k_{dp}(t-t_0)}}{k_{dp} - k_{dm}}, \text{ Eq. 3}$$

where  $C_0$  denotes the concentration at the first measured point ( $t_0$ ),  $[RNA]_0$  is the concentration of mRNA at  $t = t_0$ ; note  $C_0$  and  $t_0$  factors result from the normalisation of the eGFP concentration data. Both constructs had the same eGFP-encoding sequence. It is therefore justified to assume that the degradation time of the reporter protein is the same for both constructs. Under this assumption, we performed global fitting in which  $k_{dp}$  takes a single global value across all fitted cells (eGFP and eGFP-pApsAm D1 mRNA), while the remaining parameters were fitted individually to each cell's dataset. The data were fitted using the least-squares method with the trust-region reflective algorithm.


## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

We thank Mariusz Czarnocki-Cieciura (International Institute of Molecular and Cell Biology, Warsaw, Poland) for providing purified recombinant human CNOT7 deadenylase. We also thank Karina Kwapiszewska, Grzegorz Bubak, Jarosław Michalski and Robert Hołyst (Institute of Physical Chemistry, Polish Academy of Sciences) for their help with mRNA degradation kinetics studies. This work was supported by Virtual Research Institute Łukasiewicz Research Network—PORT Polish Center for Technology Development project “Horizon for Excellence in mRNA applications in immunOncology” [HERO] financed by the Polish Science Fund and the National Science Centre, Poland (2019/33/B/ST4/01843 to J.J. and 2018/31/B/ST5/03821 to J.K.).


1. Wolff, J. *et al.* Direct Gene Transfer into Mouse Muscle in Vivo. *Science* (1979). **247**, 1465–1468 (1990).
2. Pardi, N., Hogan, M. J., Porter, F. W. & Weissman, D. mRNA vaccines — a new era in vaccinology. *Nat. Rev. Drug Discov.* **17**, 261–279 (2018).
3. Petsch, B. *et al.* Protective efficacy of in vitro synthesized, specific mRNA vaccines against influenza A virus infection. *Nat. Biotechnol.* **30**, 1210–1216 (2012).
4. Pardi, N. *et al.* Zika virus protection by a single low-dose nucleoside-modified mRNA vaccination. *Nature* **543**, 248–251 (2017).




5. Zangi, L. *et al.* Modified mRNA directs the fate of heart progenitor cells and induces vascular regeneration after myocardial infarction. *Nat. Biotechnol.* **31**, 898–907 (2013). View Article Online  
DOI: 10.1039/D6CB00033A
6. Perez-Garcia, C. G. *et al.* Development of an mRNA replacement therapy for phenylketonuria. *Mol. Ther. Nucleic Acids* **28**, 87–98 (2022).
7. Polack, F. P. *et al.* Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *N. Engl. J. Med.* **383**, 2603–2615 (2020).
8. Baden, L. R. *et al.* Efficacy and Safety of the mRNA-1273 SARS-CoV-2 Vaccine. *N. Engl. J. Med.* **384**, 403–416 (2021).
9. Holtkamp, S. *et al.* Modification of antigen-encoding RNA increases stability, translational efficacy, and T-cell stimulatory capacity of dendritic cells. *Blood* **108**, 4009–4017 (2006).
10. Thess, A. *et al.* Sequence-engineered mRNA Without Chemical Nucleoside Modifications Enables an Effective Protein Therapy in Large Animals. *Molecular Therapy* **23**, 1456–1464 (2015).
11. Stepinski, J., Waddell, C., Stolarski, R., Darzynkiewicz, E. & Rhoads, R. E. Synthesis and properties of mRNAs containing the novel ‘anti-reverse’ cap analogs 7-methyl(3′-O-methyl)GpppG and 7-methyl (3′-deoxy)GpppG. *RNA* **7**, 1486–1495 (2001).
12. Kudla, G., Lipinski, L., Caffin, F., Helwak, A. & Zylicz, M. High Guanine and Cytosine Content Increases mRNA Levels in Mammalian Cells. *PLoS Biol.* **4**, e180 (2006).
13. Spiewla, T. *et al.* PolyA tail segmentation improves the stability of the template DNA and increases the translatability of in vitro transcribed mRNA. *Nucleic Acids Res.* **54**, gkaf1412 (2026).
14. Trepotec, Z., Geiger, J., Plank, C., Aneja, M. K. & Rudolph, C. Segmented poly(A) tails significantly reduce recombination of plasmid DNA without affecting mRNA translation efficiency or half-life. *RNA* **25**, 507–518 (2019).
15. Diken, M., Kranz, L. M., Kreiter, S. & Sahin, U. mRNA: A Versatile Molecule for Cancer Vaccines. *Current Issues in Molecular Biology* vol. 22 Preprint at <https://doi.org/10.21775/cimb.022.113> (2017).
16. Coulie, P. G., Van den Eynde, B. J., van der Bruggen, P. & Boon, T. Tumour antigens recognized by T lymphocytes: at the core of cancer immunotherapy. *Nat. Rev. Cancer* **14**, 135–146 (2014).
17. Türeci, Ö. *et al.* Targeting the Heterogeneity of Cancer with Individualized Neoepitope Vaccines. *Clinical Cancer Research* **22**, 1885–1896 (2016).
18. Tsui, N. B. Y., Ng, E. K. O. & Lo, Y. M. D. Stability of Endogenous and Added RNA in Blood Specimens, Serum, and Plasma. *Clin. Chem.* **48**, 1647–1653 (2002).
19. Geall, A. *et al.* Nonviral delivery of self-amplifying RNA vaccines. *Proceedings of the National Academy of Sciences* **109**, 14604–14609 (2012).
20. Chen, C.-Y. A., Ezzeddine, N. & Shyu, A.-B. Messenger RNA half-life measurements in mammalian cells. *Methods Enzymol.* **448**, 335–357 (2008).



21. Caulier, G., Siblino, J., Sène, L., Mauxion, F. & Séraphin, B. The CCR4–NOT complex, a multifaceted sensor of molecular signals instructing eukaryotic mRNA translation and stability. *Nucleic Acids Res.* **53**, gkaf1401 (2025). View Article Online  
DOI: 10.1093/D6CB00033A
22. Collart, M. A., Audebert, L. & Bushell, M. Roles of the CCR4-Not complex in translation and dynamics of co-translation events. *Wiley Interdiscip. Rev. RNA* **15**, e1827 (2024).
23. Raisch, T. & Valkov, E. Regulation of the multisubunit CCR4-NOT deadenylase in the initiation of mRNA degradation. *Curr. Opin. Struct. Biol.* **77**, 102460 (2022).
24. Liu, Y. *et al.* Poly(A) tail length is a major regulator of maternal gene expression during the mammalian oocyte-to-embryo transition. *bioRxiv* 2021.08.29.458052 (2021) doi:10.1101/2021.08.29.458052.
25. Dreyfus, M. & Régnier, P. The Poly(A) Tail of mRNAs: Bodyguard in Eukaryotes, Scavenger in Bacteria. *Cell* **111**, 611–613 (2002).
26. Liu, S. *et al.* Classification and function of RNA–protein interactions. *WIREs RNA* **11**, e1601 (2020).
27. Nicholson, A. L. & Pasquinelli, A. E. Tales of Detailed Poly(A) Tails. *Trends Cell Biol.* **29**, 191–200 (2019).
28. Liu, J. & Lu, F. Beyond simple tails: poly (A) tail-mediated RNA epigenetic regulation. *Trends Biochem. Sci.* **49**, 846–858 (2024).
29. Anhäuser, L., Hüwel, S., Zobel, T. & Rentmeister, A. Multiple covalent fluorescence labeling of eukaryotic mRNA at the poly(A) tail enhances translation and can be performed in living cells. *Nucleic Acids Res.* **47**, e42–e42 (2019).
30. Westerich, K. J. *et al.* Bioorthogonal mRNA labeling at the poly(A) tail for imaging localization and dynamics in live zebrafish embryos. *Chem. Sci.* **11**, 3089–3095 (2020).
31. Aditham, A. *et al.* Chemically Modified mRNAs for Highly Efficient Protein Expression in Mammalian Cells. *ACS Chem. Biol.* <https://doi.org/10.1021/acscchembio.1c00569> (2021) doi:10.1021/acscchembio.1c00569.
32. Son, S., Park, M., Kim, J. & Lee, K. ACE mRNA (Additional Chimeric Element incorporated IVT mRNA) for Enhancing Protein Expression by Modulating Immunogenicity. *Advanced Science* **11**, 2307541 (2024).
33. Liu, Y. *et al.* Remodeling of maternal mRNA through poly (A) tail orchestrates human oocyte-to-embryo transition. *Nat. Struct. Mol. Biol.* **30**, 200–215 (2023).
34. Perzanowska, O., Smietanski, M., Jemielity, J. & Kowalska, J. Chemically modified poly(A) analogs targeting PABP: structure activity relationship and translation inhibitory properties. *Chemistry – A European Journal* **n/a**, (2022).
35. Stowell, J. A. W. *et al.* Phosphorylation-dependent tuning of mRNA deadenylation rates. *Nat. Struct. Mol. Biol.* 1–8 (2025).
36. Wang, H. *et al.* Crystal structure of the human CNOT6L nuclease domain reveals strict poly(A) substrate specificity. *EMBO J.* **29**, 2566–2576 (2010).



37. Zhang, X. *et al.* PARN deadenylase is involved in miRNA-dependent degradation of TP53 mRNA in mammalian cells. *Nucleic Acids Res.* **43**, 10925–10938 (2015). Article Online  
DOI: 10.1039/D6CB00033A
38. Wolf, J. & Passmore, L. A. mRNA deadenylation by Pan2-Pan3. *Biochem. Soc. Trans.* **42**, 184–187 (2014).
39. Lim, J. *et al.* Mixed tailing by TENT4A and TENT4B shields mRNA from rapid deadenylation. *Science (1979)*. **361**, 701 LP – 704 (2018).
40. Burgers, P. M., Eckstein, F. & Hunneman, D. H. Stereochemistry of hydrolysis by snake venom phosphodiesterase. *Journal of Biological Chemistry* **254**, 7476–7478 (1979).
41. Slim, G. & Gait, M. J. Configurationally defined phosphorothioate-containing oligoribonucleotides in the study of the mechanism of cleavage of hammerhead ribozymes. *Nucleic Acids Res.* **19**, 1183–1188 (1991).
42. Maguire, S., Lohman, G. J. S. & Guan, S. A low-bias and sensitive small RNA library preparation method using randomized splint ligation. *Nucleic Acids Res.* **48**, e80–e80 (2020).
43. Wahle, E. & Winkler, G. S. RNA decay machines: deadenylation by the Ccr4-not and Pan2-Pan3 complexes. *Biochim. Biophys. Acta* **1829**, 561–570 (2013).
44. Mostafa, D. *et al.* Essential functions of the CNOT7/8 catalytic subunits of the CCR4-NOT complex in mRNA regulation and cell viability. *RNA Biol.* **17**, 403–416 (2020).
45. Harrison, B. & Zimmerman, S. B. Polymer-stimulated ligation: enhanced ligation of oligo- and polynucleotides by T4 RNA ligase in polymer solutions. *Nucleic Acids Res.* **12**, 8235–8251 (1984).
46. Tannous, B. A. Gaussia luciferase reporter assay for monitoring biological processes in culture and in vivo. *Nat. Protoc.* **4**, 582–591 (2009).
47. Karikó, K., Muramatsu, H., Ludwig, J. & Weissman, D. Generating the optimal mRNA for therapy: HPLC purification eliminates immune activation and improves translation of nucleoside-modified, protein-encoding mRNA. *Nucleic Acids Res.* **39**, e142–e142 (2011).
48. Rajivgandhi, G. *et al.* Enhanced anti-cancer activity of chitosan loaded Morinda citrifolia essential oil against A549 human lung cancer cells. *Int. J. Biol. Macromol.* **164**, 4010–4021 (2020).
49. Heiser, A. *et al.* Autologous dendritic cells transfected with prostate-specific antigen RNA stimulate CTL responses against metastatic prostate tumors. *J. Clin. Invest.* **109**, 409–417 (2002).
50. Jacobson, J. M. *et al.* Dendritic Cell Immunotherapy for HIV-1 Infection Using Autologous HIV-1 RNA: A Randomized, Double-Blind, Placebo-Controlled Clinical Trial. *J. Acquir. Immune Defic. Syndr.* **72**, 31–38 (2016).
51. Thomas, P. & Smart, T. G. HEK293 cell line: A vehicle for the expression of recombinant proteins. *J. Pharmacol. Toxicol. Methods* **51**, 187–200 (2005).
52. Warminski, M. *et al.* Amino-Functionalized 5' Cap Analogs as Tools for Site-Specific Sequence-Independent Labeling of mRNA. *Bioconjug. Chem.* **28**, 1978–1992 (2017).



53. Strzelecka, D. *et al.* Phosphodiester modifications in mRNA poly(A) tail prevent deadenylation without compromising protein expression. *RNA* **26**, 1815–1837 (2020). View Article Online  
DOI: 10.1039/D6CB00033A
54. Kalwarczyk, T. *et al.* smICA: an open source repository for mapping the concentration of fluorescently labeled molecules in living cells on the basis of confocal imaging combined with time-correlated single-photon counting. *arXiv preprint arXiv:2410.00532* (2024).



## Data Availability Statement

View Article Online  
DOI: 10.1039/D6CB00033A

The data supporting the findings of this study are available within the article and its Supplementary Information. Additional data supporting the conclusions of this work are available from the corresponding authors upon reasonable request.

Jacek Jemielity

