


ChemComm

Chemical Communications

rsc.li/chemcomm



ISSN 1359-7345


 Cite this: *Chem. Commun.*, 2025, 61, 3323

 Received 21st November 2024,
Accepted 7th January 2025

DOI: 10.1039/d4cc06176d

rsc.li/chemcomm

Studies that independently investigate [M]–C transmetalation reactions using two different metals are uncommon and yet understanding this reactivity is important to unlocking new synthetic approaches and product classes. Here, we show that the strained [Fe]–C complex, $[(\eta^6\text{-C}_5\text{Me}_4\text{-CH}_2)\text{Fe}(\text{diphosphine})]$ undergoes transmetalation with rhodium(i) and iridium(i) diolefin salts, leading to rapid Fe–C(sp³) bond cleavage and M–C(sp³) (M = Rh or Ir) bond generation.

Carbon-element (C–E) bond forming reactions are key to accessing synthetic diversity.^{1–4} Contributions to the selective generation of C–C bonds, for example, were acknowledged with the 2010 Nobel prize in chemistry.⁵ These metal-mediated transformations have had a measurable impact on the fields of drug development and discovery. The Suzuki–Miyaura coupling reaction, for example, is one of the most utilized across medicinal chemistry.⁶ Such reactions proceed in the presence of a transition metal often *via* stepwise oxidative addition, transmetalation, and reductive elimination.⁷

Transmetalation can be used to forge reactive metal–carbon ([M]–C) bonds that can be later transferred or coupled *via* reductive elimination.^{8,9} Of steps associated with cross-coupling, a detailed mechanistic understanding of transmetalation is comparatively lagging and has consequently been the subject of numerous studies.^{10–15} Intimate knowledge surrounding preferred [M]–C generation routes helps to provide a general landscape for reaction optimization.¹⁶ Electronegativity trends and by-products can sometimes be used to predict transfer propensity and the reaction outcome.

As an example of a strained [Fe]–C bond, we recently reported the preparation of an Fe(II) tucked-in complex,

Ring strain governs transmetalation behaviour at a tucked-in iron complex†

 Connor S. Durfy,[‡] Michelle Huang,[‡] Joseph A. Zurakowski,[‡] Paul D. Boyle^a
and Marcus W. Drover^{‡*}

$[(\eta^6\text{-C}_5\text{Me}_4\text{-CH}_2)\text{Fe}(\text{dnppe})]$ (dnppe = 1,2-bis(di-*n*-propylphosphino)ethane) (**1**).^{17,18} Despite the prevalence of related sandwich complexes, $[(\text{Cp}/\text{Cp}^*)_2\text{M}]$; Cp = C₅H₅[−]; Cp* = C₅Me₅[−], the reactivity of tucked-in compounds, especially those with late 3d elements, remains underexplored due to a size mismatch between the metal and L₂X₂–Cp* ring donor (compared to group 4 metals, for example).^{19,20} With **1** in hand, we wondered whether reaction with suitable metal sources would result in transmetalation, affording a programmable route towards heterometallic Cp*{Fe,M} compounds. This transformation would simultaneously enable a detailed study of Fe–C(sp³) bond cleavage and M–C(sp³) bond generation between two model organometallic molecules, helping to determine how ring strain impacts carbon-transfer chemistry, whilst informing the use of {[Fe]–C} compounds as transmetalation partners.

Herein, we investigate the transmetalation behaviour of **1** with Rh(I) and Ir(I) halides – metals known for application in a wide variety of carbon-element bond-forming cycles (Scheme 1). This reaction leads to rapid Fe–C(sp³) bond cleavage and the formation of new M–C(sp³) (M = Rh or Ir) bonds. This behavior is reversible: the addition of a diphosphine prompts regeneration of complex **1** and produces Cl–M(diphosphine). Intermolecular control reactions between [Cp*Fe(CH₃)(diphosphine)] and Cl–ML_{*n*} (L_{*n*} = diolefin or diphosphine) provide a differential outcome, resulting in clean formation of Fe–Cl and M–C species, pointing toward a heterometallic effect. These findings provide a clear example of Fe-to-Rh or – Ir hydrocarbyl transfer – and its reverse, differentiating inter-*versus* intramolecular transmetalation.

To begin, red C₆D₆ solutions of **1**¹⁷ were reacted with 0.5 equiv. of [Rh(nbd)Cl]₂ (nbd = bicyclo[2.2.1]hepta-2,5-diene) or [M(COD)Cl]₂ (M = Rh or Ir, COD = 1,5-cyclooctadiene), affording ring-opened μ-Cl heterometallic complexes $[(\eta^5\text{-C}_5\text{Me}_4\text{-CH}_2\text{-}\{M(L_n)\})\text{Fe}^{\text{II}}(\text{dnppe})(\mu\text{-Cl})]$ **2**, **3**, and **4**, respectively (Scheme 2). These reactions proceeded similarly, with a notable colour change occurring immediately upon addition – yellow brown for **2/3** and burgundy for **4**. The generation of **2–4** requires


^a Department of Chemistry, Western University, 1151 Richmond St, London, ON, N8K 3G6, Canada. E-mail: marcus.drover@uwo.ca

^b Department of Chemistry and Biochemistry, University of Windsor, 401 Sunset Ave, Windsor, ON, N9B 3P4, Canada

† Electronic supplementary information (ESI) available: Experimental details, spectroscopic data, and computational methods. CCDC 2380939–2380943. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d4cc06176d>

‡ These authors contributed equally.






Scheme 1 (A) Concept: reversible metal-to-metal transmetalation; (B) present work: transmetalation at a bridged heterometallic.

tucked-in ring-opening, giving new $M-C(sp^3)$ bonds *via* Fe-to-M alkyl transmetalation.

The formation of these products is evidenced by a shift in their $^{31}P\{^1H\}$ NMR spectrum, from $\delta_p = 89.6$ ppm (for **1**)¹⁷ to 77.4, 76.7, and 78.1 ppm, for **2–4** respectively. These chemical shifts are similar to mononuclear $[Cp^*Fe(dnppe)Cl]$ ($\delta_p = 79.4$ ppm)²¹ and suggest the formation of a dative Fe–Cl bond. Additionally



Scheme 2 Reactions of **1** with $M(I)$ ($M = Rh, Ir$) precursors to generate heterobimetallic complexes **2–4**.

supporting a $\mu-Cl$ bridge between Fe and Rh, the $^{31}P\{^1H\}$ NMR signature for **2** and **3** appears as a Rh-coupled doublet with $J_{P-Rh} = 3.1$ and 2.7 Hz, respectively.

By 1H NMR spectroscopy, desymmetrization of the Cp^*-Me protons proximal (2, 5-) and distal (3, 4-) to the $[M]-CH_2$ ($M = Rh$ or Ir) bond can be used as an additional means to support a Cp^* -bound heterometallic with $\Delta\delta \sim 1.40$ ppm. We suggest that shielding of the proximal (2, 5-) methyl sites results from adjacent metalation of an electron-rich Rh or Ir center. This shielding effect is additionally observed in the $Cp^*-CH_2-M(L_n)$ group, which shifts from $\delta_H = 2.74$ ppm in **1** to 1.13 ppm in **2**. Metallation of the $Cp^*-CH_2-M(L_n)$ is further cemented by $^{13}C\{^1H\}$ and $^1H-^{13}C\{^1H\}$ HSQC NMR spectroscopy, which for **2**, displays a Rh-coupled doublet at $\delta_C = 27.5$ ppm ($^1J_{C-Rh} = 27.2$ Hz).²² To the best of our knowledge, compounds **2–4** represent the only known $Fe(II)/Rh(I)$ or $Ir(I)$ complexes merged from a single Cp^* -derived ligand scaffold.²³ More broadly, this represents the first programmable route towards such heterometallics using a single tucked-in precursor, providing a vast space for future design.

The structures of **2–4** were confirmed by single-crystal X-ray diffraction analysis (Fig. 1). Each of the three compounds comprises a five-membered $Fe-C-C-M-Cl$ ($M = Rh$ or Ir) ring system having an envelope-type conformation, with the $\mu-Cl$ group occupying the *endo*-position. Across the series, the $Fe-M$ distance increases from $3.816(1)$ Å (**2**) to $3.863(2)$ Å (**3**) to $3.875(1)$ Å (**4**) with most other inter-ring distances staying within *ca.* 0.02 Å of one another. This can be rationalized by the larger diolefin ligand (COD vs. nbd) for $Rh(I)$ and the greater atomic radius of Ir *cf.*, Rh on going from **3** to **4**.

Given the inherent lability of metal-bound diolefin groups, we next became interested in the onwards functionalization chemistry of **2–4** with neutral L-type donor ligands, settling on diphosphines due to a strong drive for $M-P$ bond formation and ease of monitoring by $^{31}P\{^1H\}$ NMR spectroscopy (Scheme 3A). Using the nbd precursor, **2**, treatment with 1,2-bis(diphenylphosphino)ethane (dppe) at -78 °C immediately resulted in the formation of three new resonances by $^{31}P\{^1H\}$ NMR spectroscopy at $\delta_p = 79.2$ for $Fe-dnppe$ as well as 72.9 (dd, $^1J_{Rh-P} = 234.7$ Hz, $^2J_{P-P} = 26.7$ Hz) and 47.0 (dd, $^1J_{Rh-P} = 126.6$ Hz, $^2J_{P-P} = 26.7$ Hz) for the Rh-dppe component, attributed to the product **5**, $[(\eta^5-C_5Me_4-CH_2-\{Rh(dppe)\})Fe^{II}(dnppe)(\mu-Cl)]$ (Scheme 3). A *trans*-influence of the bound Cp^*CH_2 group is borne out in a marked decrease in $^1J_{Rh-P}$ coupling value from 234.7 (*trans*-Cl) to 126.6 Hz (*trans*- Cp^*CH_2).

Single crystals of **5** suitable for analysis by X-ray diffraction confirm a $\mu-Cl$ $Fe-Cl-Rh(dppe)$ complex (Scheme 3B). Complex **5** maintains the longest distance between Fe and Rh ($3.924(1)$ Å), lengthened by nearly 0.11 Å when compared to its nbd precursor **2**. Of the heterobimetallic species generated, compound **5** has the shortest $Fe-C2$ ($2.116(3)$ Å) and $Fe-Cl$ ($2.346(1)$ Å) bond lengths, suggesting the greatest “dissociation” from Rh. This point is further supported by a lengthened $Rh-C1$ bond of $2.142(3)$ for **5** vs. $2.088(2)$ Å for **2**, ascribed to a *trans*-influence of the newly installed phosphorus donor.

Solutions of **5** were found to be unstable at room-temperature, cleanly returning tucked-in complex **1** and mixtures of








Fig. 1 Molecular structures of (A) **2**, (B) **3**, and (C) **4** with ellipsoids drawn at 40% probability. Hydrogen atoms except for those on C(1) are omitted for clarity.




B. Single crystal X-ray diffraction of **5**



Scheme 3 (A) Generation of complex **5** and onwards Rh-to-Fe–C(sp³) bond exchange; (B) the molecular structure of **5** with ellipsoids drawn at 40% probability. Hydrogen atoms are omitted except for those on C(1).

[Rh(dppe)(μ-Cl)]₂ and [Rh(dppe)₂]Cl (Scheme 3A).²⁴ Given the preparative route used to access **2**: reaction of **1** with 0.5 equiv. [Rh(nbd)(μ-Cl)]₂, one might conclude that ring-opening or -closing (to return **1**) is apparently dictated by group 9 metal ligand type *i.e.*, Rh(diolefin) vs. Rh(diphosphine). This reaction is accelerated by the addition of excess dppe causing the formation of tucked-in complex **1** and [Rh(dppe)₂]Cl, which precipitates from solution (Scheme 3A) – possibly *via* the unobserved intermediacy of five-coordinate complex **6**. Examples of this elementary transformation, donor-induced transmetalation, where M = metal:




are unusual with the present case benefiting from the leaving group propensity of **1**.

To assess the favourability of this reaction the dissociation of **5** was modelled computationally. This reaction was found to be roughly thermoneutral ($\Delta G^\circ = -0.3 \text{ kcal mol}^{-1}$) having a positive value of $\Delta H^\circ = 16.8 \text{ kcal mol}^{-1}$ (see ESI†).

To explore the relationship between Fe–C bond strain and transmetalation outcome, the reactivity of the unstrained model, [Cp*Fe^{II}(dnppe)(CH₃)] with [Rh(μ-Cl)(nbd)]₂ was also tested. In this case, Fe–C(sp³) bond cleavage results to give [Cp*Fe^{II}(dnppe)(Cl)] by ³¹P{¹H} NMR spectroscopy. By ¹H NMR spectroscopy, the related Rh(i)–CH₃ compound [Rh(nbd)(μ-CH₃)₂] is not observed. However, the observation of CH₄ is consistent with its implied intermediacy. Indeed, related reactions of [Rh(diolefin)(μ-Cl)]₂ and CH₃Li, to generate bridging alkyl Rh complexes, were reported as early as 1987 by Andersen and Muettterties.^{25,26} Unlike the Rh(i) alkyl diolefin compounds **2** and **3** presented here, [Rh(COD)(μ-CH₃)₂] requires cryogenic preparation and storage. The authors note that the decomposition of this compound occurs at temperatures as low as 0 °C *via* elimination of CH₄. Relatedly, they report that efforts to synthesize the nbd analogue, “[Rh(nbd)(μ-CH₃)₂],” result in product decomposition *via* elimination of CH₄ at –30 °C, thwarting isolation.

Expanding our study to the 5d congener Ir(i), treatment of **4** with dppe was pursued (Scheme 4A). For this reaction, however, Ir–COD coordination was maintained, as evidenced by a multiplet in the ¹H NMR spectrum at $\delta_H = 3.48 \text{ ppm}$ (4H) (and the absence of free COD). Moreover, the observation of two phosphorus resonances at $\delta_P = 79.8$ and 27.1 ppm of integration 2 : 2 suggests the formation of a terminal [(η⁵-C₅Me₄-CH₂-{Ir(dppe)(COD)})Fe^{II}(dnppe)(Cl)] (**7**) (Scheme 4A) *cf.*, bridging chloride, where in the case of the latter, three distinct ³¹P NMR resonances would be expected due to local C_s-symmetry (as seen for **5**). Speaking to COD lability, single-crystal X-ray diffraction analysis of **7** revealed the μ-Cl analogue of **5** – [(η⁵-C₅Me₄-CH₂-{Ir(dppe)})Fe^{II}(dnppe)(μ-Cl)] (**8**). Chloride interaction (bridging or terminal) in **7**, however, does not dictate group 9 metal loss, which generates the five-coordinate





Scheme 4 (A) Generation of complex **7** and onwards Ir-to-Fe–C(sp³) bond exchange. (B) An unstrained model undergoes clean transmetalation.

[Ir(dppe)(COD)(Cl)] ($\delta_{\text{P}} = 34.4$ ppm) complex (along with **1**) over time in solution. Consistent with the forward direction of this process, this reaction is irreversible – treatment of **1** with [Ir(dppe)(COD)(Cl)] does not result in **7**.

As an intra- vs. intermolecular point of comparison, reactivity of the unstrained model, [Cp*Fe^{II}(dnppe)(CH₃)] with [Ir(dppe)(COD)(Cl)] was also tested (Scheme 4B). In this case, the Fe–CH₃ complex underwent clean methyl transfer to give [Ir(dppe)(COD)(CH₃)]²⁷ and [Cp*Fe^{II}(dnppe)(Cl)] as the only Fe-containing by-product. For the unstrained analogue, the intermolecular transfer of Fe–CH₃ to Ir–Cl speaks to the drive for Ir–CH₃ and Fe–Cl bond formation. For **5** and **7**, however, this driving force is offset by the stability of **1**, providing a reversal in the predicted outcome. This general reactivity trend additionally contrasts with that noted for the CpFe complex, [CpFe(CO)₂] (Cp = C₅H₅[−]) and Au–C bonds (another 5d-element), which results in Fe–C bond formation.²⁸

A family of heterometallic Fe/M (M = Rh or Ir) complexes have been systematically prepared *via* an elementary transmetalation reaction between a strained Cp*Fe tucked-in complex **1** and group 9 diolefin salts. This work establishes routes for the systematic generation of such a compositionally distinct class of Cp*{Fe,M} compound and furthers our understanding of the

factors that contribute to metal-to-metal transmetalation, an elementary reaction with direct implications for carbon-element bond formation using Fe.

The authors are grateful to Western University, the Council of Ontario Universities for a John C. Polanyi award to M. W. D., the Canadian Foundation for Innovation (LOF-212442), and the Natural Sciences and Engineering Research Council of Canada (Discovery Grant, RGPIN-2020-04480 (M. W. D.), Discovery Launch Supplement, DGECR-2020-00183), and graduate award (CGS-D/NSERC Vanier to J. A. Z.) for funding. M. H. thanks the Inorganic Chemistry Exchange (ICE) program for the opportunity to conduct research at Western University.

Data availability

The data supporting this article have been included as part of the ESI.† Crystallographic data for **2–5** and **8** have been deposited at the Cambridge Crystallographic Data Centre (CCDC 2380939–2380943†).

Conflicts of interest

There are no conflicts to declare.

References

- J. K. Stille, *Angew. Chem., Int. Ed. Engl.*, 1986, **25**, 508–524.
- I. P. Beletskaya and A. V. Cheprakov, *Chem. Rev.*, 2000, **100**, 3009–3066.
- N. Miyaura and A. Suzuki, *Chem. Rev.*, 1995, **95**, 2457–2483.
- X. Chen, *et al.*, *Angew. Chem., Int. Ed.*, 2009, **48**, 5094–5115.
- C. C. Johansson Seechurn, *et al.*, *Angew. Chem., Int. Ed.*, 2012, **51**, 5062–5085.
- D. G. Brown and J. Boström, *J. Med. Chem.*, 2016, **59**, 4443–4458.
- A. Biffis, *et al.*, *Chem. Rev.*, 2018, **118**, 2249–2295.
- X. Yan and C. Xi, *Coord. Chem. Rev.*, 2017, **350**, 275–284.
- S. C. Rasmussen, *ChemTexts*, 2021, **7**, 1.
- D. V. Partyka, *Chem. Rev.*, 2011, **111**, 1529–1595.
- C. He, *et al.*, *Angew. Chem.*, 2013, **125**, 1567–1570.
- A. A. Thomas, *et al.*, *J. Am. Chem. Soc.*, 2017, **139**, 3805–3821.
- A. A. Thomas and S. E. Denmark, *Science*, 2016, **352**, 329–332.
- M. J. Demchuk, *et al.*, *Chem. Commun.*, 2022, **58**, 68–71.
- M. L. Neidig, *et al.*, *Acc. Chem. Res.*, 2019, **52**, 140–150.
- H. Kurosawa and A. Yamamoto, *Fundamentals of Molecular Catalysis*, Elsevier Science, Amsterdam, Netherlands, 2003.
- J. A. Zurakowski and M. W. Drover, *Chem. Commun.*, 2023, **59**, 11349–11352.
- J. A. Zurakowski, *et al.*, *Chem. Sci.*, 2024, **15**, 10359–10365.
- T. J. Kealy and P. L. Pauson, *Nature*, 1951, **168**, 1039–1040.
- (a) P. J. Chirik, *Organometallics*, 2010, **29**, 1500–1517; (b) T. Ostwald, *et al.*, *Organometallics*, 2019, **38**, 829–843; (c) J. Pinkas, *et al.*, *Dalton Trans.*, 2022, **51**, 10198–10215.
- J. A. Zurakowski, *et al.*, *Inorg. Chem.*, 2023, **62**, 7053–7060.
- S. Liu and G. S. Girolami, *Organometallics*, 2021, **40**, 714–724.
- T. Shima and Z. Hou, *Organometallics*, 2009, **28**, 2244–2252.
- A. Meißner, *et al.*, *ChemPlusChem*, 2015, **80**, 169–180.
- M. A. Kulzick, *et al.*, *J. Organomet. Chem.*, 1987, **336**, 221–236.
- M. A. Kulzick, *et al.*, *J. Organomet. Chem.*, 1987, **333**, 105–118.
- M. R. Churchill and S. A. Bezman, *Inorg. Chem.*, 1973, **12**, 531–536.
- A. S. K. Hashmi and L. Molinari, *Organometallics*, 2011, **30**, 3457–3460.

