


 Cite this: *Chem. Commun.*, 2025, 61, 9326

Correction: pH-Triggered nanoreactors as oxidative stress amplifiers for combating multidrug-resistant biofilms

 Lei Huang,^{†ab} Shangming Jiang,^{†a} Bo Cai,^b Guobin Wang,^c Zheng Wang^{*bc} and Lin Wang^{*ab}

DOI: 10.1039/d5cc90180d

rsc.li/chemcomm

 Correction for 'pH-Triggered nanoreactors as oxidative stress amplifiers for combating multidrug-resistant biofilms' by Lei Huang et al., *Chem. Commun.*, 2021, 57, 4662–4665, <https://doi.org/10.1039/D1CC00247C>.

The authors regret that an incorrect image was accidentally included in Fig. 3d of the original article. An incorrect image for the agar plate photograph of *MRSA* treated with PA at a concentration of 50 $\mu\text{g mL}^{-1}$ (fourth column, first row) was used in error. The correct version of Fig. 3d is provided below. This correction does not affect the results or conclusions of this paper.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Department of Clinical Laboratory, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, 430022, China.

E-mail: lin_wang@hust.edu.cn

^b Research Center for Tissue Engineering and Regenerative Medicine, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, 430022, China. E-mail: zhengwang@hust.edu.cn

^c Department of Gastrointestinal Surgery, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, 430022, China

[†] Both authors contributed equally to the work.






Fig. 3 The antibacterial effect of CPA against MDR bacteria (*MRSA*). (a) The calcium levels in *MRSA* incubated with PBS, CaO₂, CP, and CPA for 2 h. The intracellular (b) ROS and (c) O₂^{•-} levels in *MRSA* treated with PBS, free H₂O₂, CaO₂, PA, CP, and CPA for 2 h. The final concentrations of CaO₂, CP, and CPA used in those experiments were equal to 50 µg mL⁻¹ of CaO₂. (d) The representative photographs of agar plates of *MRSA* treated with PBS, free H₂O₂, CaO₂, PA, CP, and CPA at different concentrations. (e) The relative viabilities of *MRSA* on agar plates after treatment with PBS, free H₂O₂, CaO₂, PA, CP, and CPA at different concentrations were evaluated by a colony-forming unit assay. Data shown as mean ± SD; ***p* < 0.01; N.S., not significant.

