

# PCCP

Physical Chemistry Chemical Physics

rsc.li/pccp



ISSN 1463-9076

**PAPER**

Richard M Cox *et al.*  
Thermodynamic insight into  $\text{AnO}_2^+$  bonding from  $\text{ThH}^+/\text{UH}^+$   
reactions studied by inductively coupled plasma tandem  
mass spectrometry



Cite this: *Phys. Chem. Chem. Phys.*,  
2026, **28**, 5656

# Thermodynamic insight into $\text{AnO}_2^+$ bonding from $\text{ThH}^+/\text{UH}^+$ reactions studied by inductively coupled plasma tandem mass spectrometry

Richard M Cox, \* Amanda R. Bubas and Amanda D. French

The bonding mechanisms of actinides have been a focus of fundamental research over the past few decades. In the present study, reactions of the simplest actinide-containing species,  $\text{ThH}^+$  and  $\text{UH}^+$ , with  $\text{O}_2$  and  $\text{CO}_2$  are investigated by inductively coupled plasma tandem mass spectrometry. The reactions of  $\text{ThH}^+$  and  $\text{UH}^+$  with  $\text{O}_2$  are efficient, and the reactions of  $\text{ThH}^+$  and  $\text{UH}^+$  with  $\text{CO}_2$  display reduced reaction efficiencies. For both reactions involving  $\text{CO}_2$ ,  $\text{ThO}_2^+$  and  $\text{UO}_2^+$  are observed; however, there is a clear barrier to  $\text{ThO}_2^+$  formation whereas  $\text{UO}_2^+$  forms through an exothermic, barrierless process. The experimental observations and available thermodynamic information are used to predict the outcomes of reactions involving the later  $\text{AnH}^+$ . The anticipated reaction enthalpies for Pa–Am display a clear correlation with the promotion energy of  $\text{An}^+$  to a  $6d^2$  electronic configuration,  $E_p(6d^2)$ , although a shift in the slope of the correlation of reaction enthalpies and  $E_p(6d^2)$  suggests that there is likely a change in bonding mechanism that starts with  $\text{Np}^+$ . Similar shifts have also been noted in previous studies. Beginning with  $\text{Np}^+$ , the  $6d$  orbitals become less accessible than they are for the earlier  $\text{An}^+$  as measured by  $E_p(6d^2)$ , and this accessibility of the  $6d$  orbitals may drive actinide bonding.

Received 14th November 2025,  
Accepted 12th January 2026

DOI: 10.1039/d5cp04417k

[rsc.li/pccp](http://rsc.li/pccp)

## Introduction

The actinides (An) are commonly associated with nuclear applications, but from a fundamental perspective, An compose the bottom edge of the periodic table and display complex chemical behavior that is still not well understood. Experimental investigations of the factors that guide An reactivity and bonding remain limited. Gas-phase studies offer a direct route to examine the intrinsic chemistry of the actinides (free of perturbative solvent effects) that can provide and explain trends in bonding and reactivity across the An series.<sup>1</sup>

The simplest actinide-containing species,  $\text{An-H}$ , has been studied using several gas-phase techniques. Surface ionization mass spectrometry was used for the isotopic analysis of U and Pu and led to the observation of  $\text{UH}^+$  and  $\text{PuH}^+$ ,<sup>2</sup> where the bond dissociation energy of  $\text{UH}^+$ ,  $D_0(\text{U}^+-\text{H})$  was estimated to be  $3.3 \pm 0.5$  eV. Early ion beam mass spectrometry enabled an examination of the energy dependences of the reactions of  $\text{U}^+$  with  $\text{D}_2$  and  $\text{CD}_4$  leading to the formation of  $\text{UD}^+$ .<sup>3</sup> The bond dissociation energy of  $\text{UH}^+$  was measured and reported as  $D_0(\text{U}^+-\text{H}) = 2.9 \pm 0.2$  eV. The infrared spectra of  $\text{ThH}$  and  $\text{UH}$  in noble gas matrices have also been reported, and complementary density functional theory (DFT) calculations included

in those studies were used to characterize and assign the observed bands as well as provide mechanistic insight into product formation and reaction energetics.<sup>4,5</sup>

Fourier transform ion cyclotron resonance mass spectrometry (FTICR-MS) experiments enabled the formation of  $\text{UH}^+$ ,  $\text{NpH}^+$ ,  $\text{PuH}^+$ ,  $\text{AmH}^+$ , and  $\text{CmH}^+$  as minor products in reactions of  $\text{An}^{2+}$  with various hydrocarbons.<sup>6</sup> A subsequent study presented theoretical potential energy surfaces (PES) that mostly explained the observations noted in the FT-ICR MS experiments.<sup>7</sup> Guided ion beam tandem mass spectrometry (GIBMS) was used to study the reactions of  $\text{Th}^+$  with  $\text{CH}_4$  and  $\text{H}_2\text{O}$ .<sup>8,9</sup>  $\text{ThH}^+$  was observed as a high energy product in both studies. Subsequent GIBMS studies of the reactions of  $\text{Th}^+$  and  $\text{U}^+$  with  $\text{H}_2$  led to the determination of the BDEs of  $\text{ThH}^+$  and  $\text{UH}^+$ ,  $D_0(\text{Th}^+-\text{H})^{10} = 2.45 \pm 0.07$  eV and  $D_0(\text{U}^+-\text{H})^{11} = 2.48 \pm 0.06$  eV. Experimental measurements of the electron affinities of  $\text{ThH}^-$  and  $\text{UH}^-$  accompanied by high level theoretical calculations have indicated that there is a 0.37 eV discrepancy between the theoretical and experimental  $D_0(\text{Th}^+-\text{H})$  value.<sup>12,13</sup> The reason for the discrepancy is unclear and remains an open investigation. Results from additional theoretical calculations for  $\text{AnH}^{0/+/-}$  ( $\text{An} = \text{Ac}, \text{Pa}, \text{Np}, \text{Pu}$ ) species have also been reported.<sup>12-15</sup> Most recently, the BDEs of  $\text{AnH}^+$  ( $\text{An} = \text{Th-Am}$ ) were experimentally measured by examining the kinetic energy dependence of the reaction  $\text{An}^+ + \text{CH}_4 \rightarrow \text{AnH}^+ + \text{CH}_3$ .<sup>16</sup> Table 1 summarizes the  $\text{AnH}^+$  thermochemistry published to date.

Pacific Northwest National Laboratory, Richland, WA 99352, USA.  
E-mail: richard.cox@pnl.gov



**Table 1** Experimental and theoretical bond dissociation energies (BDE) of AnH<sup>+</sup> (in eV)

$D_0(\text{An}^+-\text{H})$	$E_p(6d)^a$	Experimental <sup>b</sup>	Theoretical <sup>c</sup>	Model <sup>d</sup>
Th	0.00	$2.96 \pm 0.81$ , $2.45 \pm 0.07^d$	2.82	2.45
Pa	0.10	$2.94 \pm 0.54$	2.54	2.35
U	0.04	$2.82 \pm 0.71$ , $2.48 \pm 0.06^e$ , $2.9 \pm 0.3^f$	2.51	2.41
Np	0.00	$2.87 \pm 0.51$	2.08	2.45
Pu	1.08	$1.27 \pm 0.83$	0.94	1.37
Am	1.76	$0.92 \pm 0.74$		0.69

<sup>a</sup> Ref. 17. <sup>b</sup> Unless noted otherwise ref. 16. <sup>c</sup> FPD calculations of AnH<sup>+</sup> BDEs compiled in ref. 15. See also ref. 12–14. <sup>d</sup> Intrinsic BDE model assuming bonding between An<sup>+</sup> 6d and H 1s electrons. Presented in ref. 10. <sup>e</sup> Ref. 11. <sup>f</sup> Ref. 3.

Notably, experimental measurables, like BDEs, can be a key benchmark to help validate computational methods, and these computational methods will likely prove crucial to understanding An bonding.

A key finding of the previous experimental work is that the AnH<sup>+</sup> BDEs are correlated to the promotion energy from the ground state to the first state that populates a 6d orbital,  $E_p(6d)$ ,<sup>10,16</sup> which mirrors earlier findings that indicate that  $D_0(\text{An}^{n+}-\text{O})$  for  $n = 0-2$  are correlated to  $E_p(6d^2)$ .<sup>10,16,17</sup> The BDEs of AnF<sup>+</sup> also display some correlation to  $E_p(6d)$ ,<sup>18</sup> which follows an earlier argument by Cox, Armentrout, and de Jong using previously reported ThF<sup>+</sup>, UF<sup>+</sup>, and PuF<sup>+</sup> values.<sup>10</sup> Recent inductively coupled plasma tandem mass spectrometry (ICP-MS/MS) work has indicated that  $D_0(\text{An}^+-\text{CH}_2)$ <sup>16</sup> and  $D_0(\text{An}^+-\text{N})$ <sup>19</sup> are also correlated to  $E_p(6d^2)$ . The latter had previously been speculated by Armentrout and coworkers.<sup>20,21</sup> While this does not rule out the involvement of 5f or 7s orbitals, the 6d orbitals are clearly important to forming actinide-ligand (AnL) bonds.

Notably, the An species described in the previous paragraph have formal oxidation states (II–IV). Theoretical studies have concluded that the 5f orbitals are more active for An in higher oxidation states.<sup>22–25</sup> Classic examples include the linear OAnO<sup>+</sup> (V) species. Although the electrons of Th<sup>+</sup> do not populate the 5f orbitals in the ground state, NBO analyses have shown that 5f orbitals contribute to bonding in OThO<sup>+</sup>.<sup>22</sup> Recent work has suggested that reaction kinetics may be influenced by the participation of 5f orbitals in transient species.<sup>26</sup> In that study, the comparison of the reaction kinetics between Pu<sup>+</sup>/Nd<sup>+</sup>/Sm<sup>+</sup> + NO/CO<sub>2</sub> were used to infer 5f orbital participation in the NPuO<sup>+</sup> intermediate with corroborating theoretical potential energy surfaces (PES). Likewise, observing the reactions starting from diatomic An species (*i.e.* AnL<sup>+</sup>) may shed additional light on when the 5f orbitals are active in AnL bonding.

There are several reports of the reactions of AnO<sup>+</sup> with a variety of co-reactants.<sup>22,25,27–31</sup> Generally, AnO<sup>+</sup> species are less reactive than their An<sup>+</sup> counterparts. This may be explained by the reduced availability of electrons because two An<sup>+</sup> valence electrons are involved in the robust AnO<sup>+</sup> triple bond. Alternatively, AnO<sup>+</sup> is likely to have increased steric restrictions

compared to An<sup>+</sup> and AnH<sup>+</sup>. AnH<sup>+</sup> bonding employs one of the 6d electrons of An<sup>+</sup>. It is of interest to study the reactions of AnH<sup>+</sup> to better understand the role of An<sup>+</sup> valence electrons and assess the impact of reduced 6d electron availability upon the observed chemistry. For the simplest polyatomic, AnH<sup>+</sup>, the reactions of UH<sup>+</sup> with background gases<sup>32</sup> and nitriles<sup>33</sup> have been reported in MS<sup>n</sup> experiments by Terhorst *et al.* However, hydrides, like UH<sup>+</sup>, are a known interference for analytical ICP-MS experiments.<sup>34</sup> Here we take advantage of the inductively coupled plasma's (ICP) propensity to create hydrides in the ion source to study the reactions of ThH<sup>+</sup> and UH<sup>+</sup> with O<sub>2</sub> and CO<sub>2</sub>. We also compare the observed reactions to the analogous AnO<sub>x</sub><sup>n+</sup> ( $x = 0-1$ ,  $n = 1-2$ ) reactions to help determine the role of oxidation states in the observed chemistry. Likewise, we discuss the implications of observed thermochemical trends on An bonding.

## Experimental and theoretical methods

**CAUTION:** the An used in this study are all radioisotopes with varying activities and half-lives. All work was done within the radiological protection controls of specialized laboratories at Pacific Northwest National Laboratory.

### ICP-MS/MS

Experiments were conducted using an Agilent 8900 ICP-MS/MS located within the physical sciences facility at Pacific Northwest National Laboratory.<sup>35</sup> This instrument utilizes an ICP ion source equipped with a quartz double-pass spray chamber and 100 μL min<sup>-1</sup> perfluoroalkoxy alkane (PFA) nebulizer. Hydrides are formed within the plasma source, and the resulting ions are focused through an initial quadrupole mass filter where the reactant AnH<sup>+</sup> are selected. The reactant ions are then focused into a collision reaction cell containing an octopole ion guide which can be pressurized with a neutral reactant gas, O<sub>2</sub> (Oxarc, 99.999%) or CO<sub>2</sub> (Oxarc, 99.999%). Residual reactant and product ions are radially confined within the octopole and drift to the exit aperture. Ions are subsequently focused through a second quadrupole mass filter to identify product ions and counted at a standard electron multiplier detector.

Stock multi-element standard solutions containing 10 ng g<sup>-1</sup> of Th and U in 2% HNO<sub>3</sub> were prepared. Oxygen and carbon dioxide were used as reactant gases. The flow rates of O<sub>2</sub> and CO<sub>2</sub> were 0.09 and 0.18 mL min<sup>-1</sup>, and 0.06 and 0.12 mL min<sup>-1</sup>, respectively. Tuning parameters were optimized to provide maximum sensitivity using a 10 ng g<sup>-1</sup> <sup>232</sup>Th and <sup>238</sup>U solution, with focus on increasing hydride formation. The octopole bias was adjusted in 1 V intervals from +7 V to -78 V while keeping other cell parameters constant: octopole RF peak-to-peak voltage of 180 V, axial acceleration of 0 V, and a kinetic energy discrimination (KED, the voltage difference between the octopole bias in the collision reaction cell and the second quadrupole) of -7.0 V. Data were acquired using 3 replicates, 10 sweeps/replicate and 1 s acquisition times.



Absolute reaction cross sections ( $\sigma$ ) are calculated from the raw signal intensities using eqn (1)<sup>36</sup>

$$I = I_0 e^{-\rho \sigma l} \quad (1)$$

where  $I$  is the reactant ion intensity exiting the collision cell,  $I_0$  is the reactant ion intensity entering the collision cell,  $\rho$  is the number density of the neutral reactant in the collision cell, and  $l$  is the effective length of the collision cell. For this work  $I_0$  is estimated as the sum of all observed ions from quadrupole 2. The physical length of the collision cell in the Agilent 8900, 10 cm, is used for  $l$ , although this is not strictly accurate because the pressure gradient will extend beyond the boundaries of the collision cell. The difference is expected to be  $\leq 20\%$  and is included in the total uncertainties reported herein. Individual product ion cross sections (e.g.,  $\text{MO}^+$ ) are calculated as a percentage of the overall reaction cross section. Because the Agilent 8900 operates under multi-collision conditions, to compare the observed cross sections to the reaction collision limit, the cross sections observed at 1.5 and 3.0 mTorr for  $\text{O}_2$  and 1.4 and 2.8 mTorr for  $\text{CO}_2$  were extrapolated to zero pressure to reflect rigorous single collision conditions. Previously, we have discussed the influences on the expected number of collisions for a given neutral reactant gas and pressure.<sup>31</sup> Following this analysis, the most probable number of collisions is 2–3 at the lowest energies. Although less than ideal, the two-point extrapolation was chosen to balance instrument time and operator exposure while still executing the experiment. Absolute uncertainties in the cross sections are estimated to be  $\pm 50\%$  with relative uncertainties of  $\pm 20\%$ .

The energy in the laboratory frame is estimated from the octopole bias by eqn (2)<sup>37</sup>

$$E_{\text{Lab}} = V_p + (m/m_{\text{Ar}})(5/2)k_B T_p - V_{\text{Oct}} \quad (2)$$

where  $V_p$  is the plasma potential ( $\sim 2$  V),  $m$  is the mass of the  $\text{M}^+$  reactant ion,  $m_{\text{Ar}}$  is the mass of argon (the flow gas in the ICP),  $k_B$  is Boltzmann's constant,  $T_p$  is the ion electronic temperature entering the octopole, and  $V_{\text{Oct}}$  is the octopole bias. The temperature of the ion exiting the plasma is expected to be the plasma temperature, 8000–10 000 K. Previous work<sup>16,38,39</sup> has indicated that  $T_p$  is between 1000–10 000 K for atomic cations. For polyatomic cations, it is less clear what the electronic distribution may be. Presumably, some excess electronic energy from the atomic cations in the source is consumed to create  $\text{AnH}^+$ , and the electronic state density is likely less dense than the atomic electronic state density.<sup>10–13</sup> Consequently, the average electronic energy available is expected to be less than that available for atomic cations. An assumption is made that there are sufficient crossings between potential energy surfaces that a pathway exists between excited state reactants and ground state products so that excess electronic energy can be treated as energy available for reaction. Theoretical calculations were previously used to examine the low-lying electronic states of  $\text{ThH}^+$  and  $\text{UH}^+$ .<sup>11,13</sup> Based on this work, the electronic energy,  $E_{\text{el}}$ , is  $E_{\text{el}}(\text{ThH}^+) = 0.37$  eV and  $E_{\text{el}}(\text{UH}^+) = 0.12$  eV at 5000 K.

The energy in the center-of-mass ( $E_{\text{CM}}$ ) frame represents the kinetic energy available for a chemical reaction. The relationship between  $E_{\text{LAB}}$  and  $E_{\text{CM}}$  is described by eqn (3)<sup>36</sup>



$$E_{\text{CM}} = E_{\text{LAB}} \times M/(M + m) \quad (3)$$

where  $M$  is the mass of the neutral reactant partner,  $\text{O}_2$  (32.00 amu) or  $\text{CO}_2$  (44.01) and  $m$  is the mass of the reactant metal hydride ion,  $\text{ThH}^+$  (233.04 amu) or  $\text{UH}^+$  (239.04 amu).

## Experimental results

### $\text{AnH}^+ + \text{O}_2$

Fig. 1 displays the absolute reaction cross section of  $\text{ThH}^+$  with  $\text{O}_2$  as a function of kinetic energy. Fig. 2 displays the  $\text{UH}^+ + \text{O}_2$  cross section. Reactions (4)–(7) were observed:



Reactions (4)–(6) are observed as barrierless exothermic processes. The reaction efficiencies ( $k/k_{\text{col}}$ ) compared to the Su-Chesnavich semiclassical trajectory collision limit cross section<sup>40</sup> are  $k/k_{\text{col}}(\text{ThH}^+) = 1.52 \pm 0.76$  and  $k/k_{\text{col}}(\text{UH}^+) = 1.26 \pm 0.63$  calculated as an average from all energies  $< 1$  eV.




Fig. 1 The absolute reaction cross section of  $\text{ThH}^+ + \text{O}_2$  as a function of kinetic energy in the laboratory frame (upper x axis) and center-of-mass frame (lower x axis). The sum of the product cross sections ( $\sigma_{\text{tot}}$ ) is represented by a solid gray line. Individual products are  $\text{ThO}^+$  (red circles),  $\text{HThO}^+$  (blue squares),  $\text{Th}^+$  (green diamonds), and  $\text{ThO}_2^+$  (purple triangles). The Su-Chesnavich semi-classical trajectory ( $\sigma_{\text{traj}}$ ) collision limit<sup>40</sup> is represented by a solid black line.






Fig. 2 The absolute reaction cross section of  $\text{UH}^+ + \text{O}_2$  as a function of kinetic energy in the laboratory frame (upper x axis) and center-of-mass frame (lower x axis). The sum of the product cross sections ( $\sigma_{\text{tot}}$ ) is represented by a solid gray line. Individual products are  $\text{UO}^+$  (red circles),  $\text{HUO}^+$  (blue squares),  $\text{U}^+$  (green diamonds), and  $\text{UO}_2^+$  (purple triangles). The Su-Chesnavich semi-classical trajectory ( $\sigma_{\text{traj}}$ ) collision limit<sup>40</sup> is represented by a solid black line.

To our knowledge,  $k/k_{\text{col}}$  has not been reported for the reactions of  $\text{ThH}^+$  and  $\text{UH}^+$  with  $\text{O}_2$ . Guided ion beam tandem mass spectrometry (GIBMS) experiments have reported reaction efficiencies of  $k/k_{\text{col}} = 1.21 \pm 0.24$  for  $\text{Th}^+ + \text{O}_2$ <sup>41</sup> and  $k/k_{\text{col}} = 1.1 \pm 0.2$  for  $\text{U}^+ + \text{O}_2$ .<sup>42</sup> FTICR-MS experiments by Santos *et al.*<sup>27</sup> reported  $k/k_{\text{col}} = 0.86 \pm 0.43$  and  $k/k_{\text{col}} = 0.68 \pm 0.34$ , for the reactions of  $\text{Th}^+$  and  $\text{U}^+$  with  $\text{O}_2$ , respectively. Earlier FTICR-MS studies of the same reactions from Cornehl *et al.*<sup>43</sup> reported  $k/k_{\text{col}} = 1.12 \pm 0.45$  and  $k/k_{\text{col}} = 1.17 \pm 0.47$  for  $\text{Th}^+$  and  $\text{U}^+$ , respectively. The reaction (4) ( $\text{ThH}^+$ ) rate does not appear to be significantly hindered by the H-ligand compared to the atomic rate. A comparison of  $\text{AnH}^+$   $k/k_{\text{col}}$  to that for  $\text{An}^+$  and  $\text{AnO}^+$  is summarized in Table 2.

The product distribution is also different between  $\text{ThH}^+$  and  $\text{UH}^+$ . For  $\text{ThH}^+$  at the lowest energy, the dominant product is  $\text{ThO}^+$  (reaction (4)).  $\text{ThO}_2^+$  (reaction (6)) is the second most abundant product, while  $[\text{Th}, \text{O}, \text{H}]^+$  (reaction (5)) is a minor product. Note that previous theoretical work<sup>9,45,46</sup> indicates that  $[\text{Th}, \text{O}, \text{H}]^+$  is likely  $\text{HThO}^+$ . This structure will be adopted for future discussion of  $[\text{An}, \text{O}, \text{H}]^+$ . At slightly higher energies ( $\sim 0.5$ – $1.0$ ),  $\text{HThO}^+$  becomes the second most abundant product. For  $\text{UH}^+$  at the lowest energy, the dominant product is  $\text{UO}_2^+$  (reaction (6)).  $\text{UO}^+$  (reaction (4)) is the second most abundant product, while  $\text{HUO}^+$  (reaction (5)) is a minor product. Product branching remains relatively consistent for energies below 1 eV. Differences in the product distributions likely result from thermodynamic drivers. For example,  $D_0(\text{OU}^+-\text{O}) = 7.56 \pm 0.12 \text{ eV}^{25} \gg D_0(\text{OTh}^+-\text{O}) = 4.87 \pm 0.04 \text{ eV}^{22}$  so reaction (6) is significantly favored for  $\text{UH}^+$  compared to  $\text{ThH}^+$ . Reaction thermodynamics are discussed in greater detail below.

Table 2 A comparison of  $\text{An}^{n+}$  ( $\text{An} = \text{Th}, \text{U}; n = 1, 2$ ) and  $\text{ThL}^+$  and  $\text{UL}^+$  ( $\text{L} = \text{H}, \text{O}$ ) reaction efficiencies ( $k/k_{\text{col}}$ )<sup>a</sup>

$\text{AnL}^+ + \text{OR}$	ICP-MS/MS <sup>b</sup>	GIBMS <sup>c</sup>	FTICR-MS <sup>d</sup>
$\text{Th}^+ + \text{O}_2$		$1.21 \pm 0.24$	$0.86 \pm 0.43; 1.12 \pm 0.45$
$\text{Th}^{2+} + \text{O}_2$			$0.56 \pm 0.28$
$\text{ThH}^+ + \text{O}_2$	$1.52 \pm 0.76$		
$\text{ThO}^+ + \text{O}_2$		NR <sup>e</sup>	NR <sup>e</sup>
$\text{U}^+ + \text{O}_2$		$1.1 \pm 0.2$	$0.68 \pm 0.34; 1.17 \pm 0.47$
$\text{U}^{2+} + \text{O}_2$			$0.53 \pm 0.26$
$\text{UH}^+ + \text{O}_2$	$1.26 \pm 0.63$		
$\text{UO}^+ + \text{O}_2$		$1.16 \pm 0.23$	$0.45 \pm 0.23; 0.95 \pm 0.38$
$\text{Th}^+ + \text{CO}_2$	$1.19 \pm 0.60$	$0.88 \pm 0.18$	$0.35 \pm 0.18; 0.95 \pm 0.33$
$\text{Th}^{2+} + \text{CO}_2$			$0.55 \pm 0.28$
$\text{ThH}^+ + \text{CO}_2$	$0.49 \pm 0.25$		
$\text{ThO}^+ + \text{CO}_2$		NR <sup>e</sup>	NR <sup>e</sup>
$\text{U}^+ + \text{CO}_2$	$0.93 \pm 0.47$	$1.18 \pm 0.24$	$0.29 \pm 0.15; 1.02 \pm 0.36$
$\text{U}^{2+} + \text{CO}_2$	$0.72 \pm 0.36$		$0.38 \pm 0.17$
$\text{UH}^+ + \text{CO}_2$	$0.36 \pm 0.18$		
$\text{UO}^+ + \text{CO}_2$		$0.02 \pm 0.01^f$	$0.004 \pm 0.002; 0.002 \pm 0.001$

<sup>a</sup> Reaction efficiencies are calculated relative to the Su-Chesnavich semiclassical trajectory collision limit. See ref. 40. <sup>b</sup> This work for  $\text{AnH}^+$ . Ref. 39 for  $\text{An}^+$ .  $\text{U}^{2+}$  can be found in the SI Section. <sup>c</sup> Ref. 22, 25, 31, 41 and 42. <sup>d</sup> Ref. 27, 43 and 44. <sup>e</sup> NR = no observed reaction at lowest energy studied. <sup>f</sup> Calculated from ref. 25.

Product branching ratios are summarized in Table 3. Fig. S1–S4 also break down the branching ratios as a function of kinetic energy.

At higher energies the product  $\text{An}^+$  is observed with apparent thresholds of 1.5 eV for  $\text{Th}^+$  and 0.8 eV for  $\text{U}^+$ . This likely can be attributed to reaction (7a), which is a collision induced dissociation (CID) reaction. GIBMS work indicates that the threshold for this process should be  $\approx 2.5$  eV for both  $\text{ThH}^+$  and  $\text{UH}^+$  (see Table 1).<sup>10,11,16</sup> Alternatively, the hydroperoxyl radical could be formed through reaction (7b). Given  $D_0(\text{O}_2-\text{H}) = 2.24 \pm 0.01 \text{ eV}$ , the reaction (7b) threshold should be  $\approx 0.25$  eV for both  $\text{ThH}^+$  and  $\text{UH}^+$ . Exact assignment of the reaction is difficult because the ICP-MS/MS data is extrapolated from multicollisional conditions to “zero pressure” single collision conditions. Previous work has indicated that while this extrapolation is reasonable, the cross section may retain some multicollisional character.<sup>16,39</sup> Multi-collision conditions may lead to observable artifacts such as an

Table 3 Product branching ratios at select energies  $\leq 1$  eV. Relative uncertainty is  $\pm 10\%$



Reaction	Product	0.1 eV	0.5 eV	1 eV
$\text{ThH}^+ + \text{O}_2 \rightarrow$	$\text{ThO}^+$	0.70	0.81	0.79
	$\text{HThO}^+$	0.09	0.13	0.10
	$\text{ThO}_2^+$	0.21	0.06	0.11
$\text{UH}^+ + \text{O}_2 \rightarrow$	$\text{UO}^+$	0.31	0.26	0.32
	$\text{HUO}^+$	0.04	0.04	0.06
	$\text{UO}_2^+$	0.65	0.70	0.62
$\text{ThH}^+ + \text{CO}_2 \rightarrow$	$\text{ThO}^+$	0.45	0.47	0.47
	$\text{HThO}^+$	0.55	0.53	0.53
	$\text{ThO}_2^+$	0.00	0.00	0.00
$\text{UH}^+ + \text{CO}_2 \rightarrow$	$\text{UO}^+$	0.28	0.31	0.40
	$\text{HUO}^+$	0.47	0.51	0.60
	$\text{UO}_2^+$	0.25	0.18	0.00



apparent threshold that is shifted to lower energies relative to the thermodynamic threshold. Furthermore,  $\text{AnH}^+$  can be partially activated upon collision storing some energy within rovibrational modes so that subsequent collisions may remove the ligand at lower energies. By contrast, competition with the more favorable reactions (4)–(6) may also delay the onset of reaction (7b). Consequently,  $\text{An}^+$  may be formed through either mechanism; nevertheless, the observed thresholds in Fig. 1 and 2 are consistent with  $D_0(\text{Th}^+-\text{H}) = 2.45 \pm 0.07$  eV and  $D_0(\text{U}^+-\text{H}) = 2.48 \pm 0.06$  eV reported in GIBMS studies.<sup>10,11</sup>

### $\text{AnH}^+ + \text{CO}_2$

Fig. 3 and 4 display the absolute reaction cross section of  $\text{ThH}^+$  and  $\text{UH}^+$  with  $\text{CO}_2$  as a function of kinetic energy. Reactions (8)–(11) were observed:



The observed reaction efficiencies are  $k/k_{\text{col}}(\text{ThH}^+) = 0.49 \pm 0.25$  and  $k/k_{\text{col}}(\text{UH}^+) = 0.36 \pm 0.18$ . Like  $\text{AnH}^+ + \text{O}_2$ , the reaction of  $\text{AnH}^+$  with  $\text{CO}_2$  has not been previously reported. The reactions of  $\text{An}^+ + \text{CO}_2$  have been previously reported for ICP-MS/MS, GIBMS, and FTICR-MS.<sup>25,27,30,31,39</sup> Reaction efficiencies are




Fig. 4 The absolute reaction cross section of  $\text{UH}^+ + \text{CO}_2$  as a function of kinetic energy in the laboratory frame (upper x axis) and center-of-mass frame (lower x axis). The sum of the product cross sections ( $\sigma_{\text{tot}}$ ) is represented by a solid gray line. Individual products are  $\text{UO}^+$  (red circles),  $\text{HUO}^+$  (blue squares),  $\text{U}^+$  (green diamonds), and  $\text{UO}_2^+$  (purple triangles). The Su–Chesnavich semi-classical trajectory ( $\sigma_{\text{traj}}$ ) collision limit<sup>40</sup> is represented by a solid black line.

listed in Table 2. For the atomic cations, ICP-MS/MS,<sup>39</sup> GIBMS,<sup>31</sup> and FTICR-MS<sup>43</sup> reported values agree very well with the exception of the FTICR-MS efficiencies reported by Santos *et al.*<sup>27</sup> The  $\text{AnH}^+$  reaction efficiencies are significantly reduced from the atomic reaction efficiencies. Table 2 also lists  $k/k_{\text{col}}$  for  $\text{AnO}^+$  reported from GIBMS and FTICR-MS experiments.<sup>25,27,31,43,44</sup>

While  $k/k_{\text{col}}(\text{AnH}^+)$  is significantly reduced from  $k/k_{\text{col}}(\text{An}^+)$ ,  $k/k_{\text{col}}(\text{AnH}^+) \gg k/k_{\text{col}}(\text{AnO}^+)$ , see Table 2. For  $\text{ThO}^+ + \text{CO}_2$ ,  $D_0(\text{OTh}^+-\text{O}) = 4.87 \pm 0.04$  eV<sup>22</sup> <  $D_0(\text{OC}-\text{O}) = 5.45$  eV,<sup>47</sup> so no reaction is expected. However,  $D_0(\text{OU}^+-\text{O}) = 7.56 \pm 0.12$  eV<sup>25</sup> >>  $D_0(\text{OC}-\text{O}) = 5.45$  eV, so thermodynamics do not explain the low reaction efficiency. Note also that by GIBMS,<sup>31</sup> which measures the reaction energy dependence, the observed reaction threshold ( $E_0$ ) for  $\text{ThO}^+ + \text{CO}_2$  did not match the thermodynamic threshold, which indicates the presence of some barrier to reaction.

Unlike  $\text{AnH}^+ + \text{O}_2$ , the primary reaction observed with  $\text{CO}_2$  was reaction (9). For  $\text{ThH}^+$ ,  $\text{HThO}^+$  was slightly favored by  $\text{HThO}^+/\text{ThO}^+ = 55/45$  for all energies  $\leq 1$  eV. For  $\text{UH}^+$ , the branching is  $\text{HUO}^+/\text{UO}^+ = 60/40$  (note  $\text{UO}_2^+$  has been removed for two energies, see Table 3) for all energies  $\leq 1$  eV. Reaction (10) was observed for  $\text{UH}^+$  at the lowest energies, but this product was no longer observed by 1 eV with a rapid decline occurring at  $\approx 0.4$  eV. A slight increase in the reaction (8) cross section occurs in this region, which may suggest a connection between the two channels, possibly loss of an O ligand from  $\text{AnO}_2^+$ , but the energy resolution in Fig. 4 is not sufficient to definitively establish this argument. Reaction (10) is not observed until higher energies for  $\text{ThH}^+$  with an apparent threshold of 2 eV. Reaction (11) is presumably a CID reaction, although the formation of  $\text{HCO}_2$  is conceivably possible.




Fig. 3 The absolute reaction cross section of  $\text{ThH}^+ + \text{CO}_2$  as a function of kinetic energy in the laboratory frame (upper x axis) and center-of-mass frame (lower x axis). The sum of the product cross sections ( $\sigma_{\text{tot}}$ ) is represented by a solid gray line. Individual products are  $\text{ThO}^+$  (red circles),  $\text{HThO}^+$  (blue squares),  $\text{Th}^+$  (green diamonds), and  $\text{ThO}_2^+$  (purple triangles). The Su–Chesnavich semi-classical trajectory ( $\sigma_{\text{traj}}$ ) collision limit<sup>40</sup> is represented by a solid black line.



Apparent thresholds of 1.5 and 1 eV were observed for  $\text{ThH}^+$  and  $\text{UH}^+$ , respectively. Despite the possible multicollisional character retained in the reaction (11) cross sections, these observed thresholds are likely consistent with the established BDEs of  $\text{ThH}^+$  and  $\text{AnH}^+$ .

## Discussion

### Comparison of reaction rates

The reaction efficiencies of  $\text{An}^{n+}$  ( $\text{An} = \text{Th}, \text{U}; n = 1, 2$ ) and  $\text{AnL}^+$  ( $\text{L} = \text{H}, \text{O}$ ) with  $\text{O}_2$  and  $\text{CO}_2$  from ICP-MS/MS,<sup>39</sup> GIBMS,<sup>22,25,31,41,42</sup> and FTICR-MS<sup>27,43,44</sup> reactions are listed in Table 2. For these reactions, the precursor ion includes  $\text{An}$  in nominal oxidation states of I–III. ICP-MS/MS data available for the reaction  $\text{U}^{2+} + \text{CO}_2$  is also included in Table 2. The cross section for this reaction can be found in Fig. S5 in the SI Section.

In general, the observed ICP-MS/MS  $k/k_{\text{col}}$  for atomic cations matches reasonably well with the GIBMS measurements within experimental uncertainty. Likewise, the observed efficiencies from ICP-MS/MS are consistent with the Cornehl *et al.* FTICR-MS<sup>43</sup> results. The FTICR-MS results from Gibson and coworkers are an average factor of  $3 \pm 1$  times lower than the GIBMS and ICP-MS/MS values, Table 3. The difference between GIBMS and FTICR-MS  $k/k_{\text{col}}$  have been noted previously.<sup>31</sup> Discrepancies between the reaction efficiencies reported by ICP-MS/MS, GIBMS, and FTICR-MS can likely be attributed to differences in the starting electronic energy distributions and collision conditions achieved by each of the methods.

The FTICR-MS work from Gibson and coworkers<sup>27,44</sup> are the most complete for  $\text{An}$  (I–III) starting oxidation states. For the reactions  $\text{ThO}^+ + \text{O}_2/\text{CO}_2$ , no reaction was observed. This result is not surprising because both reactions are endothermic as confirmed by GIBMS experiments.<sup>22,31</sup> All other reactions were observed by FTICR-MS. For  $\text{An}^{n+} + \text{O}_2$  ( $n = 1, 2$ ) the reaction efficiency decreases slightly with increasing oxidation state (although with overlapping uncertainty). For  $\text{UO}^+ + \text{O}_2$ , the reaction efficiency also decreases slightly but is still within uncertainty of both  $\text{U}^{n+}$  ( $n = 1, 2$ ) reactions. By contrast, the efficiency of  $\text{UO}_x^+$  ( $x = 0, 1$ ) +  $\text{O}_2$  measured by GIBMS are identical within experimental uncertainty.<sup>42</sup> The FTICR-MS reaction trend for  $\text{An}^{n+}$  ( $n = 1, 2$ ) +  $\text{CO}_2$  appears to be reversed, yet the efficiency for  $\text{UO}^+ + \text{CO}_2$  is significantly reduced compared to  $\text{U}^{n+} + \text{CO}_2$ . This result is echoed in the GIBMS report.<sup>25</sup>

At present, a complete reaction set is not available for ICP-MS/MS. Assuming that the ICP-MS/MS reaction efficiencies are similar to those reported by GIBMS, then the reaction efficiency may increase slightly for  $\text{AnH}^+ + \text{O}_2$  compared to  $\text{An}^+ + \text{O}_2$  but would be within the combined experimental uncertainties. A similar result may be expected for  $\text{An}^{2+}$  given the historical differences between FTICR-MS, GIBMS, and ICP-MS/MS. Reaction enthalpies calculated from  $\text{AnO}_x^{n+}$  ( $n = 0-2; x = 1, 2$ ) values compiled by Marçalo and Gibson<sup>17</sup> indicate that all the  $\text{O}_2$  reactions in Table 2 will have similar thermodynamic drivers (except  $\text{ThO}^+ + \text{O}_2$ ). Likewise, Table 4 indicates that the thermodynamics associated with  $\text{AnH}^+$  reactions with  $\text{O}_2$  will be similar. For the  $\text{CO}_2$  reactions in Table 2, there is a substantial difference between  $\text{An}^+ + \text{CO}_2$  and  $\text{AnH}^+ + \text{CO}_2$  reaction efficiencies. Table 4 indicates  $\approx 0.5$  eV difference in reaction enthalpies.

The thermodynamics of the reactions observed in this work are discussed in greater detail in the next section. Table S1 also compiles the thermodynamic information for the reactions listed in Table 2. For  $\text{An} \geq \text{Np}^+$ , there is a significant difference in thermodynamics between oxidation states. Marçalo and Gibson tie this difference to  $E_p(6d^2)$ .<sup>17</sup> Because  $E_p(6d^2)$  are a rough measure of the energy of the 6d orbitals relative to the 5f orbitals populated in most  $\text{An}^{n+}$  ground states, the difference between reactivity observed in Table 2 may be a function of the accessibility of the 6d orbitals. Conversely, the difference between  $\text{An}$  reactivity with  $\text{O}_2$  and  $\text{CO}_2$  follow similar trends regardless of the  $\text{An}$  oxidation state. The reduced reactivity of  $\text{CO}_2$  has been discussed in greater detail previously but has been tied to the weak interaction of a late intermediate along the potential energy surface.<sup>26,31,39,48</sup>

### $\text{An}^+$ thermochemistry

Most observed reactions are clearly exothermic reactions that seemingly will only provide upper or lower limits to BDEs. Nevertheless, the reactions here may refine existing or provide new thermodynamic values. The expected reaction enthalpies are discussed below.

The reaction enthalpy ( $\Delta_r H_0^\circ$ ) for the ligand exchange reactions, reactions (4) and (8), are:

$$\Delta_r H_0^\circ(4) = D_0(\text{An}^+ - \text{H}) + D_0(\text{O} - \text{O}) - D_0(\text{An}^+ - \text{O}) - D_0(\text{O} - \text{H}) \quad (12)$$

Table 4 Bond dissociation energies of X–L and LX–O ( $X = \text{C}, \text{O}, \text{An}^+, L = \text{H}, \text{O}$ ) and reaction enthalpies in eV

X	$D_0(\text{X}-\text{H})^a$	$D_0(\text{X}-\text{O})^b$	$D_0(\text{OX}-\text{H})^c$	$D_0(\text{OX}-\text{O})^b$	$\Delta_r H_0^\circ(4)^d$	$\Delta_r H_0^\circ(8)^e$	$\Delta_r H_0^\circ(6)^f$	$\Delta_r H_0^\circ(10)^g$
C	3.53 <sup>c</sup>	11.09 <sup>c</sup>	1.81	5.45 <sup>c</sup>				
O	4.44 <sup>c</sup>	5.11 <sup>c</sup>	2.24					
Th <sup>+</sup>	2.45 ± 0.07 <sup>h</sup>	8.57 ± 0.14 <sup>i</sup>		4.87 ± 0.04 <sup>j</sup>	−5.45 ± 0.16	−2.48 ± 0.16	−5.88 ± 0.16	2.02 ± 0.16
Pa <sup>+</sup>	2.94 ± 0.54	8.29 ± 0.52		8.08 ± 0.30	−4.68 ± 0.75	−1.71 ± 0.75	−8.33 ± 0.81	−0.43 ± 0.81
U <sup>+</sup>	2.48 ± 0.06 <sup>k</sup>	8.01 ± 0.13 <sup>l</sup>		7.56 ± 0.12 <sup>m</sup>	−4.87 ± 0.15	−1.90 ± 0.15	−7.99 ± 0.19	−0.09 ± 0.19
Np <sup>+</sup>	2.87 ± 0.51	7.88 ± 0.10		6.32 ± 0.23	−4.34 ± 0.52	−1.37 ± 0.52	−6.22 ± 0.57	1.68 ± 0.57
Pu <sup>+</sup>	1.27 ± 0.83	6.75 ± 0.20		5.28 ± 0.39	−4.81 ± 0.85	−1.84 ± 0.85	−5.64 ± 0.94	2.26 ± 0.94
Am <sup>+</sup>	0.92 ± 0.74	5.80 ± 0.29		4.25 ± 0.58	−4.21 ± 0.79	−1.24 ± 0.79	−4.02 ± 0.98	3.88 ± 0.98

<sup>a</sup> Ref. 16 unless indicated otherwise. <sup>b</sup> Ref. 17 unless indicated otherwise. <sup>c</sup> NIST Chemistry Webbook unless indicated otherwise. <sup>d</sup> See eqn (12). <sup>e</sup> See eqn (13). <sup>f</sup> See eqn (16). <sup>g</sup> See eqn (17). <sup>h</sup> Ref. 10. <sup>i</sup> Ref. 41. <sup>j</sup> Ref. 22. <sup>k</sup> Ref. 11. <sup>l</sup> Ref. 42. <sup>m</sup> Ref. 25.



$$\Delta_r H_0^\circ(8) = D_0(\text{An}^+-\text{H}) + D_0(\text{OC}-\text{O}) - D_0(\text{An}^+-\text{O}) - D_0(\text{OC}-\text{H}) \quad (13)$$

Existing thermochemical values are listed in Table 4. Both reaction (4) and (8) are quite exothermic.  $\Delta_r H_0^\circ(4) = -5.45$  and  $-4.87$  eV for  $\text{ThH}^+$  and  $\text{UH}^+$ , respectively, while  $\Delta_r H_0^\circ(8) = -2.48$  and  $-1.90$  eV. For the other An, the more discerning reaction (8) should be exothermic if  $D_0(\text{An}^+-\text{O}) \geq D_0(\text{An}^+-\text{H}) + 3.64$  eV. From  $D_0(\text{An}^+-\text{H})$  listed in Table 4, reaction (8) should be exothermic for  $\text{An}^+ = \text{Th}^+ - \text{Am}^+$ .

For reactions (5) and (9), the reaction enthalpies are:

$$\Delta_r H_0^\circ(5) = D_0(\text{O}-\text{O}) - D_0(\text{HAn}^+-\text{O}) \quad (14)$$

$$\Delta_r H_0^\circ(9) = D_0(\text{OC}-\text{O}) - D_0(\text{HAn}^+-\text{O}) \quad (15)$$

Consequently, reactions (5) and (9) are exothermic if  $D_0(\text{HAn}^+-\text{O}) \geq D_0(\text{O}-\text{O})/D_0(\text{OC}-\text{O})$ . Fig. 3 and 4 indicate that  $D_0(\text{HAn}^+-\text{O}) \geq D_0(\text{OC}-\text{O}) = 5.45$  eV for both  $\text{Th}^+$  and  $\text{U}^+$ . The branching ratios may provide additional information to refine  $D_0(\text{HAn}^+-\text{O})$ . Assuming that the primary driver to the branching ratios is the reaction enthalpy, then  ${}^\circ\Delta_r H_0^\circ(9) \leq {}^\circ\Delta_r H_0^\circ(8) = -2.48$  and  $-1.90$  eV for  $\text{ThH}^+$  and  $\text{UH}^+$ . This is true for  $D_0(\text{HTh}^+-\text{O}) \geq 7.93$  and  $D_0(\text{HU}^+-\text{O}) \geq 7.38$  eV. Because reaction dynamics are controlled by more than the thermodynamic driver, the BDEs may not be this large. Nevertheless, comparison of theoretical bond distances<sup>9</sup>  $r(\text{Th}^+-\text{O}) = 1.79$  Å and  $r(\text{HTh}^+-\text{O}) = 1.80$  Å indicates that both the  $\text{ThO}^+$  and  $\text{HThO}^+$  BDEs are likely of similar strength, consistent with  $D_0(\text{HTh}^+-\text{O}) \geq 7.93$  eV.

For reaction (6) and (10), the reaction enthalpies are:

$$\Delta_r H_0^\circ(6) = D_0(\text{An}^+-\text{H}) + D_0(\text{O}-\text{O}) - D_0(\text{OAn}^+-\text{O}) - D_0(\text{An}^+-\text{O}) \quad (16)$$

$$\Delta_r H_0^\circ(10) = D_0(\text{An}^+-\text{H}) + D_0(\text{OC}-\text{O}) + D_0(\text{C}-\text{O}) - D_0(\text{OAn}^+-\text{O}) - D_0(\text{An}^+-\text{O}) - D_0(\text{C}-\text{H}) \quad (17)$$

$\Delta_r H_0^\circ(10) = 2.02 \pm 0.16$  and  $-0.09 \pm 0.19$  eV for  $\text{ThH}^+$  and  $\text{UH}^+$ , respectively. The apparent threshold observed for reaction (10) in Fig. 3 is consistent with  $\Delta_r H_0^\circ(10) = 2.02 \pm 0.16$ . Likewise, the observation of a barrierless exothermic reaction for reaction (10) in Fig. 4 is also consistent with  $\Delta_r H_0^\circ(10) = -0.09 \pm 0.19$  eV. Notably, eqn (17) requires that  $D_0(\text{OAn}^+-\text{O}) + D_0(\text{An}^+-\text{O}) - D_0(\text{An}^+-\text{H}) \geq 13.01$  eV for reaction (10) to be exothermic.

Table 4 indicates that this may only be true for  $\text{PaH}^+$  and  $\text{UH}^+$ , although the high uncertainty in  $\Delta_r H_0^\circ(10) = -0.43 \pm 0.81$  eV may indicate otherwise for  $\text{PaH}^+$ . Observation of the energy dependence of reaction (10) may refine some of the uncertainty in the reported  $\text{Pa}^+$  BDEs.

There is a noticeable change in the reaction (10) thermochemistry after  $\text{UH}^+$ , as detailed in Table 4. While it has been clearly demonstrated that the 5f orbitals contribute to bonding in linear  $\text{OAnO}^+$ , the primary participants in bonding are the 6d orbitals. Promotion energies to a state with one unpaired 6d electron are nearly equivalent in the early series  $E_p(6d)$   $\text{Th}^+ \approx E_p(6d)$   $\text{Pa}^+ \approx E_p(6d)$   $\text{U}^+ \approx E_p(6d)$   $\text{Np}^+ \ll E_p(6d)$




Fig. 5 The correlation between  $E_p(6d^2)$  and the reaction (10) enthalpy calculated using the values found in Table 4 and eqn (17). The solid line represents the least square linear regression trend line  $(-0.5 + (1.3 \pm 0.3)E_p(6d^2), r^2 = 0.87)$ .

$\text{Pu}^+ \ll E_p(6d)$   $\text{Am}^+$ . For  $E_p(6d^2)$ ,  $E_p(6d^2)$   $\text{Th}^+ < E_p(6d^2)$   $\text{Pa}^+ \approx E_p(6d^2)$   $\text{U}^+ < E_p(6d^2)$   $\text{Np}^+ \ll E_p(6d^2)$   $\text{Pu}^+ \ll E_p(6d^2)$   $\text{Am}^+$ . This appears to lead to a shift in the reaction (10) enthalpy after U to a new trend as observed in Fig. 5. Note that  $\text{Th}^+$  is excluded from this analysis because it lacks the four valence electrons to form strong dioxide bonds.

Promotion energies,  $E_p(6d)$  and  $E_p(6d^2)$ , to some extent represent the accessibility of the 6d orbitals for bonding. Comparing these promotion energies to the  $\Delta_r H_0^\circ(10)$  indicates how influential the 6d orbitals may be in forming  $\text{OAnO}^+$  bonds. Assuming similar driving forces across the An series to  $\text{OAnO}^+$  bonding, the trend should be linear. Fig. 5 does indicate a relatively linear correlation, yet a clear difference between the early and latter An is observed. This mirrors observations in other single ligand systems that may have deviations in observed trends between  $\text{Np}^+$  and  $\text{Pu}^+$ .<sup>16,19</sup> This trend may suggest that there is a difference in bonding mechanism (*i.e.* orbital participation) for  $\text{NpO}_2^+ - \text{AmO}_2^+$ . Considering only spatial overlap as the driving force to molecular orbital formation, Fig. 5 is consistent with 5f orbital participation where the 5f orbitals are spatially larger for the early An. Armentrout and coworkers<sup>22,25</sup> have demonstrated 5f orbital participation for  $\text{ThO}_2^+$  and  $\text{UO}_2^+$ . 5f orbital participation in hydrocarbon activation in the early An has also been argued previously to interpret FTICR-MS studies of the reactions of  $\text{An}^+$  with hydrocarbons of varying lengths.<sup>49</sup> Nevertheless, 5f orbital participation in the early An cannot explain the trend observed in Fig. 5. A natural bond order analysis of  $\text{AnO}_2^+$  ( $\text{An} = \text{U} - \text{Am}$ ) presented by Feng, Glendening, and Peterson<sup>50</sup> indicates that the 5f and 6d orbital participation in both the  $\sigma$  and  $\pi$  bonding orbitals remains consistent across the four An at 50% 6d and 50% 5f (df hybrid orbital) and 50% 6d and 35% 5f ( $\text{pd}^3\text{f}^2$  hybrid orbital), respectively. Computational studies<sup>51</sup> utilizing B3LYP indicate two  $\sigma$  and two  $\pi$  bonding orbitals for neutral  $\text{AnO}_2$ .  $\sigma_g$  and  $\pi_g$  orbitals were characterized as having major



contributions from the O 2p orbitals with minor contributions from the An 6d/7s orbitals. By contrast the  $\sigma_u$  and  $\pi_u$  orbitals were characterized by major An 5f and minor O 2p orbital contributions. For the early An, the gerade orbitals are lower in energy, but this shifts between Am and Cm. The shift in the relative energy between the 5f and 6d orbitals is possibly the driving force to the shift in bonding. Between  $\text{Np}^+$  and  $\text{Pu}^+$  is where the 6d orbitals become less accessible. This concept can be observed in Table 1 where the  $E_p(6d)$  values are listed and Table S2 where  $E_p(6d^2)$  are listed. For both promotion energies, there is a clear escalation in energies from  $\text{Np}^+$  to  $\text{Pu}^+$ . Since the 5f/6d orbital participation does not seem to vary significantly according to theoretical observations, it is plausible that how the participation is incorporated shifts (*i.e.* orbital overlap *vs.* energy driven degeneracy).

An alternate explanation could be that  $\text{NpO}_2^+$  is unusual. Removing the  $\text{NpO}_2^+$  from the trend in Fig. 5 ( $r^2 = 0.87$ ) results in an improved linear correlation (Fig. S6,  $r^2 = 0.99$ ). This correlation excluding  $\text{NpO}_2^+$  is stronger than both of the linear trends displayed in Fig. S7 ( $r^2 = 0.95$  for both trends).  $\text{NpO}_2^+$  may be an outlier because the electronic configuration of  $\text{Np}^+$  is unusual for the early An series and has a  $^7L_5$  ( $5f^4 6d 7s$ ) ground level with a low lying  $^5I_4$  ( $5f^4 7s^2$ , 0.003 eV) level.<sup>52</sup>  $\text{Pa}^+$  ( $n = 4$ ) and  $\text{U}^+$  ( $n = 5$ ) both have  $5f^{n-2} 7s^2$  ground configurations, although the  $5f^{n-2} 6d 7s$  configurations are very low lying at 0.10 and 0.04 eV, respectively.  $\text{Pu}^+$  ( $n = 7$ ) and  $\text{Am}^+$  ( $n = 8$ ) both have  $5f^{n-1} 7s$  ground configurations. The 5f, 6d, and 7s orbitals of ground state  $\text{Np}^+$  are occupied, which may suggest that they are also similar in energy. This energy similarity may drive a bonding mechanism (orbital participation) for  $\text{Np}^+$  that is different from the other  $\text{An}^+$ , although theoretical calculations indicate minimal differences in the bonding orbital contribution.<sup>50</sup>  $\text{Np}^+$  also has a low lying  $j = (5/2, 1/2)$   $5f^3 7s$  level at 0.01 eV, so if the ground electronic configuration is related to the observed trend in Fig. 5, then  $\text{Np}^+$  can be grouped with both  $\text{Pa}^+/\text{U}^+$  and  $\text{Pu}^+/\text{Am}^+$ . The experimental uncertainty limits the conclusions that can be drawn because the trend excluding  $\text{NpO}_2^+$  (Fig. S6) and the split trend (Fig. S7) are both reasonable. Additional experimental work is necessary to provide support for the validity of either argument.

## Conclusion

The reactions of  $\text{AnH}^+ + \text{O}_2/\text{CO}_2$  ( $\text{An}^+ = \text{Th}, \text{U}$ ) may provide additional insight into An bonding. Observed reactions, in particular reaction (8), are consistent with thermochemistry derived from previous reports.<sup>16,17</sup> Extending these results to the additional An in the series indicates that there may be a change in bonding mechanism that starts at  $\text{Np}^+$ , Fig. 5. Notably, at  $\text{Np}^+$  the 6d orbitals become less accessible than the earlier  $\text{An}^+$  as measured by  $E_p(6d^2)$ . Computational studies,<sup>50</sup> however, suggest that there is minimal change in the orbitals involved in  $\text{AnO}_2^+$  bonding. Because the 5f orbitals contract with increasing atomic number, the consistent composition of the An bonding orbitals may be indicative of the

newer concept of energy driven covalency.<sup>53</sup> This concept is far from proven, but observed shifts in bonding in gas phase experiments<sup>16,19</sup> are consistent with an energy driven covalency model. Although much of the focus of An chemistry has been on when and why the 5f orbitals participate in bonding, these may be the wrong questions to ask. It appears that there may be a fundamental shift in bonding mechanism between early and latter An. Additional studies should be focused on understanding that difference.

## Conflicts of interest

There are no conflicts to declare.

## Data availability

All data used within this document are available within this document, the accompanying supplementary information (SI), or the manuscripts referenced in this document. Supplementary information: tabulated thermochemistry values, product branching ratios as a function of kinetic energy, kinetic energy dependence absolute reaction cross section of  $\text{U}^{2+} + \text{CO}_2$ . See DOI: <https://doi.org/10.1039/d5cp04417k>.

## Acknowledgements

The experimental data was collected with support from the Open Call Initiative under the Laboratory Directed Research and Development (LDRD) Program at Pacific Northwest National Laboratory (PNNL). Interpretation of this experimental data and writing this manuscript was supported by a U.S. Department of Energy, Office of Science, Early Career Research Program award in the Basic Energy Sciences, Division of Chemical Sciences, Geosciences, and Biosciences, Heavy Element Chemistry program under FWP 81364. PNNL is a multi-program national laboratory operated for the U.S. Department of Energy (DOE) by Battelle Memorial Institute under Contract No. DE-AC05-76RL01830.

## References

- 1 M. C. Heaven and K. A. Peterson, Probing Actinide Bonds in the Gas Phase: Theory and Spectroscopy, in *Experimental and Theoretical Approaches to Actinide Chemistry*, ed. J. K. Gibson and W. A. de Jong, John Wiley & Sons Ltd, Oxford, 2018.
- 2 P. E. Moreland, D. J. Rokop and C. M. Stevens, Mass-Spectrometric Observations of Uranium and Plutonium Monohydrides Formed by Ion–Molecule Reaction, *Int. J. Mass Spectrom. Ion Phys.*, 1970, **5**(1), 127–136.
- 3 P. B. Armentrout, R. V. Hodges and J. L. Beauchamp, Endothermic Reactions of Uranium Ions with  $\text{N}_2$ ,  $\text{D}_2$ , and  $\text{CD}_4$ , *J. Chem. Phys.*, 1977, **66**(10), 4683.
- 4 P. F. Souter, G. P. Kushto, L. Andrews and M. Neurock, Experimental and Theoretical Evidence for the Isolation of



- Thorium Hydride Molecules in Argon Matrices, *J. Phys. Chem. A*, 1997, **101**(7), 1287–1291.
- 5 X. Wang, L. Andrews and L. Gagliardi, Infrared Spectra of ThH<sub>2</sub>, ThH<sub>4</sub>, and the Hydride Bridging ThH<sub>4</sub>(H<sub>2</sub>)<sub>x</sub> (x = 1–4) Complexes in Solid Neon and Hydrogen, *J. Phys. Chem. A*, 2008, **112**(8), 1754–1761.
  - 6 J. Marçalo, M. Santos and J. K. Gibson, Gas-phase Reactions of Doubly Charged Actinide Cations with Alkanes and Alkenes-Probing the Chemical Activity of 5f Electrons from Th to Cm, *Phys. Chem. Chem. Phys.*, 2011, **13**(41), 18322–18329.
  - 7 E. Di Santo, M. Santos, M. C. Michelini, J. Marçalo, N. Russo and J. K. Gibson, Gas-Phase Reactions of the Bare Th<sup>2+</sup> and U<sup>2+</sup> Ions with Small Alkanes, CH<sub>4</sub>, C<sub>2</sub>H<sub>6</sub>, and C<sub>3</sub>H<sub>8</sub>: Experimental and Theoretical Study of Elementary Organoactinide Chemistry, *J. Am. Chem. Soc.*, 2011, **133**(6), 1955–1970.
  - 8 R. M. Cox, P. B. Armentrout and W. A. de Jong, Activation of CH<sub>4</sub> by Th<sup>+</sup> as Studied by Guided Ion Beam Mass Spectrometry and Quantum Chemistry, *Inorg. Chem.*, 2015, **54**(7), 3584–3599.
  - 9 R. M. Cox and P. B. Armentrout, Activation of Water by Thorium Cation: A Guided Ion Beam and Quantum Chemical Study, *J. Am. Soc. Mass Spectrom.*, 2019, **30**(10), 1835–1849.
  - 10 R. M. Cox, P. B. Armentrout and W. A. de Jong, Reactions of Th<sup>+</sup> + H<sub>2</sub>, D<sub>2</sub>, and HD Studied by Guided Ion Beam Tandem Mass Spectrometry and Quantum Chemical Calculations, *J. Phys. Chem. B*, 2016, **120**, 1601–1614.
  - 11 W.-J. Zhang, M. Demireva, J. Kim, W. A. de Jong and P. B. Armentrout, Reactions of U<sup>+</sup> with H<sub>2</sub>, D<sub>2</sub>, and HD Studied by Guided Ion Beam Tandem Mass Spectrometry and Theory, *J. Phys. Chem. A*, 2021, **125**(36), 7825–7839.
  - 12 G. F. de Melo, M. Vasiliu, M. Marshall, Z. Zhu, B. A. Tufekci, S. M. Ciborowski, M. Blankenhorn, R. M. Harris, K. H. Bowen and D. A. Dixon, Experimental and Computational Description of the Interaction of H and H<sup>−</sup> with U, *J. Phys. Chem. A*, 2022, **126**(27), 4432–4443.
  - 13 M. Vasiliu, K. A. Peterson, M. Marshall, Z. Zhu, B. A. Tufekci, K. H. Bowen and D. A. Dixon, Interaction of Th with H<sup>0/+−</sup>: Combined Experimental and Theoretical Thermodynamic Properties, *J. Phys. Chem. A*, 2022, **126**(2), 198–210.
  - 14 G. F. de Melo and D. A. Dixon, Protactinium and Actinium Monohydrides: A Theoretical Study on Their Spectroscopic and Thermodynamic Properties, *J. Phys. Chem. A*, 2022, **126**(36), 6171–6184.
  - 15 G. F. de Melo and D. A. Dixon, Energetic and Electronic Properties of NpH<sup>0/+−</sup> and PuH<sup>0/+−</sup>, *J. Phys. Chem. A*, 2023, **127**(14), 3179–3189.
  - 16 A. R. F. Bubas, D. Amanda, K. M. Melby, M. J. Rodriguez and R. M. Cox, The Balance of Orbital Overlap and Orbital Energy in the Activation of Methane by Actinide Cations: Insights from Inductively Coupled Plasma Tandem Mass Spectrometry, *Inorg. Chem. Front.*, 2025, **12**, 1503–1516.
  - 17 J. Marçalo and J. K. Gibson, Gas-Phase Energetics of Actinide Oxides: An Assessment of Neutral and Cationic Monoxides and Dioxides from Thorium to Curium, *J. Phys. Chem. A*, 2009, **113**(45), 12599–12606.
  - 18 J. K. Gibson, Bond Dissociation Energies Reveal the Participation of d Electrons in f-Element Halide Bonding, *J. Phys. Chem. A*, 2022, **126**(2), 272–285.
  - 19 R. M. Cox, A. R. Bubas, K. M. Melby, A. D. French, M. J. Rodriguez, M. P. Prange, N. Govind, K. A. Peterson and P. B. Armentrout, The Reactions of An<sup>+</sup> (An<sup>+</sup> = Th<sup>+</sup>, U<sup>+</sup>–Am<sup>+</sup>) + NO Reactions Observed by Inductively Coupled Plasma Tandem Mass Spectrometry, 2025, in progress.
  - 20 R. M. Cox, A. Kafle, P. B. Armentrout and K. A. Peterson, Bond Energy of ThN<sup>+</sup>: A Guided Ion Beam and Quantum Chemical Investigation of the Reactions of Thorium Cation with N<sub>2</sub> and NO, *J. Chem. Phys.*, 2019, **151**(3), 034304.
  - 21 A. R. Bubas, A. Kafle, B. C. Stevenson and P. B. Armentrout, The Bond Energy of UN<sup>+</sup>: Guided Ion Beam Studies of the Reactions of U<sup>+</sup> with N<sub>2</sub> and NO, *J. Chem. Phys.*, 2024, **160**, 164305.
  - 22 P. B. Armentrout and K. A. Peterson, Guided Ion Beam and Quantum Chemical Investigation of the Thermochemistry of Thorium Dioxide Cations: Thermodynamic Evidence for Participation of f Orbitals in Bonding, *Inorg. Chem.*, 2020, **59**(5), 3118–3131.
  - 23 K. S. Otte, J. E. Niklas, C. M. Studvick, C. L. Montgomery, A. R. C. Bredar, I. A. Popov and H. S. La Pierre, Proton-Coupled Electron Transfer at the Pu<sup>5+/4+</sup> Couple, *J. Am. Chem. Soc.*, 2024, **146**, 21859–21867.
  - 24 J. E. Niklas, K. S. Otte, C. M. Studvick, S. Roy Chowdhury, B. Vlasisavljevich, J. Bacsá, F. Kleemiss, I. A. Popov, H. S. La Pierre and A. Tetrahedral, Neptunium(v) Complex, *Nat. Chem.*, 2024, **16**(9), 1490–1495.
  - 25 A. R. Bubas, W.-J. Zhang, P. B. Armentrout and A. Guided Ion Beam Investigation, of UO<sub>2</sub><sup>+</sup> Thermodynamics and f orbital Participation: Reactions of U<sup>+</sup> + CO<sub>2</sub>, UO<sup>+</sup> + O<sub>2</sub>, and UO<sup>+</sup> + CO, *J. Chem. Phys.*, 2023, **159**(24), 244305.
  - 26 R. M. Cox, A. R. Bubas, K. M. Melby, A. D. French, M. J. Rodriguez, M. P. Prange and N. Govind, The Curious Case of Pu<sup>+</sup>: Insight on 5f Orbital Activity from Inductively Coupled Plasma Tandem Mass Spectrometry (ICP-MS/MS) Reactions, *Inorg. Chem.*, 2024, **63**(43), 20617–20624.
  - 27 M. Santos, J. Marçalo, A. P. D. Matos, J. K. Gibson and R. G. Haire, Gas-Phase Oxidation Reactions of Neptunium and Plutonium Ions Investigated via Fourier Transform Ion Cyclotron Resonance Mass Spectrometry, *J. Phys. Chem. A*, 2002, **106**, 7190–7194.
  - 28 M. Santos, A. P. d Matos, J. Marçalo, J. K. Gibson, R. G. Haire, R. Tyagi and R. M. Pitzer, Oxidation of Gas-Phase Protactinium Ions, Pa<sup>+</sup> and Pa<sup>2+</sup>: Formation and Properties of PaO<sub>2</sub><sup>2+</sup>(g), Protactinyl, *J. Phys. Chem. A*, 2006, **110**, 5751–5759.
  - 29 M. Santos, J. Marçalo, J. P. Leal, A. P. D. Matos, J. K. Gibson and R. G. Haire, FTICR-MS Study of the Gas-phase Thermochemistry of Americium Oxides, *Int. J. Mass Spectrom.*, 2003, **228**, 457–465.
  - 30 P. B. Armentrout and J. L. Beauchamp, Reactions of U<sup>+</sup> and UO<sup>+</sup> with O<sub>2</sub>, CO, CO<sub>2</sub>, COS, CS<sub>2</sub> and D<sub>2</sub>O, *Chem. Phys.*, 1980, **50**, 27–36.



- 31 R. M. Cox, K. Harouaka, M. Citir and P. B. Armentrout, Activation of CO<sub>2</sub> by Actinide Cations (Th<sup>+</sup>, U<sup>+</sup>, Pu<sup>+</sup>, and Am<sup>+</sup>) as Studied by Guided Ion Beam and Triple Quadrupole Mass Spectrometry, *Inorg. Chem.*, 2022, 8168–8181.
- 32 J. G. Terhorst, T. A. Corcovilos and M. J. van Stipdonk, Conversion of a UO<sub>2</sub><sup>2+</sup> Precursor to UH<sup>+</sup> and U<sup>+</sup> Using Tandem Mass Spectrometry to Remove Both “yl” Oxo Ligands, *J. Am. Soc. Mass Spectrom.*, 2023, 34(11), 2439–2442.
- 33 J. G. Terhorst, T. A. Corcovilos, S. J. Lenze and M. J. van Stipdonk, Synthesis of Organo-Uranium(II) Species in the Gas-Phase using Reactions Between [UH]<sup>+</sup> and Nitriles, *Dalton Trans.*, 2025, 54(1), 231–238.
- 34 K. P. Hobbs, A. D. French, K. M. Melby, E. J. Bylaska, K. Harouaka, R. M. Cox, I. J. Arnquist and C. L. Beck, Assessing Gas-Phase Ion Reactivity of 50 Elements with NO and the Direct Application for <sup>239</sup>Pu in Complex Matrices Using ICP-MS/MS, *Anal. Chem.*, 2024, 96(15), 5807–5814.
- 35 K. Harouaka, C. Allen, E. Bylaska, R. M. Cox, G. C. Eiden, M. L. D. Vacri, E. W. Hoppe and I. J. Arnquist, Gas-Phase Ion-Molecule Interactions in a Collision Reaction Cell with Triple Quadrupole-Inductively Coupled Plasma Mass Spectrometry: Investigations with N<sub>2</sub>O as the Reaction Gas, *Spectrochim. Acta, Part B*, 2021, 186, 106309.
- 36 P. B. Armentrout, Mass Spectrometry – Not Just a Structural Tool: The Use of Guided Ion Beam Tandem Mass Spectrometry to Determine Thermochemistry, *J. Am. Soc. Mass Spectrom.*, 2002, 13(5), 419–434.
- 37 N. Yamada, Kinetic Energy Discrimination in Collision/Reaction Cell ICP-MS: Theoretical Review of Principles and Limitations, *Spectrochim. Acta, Part B*, 2015, 110, 31–44.
- 38 A. R. Bubas, R. M. Cox, K. M. Melby, A. D. French, M. J. Rodriguez, M. P. Prange, N. Govind and P. B. Armentrout, The Reactions of Ln<sup>+</sup> (Ln<sup>+</sup> = Ce<sup>+</sup>, Pr<sup>+</sup>, Nd<sup>+</sup>, Sm<sup>+</sup>, Eu<sup>+</sup>) + NO Reactions Observed by Inductively Coupled Plasma Tandem Mass Spectrometry, *J. Chem. Phys.*, 2025, submitted.
- 39 R. M. Cox, K. M. Melby, A. D. French and M. J. Rodriguez, f-Block Reactions of Metal Cations with Carbon Dioxide Studied by Inductively Coupled Plasma Tandem Mass Spectrometry, *Phys. Chem. Chem. Phys.*, 2024, 26(1), 209–218.
- 40 T. Su and M. T. Bowers, Ion-Polar Molecule Collisions: The Effect of Ion Size on Ion-Polar Molecule Rate Constants: The Parameterization of the Average Dipole Orientation Theory, *Int. J. Mass Spectrom. Ion Phys.*, 1973, 12, 347–356.
- 41 R. M. Cox, M. Citir, P. B. Armentrout, S. R. Battey and K. A. Peterson, Bond Energies of ThO<sup>+</sup> and ThC<sup>+</sup>: A Guided Ion Beam and Quantum Chemical Investigation of the Reactions of Thorium Cation with O<sub>2</sub> and CO, *J. Chem. Phys.*, 2016, 144, 184309.
- 42 W. Zhang, A. R. E. Hunt, M. Demireva, J. Kim, K. A. Peterson and P. B. Armentrout, Bond Energies of UO<sup>+</sup> and UC<sup>+</sup>: Guided Ion Beam and Quantum Chemical Studies of the Reactions of Uranium Cation with O<sub>2</sub> and CO, *Isr. J. Chem.*, 2023, e202300026.
- 43 H. H. Cornehl, R. Wesendrup, M. Diefenbach and H. Schwarz, A Comparative Study of Oxo-Ligand Effects in the Gas-Phase Chemistry of Atomic Lanthanide and Actinide Cations, *Chem. – Eur. J.*, 1997, 3(7), 1083–1090.
- 44 J. K. Gibson, R. G. Haire, M. Santos, J. Marcalo and A. Pires de Matos, Oxidation Studies of Dipositive Actinide Ions, An<sup>2+</sup> (An = Th, U, Np, Pu, Am) in the Gas Phase: Synthesis and Characterization of the Isolated Uranyl, Neptunyl, and Plutonyl Ions UO<sub>2</sub><sup>2+(g)</sup>, NpO<sub>2</sub><sup>2+(g)</sup>, and PuO<sub>2</sub><sup>2+(g)</sup>, *J. Phys. Chem. A*, 2005, 109(12), 2768–2781.
- 45 G. Mazzone, M. d. C. Michelini, N. Russo and E. Sicilia, Mechanistic Aspects of the Reaction of Th<sup>+</sup> and Th<sup>2+</sup> with Water in the Gas Phase, *Inorg. Chem.*, 2008, 47(6), 2083–2088.
- 46 J. Zhou and H. B. Schlegel, Ab Initio Molecular Dynamics Study of the Reaction between Th<sup>+</sup> and H<sub>2</sub>O, *J. Phys. Chem. A*, 2010, 114(33), 8613–8617.
- 47 R. D. Johnson III, NIST Computational Chemistry Comparison and Benchmark Database. <https://cccbdb.nist.gov/> (accessed April 12, 2018).
- 48 P. B. Armentrout and R. M. Cox, Potential Energy Surface for Reaction of Sm<sup>+</sup> + CO<sub>2</sub> → SmO<sup>+</sup> + CO: Guided Ion Beam and Theoretical Studies, *Phys. Chem. Chem. Phys.*, 2017, 19, 11075–11088.
- 49 J. K. Gibson, R. G. Haire, J. Marçalo, M. Santos, A. Pires de Matos, M. K. Mroziak, R. M. Pitzer and B. E. Bursten, Gas-Phase Reactions of Hydrocarbons with An<sup>+</sup> and AnO<sup>+</sup> (An = Th, Pa, U, Np, Pu, Am, Cm): The Active Role of 5f Electrons in Organoprotactinium Chemistry, *Organometallics*, 2007, 26(16), 3947–3956.
- 50 R. Feng, E. D. Glendening and K. A. Peterson, Actinyl Cation-Cation Interactions in the Gas Phase: An Accurate Thermochemical Study, *Phys. Chem. Chem. Phys.*, 2019, 21(15), 7953–7964.
- 51 A. Kovacs, P. Pogany and R. J. M. Konings, Theoretical Study of Bond Distances and Dissociation Energies of Actinide Oxides AnO and AnO<sub>2</sub>, *Inorg. Chem.*, 2012, 51(8), 4841–4849.
- 52 J. E. Sansonetti and W. C. Martin, NIST Standard Reference Database 108. <https://www.nist.gov/pml/handbook-basic-atomic-spectroscopic-data>.
- 53 J. Su, E. R. Batista, K. S. Boland, S. E. Bone, J. A. Bradley, S. K. Cary, D. L. Clark, S. D. Conradson, A. S. Ditter, N. Kaltsoyannis, J. M. Keith, A. Kerridge, S. A. Kozimor, M. W. Löble, R. L. Martin, S. G. Minasian, V. Mocko, H. S. La Pierre, G. T. Seidler, D. K. Shuh, M. P. Wilkerson, L. E. Wolfsberg and P. Yang, Energy-Degeneracy-Driven Covalency in Actinide Bonding, *J. Am. Chem. Soc.*, 2018, 140(51), 17977–17984.

