

PCCP

Physical Chemistry Chemical Physics

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: R. Veillet, O. Venot, F. Citrangolo Destro, T. Fages, R. Fournet, P. Glaude, R. BOUNACEUR and B. Sirjean, *Phys. Chem. Chem. Phys.*, 2025, DOI: 10.1039/D4CP04873C.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Cite this: DOI: 00.0000/xxxxxxxxxx

Received Date
Accepted Date

DOI: 00.0000/xxxxxxxxxx

A theoretical study of CH_3SH pyrolysis kinetics[†]Roméo Veillet,^{*a} Olivia Venot,^a Fabiola Citrangolo Destro,^b Timothée Fages,^b René Fournet,^b Pierre-Alexandre Glaude,^b Roda Bounaceur,^b and Baptiste Sirjean^b

The decomposition mechanism of methanethiol (CH_3SH) under pyrolysis conditions has been investigated using electronic structure calculations at the CCSD(T)-F12/cc-pVTQ ∞ Z-F12//CCSD(T)-F12/cc-pVDZ-F12 level of theory for several potential energy surfaces such as $\text{CH}_3\text{SH} + \text{SH}$, $\text{CH}_3\text{SH} + \text{CH}_3$, $\text{CH}_2\text{S} + \text{SH}$, $\text{CH}_2\text{S} + \text{CH}_3$ and $\text{CS} + \text{SH}$. Pressure dependent reactions rates for the associated reactions have been deduced from energies, frequencies and geometries calculations by solving the master equation for each surface. Complex behavior is found for these systems with very low to submerged energy barriers for the $\text{CH}_2\text{S} + \text{SH}$ addition on the S atom, mainly due to multireference effects and Van der Waals complex formation. The former has been explored in details with CASPT2-F12/cc-pVTZ-F12 calculations up to an active space size of 13 electrons in 15 orbitals, and the latter was accounted for with phase space theory. Comparison with available experimental data shows that the resulting mechanism robustly predicts the abundance of the main species over the full temperature range.

1 Introduction

High temperature kinetics of sulfur-bearing compounds in gas phase are relevant to a wide span of applications, ranging from practical industrial applications of combustion, up to theoretical research on photochemistry and atmospheric chemistry^{1–3}. In combustion applications, the Claus process is the standard method to purify natural gas and extract sulfur from gases containing hydrogen sulfide, such as sour shale gas, to produce elemental sulfur for the chemical industry⁴. This process involves the oxidation of hydrogen sulfide (H_2S), which is the main sulfur-bearing species in natural gas, but other contaminants such as methanethiol CH_3SH can also be present in concentrations sufficient to impair sulfur yields⁵. The highly exothermic nature of this oxidation leads to high temperatures in the reactor and to the thermal decomposition of these sulfur-bearing compounds. Research for optimizing this process has also been increasing in popularity due to the recent interest in biogas production as a renewable energy source. This type of gas also contains high amounts of sulfur compounds in the form of H_2S and CH_3SH , that needs to be removed for its use in gas turbines^{6,7}.

The kinetics of CH_3SH have also been previously explored by atmospheric chemists, because of its crucial role in the sulfur cycle on Earth. It involves sulfur emissions from organic sulfur compounds of living organisms in the oceans, released in the form of

gases such as CH_3SH and CH_3SCH_2 (dimethylsulfide or DMS), the latter constituting its main form in the atmosphere^{8,9}. These molecules have a very short residence time in the atmosphere (about 1 day) because they are quite rapidly oxidized by atmospheric molecular oxygen, which promptly reacts with the CH_3S and CH_3SCH_2 radicals produced by the reaction of CH_3SH and DMS with other radicals produced by photochemistry such as the hydroxyl radical (OH)^{10–13}.

These reactions happen on Earth at relatively low temperatures, but the need for high temperature data has been increasing these past few years in the field of astrophysics, in particular in exoplanets atmospheric studies, due to the recent observational evidence of high-temperature photochemistry. Since the launch of the James Webb Space Telescope in December 2021, the exoplanet atmospheric chemistry community has reported multiple detection of sulfur, notably in the warm atmospheres of planets such as WASP-39 b and WASP-107 b^{14,15}. The observations of these 2 planets constitute the very first detections of sulfur element in exoplanet atmospheres. Surprisingly, sulfur dioxide (SO_2), the highly oxidized state of sulfur, has been detected, whereas its presence was not expected according to chemical equilibrium predictions. The presence of SO_2 has been explained through a pathway initiated by photochemistry^{14,15}. Multiple sulfur kinetic networks have been used to try to model the atmosphere of these planets and reproduce the observational data^{14,16–19}, but these networks only include sulfur chemistry as an S/O/H system, although in hydrogen dominated atmospheres such as the ones expected for WASP-39 b and WASP-107 b, carbon and nitrogen are also present in higher concentrations than

^a Université Paris Cité and Univ Paris Est Creteil, CNRS, LISA, F-75013 Paris, France^b Université de Lorraine, CNRS, LRGP, F-54000 Nancy, France

† Supplementary Information available: [details of any supplementary information available should be included here]. See DOI: 10.1039/cXCP00000x/



sulfur, in the form of CH_4 and NH_3 . Photochemistry of these two species could couple to photochemistry of sulfur compounds through C/H/S species, of which CH_3SH is the simplest representative. To our knowledge, this coupling remains to this day an unexplored area of sulfur chemistry in exoplanets and has not been included in any chemical network used to model these atmospheres. Hence, this possibility should at least be explored and properly modeled to improve our confidence in the predictions of these networks.

This work aims to develop and validate a detailed kinetic model for the pyrolysis of CH_3SH , using high-level *ab initio* calculations, such as CCSD(T)-F12 and CASPT2-F12, to calculate the rate constants of key reactions of the mechanism. First, we discuss in Sect. 2 how we identify the most sensitive reactions for the validation of the kinetic model against pyrolysis speciation data. Then, we present in Sect. 3.3 the results of our calculations and simulate experimental pyrolysis data from the literature with this kinetic model in Sect. 3.4. The final section concludes with a discussion of the implications of this work.

2 Methodology

Detailed kinetic models for the pyrolysis of CH_3SH are scarce in the literature. Recently, Colom-Diaz et al. (2021)²⁰ proposed a mechanism validated against their experimental data measured in a jet stirred reactor. This model is based on a previous work from the group²¹ that models CH_3SH oxidation and validates it against species abundance profiles obtained in a plug flow reactor. Other works on C/S species such as CS_2 exist, for example in Glarborg et al. (2014)²², but usually without including C/H/S species. For C/H/S species, Alzueta et al. (2019)²¹ and Colom-Diaz et al. (2021)²⁰ proposed a mechanism used to simulate their experiments from CH_3SH combustion and pyrolysis in flow and jet-stirred reactors. If their model is able to capture the oxidation behavior of CH_3SH , the pyrolysis mechanism still needs to be improved. However, given the lack of available data on the rate constants of reactions involving C/H/S species, the accurate modeling of CH_3SH pyrolysis and especially its decomposition products over wide ranges of temperatures is a very complex and delicate task.

In consequence, we chose to build the C/H/S mechanism from the scratch, using exhaustive analogies with equivalent oxygenated species to build a strong core for the mechanism. This applied to CH_3SH and its primary pyrolysis products such as CH_2S , CS_2 and their corresponding radicals. Enthalpy differences were considered in the activation energies of these reactions when the original or new reaction was implemented in the endothermic direction. For example when implementing the $\text{CH}_3\text{SH} + \text{HS}_2 \rightarrow \text{CH}_2\text{SH} + \text{H}_2\text{S}_2$ reaction, because the analog reaction $\text{CH}_3\text{OH} + \text{HO}_2 \rightarrow \text{CH}_2\text{OH} + \text{H}_2\text{O}_2$ was already implemented in the endothermic direction in our C/H/O/N core, we subtracted its enthalpy difference at 1000 K (about 9.9 kcal/mol) from its activation energy (18.8 kcal/mol) and added the enthalpy difference of the analogous sulfur reaction (26.4 kcal/mol) to yield its activation energy (35.3 kcal/mol). This was done systematically to avoid bias in the activation energy when the reaction enthalpy differs too much from those of the analog reaction

available. Contrary to CH_3SH chemistry, H_2S and $\text{C}_0\text{-C}_2$ chemistry were included based on extensive validations done against literature data. This was done starting from a full C/H/O/N mechanism from a previous work²³, in which we integrated an S/H/O sub-mechanism from Stagni et al. (2022)²⁴ without replacing our H/O submechanism. The references for the C/H/O reactions used for the C/H/S analogies are directly taken from the C/H/O core mechanism of Veillet et al. (2024)²³, and is based on the work of Burke et al. (2016)²⁵. This resulted in over 120 analogies, which can be found in the text file of the mechanism in Supplementary Data. For thermodynamic data, we used NASA polynomials from Glarborg et al. (2014)²² for C/S and O/C/S species, from Colom-Diaz et al. (2021)²⁰ and Sirjean et al. (2017)²⁶ for C/H/S species and Stagni et al. (2022)²⁴ for S/H and S/H/O species. Available experimental data on CH_3SH pyrolysis was then simulated with the resulting kinetic model and compared with its predictions. Formation pathway and sensitivity analysis was performed to target key reactions and further improve the model by computing their corresponding rates with state of the art *ab initio* methods.

3 Results


3.1 Target Experimental Data

After building the core of the C/H/S kinetic mechanism on these analogies, we simulated experimental data from the literature using the open source kinetic solver Cantera²⁷. The simulated conditions were taken from the pyrolysis experiments of Alzueta et al. (2019)²¹, listed as Set 1 in their work, and performed at atmospheric pressure between 800 K to 1400 K in a turbulent flow reactor with a temperature-dependent residence time of $\frac{194.6}{T[\text{K}]}$ seconds. The input gas mixture was composed of 983 ppm of CH_3SH and 42 ppm of O_2 , resulting in an effective oxidizer-to-fuel equivalence ratio of 0.014. The remaining bath gas used for dilution was N_2 . At the exit of the flow reactor, the abundance of CH_3SH , CH_4 , CO , CS_2 , H_2 , H_2S and SO_2 were measured. This data has been used to validate our mechanism performances and suggest improvements with *ab initio* calculations by looking at the discrepancies between experimental data and our modeling. More specifically, we ran a sensitivity and formation pathway analysis using Cantera to identify key reactions involved in the formation of major species. Simulations were performed with an ideal plug flow reactor and fixed temperature, as the high dilution and temperature control should prevent deviations from the target temperature.

3.2 Decomposition mechanism

Results from sensitivity and formation pathway analysis in these conditions around 1400 K suggested the decomposition mechanism shown in Fig. 1. In the first step of the decomposition, CH_3SH gets thermally dissociated into CH_3 and SH radicals. In our mechanism, this reaction has been approximated by analogy with the pressure dependent $\text{CH}_3\text{OH} \rightarrow \text{CH}_3 + \text{OH}$ reaction from Jasper et al. (2007)²⁸. As detailed in Sect. 2, this means that the pre-exponential and temperature exponent parameters of the Arrhenius expression were assumed identical with $\text{CH}_3\text{OH} \rightarrow \text{CH}_3 + \text{OH}$ but the activation energy was mod-



Fig. 1 Thermal decomposition mechanism for CH_3SH .

ified by adding the reaction enthalpy difference to $\text{CH}_3\text{SH} \rightarrow \text{CH}_3 + \text{SH}$, which was computed from each species NASA polynomials in the thermochemical data. This assumes that the enthalpy and entropy difference of the loose transition state to the reactant, the geometry of corresponding dividing surface, its rotational and vibrational partition functions and their temperature dependence are all identical between these reactions, with the exception of the translational mode in the reaction coordinate which is assumed to be the only varying parameter resulting in the previously described modification of the activation energy. The parameters for the pressure dependence of the reaction rate are also assumed to be identical to $\text{CH}_3\text{OH} \rightarrow \text{CH}_3 + \text{OH}$ with this approach, resulting in the same Troe parameters ($A = -0.4748$, $T_3 = 35580$ K, $T_1 = 1116$ K, $T_2 = 9023$ K) and Arrhenius coefficients for high pressure ($A = 2.084 \times 10^{18} \text{ s}^{-1}$, $n = -0.615$, $E = 74000 \text{ cal/mol}$) and low pressure ($1.500 \times 10^{43} \text{ cm}^3 \text{ mol}^{-1} \text{ s}^{-1}$, $n = -6.995$, $E = 79000 \text{ cal/mol}$). These assumptions are strong but seem reasonable due to the very similar geometry between CH_3OH and CH_3SH , and their validity were later confirmed by their good agreement with available experimental data and subsequent theoretical calculations (VRC-TST and ILT) that were beyond the scope of this work and will be the focus of a future publication. Briefly, these results showed that VRC-TST and our analogy usually agreed at least qualitatively, staying within the same order of magnitude in the whole P-T range studied (1000 - 10^{-6} bar, 200 - 5000 K), with up to a factor of 7 in the fall-off region, and significantly better agreement for low temperatures (below 1000 K). For high temperatures, ILT showed similar agreement but disagrees with our analogy and VRC-TST for low temperatures, predicting a rate constant up to 5 orders of magnitude lower at 300 K and an unphysical reverse rate for the recombination reaction $\text{CH}_3 + \text{SH} \rightarrow \text{CH}_3\text{SH}$ ($10^{16} \text{ cm}^3 \text{ mol}^{-1} \text{ s}^{-1}$) in the high pressure limit, which hints towards numerical issues in the procedure. The approximate nature of this analogy should therefore be kept in mind when using the present rate constant far from the experimental conditions for which its accuracy has been confirmed.

Following the CH_3SH dissociation, the CH_3 and SH radicals produced then react with CH_3SH through H-abstraction reactions,

mainly $\text{CH}_3\text{SH} + \text{CH}_3 \rightarrow \text{CH}_3\text{S} + \text{CH}_4$ and $\text{CH}_3\text{SH} + \text{SH} \rightarrow \text{CH}_3\text{S} + \text{H}_2\text{S}$, as CH_3S has a lower energy than its CH_2SH isomer, contrary to their CH_3O and CH_2OH analogs. Addition on the S lone pair has been investigated, but no stable structure was found. Resulting radicals CH_3S and CH_2SH can then undergo a β -scission through $\text{CH}_3\text{S} \rightarrow \text{CH}_2\text{S} + \text{H}$ and $\text{CH}_2\text{SH} \rightarrow \text{CH}_2\text{S} + \text{H}$ or react with CH_3 and SH radicals through the disproportionation reactions $\text{CH}_3\text{S} + \text{CH}_3 \rightarrow \text{CH}_2\text{S} + \text{CH}_4$, $\text{CH}_2\text{SH} + \text{CH}_3 \rightarrow \text{CH}_2\text{S} + \text{CH}_4$, $\text{CH}_3\text{S} + \text{SH} \rightarrow \text{CH}_2\text{S} + \text{H}_2\text{S}$ and $\text{CH}_2\text{SH} + \text{SH} \rightarrow \text{CH}_2\text{S} + \text{H}_2\text{S}$. The latter reactions are highly favored by reaction enthalpies at 1000 K (calculated based on thermodynamic data from our mechanism) of -56.4, -68.1, -42.0, and -53.6 kcal/mol respectively. For $\text{CH}_2\text{SH} + \text{CH}_3$ and $\text{CH}_3\text{S} + \text{CH}_3$, the $\text{C}_2\text{H}_4 + \text{H}_2\text{S}$ exit channel was also considered, but requires multiple isomerization and is not favorable in terms of entropy (respectively -13.4 and -4.0 cal/mol/K for the $\text{C}_2\text{H}_4 + \text{H}_2\text{S}$ exit channel in comparison to -16.3 and -6.9 cal/mol/K for the $\text{CH}_2\text{S} + \text{CH}_4$ exit channel at 1000 K), while being about the same in enthalpy (1.4 kcal/mol lower for $\text{C}_2\text{H}_4 + \text{H}_2\text{S}$). For the $\text{CH}_3\text{S} + \text{SH}$ and $\text{CH}_2\text{SH} + \text{SH}$ pathways, the other possible exit channel is $\text{CH}_4 + \text{S}_2$, which has an enthalpy lower by 10.1 kcal/mol, but no transition state was found for this reaction on the triplet surface.

These reactions all create the intermediate species CH_2S , that can then undergo either an H-abstraction through $\text{CH}_2\text{S} + \text{CH}_3 \rightarrow \text{CHS} + \text{CH}_4$, $\text{CH}_2\text{S} + \text{SH} \rightarrow \text{CHS} + \text{H}_2\text{S}$ or an addition on each side of the C-S double bond with the CH_3 and SH radicals. Formation and consumption of CH_2S is found to be a major kinetic bottleneck that controls both the relative and absolute proportions of the major pyrolysis end products. The description of its reactions with CH_3 and SH is thus crucial, as the pressure dependence of the branching ratios between the different exit channels heavily depends on the enthalpy difference for each channel. This effect cannot be easily corrected for due to the vastly different behavior of SH radicals in comparison to OH radicals in regard to double bond addition, as shown by Degirmenci et al. (2016)²⁹. Radicals with an unpaired electron localized on the sulfur atom also are subject to way lower barriers and higher enthalpy differences when the double bond also involves an S atom. As a result, OH and CH_3 additions on CH_2O are usually minor reaction pathways in comparison to the H-abstraction and are supposed to approach the high pressure limit for its rate constant. However, this may not be true on the potential energy surface of the S analog, especially at the high temperatures imposed during pyrolysis which favors the dissociation of the adduct. This is especially crucial because if the H-abstraction dominates, the formation of CHS will lead to end products such as CS_2 , which is formed almost exclusively from CHS through the CS intermediate species by the reactions $\text{CHS} + \text{CH}_3 \rightarrow \text{CS} + \text{CH}_4$, $\text{CHS} + \text{SH} \rightarrow \text{CS} + \text{H}_2\text{S}$ and then by the ipso-addition $\text{CS} + \text{SH} \rightarrow \text{CS}_2 + \text{H}$. Otherwise, if the addition dominates, other exit channels such as $\text{CH}_2\text{S} + \text{SH} \rightarrow \text{CH}_3 + \text{S}_2$ or $\text{CH}_2\text{S} + \text{CH}_3 \rightarrow \text{C}_2\text{H}_4 + \text{SH}$ leads to products with a broken C-S bond and may prevent the formation of CS_2 .






Fig. 2 Potential energy surface of the $\text{CH}_3\text{SH} + \text{SH}$ reaction at the CCSD(T)-F12/CBS//CCSD(T)-F12/cc-pVDZ-F12 level of theory.

3.3 Ab initio calculations

To improve the core C/H/S mechanism that was built using analogies, we identified five crucial potential energy surfaces from key reactions mentioned in the previous section: $\text{CH}_3\text{SH} + \text{SH}$, $\text{CH}_3\text{SH} + \text{CH}_3$, $\text{CH}_2\text{S} + \text{SH}$, $\text{CH}_2\text{S} + \text{CH}_3$, and $\text{CS} + \text{SH}$. For each of them, all the geometries and vibrational frequencies were computed at the CCSD(T)-F12/cc-pVDZ-F12 level of theory using the Molpro software^{30–32}. Energies were extrapolated to the CCSD(T)-F12 complete basis set limit from cc-pVTZ-F12 and cc-pVQZ-F12 energy calculations. Pressure-dependent phenomenological rate constants were then deduced using the Master Equation solver MESS³³. Internal rotors were corrected from the partition function using the 1D hindered rotor model with scans at the B2PLYPD3/cc-pVDZ level of theory with the Gaussian 16 software. Thermodynamical data in the NASA 7 polynomial format were computed using the atomization method with the Thermrot³⁴ software and energies/frequencies/geometries as previously mentioned. For reactions involving a H-atom transfer, tunneling corrections were applied using an asymmetric Eckart potential. Computed geometries, vibrational frequencies, torsional potentials, and the parameters used to solve the master equation are available in supplemental information in the form of MESS and Thermrot inputs. We detail hereafter the calculations for each reaction.

3.3.1 $\text{CH}_3\text{SH} + \text{SH}$ and $\text{CH}_3\text{SH} + \text{CH}_3$

Because of the importance and sensitivity of the first H-abstractions with CH_3SH , we decided to compute the full potential energy surface for $\text{CH}_3\text{SH} + \text{SH}$ and $\text{CH}_3\text{SH} + \text{CH}_3$. Two different exit channels were identified for each surface: $\text{CH}_3\text{S} + \text{H}_2\text{S}$, $\text{CH}_2\text{SH} + \text{H}_2\text{S}$ for $\text{CH}_3\text{SH} + \text{SH}$, and $\text{CH}_3\text{S} + \text{CH}_4$, $\text{CH}_2\text{SH} + \text{CH}_4$ for $\text{CH}_3\text{SH} + \text{CH}_3$, each corresponding to a different H-abstraction site. The corresponding potential energy surfaces are shown in Figs. 2 and 3.

For $\text{CH}_3\text{SH} + \text{SH}$, the formation of two Van der Waals complexes was found to have a significant impact on the kinetics at low temperatures. Indeed, this Van der Waals interaction results in an almost submerged barrier for $\text{CH}_3\text{SH} + \text{SH} \rightarrow \text{CH}_3\text{S} + \text{H}_2\text{S}$, which features a barrier height lower than the depth of the pre-reactive complex that crucially affects the perceived




Fig. 3 Potential energy surface of the $\text{CH}_3\text{SH} + \text{CH}_3$ reaction at the CCSD(T)-F12/CBS//CCSD(T)-F12/cc-pVDZ-F12 level of theory.




Fig. 4 Computed rate coefficients for the $\text{CH}_3\text{SH} + \text{SH}$ abstractions. Analogies with the $\text{CH}_3\text{OH} + \text{OH}$ equivalents from Xu et al. (2007)³⁵ are shown in dashed lines.

barrier height when including the effect of tunneling. To account for the formation of this pre-reactive complex, phase space theory, implemented in MESS code³³ was used for the barrierless formation of the entrance channel complex, based on a 1-dimensional potential between the centers of mass of the two reactants calculated at the B2PLYPD3/aug-CC-pVTZ level of theory. The resulting rate constants for these reactions are shown in Fig. 4, compared to the previously obtained values from the analog reaction $\text{CH}_3\text{OH} + \text{OH}$. The effect of bond dissociation energy differences between $\text{CH}_3\text{S-H}$ (-86.8 kcal/mol) and $\text{CH}_3\text{O-H}$ (-105.2 kcal/mol) are clearly seen for the reaction $\text{CH}_3\text{SH} + \text{SH} \rightarrow \text{CH}_3\text{S} + \text{H}_2\text{S}$. The exit channel $\text{CH}_3\text{S} + \text{H}_2\text{S}$ is highly favored over $\text{CH}_2\text{SH} + \text{H}_2\text{S}$ at low temperatures, although at high temperatures (1000 - 2000 K) both reactions contributes significantly to the total abstraction rate constant. This behavior is clearly in contrast with their oxygenated analog, where $\text{CH}_2\text{OH} + \text{OH}$ is always the preferred exit channel for all the plotted temperature range. Differences in the slope of the curves also show variations in the activation energy between the two systems, particularly for the reaction yielding CH_2SH . For the similar $\text{CH}_3\text{SH} + \text{CH}_3$ H-abstraction reactions, the entrance channel complex had a negligible energy in comparison to the activation energy (0.5 kcal/mol at the CCSD(T)-F12/CBS//CCSD(T)-F12/cc-pVDZ-F12 level of theory). Thus, for these reactions the rate constant were com-






Fig. 5 Computed rates coefficients for the $\text{CH}_3\text{SH} + \text{CH}_3$ H-abstractions. Analogies with the $\text{CH}_3\text{OH} + \text{CH}_3$ equivalents from Alecu et al. (2011)³⁶ are shown in dashed lines.

puted using Thermrot³⁴, and only considering the reactants, the transition states and the products. The resulting rate constants for this potential energy surface are shown in Fig. 5, as well as the rate constants determined by analogy with the $\text{CH}_3\text{OH} + \text{CH}_3$ equivalents. We observe the same behavior as previously stated with $\text{CH}_3\text{SH} + \text{SH}$, with the exit channel $\text{CH}_3\text{S} + \text{CH}_4$ dominating at low temperatures while $\text{CH}_2\text{SH} + \text{CH}_4$ only significantly contributes at very high temperatures. However, its amplitude is different, as the ratio between each corresponding rate at 500 K is around one order of magnitude for the $\text{CH}_3\text{SH} + \text{SH}$ surface, and around three orders of magnitude for the $\text{CH}_3\text{SH} + \text{CH}_3$ surface. The comparison with the oxygen analog also shows the inverse behavior between CH_3SH and CH_3OH , the latter favoring the CH_2OH exit channel. This has a huge impact on the accuracy of the analogy for the $\text{CH}_3\text{SH} + \text{CH}_3 \rightarrow \text{CH}_3\text{S} + \text{CH}_4$ reaction, but surprisingly has little impact on the rate constant of $\text{CH}_3\text{SH} + \text{CH}_3 \rightarrow \text{CH}_2\text{SH} + \text{CH}_4$.

3.3.2 $\text{CH}_2\text{S} + \text{SH}$ and $\text{CH}_2\text{S} + \text{CH}_3$

As explained in Sect. 2, the reactions of CH_2S with SH and CH_3 and the competition between addition and H-abstraction play a crucial role in controlling the products. In consequence, we decided to compute the full potential energy surface for both cases.

For $\text{CH}_2\text{S} + \text{SH}$, we considered the H-abstraction to $\text{CHS} + \text{H}_2\text{S}$ and the additions on the double bond, both on the sulfur atom, forming $\text{CH}_2\text{S}_2\text{H}$ and on the carbon atom, forming HSCH_2S . We also considered isomerizations both by H and SH transfers, adding the respective pathways $\text{CH}_2\text{S}_2\text{H} \rightarrow \text{CH}_3\text{S}_2$ and $\text{HSCH}_2\text{S} \rightarrow \text{CH}_2\text{S}_2\text{H}$. For the exit channels, we considered the β -scissions $\text{CH}_3\text{S}_2 \rightarrow \text{CH}_3 + \text{S}_2$ and $\text{HSCH}_2\text{S} \rightarrow \text{SCHSH} + \text{H}$. Due to the barrierless behavior of the $\text{CH}_3 + \text{S}_2$ exit channel, its inclusion required using multireference methods. This was done using Phase Space Theory together with a C-S bond scan at $\text{CASPT2}(11e,8o)/\text{cc-pVDZ}/\text{CASSCF}(11e,8o)/\text{cc-pVDZ}$ level of theory. The contributing determinants to the wavefunction during this relaxed scan are shown in Fig. ???. The resulting full potential energy surface is shown in Fig. 6.

The first issue was the transition state TS_2 of the addition on

the sulfur atom, which has an energy lower than the entry channel by 4.2 kcal/mol. When doing the intrinsic reaction coordinate scan to search for the geometry of the Van der Waals complex W_1 at the $\text{B2PLYPD3}/\text{cc-pVDZ}$ level of theory, we noticed the formation of two Van der Waals complexes with a peculiar geometry, showed in Fig. 7. The first one (VdW 1) is found to have a very close geometry to the transition state, with a S-S distance of 2.92 Å in comparison to 2.67 Å for the transition state. The S-H bond of the SH radical is not in the same plane as the CH_2S molecule, and this structure seems almost covalently bound given the orientation of the SH fragment. The second one (VdW2, W_1 in Fig. 6) is more in line with the expected geometry for such a system, with SH and CH_2S in the same plane, and the H atom oriented towards the S atom of CH_2S and a more distant position of the SH fragment. The first one was found to have a lower energy in our preliminary $\text{B2PLYPD3}/\text{cc-pVDZ}$ calculations. We also tried comparing the energies of the two geometries with $\text{CCSD}/\text{cc-pVDZ}$ level of theory to confirm this relationship, but we obtained the opposite result, with no Van der Waals complex with similar geometry to the first one and the second one behind the configuration minimizing the energy of the system. To try to understand the discrepancies between these results, we also tried to optimize the geometries with Gaussian counterpoise correction of the basis set superposition error, but we did not find significant contribution to justify these changes. To account for the Van der Waals complex formation, we therefore decided to use phase space theory with the second geometry complex based on a 1-dimensional potential between the centers of mass of the two reactants calculated at the $\text{B2PLYPD3}/\text{aug-CC-pVDZ}$ level of theory. Moreover, the subsequent calculations we performed at the $\text{CCSD}(\text{T})\text{-F12/CBS}/\text{CCSD}(\text{T})\text{-F12}/\text{cc-pVDZ-F12}$ level of theory exhibited convergence issues for both the transition state of the addition on the sulfur atom and the Van der Waals complex (TS_2 and W_1 in Fig. 6 respectively), which forced us to investigate on the matter. For these calculations, the T_1 diagnostic³⁷ was found to be above 0.05, which induced important oscillations in the CCSD iterative solver of Molpro, that were not solvable with a level shift nor incrementing the number of maximum steps. Therefore, given this extreme contribution of the triplets to the total energy, we used CASPT2 calculations. To determine the relevant active space size, we first performed a single point energy calculation in CASSCF with all the valence electrons included and 3 unoccupied valence orbitals, which resulted in 19 electrons in 13 orbitals, and looked at the occupation of the lower energy orbitals. From there, it was found that the LUMO was occupied at around 10%, and when looking at the Slater determinants composition of the wavefunction, this occupation was mainly caused by excited electron pair occupying this level, which contributed with a negative amplitude of around -0.15. Other determinants formed by the unpairing of electrons were also found to contribute the LUMO occupation, but also for the next orbital (LUMO + 1), which was occupied at around 3 %. As the involved electron pairs were the two doubly occupied orbitals with the highest energy, we decided to include these orbitals in the active space, as well as the singly occupied orbital and the next two unoccupied orbitals, which resulted in an active space of 5 electrons in 5 orbitals.






Fig. 6 Potential energy surface of the $\text{CH}_2\text{S} + \text{SH}$ reaction at the CCSD(T)-F12/CBS//CCSD(T)-F12/cc-pVDZ-F12 level of theory.




Fig. 7 B2PLYPD3/cc-pVDZ geometries of the two Van der Waals complexes found for the addition reaction $\text{CH}_2\text{S} + \text{SH} \rightarrow \text{CH}_2\text{S}_2\text{H}$. The first one is the minimum at this level of theory, while at the CCSD/cc-pVDZ level, the second one is the minimum. CCSD(T)-F12/cc-pVDZ-F12 calculations on this second geometry indicated a T1 diagnostic above 0.05 that prevented convergence of the wavefunction.

We performed a relaxed scan of the S-S bond length from 2 to 11 Å in CASPT2(5e,5o)/cc-pVDZ to capture the full Van der Waals interaction. We also recomputed the same scan from 2 to 3.6 Å with both CASPT2(5e,5o)-F12/cc-pVTZ-F12//CASPT2(5e,5o)/cc-pVDZ and CCSD(T)-F12/cc-pVDZ-F12 level of theories to focus on the transition state position. The results of this comparison are shown in Figs. 8 and 9. The position of the two minima on the potential energy surface is clearly visible on the full scan with CASPT2(5e,5o)/cc-pVDZ, but contrary to B2PLYPD3/cc-pVDZ, CCSD(T)-F12/cc-pVDZ-F12 and also CASPT2(5e,5o)-F12/cc-pVTZ-F12//CASPT2(5e,5o)/cc-pVDZ calculations, the second geometry with both fragments in the same plane (VdW 2) is of lower energy. However, this seems to be caused by an insufficient basis set size, as the energy calculations with CASPT2(5e,5o)-F12/cc-pVTZ-F12 using geometries from CASPT2(5e,5o)/cc-pVDZ calculations results in this geometry (VdW 2) being higher in energy than the out-of-plane geom-




Fig. 8 Relaxed scan of the S - S bond length in the $\text{CH}_2\text{S} + \text{SH}$ potential energy surface with CASPT2(5e,5o)/cc-pVDZ level of theory.

etry (VdW 1). Also with the CASPT2(5e,5o)-F12/cc-pVTZ-F12 level of theory, the transition state seems to be of higher energy than the $\text{CH}_2\text{S} + \text{SH}$ asymptote, which is not true anymore with all the other levels of theory previously mentioned. This can be seen on the second, finer close-range scan that was performed in Fig. 9, together with the transition state position that is very sensitive to the level of theory used. Another notable difference between CASPT2(5e,5o)/cc-pVDZ and CCSD(T)-F12/cc-pVDZ-F12 geometry optimization is that the former has available analytical gradients in Molpro, while the second uses the analytical gradient from the Alaska program³⁸. Different options for computing the numerical gradients were tested (finer step size, tighter convergence criteria, different coordinate system...) and the energy and position of the transition state seem to be quite dependent on these parameters, indicating that the flat geometry of the potential energy surface might result in a loose transition state and convergence to geometries far from its real one. However, for the default parameters, the relatively good agreement with geometries from CASPT2(5e,5o)/cc-pVDZ analytical gradient seems to indicate that the precision of the results from numerical gradients




Fig. 9 Relaxed scan of the S-S bond length in the $\text{CH}_2\text{S} + \text{SH}$ potential energy surface with two level of theory : CCSD(T)-F12/cc-pVDZ-F12 (yellow) and CASPT2(5e,5o)-F12/cc-pVTZ-F12 //CASPT2(5e,5o)/cc-pVDZ (blue). Starting from 3 Å, oscillations in the coupled cluster iterative solver prevents the wavefunction from converging.

might be sufficient for the purposes of this kinetic study.

For the reaction rate calculations, we therefore used the CCSD(T)-F12/cc-pVDZ-F12 geometries and energies, while using the coplanar Van der Waals complex geometry obtained (VdW 2) for its Phase Space Theory treatment. The resulting branching ratio for each exit channel and the rates at 1 bar are shown in Figs. 10 and 11 respectively.




Fig. 10 Computed branching ratio for the $\text{CH}_2\text{S} + \text{SH}$ potential energy surface in Fig. 6.

On the full pressure and temperature range (10^{-6} - 100 bar, 300 - 2500 K), we see that mainly two exit channels are relevant: the H-abstraction $\text{CHS} + \text{H}_2\text{S}$ and the addition on the sulfur atom yielding $\text{CH}_2\text{S}_2\text{H}$. At high pressures and low temperatures, stabilization of the HSCH_2S radical also contributes significantly to the total reaction rate. At 1 bar, which corresponds to the pressure in the experiments of Colom-Diaz et al. (2021)²⁰, we see that the H-abstraction dominates above 1000 K, while under this value, the addition on the sulfur atom dominates, stabilizing on the $\text{CH}_2\text{S}_2\text{H}$ radical, but also to a lesser extent on the HSCH_2S radical, which contributes increasingly at higher temperatures.




Fig. 11 Computed rate coefficients at 1 bar for the $\text{CH}_2\text{S} + \text{SH}$ potential energy surface. H-abstraction is shown in black, additions in red, and β -scission in blue.

The addition on the sulfur atom followed by the isomerization and stabilization to CH_3S_2 , however, is negligible. β -scission exit channels such as $\text{CH}_3 + \text{S}_2$ are never the favored reaction, but can possibly occur in multiple steps after the addition at low temperatures (< 1000 K). For the $\text{SCHSH} + \text{H}$ exit channels, its high energy transition state prevents it from any significant contribution, and thus is assumed to be negligible.

For the $\text{CH}_2\text{S} + \text{CH}_3$ potential energy surface, we considered the H-abstraction yielding to $\text{CHS} + \text{CH}_4$ (TS_3, P_2) and both the addition on the carbon atom to $\text{CH}_3\text{CH}_2\text{S}$ (TS_1, W_1) and the addition on the sulfur atom to CH_2SCH_3 (TS_2, W_2). We also considered the isomerizations by H atom transfer $\text{CH}_3\text{CH}_2\text{S} \rightarrow \text{CH}_2\text{CH}_2\text{SH}$ (TS_5) and CH_3 fragment transfer $\text{CH}_2\text{SCH}_3 \rightarrow \text{CH}_3\text{CH}_2\text{S}$ (TS_4). The β -scission exit channel to $\text{C}_2\text{H}_2 + \text{SH}$ (TS_6, P_3) was also considered. The resulting potential energy surface is shown in Fig. 12. Branching ratios and reaction rates are shown in Figs. 13 and 14 respectively. For the full pressure-temperature range computed, we can see that mainly three exit channels dominate : either the β -scission $\text{C}_2\text{H}_4 + \text{SH}$ at low temperatures and low pressures, the H-abstraction to $\text{CHS} + \text{CH}_4$ at high temperatures and low pressures, or the addition on the sulfur atom to CH_2SCH_3 at high pressures and low temperatures. At 1 bar, we see that the H-abstraction to $\text{CHS} + \text{CH}_4$ only becomes dominant above 1600 K, while under 1200 K the addition on the sulfur atom to CH_2SCH_3 dominates. Between these temperatures, the exit channel to the β -scission $\text{C}_2\text{H}_4 + \text{SH}$ is the major process that happens directly from the $\text{CH}_2\text{S} + \text{SH}$ input channel. In comparison to the $\text{CH}_2\text{S} + \text{SH}$ reaction rates, we can also see that the reactivity is a bit lower as the reaction rates only exceed 10^{12} $\text{cm}^3/\text{mol/s}$ for the H-abstraction above 1700 K, but the reaction enthalpy is also greater, which should result in slower reversed reactions.

3.3.3 CS + SH

Due to the formation of CHS radicals and their abstraction to CS , the next crucial reaction towards the formation of CS_2 is $\text{CS} + \text{SH}$. This reaction has been investigated with the same methodology described in Sect. 3.3, and the full resulting potential energy surface is shown in Fig. 15. We identified one addition site on




Fig. 12 Potential energy surface of the $\text{CH}_2\text{S} + \text{CH}_3$ reaction at the CCSD(T)-F12/CBS//CCSD(T)-F12/cc-pVDZ-F12 level of theory.




Fig. 13 Computed branching ratio for the $\text{CH}_2\text{S} + \text{CH}_3$ potential energy surface in Fig. 12.




Fig. 14 Computed rate coefficients at 1 bar for the $\text{CH}_2\text{S} + \text{CH}_3$ potential energy surface. H-abstraction is shown in black, additions in red, and β -scission in blue.

the carbon atom to form the SCSH radical (W_1 through TS_3), which can then undergo a β -scission to the $\text{CS}_2 + \text{H}$ products (P_2 through TS_2) to form the ipso-addition $\text{CS} + \text{SH} \rightarrow \text{CS}_2 + \text{H}$. This radical can also undergo an isomerization to the more stable SCHS radical (W_2 through TS_4) due to a resonance between the S atoms bearing the unpaired electron. This radical can also undergo a β -scission to the products $\text{CS}_2 + \text{H}$ (P_2 through TS_1). The evaluation of the energy of the addition transition state revealed a very low barrier, under the energy of the $\text{CS} + \text{SH}$ products. The formation of a Van der Waals complex with a linear geometry was identified (W_1) and treated with Phase Space Theory. The electronic energy of this complex was found to be about 0.5 kcal/mol under the transition state energy, but when adding the zero point energy of both structures their energy difference becomes lower than the computation accuracy (computed difference of 0.02 kcal/mol). The corresponding branching ratios and reaction rates for this potential energy surface are shown in Figs. 16 and 17 from the $\text{CS} + \text{SH}$ perspective and in Figs. 18 and 19

from the $\text{CS}_2 + \text{H}$ perspective, respectively. For the $\text{CS} + \text{SH}$ entry point, there is only one relevant pathway, which is the ipso-addition $\text{CS} + \text{SH} \rightarrow \text{CS}_2 + \text{H}$ that has a branching ratio of one over the whole range of pressures and temperatures used for the calculation. This results from its reaction rate being over 6 orders of magnitude higher than the two other reactions that stabilize the radicals SCSH and SCHS, and is itself due to a combination of two factors. First, the rate for this reaction is effectively restricted by the collision limit, as the limiting step is the formation of the Van der Waals complex due to the energy barrier of the transition state for the addition being lower than the $\text{CS} + \text{SH}$ energy. The energy difference between this barrier and the complex is so low that it cannot be stabilized even at low temperatures and high pressures, given that it is lower than thermal energy even at 200 K. Second, the β -scission transition state is lower than the $\text{CS} + \text{SH}$ energy by around 19 kcal/mol and the low well depth of about 18 kcal/mol under this transition state prevents the resulting radical to be stabilized at any pressure explored here. Conversely, if we look at $\text{CS}_2 + \text{H}$ as the entrance channel, the $\text{SCSH} \rightarrow$




Fig. 15 Potential energy surface of the CS + SH reaction at the CCSD(T)-F12/CBS//CCSD(T)-F12/cc-pVDZ-F12 level of theory.




Fig. 16 Computed branching ratio for the CS + SH potential energy surface in Fig. 15.




Fig. 18 Computed branching ratio for the CS + SH potential energy surface in Fig. 15 from the perspective of the CS2 + H entrance channel.




Fig. 17 Computed rate coefficients at 1 bar for the CS + SH potential energy surface. Addition is shown in red, ipso-addition in blue. The dashed line also requires an isomerization.




Fig. 19 Computed rate coefficients at 1 bar for the CS2 + H potential energy surface. Addition is shown in red, ipso-addition in blue. The dashed line also requires an isomerization.

CS + SH β -scission has to overcome the reverse reaction enthalpy, which creates a region at low temperatures (< 1000 K at atmo-

spheric pressure) where the addition product SCSH is stabilized, while the ipso-addition $\text{CS}_2 + \text{H} \rightarrow \text{CS} + \text{SH}$ only dominates at

high temperatures. The other addition to SCHS, while favorable in enthalpy in comparison to SCSH, is never dominant because of the higher energy of transition state for this pathway.

3.4 Comparison with experimental data

From the calculations of Sect. 3.3, we included in the mechanism every rate constant showed in Figs. 4, 5, 14, 11 and 17 in addition to each dominant pathway from other entrance channels that didn't correspond to a reverse reaction of these. The performance of the mechanism developed is tested against the experimental data on CH_3SH pyrolysis of Alzueta et al. (2019)²¹ on Fig. 20, and the corresponding sensitivity and rate of production analysis for $T = 1400$ K are given in Figs ?? and ?? respectively. For CH_3SH consumption, the mechanism perfectly reproduces the experimental data mainly thanks to the analogy used for the CH_3SH thermal decomposition $\text{CH}_3\text{SH} \rightarrow \text{CH}_3 + \text{SH}$, which used the differences in reaction enthalpies to compute the activation energy. This major initiation reaction is, as expected, the most sensitive reaction for every species. The production of CH_4 is in good agreement with the experimental data, as the mechanism correctly reproduces the shape of the temperature dependence, but underestimates its abundance by about 100 ppm over 1000 K. Fig. ?? shows that CH_4 is mainly produced by the reaction $\text{CH}_3 + \text{H}_2\text{S} \rightarrow \text{CH}_4 + \text{SH}$, although it is not a sensitive reaction. The most sensitive reaction excluding the initiation is $\text{CH}_3 + \text{CH}_3\text{SH} \rightarrow \text{CH}_3\text{S} + \text{CH}_4$, which has been computed in Sect. 3.3. The reevaluation of this rate constant is the main cause of the improvement of the simulations over the Colom-Diaz et al. (2021)²⁰ mechanism. As we already extensively investigated both the abstractions and the additions on CH_2S , we know it is not the cause of CH_4 underestimated abundance. Fig. ?? gives an idea of the species to which the remaining carbon is distributed. Among the species CH_2S , CS_2 , C_2H_6 , C_2H_4 and C_2H_2 that all bear carbon atoms, CH_2S and C_2H_6 are intermediate species that do not contribute to the final carbon balance. The overabundance of the CS_2 is also too small to fully explain the difference, and does not correspond to the same temperature range as the under-abundance of CH_4 , which it cannot explain between 1000 and 1200 K. Therefore, the issue probably lies in C_2H_4 and C_2H_2 , which have significant abundances of 80 and 60 ppm respectively at the end of the simulation. As these species contain 2 carbon atoms, together they represent a potential of 280 ppm of CH_4 if hydrogen is supplied from another source. As previously seen in Sect. 3.3 and observed by Degirmenci et al. (2016)²⁹, the addition of SH radicals on double bonds is highly favored in comparison to OH, and could potentially be the cause of C_2H_4 and C_2H_2 decomposition to CH_4 above 1000 K. Further study of the full potential energy surface of $\text{C}_2\text{H}_4 + \text{SH}$ and $\text{C}_2\text{H}_2 + \text{SH}$ with both H-abstraction and addition considered would be needed, but is out of the scope of this work. The production of CS_2 matches also quite well the experimental data, despite overestimating its abundance by around 50 ppm over 1250 K. Its main production pathway is the reaction $\text{CS} + \text{SH} \rightarrow \text{CS}_2 + \text{H}$ and to a lesser extent the reaction $\text{CS} + \text{S}_2 \rightarrow \text{CS}_2 + \text{S}$. The most sensitive reactions besides the initiation are the H-abstraction $\text{CH}_2\text{S} + \text{SH} \rightarrow$

$\text{CHS} + \text{H}_2\text{S}$ and the ipso-addition $\text{CS} + \text{SH} \rightarrow \text{CS}_2 + \text{H}$, which we computed in Sect. 3.3, and believe to be the cause of improvement over the Colom-Diaz et al. (2021)²⁰ mechanism. Improvement in the CS_2 abundance above 1200 K could probably be achieved by considering other reactions of CS, which is very reactive in contrast to CO despite being a closed-shell molecule. This could include the study of the full $\text{CS} + \text{CH}_3$ potential energy surface, or reactions with other abundant closed-shell species such as CH_3SH , H_2 , H_2S and CH_4 . The abundance of H_2S is also well reproduced with a slight underestimation similar to CH_4 , although its magnitude is closer to 50 ppm. At 1400 K, H_2S seems to approach thermochemical equilibrium, with production mainly comes from H-abstractions on CH_2S and CH_3SH through the reactions $\text{CH}_2\text{S} + \text{SH} \rightarrow \text{CHS} + \text{H}_2\text{S}$ and $\text{CH}_3\text{SH} + \text{SH} \rightarrow \text{CH}_3\text{S} + \text{H}_2\text{S}$. On the opposite, destruction reactions are mainly H-abstractions on H_2S by the H and CH_3 radicals through the reactions $\text{H}_2\text{S} + \text{H} \rightarrow \text{H}_2 + \text{SH}$ and $\text{H}_2\text{S} + \text{CH}_3 \rightarrow \text{CH}_4 + \text{SH}$. Sensitive reactions are mainly H-abstraction on CH_3SH by SH radicals for both channels $\text{CH}_3\text{SH} + \text{SH} \rightarrow \text{CH}_3\text{S} + \text{H}_2\text{S}$ and $\text{CH}_3\text{SH} + \text{SH} \rightarrow \text{CH}_2\text{SH} + \text{H}_2\text{S}$, and to a lesser extent the non-thermal addition/isomerisation/beta-scission reaction $\text{CH}_2\text{S} + \text{SH} \rightarrow \text{CH}_3 + \text{S}_2$ that were all computed in Sect. 3.3. The massive improvement over the Colom-Diaz et al. (2021)²⁰ mechanism is therefore believed to be mainly due to the accurate computation of the H-abstraction reactions on CH_3SH by SH radicals with the treatment of the Van der Waals complex in Sect. 3.3. The abundance of H_2 is almost perfectly reproduced. In comparison, the Colom-Diaz et al. (2021)²⁰ mechanism overestimates H_2 by a factor of 2, and underestimates H_2S and CH_4 by more than a factor of 3. Its formation mainly involves H-abstraction on H_2S , CH_2S and CH_3SH by the H atom through the reactions $\text{H}_2\text{S} + \text{H} \rightarrow \text{H}_2 + \text{SH}$, $\text{CH}_2\text{S} + \text{H} \rightarrow \text{CHS} + \text{H}_2$ and $\text{CH}_3\text{SH} + \text{H} \rightarrow \text{CH}_3\text{S} + \text{H}_2$. When looking at the most sensitive reactions, the H-abstraction on CH_3SH is the most significant, along with the thermal decomposition of the CH_3S radical through the beta-scission $\text{CH}_3\text{S} \rightarrow \text{CH}_2\text{S} + \text{H}$. This reaction has seen its activation energy adjusted by preliminary B2PLYPD3/cc-pVDZ calculations, which are believed to be a source of improvement over the Colom-Diaz et al. (2021)²⁰ mechanism. As some residual oxygen was left in the experimental conditions, CO was also measured in the original data set, and although the temperature dependence doesn't match the experimental data perfectly, the order of magnitude is well reproduced with the present mechanism. This species is produced exclusively through the thermal decomposition of HCO through the reaction $\text{HCO} \rightarrow \text{CO} + \text{H}$, which seems to be produced by the oxidation mechanism through the addition of O_2 and its non-thermal decomposition to CH_2O through the reaction $\text{CH}_2\text{S} + \text{O}_2 \rightarrow \text{CH}_2\text{O} + \text{SO}$ and the H-abstraction of CH_2O through the reaction $\text{CH}_2\text{O} + \text{SH} \rightarrow \text{HCO} + \text{H}_2\text{S}$. These reactions were also analogies with activation energies approximated thanks to B2PLYPD3/cc-pVDZ calculations. Further work would be needed on the oxidation mechanism, but such a study is outside the scope of this work.






Fig. 20 Comparison between present work (yellow) and Colom-Diaz et al. (2021)²⁰ (blue) mechanisms on experimental CH₃SH pyrolysis data from Alzueta et al. (2019)²¹.



Conclusions

We presented an extensive theoretical study of the kinetics of CH_3SH pyrolysis. A reaction mechanism was built upon available reaction sub-mechanisms from the literature for H/O/S and C/S kinetics, and using analogies with oxygenated molecules for the basic C/H/S coupling. Sensitivity and rates of production analysis was then performed to determine key reactions $\text{CH}_3\text{SH} + \text{SH}$, $\text{CH}_3\text{SH} + \text{CH}_3$, $\text{CH}_2\text{S} + \text{SH}$, $\text{CH}_2\text{S} + \text{CH}_3$, $\text{CS} + \text{SH}$ for which theoretical reaction rates were calculated. Pressure dependence and pre-reactive complex formation were treated using master equation and phase space theory methods together with a CCSD(T)-F12/CBS//CCSD(T)-F12/cc-pVDZ-F12 level of theory. Multireference effects were also investigated using CASPT2 calculations. Additions on CH_2S are found to have a significant contribution in contrast to the case of CH_2O , due to the very low energy of the transition states. This often causes transition states with negative energies relative to the enthalpy asymptotes and create cases where pre-reactive complex treatment is necessary. Available experimental data have been compared to kinetics simulations, and the developed mechanism exhibits a robust behavior that correctly reproduces the experimental data.

Conflicts of interest

There are no conflicts to declare.

Data availability

The data supporting this article have been included as part of the Supplementary Information that contains all theoretical data as inputs of MESS and THEMROT codes, the kinetic mechanism in CHEMKIN and CANTERA formats, and supplementary Figures S1 to S4.

Acknowledgements

This project is funded by the ANR project 'EXACT' (ANR-21-CE49-0008-01). In addition, O.V. acknowledges funding from the Centre National d'Études Spatiales (CNES). High performance computing resources were provided by IDRIS under the allocation AD010812434R3 made by GENCI and also by the EXPLOR center hosted by the University of Lorraine.

Notes and references

- 1 J. S. Eow, *Environmental progress*, 2002, **21**, 143–162.
- 2 H. Bian, B. Xu, H. Zhang, Q. Wang, H. Zhang, S. Zhang and D. Xia, *International Journal of Quantum Chemistry*, 2019, **119**, e25822.
- 3 D. J. Kieber, J. Jiao, R. P. Kiene and T. S. Bates, *Journal of Geophysical Research: Oceans*, 1996, **101**, 3715–3722.
- 4 N. J. Nabikandi and S. Fatemi, *Journal of Industrial and Engineering Chemistry*, 2015, **30**, 50–63.
- 5 R. K. Rahman, A. G. Raj and S. Ibrahim, Abu Dhabi International Petroleum Exhibition and Conference, 2017, p. D021S052R004.
- 6 W. Burgers, P. Northrop, H. Kheshgi and J. Valencia, *Energy Procedia*, 2011, **4**, 2178–2184.
- 7 E. Dumont, *International Journal of Energy and Environment*, 2015, **6**, 479–498.
- 8 B. Nguyen, B. Bonsang and A. Gaudry, *Journal of Geophysical Research: Oceans*, 1983, **88**, 10903–10914.
- 9 T. Bates, B. Lamb, A. Guenther, J. Dignon and R. Stoiber, *Journal of Atmospheric Chemistry*, 1992, **14**, 315–337.
- 10 M. Chin, D. J. Jacob, G. M. Gardner, M. S. Foreman-Fowler, P. A. Spiro and D. L. Savoie, *Journal of Geophysical Research: Atmospheres*, 1996, **101**, 18667–18690.
- 11 D. Davis, G. Chen, P. Kasibhatla, A. Jefferson, D. Tanner, F. Eisele, D. Lenschow, W. Neff and H. Berresheim, *Journal of Geophysical Research: Atmospheres*, 1998, **103**, 1657–1678.
- 12 G. S. Tyndall and A. Ravishankara, *International Journal of Chemical Kinetics*, 1991, **23**, 483–527.
- 13 J. Chen, T. Berndt, K. H. Møller, J. R. Lane and H. G. Kjaergaard, *The Journal of Physical Chemistry A*, 2021, **125**, 8933–8941.
- 14 S.-M. Tsai, E. K. Lee, D. Powell, P. Gao, X. Zhang, J. Moses, E. Hébrard, O. Venot, V. Parmentier, S. Jordan *et al.*, *Nature*, 2023, **617**, 483–487.
- 15 A. Dyrek, M. Min, L. Decin, J. Bouwman, N. Crouzet, P. Mollière, P.-O. Lagage, T. Konings, P. Tremblin, M. Güdel *et al.*, *Nature*, 2024, **625**, 51–54.
- 16 S.-M. Tsai, M. Malik, D. Kitzmann, J. R. Lyons, A. Fateev, E. Lee and K. Heng, *The Astrophysical Journal*, 2021, **923**, 264.
- 17 J. I. Moses, C. Visscher, J. J. Fortney, A. P. Showman, N. K. Lewis, C. A. Griffith, S. J. Klippenstein, M. Shabram, A. J. Friedson, M. S. Marley *et al.*, *The Astrophysical Journal*, 2011, **737**, 15.
- 18 P. B. Rimmer and S. Rugheimer, *Icarus*, 2019, **329**, 124–131.
- 19 B. Drummond, P. Tremblin, I. Baraffe, D. S. Amundsen, N. J. Mayne, O. Venot and J. Goyal, *Astronomy & Astrophysics*, 2016, **594**, A69.
- 20 J. Colom-Díaz, M. Alzueta, Z. Zeng, M. Altarawneh and B. Dlugogorski, *Fuel*, 2021, **283**, 119258.
- 21 M. U. Alzueta, R. Pernía, M. Abián, Á. Millera and R. Bilbao, *Combustion and Flame*, 2019, **203**, 23–30.
- 22 P. Glarborg, B. Halaburt, P. Marshall, A. Guillory, J. Troe, M. Thellefsen and K. Christensen, *The Journal of Physical Chemistry A*, 2014, **118**, 6798–6809.
- 23 R. Veillet, O. Venot, B. Sirjean, R. Bounaceur, P.-A. Glaude, A. Al-Refaie and E. Hébrard, *Astronomy & Astrophysics*, 2024, **682**, A52.
- 24 A. Stagni, S. Arunthanayothin, L. Pratali Maffei, O. Herbinet, F. Battin-Leclerc and T. Faravelli, *Chemical Engineering Journal*, 2022, **446**, 136723.
- 25 U. Burke, W. K. Metcalfe, S. M. Burke, K. A. Heufer, P. Dagaut and H. J. Curran, *Combustion and Flame*, 2016, **165**, 125–136.
- 26 B. Sirjean, J.-C. Lizardo-Huerta, L. Verdier, R. Fournet and P.-A. Glaude, *Proceedings of the Combustion Institute*, 2017, **36**, 499–506.
- 27 D. G. Goodwin, H. K. Moffat, I. Schoegl, R. L. Speth and B. W. Weber, *Cantera: An Object-oriented Software Toolkit for*



- Chemical Kinetics, Thermodynamics, and Transport Processes*, <https://www.cantera.org>, 2023, Version 3.0.0.
- 28 A. W. Jasper, S. J. Klippenstein, L. B. Harding and B. Ruscic, *The Journal of Physical Chemistry A*, 2007, **111**, 3932–3950.
- 29 I. Degirmenci and M. L. Coote, *The Journal of Physical Chemistry A*, 2016, **120**, 1750–1755.
- 30 H.-J. Werner, P. J. Knowles, G. Knizia, F. R. Manby and M. Schütz, *Wiley Interdisciplinary Reviews: Computational Molecular Science*, 2012, **2**, 242–253.
- 31 H.-J. Werner, P. J. Knowles, F. R. Manby, J. A. Black, K. Doll, A. Heßelmann, D. Kats, A. Köhn, T. Korona, D. A. Kreplin *et al.*, *The Journal of chemical physics*, 2020, **152**, 144107.
- 32 H.-J. Werner, P. J. Knowles, P. Celani, W. Györffy, A. Hesselmann, D. Kats, G. Knizia, A. Köhn, T. Korona, D. Kreplin, R. Lindh, Q. Ma, F. R. Manby, A. Mitrushenkov, G. Rauhut, M. Schütz, K. R. Shamasundar, T. B. Adler, R. D. Amos, S. J. Bennie, A. Bernhardsson, A. Berning, J. A. Black, P. J. Bygrave, R. Cimiraglia, D. L. Cooper, D. Coughtrie, M. J. O. Deegan, A. J. Dobbyn, K. Doll, M. Dornbach, F. Eckert, S. Erfort, E. Goll, C. Hampel, G. Hetzer, J. G. Hill, M. Hodges, T. Hrenar, G. Jansen, C. Köpl, C. Kollmar, S. J. R. Lee, Y. Liu, A. W. Lloyd, R. A. Mata, A. J. May, B. Mussard, S. J. McNicholas, W. Meyer, T. F. Miller III, M. E. Mura, A. Nicklass, D. P. O'Neill, P. Palmieri, D. Peng, K. A. Peterson, K. Pflüger, R. Pitzer, I. Polyak, M. Reiher, J. O. Richardson, J. B. Robinson, B. Schröder, M. Schwilk, T. Shiozaki, M. Sibaev, H. Stoll, A. J. Stone, R. Tarroni, T. Thorsteinsson, J. Toulouse, M. Wang, M. Welborn and B. Ziegler, *MOLPRO*, version 2022.1, a package of *ab initio* programs, see <https://www.molpro.net>.
- 33 Y. Georgievskii, J. A. Miller, M. P. Burke and S. J. Klippenstein, *The Journal of Physical Chemistry A*, 2013, **117**, 12146–12154.
- 34 J. Lizardo-Huerta, B. Sirjean, R. Bounaceur and R. Fournet, *Physical Chemistry Chemical Physics*, 2016, **18**, 12231–12251.
- 35 S. Xu and M.-C. Lin, *Proceedings of the Combustion Institute*, 2007, **31**, 159–166.
- 36 I. Alecu and D. G. Truhlar, *The Journal of Physical Chemistry A*, 2011, **115**, 14599–14611.
- 37 T. J. Lee and P. R. Taylor, *International Journal of Quantum Chemistry*, 1989, **36**, 199–207.
- 38 R. Lindh, *Theoretica chimica acta*, 1993, **85**, 423–440.



Data availability

The data supporting this article have been included as part of the Supplementary Information that contains all theoretical data as inputs of MESS and THEMROT codes, the kinetic mechanism in CHEMKIN and CANTERA formats, and supplementary Figures S1 to S4.