



Cite this: *Phys. Chem. Chem. Phys.*,  
2025, 27, 6810

# Correction: Achieving 9% EQE in light-emitting electrochemical cells via a TADF-sensitized fluorescence strategy


Zeyang Zhou,<sup>ab</sup> Qingda Chang,<sup>ab</sup> Rui Chen,<sup>ab</sup> Pengfei Jin,<sup>ab</sup> Baipeng Yin,<sup>\*a</sup>  
Chuang Zhang<sup>\*a</sup> and Jiannian Yao<sup>ac</sup>

DOI: 10.1039/d5cp90050f

rsc.li/pccp

Correction for 'Achieving 9% EQE in light-emitting electrochemical cells via a TADF-sensitized fluorescence strategy' by Zeyang Zhou et al., *Phys. Chem. Chem. Phys.*, 2024, **26**, 24498–24505, <https://doi.org/10.1039/D4CP02801E>.

The molecular structure of 4CzPN-*t*Bu was displayed incorrectly in Fig. 1 in the originating article; the correct molecule is shown here.



**Fig. 1** (a) Schematics of the operation mechanism of LECs after the formation of the electric double layers (EDLs), p-doped, n-doped, and intrinsic zones. The red and blue arrows represent the energy transfer processes with and without a sensitizer, respectively. (b) Chemical structure of emitter 4CzPN-*t*Bu and sensitizer DMAC-DPS. (c) Cyclic voltammograms of mix-host PVK:OXD-7, sensitizer DMAC-DPS, and emitter 4CzPN-*t*Bu. (d) EL spectrum measured from LECs with and without DMAC-DPS. (e) Current density (A m<sup>-2</sup>)-voltage (V)-luminance (cd m<sup>-2</sup>) curve of LECs with and without DMAC-DPS.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

<sup>a</sup> Key Laboratory of Photochemistry, Beijing National Laboratory for Molecular Sciences, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100190, China.  
E-mail: yinbaipeng@iccas.ac.cn, zhangc@iccas.ac.cn

<sup>b</sup> University of Chinese, Academy of Sciences, Beijing 100049, China

<sup>c</sup> Institute of Molecular Engineering Plus, Fuzhou University, Fuzhou 350108, China

