


Environmental Science Processes & Impacts

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: C. Abdel-Nour, S. Rossignol, B. Samia, M. Bou-Saad, N. K. Dieng, S. Lebarillier, L. Asia, A. Monod and P. WONG-WAH-CHUNG, *Environ. Sci.: Processes Impacts*, 2025, DOI: 10.1039/D5EM00203F.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Hydroxyl radical-initiated degradation kinetics of organic pollutants in surfactant-rich environments

View Article Online
DOI: 10.1039/C5EM00203F

Carole Abdel Nour^a, Stéphanie Rossignol^{*a}, Boulos Samia^a, Maria Bou Saad^a, Ndeye Khoyane Dieng^a, Stéphanie Lebarillier^a, Laurence Asia^a, Anne Monod^a, Pascal Wong-Wah-Chung^a

^aAix Marseille Univ, CNRS, LCE, Marseille, France

^{*}Corresponding author

Keywords: surfactants, hydroxyl radical, kinetic rate constants, organic pollutants, sodium dodecyl sulfate, rhamnolipids, carbamazepine, phenanthrene

Abstract

Surfactant-rich aqueous media are common in natural environments. The sea surface microlayer and the sea spray droplets are good examples and are also frequently markedly enriched in organic pollutants. This study focuses on the degradation kinetics of organic pollutants initiated by the hydroxyl radical in such surfactant-rich environments. The apparent second-order rate constants of the reaction for the hydroxyl radical with two model pollutants, carbamazepine and phenanthrene, were determined in the presence of sodium dodecyl sulfate or rhamnolipids at concentrations below and above their critical micelle concentration (CMC). The results show that the apparent rate constant remains unaffected below the CMC, unless additional reactive processes are induced. From the results obtained above the CMC, second-order rate constants were derived for the reaction of the hydroxyl radical with the pollutant trapped in the surfactant micelles. A rather similar decrease of about a factor of 3 in comparison to pre-micellar conditions was obtained for all three experimental conditions tested. This work appears to provide a suitable foundation for evaluating the impact of such surfactant-rich environments on the half-lives and fates of organic pollutants and a preliminary indication of how reaction kinetics may be modified in various organized surfactant structures.

Environmental Significance Statement

Photo-induced processes represent a significant pathway for the degradation of most organic pollutants. Studying the kinetics of these processes is of paramount importance for the evaluation of the lifetime of these pollutants in aqueous surface and atmospheric compartments. Here, we expended the existing knowledge on this subject by investigating the degradation kinetics of pollutants in surfactant-rich aqueous media. Indeed, organic pollutants are known to be enriched in such media, e.g. in the sea surface microlayer. This work shows that surfactants can increase the pollutants' lifetime against hydroxyl radical attack through both radical quenching and micelle trapping. It provides a quantitative assessment of the decrease in reaction rate constants, enabling this modified kinetics to be taken into account in models.

1. Introduction

Assessing the fate of pollutants in the environment is a crucial task in determining their impacts on ecosystems and human health. The pollutant's half-life and the structures of its degradation products are essential data in this context, contributing to accurate risk assessment by evaluating the pollutant's persistence and the potential adverse effects resulting from its transformation.

However, the chemical processes involved may depend on both the pollutant and the environment, introducing a complexity that is often underestimated in risk assessment. An example of this is the observed slower degradation rate by gas phase hydroxyl radicals of pesticides sorbed onto/into atmospheric particles compared to gas phase pesticides. The difference is significant as this slower rate would explain the unexpected long-range atmospheric transport of some of these compounds.^{1–4} Another example is the inhibiting or amplifying effect of different organic coatings on the kinetics of the heterogeneous reaction of polycyclic aromatic hydrocarbons (PAHs) with ozone at the air-water interface.^{5,6}

In natural surface waters, the fate of organic pollutants is strongly linked to light-initiated degradation processes. This photochemistry is most often studied in the homogeneous dilute phase. Kinetic rate constants are inferred from laboratory photodegradation experiments and are then used to predict the pollutant half-lives in a given body of water, taking into account variations in light flux and concentrations of organic and inorganic matter.^{7–9} An interesting application of this is the recent study by Rosso and Vione¹⁰ in which they modelled half-lives of emerging pollutants in a context of evaporative surface water concentration in arid and semi-arid environments predicting modifications in the involvement of the different degradation pathways (direct photolysis, triplet sensitized reactions, etc.). This highly instructive study nevertheless assumes that reaction rate constants remain unchanged whatever the concentration factor. If this assumption seems justified in the range examined, i.e. a water depth decreasing to one meter, it would appear to be reassessed when approaching a depth of zero meters.

The presence of organic compounds with surface-active properties (of biogenic or anthropogenic origin) could, for instance, lead to the formation of micelles or related organized structures (monolayers, vesicles, ...) in which the organic pollutants can partition, due to hydrophobic interactions.^{11–20} This could potentially modify reaction kinetics and/or preferential pathways.^{18,21–23} This type of organic-rich medium, with a significant fraction of surface-active compounds, is not uncommon in the environment. The sea surface microlayer (SML) for instance is enriched in compounds with surface-active properties.^{24–26} Those biogenic surface-active compounds, such as carbohydrates, proteins, lipids, arise from the water column through adsorption at the surface of air bubbles produced mainly by breaking waves.²⁷ Visible slicks (with at least a monolayer of organic matter at the air-water interface) are stable for wind speeds of up to 4 m.s⁻¹ and a SML can exist up to wind speeds of 6–10 m.s⁻¹. The SML is thus almost ubiquitous.²⁸ Sea spray droplets, emitted into the atmosphere when the plumes of air bubbles generated by breaking waves burst at the ocean surface, are also enriched in organic matter, including compounds with surface-active properties, compared to bulk sea water^{29–32} and individual concentrations can increase upon droplet evaporation and formation of sea spray aerosol (SSA).

It is also important to note that concentrations of organic pollutants are likely to be higher in these organic-rich media than in the more studied diluted phase. Rosso and Vione¹⁰ logically supposed increased organic pollutants concentrations in evaporative waters. Organic pollutants, especially the most hydrophobic ones (PCBs, aliphatic and aromatic hydrocarbons, PFASs, ...), are found to be enriched in the SML^{33–38} with enrichment factors compared to the underlying water up to 950 (value for anthracene in a harbor³⁸), and can include anthropogenic surfactants.^{39,40} Moreover, it has been shown that these pollutants can be transferred in SSA, with enrichment factors compared to bulk sea water increasing as the SSA size decreases, particularly in the submicron range.^{37,38,41,42} In these surfactant-rich environments, organic pollutants can partition into micelles (or related structures) and/or into surfactant monolayers or multilayers at the water-air interface.^{43–45} Kinetic and photochemical pathways may then differ significantly from those observed in the diluted phase.^{22,46}

The aim of this work is thus to investigate the photochemical fate of organic pollutants in surfactant-rich natural environments, such as SML or evaporative water droplets from sea spray. This study focuses on the degradation kinetics of two pollutants with significantly different water solubilities and octanol-water partition coefficient (K_{ow}), carbamazepine and phenanthrene ($\log K_{ow}$ of 2.45^{47–49} and 4.46–4.65^{50,51} respectively). Here, we studied hydroxyl radical-initiated degradation in the presence of either sodium dodecyl sulfate (SDS), a well-characterized anionic surfactant that can also be found in the environment, or rhamnolipids (RLs), a biosurfactant that may better represent surfactants of biogenic origin encountered in environmental samples. Structures of pollutants and surfactants are presented in Figure 1. The apparent rate constants of reaction between the pollutant and hydroxyl radicals were determined following the competition kinetics method, using the reaction of terephthalic acid (TPA) with hydroxyl radicals as the reference.




Figure 1: Structures of (a) carbamazepine (CBZ), (b) phenanthrene (Phe), (c) sodium dodecyl sulfate (SDS), (d) mono-rhamnolipids, and (e) di-rhamnolipids (RLs). The chain length (n+6) of the commercial mixture of mono- and di-rhamnolipids used in this study is unknown but was assumed to be ten carbon atoms (n = 4) for micellar volume calculations (see SI for details).

2. Methods

2.1. Chemicals

Carbamazepine (CBZ, 99% purity), phenanthrene (Phe, 99.5% purity), terephthalic acid (TPA, 98% purity), fluoranthene (Fluo, 98% purity), sodium dodecyl sulfate (SDS, >99% purity), rhamnolipids biosurfactants from *Pseudomonas aeruginosa* (RLs, 90% purity), acetonitrile (HPLC grade, ≥99.9% purity), hexane (≥99.9% purity), dichloromethane (≥99.8% purity, amylene as stabilizer) and dimethylformamide (≥99% purity) were purchased from Sigma-Aldrich (St-Louis, USA) and used without further purification. Non stabilized hydrogen peroxide (H₂O₂, 30 wt% solution in water) was provided by Thermo Fisher Scientific. Potassium hydroxide pellets (KOH, >85%) was purchased from Carlo Erba Reagents (Milano, Italy) and a stock solution was prepared every month at 0.1 M. Stock solutions of Fluo were prepared in hexane at a concentration of 50 mg L⁻¹ in a 100 mL amber glass bottle and stored at 4°C. All aqueous solutions were prepared with ultrapure water (Merck, Direct-Q® 5 UV, 18.2 MΩ cm).

2.2. Determination of the critical micelle concentration of SDS and RLs

View Article Online
DOI: 10.1039/D5EM00203F

The critical micelle concentration (CMC) of SDS was determined using conductivity measurements at room temperature. The conductivity ($\mu\text{S cm}^{-1}$) of a series of SDS solutions with various concentrations was measured using a conductivity meter Multi 340i and a measuring cell TetraCon 325 from WTW®. Conductivity was plotted as a function of surfactant concentration. Since free surfactant molecules have a different molar conductivity to micelle surfactant molecules, the CMC is identified as the point on plot where the slope changes. The CMC of RLs was determined using the pendant drop technique. The surface tension (ST) of RLs solutions was measured with a drop shape analyzer DSA-30 from Krüss. The shape of the drop hanging from a needle is determined by the balance of forces which includes the surface tension of the liquid. The surface tension can be related to the drop shape by the equation of Young-Laplace. The accuracy of the measurements is $\pm 0.1 \text{ mN m}^{-1}$. Prior to each experiment, the instrument was calibrated by measuring the surface tension of pure water. The CMC is determined by measuring the ST of a series of RLs solutions with different concentrations. Above the CMC, ST is independent of the surfactant concentration. ST was plotted as a function of the decimal logarithm of the surfactant concentration. The CMC is identified as the point of intersection between the regression straight line of the linearly dependent region and the straight line passing through the plateau.

2.3. CBZ degradation experiments

The CBZ photodegradation experiments were carried out in a photoreactor equipped with a xenon arc lamp (300 W - LOT Quantum Design, France). The lamp outlet is equipped with a water filter that eliminates infrared radiation ($\lambda > 800 \text{ nm}$) and helps reduce the thermal load. The horizontal beam is deflected at a 90° angle by a dichroic mirror before passing through an atmospheric attenuation filter that removes ultraviolet radiation below 290 nm before reaching the water surface in the reactor. The resulting lamp spectrum is provided in Supplementary Information (SI-Figure S1). The reactor has a volume of approximately 100 mL, and the experiments took place in 50 mL of the reaction mixture. Throughout all experiments, the reactor was hermetically closed with a cap with a quartz window and was equipped with a double wall that allowed water circulation and regulation of the temperature at $(20 \pm 1)^\circ\text{C}$. Homogeneity was ensured by magnetic stirring. Sampling was performed through a side sampling port.

Stock solutions of CBZ and TPA were prepared at $1.6 \times 10^{-4} \text{ M}$ and $4.8 \times 10^{-5} \text{ M}$ respectively using ultrapure water (Merck, Direct-Q® 5 UV, $18.2 \text{ M}\Omega \text{ cm}$) and stored in the dark at 4°C . Working solutions were prepared before each experiment by dilution of stock solutions in ultrapure water to concentrations of $2 \times 10^{-5} \text{ M}$ for each reagent and the pH was adjusted to 8.2 using a 0.1 M KOH solution. This pH was chosen because it is representative of marine environments and because it implies the predominance of the doubly deprotonated form of TPA ($\text{pK}_{\text{a}_{\text{TPA}}} = 3.51$ and 4.82), strongly limits the partitioning of TPA in surfactant micelles. Hydroxyl radicals were produced by H_2O_2 photolysis. The concentration of H_2O_2 and surfactant (SDS or RLs) added in each experiment was estimated to achieve a degradation of about 70-80% of CBZ within 4 hours of irradiation, using a kinetic model based on the known reaction rate constants of CBZ, TPA and SDS towards $\cdot\text{OH}$ radicals. A summary of the initial conditions of all the experiments performed with CBZ is provided in the Supplementary Information (SI-Tables S1 and S2). Aliquots of the reaction mixture were taken from the reactor for CBZ and TPA quantification right before the irradiation was turned on and after 30, 60, 120, 180 and 240 minutes.

2.4. Phe degradation experiments

The Phe photodegradation experiments were carried out using an irradiation chamber equipped with 3 UV fluorescent tubes (CLEO Compact S 25W from ISOLde). The spectral irradiance at the center of the chamber is provided in the Supplementary Information (SI-Figure S2). The chamber is equipped

with a ventilation system to maintain a constant temperature of $(25 \pm 2)^\circ\text{C}$ inside the box throughout the experiments.

The reactor was a glass crystallizer on which a quartz plate was sealed with a Teflon gasket prior each experiment. The crystallizer was filled with 200 mL of the reaction mixture, thereby ensuring that the headspace volume was minimized to limit the volatilization of Phe. Homogeneity was ensured by magnetic stirring.

A stock solution of both Phe and TPA at concentrations of 5.6×10^{-6} and 3×10^{-5} M respectively was prepared in one liter of ultrapure water every two weeks. This solution was left under magnetic stirring for 2 days in the dark at room temperature to maximize the solubilization of phenanthrene. Prior each experiment, 200 mL of the stock solution were filtered into the reactor through a hydrophilic PTFE filter (0.22 μm Clarify-PTFE (hydrophilic) syringe filters from Phenomenex). The pH was then adjusted to 8.2 using a 0.1 M KOH solution. The concentration of H_2O_2 and surfactant (SDS or RLs) added in each experiment was estimated to achieve a degradation of about 70-80% of Phe within 4 hours of irradiation, using a kinetic model based on the known reaction rate constants of Phe, TPA and SDS towards $\cdot\text{OH}$ radical. A summary of the initial conditions of all the experiments performed with Phe is provided in the Supplementary Information (SI-Tables S3 and S4). Aliquots of the reaction mixture were taken from the reactor for Phe and TPA quantification right before the irradiation and after 30, 60, 120, 180 and 240 minutes of exposure.

2.5. Analytical methods

The analyses of CBZ and TPA were carried out by high performance liquid chromatography using a UHPLC PerkinElmer FX 10 equipped with a 330-diode array UV-visible detector (Flexar PDA Plus model) and a 410 automatic injector. The chromatographic separation was carried out on a Brownlee SPP RP-amide column (150 mm x 2.1 mm; 2.7 μm) maintained at 30°C . The volume of reaction mixture injected was 10 μL . Analytes were separated with a mixture of acetonitrile/water with 0.1% formic acid at a flow rate of 0.5 mL min^{-1} using the following gradient: from 5% to 50 % acetonitrile within 20 min and then back to initial conditions within 5 min. TPA and CBZ were detected at $\lambda = 241$ and 285 nm respectively. Under these conditions, their retention times were 8 and 20 min respectively. The data was processed by the Chromera software.

For Phe extraction and concentration (SI-Figure S3), aliquots of 5 mL of the reaction mixture were sampled and Fluo (20 μL of the stock solution), used as an extraction surrogate, was introduced. The aliquots were then extracted 4 times, adding 5 mL of dichloromethane and agitating on a rotary agitator for 7 min at each cycle. The 20 mL of dichloromethane were then evaporated to 1 mL, 100 μL of dimethylformamide were added and the remaining dichloromethane was evaporated. 900 μL of acetonitrile were then added and the 1 mL solutions were filtered onto a 0.22 μm PTFE filter before chromatographic analysis. In the presence of SDS in the reaction mixture, 200 mg of KCl was introduced after Fluo addition and prior to dichloromethane extractions to precipitate SDS and avoid excessive foaming. The global Phe recovery was estimated to be $(95 \pm 4) \%$ (SI-Table S5).

The analysis of Phe and Fluo was carried out using a UHPLC PerkinElmer FX6a coupled to an A-30 fluorescence detector Atlas. An Agilent Chromsep pursuit 5PAH column (250 mm x 4.6 mm; 5 μm) equipped with a precolumn CP 28151 was used and maintained at 30°C . The injected volume was 10 μL . Analytes were separated with a mixture of acetonitrile/water (80:20, v/v) with 0.1% formic acid at a flow rate of 1.2 mL min^{-1} using an isocratic elution mode. Detection of Phe and Fluo were performed at $\lambda_{\text{ex}}/\lambda_{\text{em}}$ of 248/375 and 280/462 nm, respectively. The data was processed by the software Waters Empower®.

2.6. Kinetics data treatment

View Article Online
DOI: 10.1039/D5EM00203F

Apparent second order reaction rate constants between the pollutant and the hydroxyl radicals, $k_{app(pollutant,HO^{\cdot})}$, were determined following the competition kinetics method.⁵² TPA was selected as the reference compound.^{53,54} TPA being doubly deprotonated at pH 8.2 ($pK_{a,TPA} = 3.51$ and 4.82), negligible partitioning into the negatively charged micelles of surfactant is supposed. Its reaction rate constant with hydroxyl radicals, $k_{(TPA,HO^{\cdot})}$, has been reported to be $(4.4 \pm 0.1) \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$.⁵⁴ The apparent second order rate constant for the reaction between the pollutant and hydroxyl radicals, $k_{app(pollutant,HO^{\cdot})}$, was determined using the following equation (see SI for details):

$$\ln \frac{[pollutant]_t}{[pollutant]_0} = \frac{k_{app(pollutant,HO^{\cdot})}}{k_{(TPA,HO^{\cdot})}} \ln \frac{[TPA]_t}{[TPA]_0} \quad \text{Eq. 1}$$

where $[pollutant]_0$ and $[TPA]_0$ are the initial concentrations and $[pollutant]_t$ and $[TPA]_t$ the concentrations remaining in the solution at an irradiation time t . The ratio $k_{app(pollutant,HO^{\cdot})}/k_{(TPA,HO^{\cdot})}$ correspond to the slope of the linear regression obtained by monitoring pollutant and TPA concentrations over time, from which $k_{app(pollutant,HO^{\cdot})}$ was inferred. The uncertainty for each $k_{app(pollutant,HO^{\cdot})}$ value was estimated based on the standard deviation of replicate measurements (SI-Tables S1 to S4) and the uncertainty associated with the $k_{(TPA,HO^{\cdot})}$ value.

2.7. Adaptation of the micellar pseudophase model

The pseudophase model provides a theoretical approach to describe reaction kinetics in heterogeneous systems. In this model^{23,55,56}, the total available volume of micelles of a given surfactant S (V_{Sm}) is assimilated to a conventional reaction medium. Here, as the hydroxyl radical is a transient species, this model was adapted (see SI for details) assuming that the hydroxyl radical is in the aqueous phase and can react either with the pollutant P solubilized in the water phase or with the pollutant entrapped in the micellar pseudophase. The apparent second-order reaction rate constant for the reaction of a pollutant with hydroxyl radicals, $k_{app(pollutant,HO^{\cdot})}$, can then be expressed as a function of the true second-order reaction rate constant of hydroxyl radicals with water solubilized pollutant, $k_{(w-pollutant,HO^{\cdot})}$, the true second-order reaction rate constant of the reaction of hydroxyl radicals with the pollutant molecules trapped into the micelles of surfactant, $k_{(Sm-pollutant,HO^{\cdot})}$, the partition coefficient of the pollutant between the surfactant micelles and water, $K_{mw(pollutant,S)}$, and the total micellar volume, V_{Sm} (equation 2):

$$k_{app(pollutant,HO^{\cdot})} = \frac{k_{(w-pollutant,HO^{\cdot})} + k_{(Sm-pollutant,HO^{\cdot})} \cdot K_{mw(pollutant,S)} \cdot V_{Sm}}{1 + K_{mw(pollutant,S)} \cdot V_{Sm}} \quad \text{Eq. 2}$$

The experimental values of $k_{app(pollutant,HO^{\cdot})}$ were plotted as a function of V_{Sm} and fitted using Eq. 2, allowing the estimation of the three equation parameters $k_{(w-pollutant,HO^{\cdot})}$, $k_{(Sm-pollutant,HO^{\cdot})}$ and $K_{mw(pollutant,S)}$.

3. Results and Discussion

3.1. Determination of the critical micelle concentrations

The CMC of SDS in ultrapure water was found to be 2300 mg L^{-1} ($\approx 8.0 \text{ mM}$) (SI-Figure S4 and SI-Table S6). This value is in the range of those reported in the literature, which generally fall in the $7.9\text{--}8.3 \text{ mM}$ range at 25°C .^{57–60} The presence of CBZ ($2 \times 10^{-5} \text{ M}$), H_2O_2 (1 M) and KOH (for pH adjustment at 8.2) induced no significant change, with a mean CMC value of $2192 \pm 223 \text{ mg L}^{-1}$. This latter value has been used to estimate the total volume of SDS micelles in the framework of the micellar pseudophase model.

RLs surfactants used in this study were shown to reduce the surface tension of water from 72 to 32 mN m⁻¹ at room temperature. The CMC value of the RLs solution in ultrapure water was about 145 mg L⁻¹ (SI-Figure S5). This value has been used to estimate the total volume of RLs micelles in the framework of the micellar pseudophase model. The previously reported CMC values range from 1.6 to 400 mg.L⁻¹.^{61–63} This broad variation in CMCs values is likely due to the fact that RLs are produced by different strains under varying production conditions, leading to different proportions of mono- and di-rhamnolipids. The extraction and purification methods used to isolate rhamnolipids from microbial cultures can also affect the purity and composition of the rhamnolipid samples, potentially leading to differences in reported CMC values.

3.2. Degradation kinetics in the presence of SDS

3.2.1. CBZ degradation kinetics

The photochemical stability of CBZ in ultrapure water at a pH adjusted to 8.2 was validated under our experimental conditions (SI-Figure S6). In a preliminary series of experiments, slower CBZ degradation rates were observed in the presence of increasing concentrations of SDS for a given initial H₂O₂ concentration (SI-Figure S7), reflecting the quenching of ·OH radical by SDS molecules ($k_{(SDS,HO^{\cdot})} = 8.2 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ ⁶⁴). Nevertheless, this simple approach suffers from reproducibility issues, which make the determination reaction rate constants unreliable. To encompass this, the competition kinetic method was used to infer the second order apparent rate constant for the reaction of ·OH radicals with CBZ, $k_{app}(CBZ,HO^{\cdot})$, for each experimental condition. Figure 2(a) shows the results obtained for CBZ in the presence of increasing concentrations of SDS.




Figure 2: Apparent second-order rate constant for the reaction of hydroxyl radicals with (a) CBZ and (b) Phe as a function of SDS concentration. Error bars correspond to the uncertainty estimated based on the standard deviation of replicate measurements and the uncertainty associated with the $k_{(TPA,HO^{\cdot})}$ value. Vertical dashed line indicates the CMC of SDS.

The rate constant value obtained in the absence of surfactant, $(7.39 \pm 0.24) \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ is within the range of previously reported values, which vary from 2.05 to $18 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$.^{65–67} Discrepancies may arise from temperature, pH dependencies and the determination method employed. The presence of SDS at concentrations below the CMC does not significantly affect the value of the apparent reaction rate constant, indicating that SDS monomers can inhibit the degradation of CBZ but only by acting as a hydroxyl radical quencher. Conversely, above the CMC, a significant decrease in the value of the apparent reaction rate constant is observed. This decrease is not linear and tends to reach a plateau at high SDS concentrations. According to previous studies, the neutral CBZ molecules are assumed to partition into the apolar SDS micelle core.^{68,69} The total volume of the SDS micelles, V_{SDSm} , was estimated assuming a micellar radius of 1.84 nm and a number of monomers per micelle of 64⁷⁰ (see SI for details). V_{SDSm} increases linearly with the SDS concentration above the CMC (Figure S8(a)).

The experimental values of $k_{\text{app}}(\text{CBZ}, \text{HO}^\bullet)$ were plotted as a function of V_{SDSm} and fitted using Eq. 2. (Figure 3(a)). The resulting estimated values of the three equation parameters $k_{(\text{w-CBZ}, \text{HO}^\bullet)}$, $k_{(\text{SDSm-CBZ}, \text{HO}^\bullet)}$ and $K_{\text{mw}}(\text{CBZ}, \text{SDS})$ were $(7.25 \pm 0.54) \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$, $(1.88 \pm 0.42) \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ and 489 ± 171 , respectively (Table 1). CBZ shielding by SDS micelles thus results in a 74 % decrease of the second-order rate constant of its reaction with hydroxyl radicals. This reaction rate constant decrease leads to an increase in the half-life of CBZ towards HO^\bullet reaction by a factor of 3.8. The estimated value of $\log K_{\text{mw}}(\text{CBZ}, \text{SDS})$ is 2.69, with an uncertainty range of 2.50 to 2.82. This value is within the range of the published CBZ octanol-water partition coefficient ($\log K_{\text{ow}, \text{CBZ}}$), which varies from 1.51 to 2.77 for calculated values, with a commonly used experimental value of 2.45.^{47–49} Using this $\log K_{\text{mw}}(\text{CBZ}, \text{SDS})$ value, the molar fraction of CBZ trapped in SDS micelles was inferred (see SI for details) and plotted as a function of SDS concentration (Figure 3(b)). This figure shows that, even for a fairly water-soluble compound such as CBZ, the molar fraction of molecules trapped in SDS micelles rapidly becomes non-negligible above the CMC of SDS, reaching a value of 0.5 at twice the CMC (corresponding to a V_{SDSm} of 2 ml L^{-1}).




Figure 3: (a) Apparent second-order reaction rate constant measured for the reaction of hydroxyl radicals with CBZ as a function of estimated total SDS micelle volume fitted with Eq. 2. (b) Corresponding estimated molar fraction of CBZ trapped into SDS micelles as a function of estimated total SDS micelle volume.

3.2.2. Phe degradation kinetics

The experimental conditions used for Phe degradation experiments were determined so that the direct photolysis as well as the volatilization of Phe can be considered negligible (SI-Figure S9). The evolution of the measured apparent rate constant of the reaction between hydroxyl radicals and Phe as a function of SDS concentration is shown on Figure 2(b). The reaction rate constant value obtained without SDS, $(1.22 \pm 0.16) \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$, is in good agreement with the lowest ones previously reported that range from 1.24×10^{10} to $2.3 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$.^{71,72}

In contrast to the results obtained for CBZ, the apparent second-order reaction rate constant for Phe increased with SDS concentration below the CMC and reached a maximum of $(1.31 \pm 0.36) \times 10^{11} \text{ M}^{-1} \text{ s}^{-1}$ close to the CMC. This increase causes the rate constant to exceed the classical diffusion-controlled limit value observed for bimolecular reactions in the aqueous phase at ambient temperature, which is around $10^{10} \text{ M}^{-1} \text{ s}^{-1}$.⁷³ The diffusion-controlled limit is due to molecular agitation in a medium at a certain temperature, and to the fact that the reaction cannot occur faster than the reactive molecules colliding. Molecular diffusion depends on the medium properties, including viscosity, the size of the reactants, and temperature. Here, the diffusion limited rate constant was determined to be approximately $8.5 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ (see SI for calculation details). The reaction of Phe with hydroxyl radicals in the absence of surfactant is thus probably limited by diffusion. Since the actual rate constant of the reaction cannot exceed the diffusion rate constant, parallel processes must be involved in the presence of SDS. Enhanced direct photolysis is excluded as Phe stability was validated in the presence of SDS (SI-Figure S9). This increase in the apparent second-order reaction rate constant corresponds to a decrease in the Phe half-life by a factor of almost 11. Complementary experiments carried out in the presence of RLs instead of SDS at concentrations below the CMC show no significant influence on the apparent rate constant (SI-Figure S10). This suggests the existence of specific side processes resulting from the simultaneous presence of both Phe and SDS, as discussed in section 3.4.

Figure 2b shows that above the CMC, as for CBZ, the apparent reaction rate constant of Phe decreases. Figure 4(a) shows the plot of the apparent reaction rate constant values as a function of total SDS micelle volume in the SDS concentration range of 2000-7000 mg L⁻¹, fitted with Eq. 2. The resulting estimated values of the three equation parameters $k_{(w-Phe,HO^{\bullet})}$, $k_{(SDSm-Phe,HO^{\bullet})}$ and $K_{mw(Phe,SDS)}$ were $(1.31 \pm 0.20) \times 10^{11} \text{ M}^{-1} \text{ s}^{-1}$, $(4.17 \pm 0.40) \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$ and $(1.1 \pm 10.9) \times 10^5$, respectively (Table 1). The $k_{(w-Phe,HO^{\bullet})}$ value is irrelevant as it corresponds to the apparent reaction rate constant for water solubilized Phe in the presence of SDS concentration just below the CMC, thus including side degradation processes (discussed in section 3.4). This fit also suffers from a large uncertainty in the $K_{SDSm-w,Phe}$ value, probably due to a lack of data in the 0-0.9 range for the mole fraction of Phe trapped in SDS micelles (Figure 4(b)), making it statistically insignificant. Nevertheless, the corresponding estimated log $K_{mw(Phe,SDS)}$ value, 5.06, is only slightly above the values reported in the literature for the log K_{ow} of Phe, which ranges from 4.46 to 4.65.^{50,51} The decrease in the apparent reaction rate constant observed above the CMC of SDS is consistent with the results obtained for CBZ, i.e. with a shielding effect of micelle trapping. The estimated value of the second-order rate constant for the reaction of hydroxyl radicals with Phe molecules trapped in SDS micelles, $(4.17 \pm 0.40) \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$, corresponds to a 68% decrease compared to the pre-CMC value. It is thus of the same order as that observed for CBZ (74%). Nevertheless, this $k_{(SDSm-Phe,HO^{\bullet})}$ value remains significantly higher than the reaction rate constant obtained without any surfactant addition $((1.22 \pm 0.16) \times 10^{10} \text{ M}^{-1} \text{ s}^{-1})$. This suggests that, if micelle trapping effectively protects Phe from the hydroxyl radical attack, this effect does not fully offset the

increase in reactivity induced by free SDS monomers and does not significantly affect degradation processes occurring in parallel with the hydroxyl radical attack.




Figure 4: (a) Apparent second-order reaction rate constant measured for the reaction of hydroxyl radicals with Phe as a function of estimated total SDS micelle volume fitted with Eq. 2, for SDS concentrations in the range of 2000 - 7000 mg L⁻¹. (b) Corresponding estimated molar fraction of Phe trapped into SDS micelles as a function of estimated total SDS micelle volume.

3.3. Degradation kinetics of CBZ in the presence of RLs

Figure 5 shows the evolution of the experimental values of the apparent second-order rate constant of the reaction of CBZ with hydroxyl radicals in the presence of RLs. The general trend appears to be quite similar to that obtained with SDS, with a significant decrease in the apparent reaction rate constant at RLs concentrations above the CMC. For RLs concentrations below the CMC, the rate constant value determined in the absence of RLs ($(7.39 \pm 0.24) \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$) is higher than the values obtained at 10 and 80 mg L⁻¹ of RLs ($(6.77 \pm 0.26) \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ and $(6.80 \pm 0.16) \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ respectively), while these two latter are not significantly different. Whereas SDS clearly acts only as a quencher of hydroxyl radicals in the pre-micellar region, these results could suggest interactions of RLs with CBZ under similar conditions. Another explanation could be that RLs start to form aggregates at concentrations below the determined CMC, which was established by surface tension measurements. In any case, the dataset is too small to support any firm conclusion and the effect remains very limited compared to the decrease in the rate constant above the CMC. In this post-micellar region, CBZ behaves similarly as with SDS, partitioning into micelles with corresponding shielding effects. At pH 8.2, RLs are in their deprotonated form (pKa ranging from 4.28 to 5.50 depending on concentration⁷⁴) and are assumed to form micelles only, with negligible presence of larger structures such as of lamellae or vesicles.^{75,76} The total volume of RLs micelles, V_{RLSm} , was approximated using an estimated mean molecular weight of 577 g mol⁻¹, a micellar radius of 2.4 nm and a number of monomers per micelle of 25^{15,76} (see SI for details). The total volume of RLs micelles, V_{RLSm} , increases linearly with RLs

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

concentration above the CMC (SI-Figure 9(b)). The experimental values of $k_{app}(CBZ,HO^{\bullet})$ were plotted as a function of V_{RLsm} and fitted using Eq. 2. (Figure 6(a)). The resulting estimated values of the three equation parameters $k_{(w-CBZ,HO^{\bullet})}$, $k_{(RLsm-CBZ,HO^{\bullet})}$ and $K_{mw}(CBZ,RLs)$ were $(6.86 \pm 0.24) \times 10^9 M^{-1} s^{-1}$, $(2.67 \pm 0.42) \times 10^9 M^{-1} s^{-1}$ and 2037 ± 1162 , respectively (Table 1). CBZ shielding by RLs micelles thus results in a 61 % decrease of the second-order rate constant of its reaction with hydroxyl radicals. It is interesting to note that the rate constant values obtained for CBZ trapped in micelles of SDS and RLs are not significantly different. The estimated value of $\log K_{mw}(CBZ,RLs)$ is 3.31, with an uncertainty range of 2.94 to 3.51. This value is slightly higher than both the value obtained for SDS micelles and the $\log K_{ow}$ of CBZ (2.69 and 1.51-2.77⁴⁷⁻⁴⁹ respectively). This may indicate a higher propensity of RLs micelles to trap CBZ compared to SDS ones. Using this $\log K_{mw}(CBZ,RLs)$ value, the molar fraction of CBZ trapped in RLs micelles was plotted as a function of RLs concentration (Figure 6(b)). As with SDS, the molar fraction of CBZ molecules trapped in RLs micelles rapidly becomes non-negligible as the concentration of RLs increases. It should be pointed out that the value determined here as the CMC of RLs ($145 mg L^{-1}$) is similar to those obtained in most studies^{15,63}, but has previously been shown to actually represent the critical aggregation concentration (CAC) above which only premicellar aggregates are formed, the true CMC occurring at a higher RLs concentration ($7.5 mM$ for monorhamnolipids, equivalent to $3.8 g L^{-1}$).⁷⁶ Molecular dynamic simulations performed in this CAC-CMC range by Eismín *et al.*⁷⁶ predict loosely packed premicelles into which water can significantly penetrate. This suggests that fully formed RLs micelles could have a higher shielding effect than the one quantified here. Nevertheless, our value seems relevant for environmental conditions with surfactant concentrations just above CMC.




Figure 5: Apparent second-order rate constant for the reaction of hydroxyl radicals with CBZ as a function of RLs concentration. Error bars correspond to the uncertainty estimated based on the standard deviation of replicate measurements and the uncertainty associated with the $k_{(TPA,HO^{\bullet})}$ value. Vertical dashed line indicates the CMC of RLs.




Figure 6: (a) Apparent second-order reaction rate constant measured for the reaction of hydroxyl radicals with CBZ as a function of estimated total RLS micelle volume fitted with Eq. 2. (b) Corresponding estimated molar fraction of CBZ trapped into RLS micelles as a function of estimated total RLS micelle volume.

3.4. Discussion

3.4.1. Influence of surfactants below the CMC

The presence of SDS or RLS concentrations below the CMC induces no significant modification in the second-order rate constant of the reaction of hydroxyl radical with the two model pollutants chosen, CBZ or Phe, except in the case of Phe in the presence of SDS. This suggests that, in many cases, solubilized surfactants monomers can reduce the half-life of organic pollutants relative to $^{\bullet}OH$ reactions by acting solely as $^{\bullet}OH$ radical quenchers. This effect alone can have a dramatic influence on the half-life of pollutants in organic-rich environments.¹⁰ However, the notable ~11-fold increase in the apparent second-order rate constant of the reaction of Phe with increasing SDS concentrations below the CMC suggests that the situation can sometimes be more complex, requiring consideration of side processes. Here, the hypothesis that additional reactive species are produced during SDS degradation must be considered. Degradation of SDS molecules initiated by hydroxyl radicals has been shown to lead to the formation of peroxy (ROO^{\bullet}) and sulfate ($SO_4^{\bullet-}$) radicals,⁷⁷ both being able to react with organics. Given the second-order rate constant values determined in this study for the reaction of hydroxyl radicals with CBZ and Phe and those from previous studies for the reaction of sulfate radicals with TPA, CBZ and Phe, the expected $k_{HO^{\bullet}} / k_{SO_4^{\bullet-}}$ ratio (SI-Table S7) is only slightly higher for Phe than for CBZ (ratios ranging from 4 to 9 and from 3 to 4, respectively), while it is significantly higher for TPA (ratio of 26).^{65,78–81} Compared to TPA, both CBZ and Phe should therefore appear affected by the formation of sulfate radicals, albeit to a slightly lesser extent in the case of CBZ. The fact that the reaction rate constant for CBZ appears unaffected by the hypothetical formation of sulfate radicals,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

suggests that additional side processes contribute to the increasing rate constants observed for Phe with SDS concentration.

Table 1: Second-order rate constants previously reported and determined in this study for the reaction of CBZ and Phe with hydroxyl radicals, octanol-water partitioning coefficients previously reported and surfactant micelles-water partitioning coefficients determined in this study.

HO [•] reaction second order rate constant (M ⁻¹ s ⁻¹)	CBZ		Phe	
	Free pollutant - Previous studies	3.1 to 18 × 10 ⁹ ^{65,66}	1.24 to 2.3 × 10 ¹⁰ ^{71,72}	
	Free pollutant without surfactants - This study ^a	(7.39 ± 0.24) × 10 ⁹ ^a	(1.22 ± 0.16) × 10 ¹⁰ ^a	
	Free pollutant with SDS below CMC - This study	(7.25 ± 0.54) × 10 ⁹	(1.31 ± 0.20) × 10 ¹¹ ^b	
	Free pollutant with RLs below CMC - This study	(6.86 ± 0.24) × 10 ⁹	(1.13 ± 0.23) × 10 ¹⁰ ^c	
	Pollutant trapped in SDS micelles - This study	(1.88 ± 0.42) × 10 ⁹	(4.17 ± 0.40) × 10 ¹⁰ ^b	
	Pollutant trapped in RLs micelles - This study	(2.67 ± 0.42) × 10 ⁹	nd	
K _{ow} (log K _{ow}) - Previous studies		32 - 589 (1.51 - 2.77) ⁴⁷⁻⁴⁹	2.88 × 10 ⁴ - 4.47 × 10 ⁴ (4.46 - 4.65) ^{50,51}	
K _{mw(SDS)} (log K _{mw(SDS)}) - This study		489 ± 171 (2.69)	(1.1 ± 10.9) × 10 ⁵ (5.06)	
K _{mw(RLs)} (log K _{mw(RLs)}) - This study		2037 ± 1162 (3.31)	nd	

^a Determined in ultrapure water at pH 8.2 adjusted with the 1 M KOH solution
^b These values involve side processes, see text for details
^c This value represents the arithmetic mean of all measurements obtained below the RLs' CMC in the series of experiments presented in SI-Figure S1 with the corresponding standard deviation
nd: not determined

3.4.2. Influence of surfactants above the CMC

Above the CMC, a similar trend is observed in the evolution of the apparent second-order rate constant for the three experimental conditions tested. The partitioning of the pollutant in the micelles of surfactant decreases its apparent reaction rate constant with hydroxyl radicals, according to previous studies.^{82,83} Here, the results show that this decrease reaches a plateau when the micellar volume becomes sufficient to tend to trap all the available molecules of pollutant. Interestingly, the magnitude of this decrease is quite similar for both model pollutants and model surfactants, with a reduction by a factor of about 3 (3.9, 3.1 and 2.6 for CBZ with SDS, Phe with SDS and CBZ with RLs respectively). A similar extend was found by Bansal et al.⁸³ for the degradation of benzene initiated by H₂O₂ photolysis in solutions of three different surfactants, with second-order reaction rate constants with hydroxyl radicals decreasing from 8.2 to 2.6-3.0 × 10⁹ M⁻¹ s⁻¹. SDS has also been shown by Faust and Abbatt⁸⁴ to slow down the rate constant of the heterogeneous reactions of a highly hydrophilic compound (tricarballic acid) solubilized in aqueous droplets with gas phase hydroxyl radicals. Despite the heterogeneous nature of the chemistry involved in this case, there again the decrease in the reaction rate was found to fall within the same order of magnitude, with a decrease of about 60% in the presence of a SDS monolayer at the water surface.⁸⁴ The consistency of all these results provides an interesting starting point for considering such modified kinetics in the assessment of the fate of organic pollutants in surfactant-rich environments.

In addition, the results presented here indicate that the generally easily available K_{ow} value could be employed to assess the molar fraction of a pollutant that partitions into a given volume of micelles, even for relatively water-soluble compounds. A limitation may be the required absence of ionic or other interactions between the hydrophilic head group of the surfactant and the pollutant, as it was the case in this study with nonionic pollutants and anionic surfactants. For hydrophobic compounds, some K_{ow}-K_{mw} relationships can be found in the literature.^{16,85}

These findings may also be used as a preliminary indication of how reaction kinetics may be modified in other surfactant-organized structures, including surfactant mono- or multilayers at water-air interfaces (e.g. SML or atmospheric droplets) and vesicles and lamellae for media with higher surfactant concentrations.

4. Conclusion

View Article Online
DOI: 10.1039/D5EM00203F

This work was focused on the determination of the kinetics of the reaction of the hydroxyl radical with pollutants in surfactant rich media. It was shown that, in the absence of side processes, the presence of monomers of the two tested surfactants, SDS (a well-known anionic surfactant) and RLs (a biogenic surfactant), at concentrations below the CMC does not significantly affect the value of the second-order reaction rate constant. In these concentration ranges, the pollutant degradation can nevertheless be slowed down by the quenching of hydroxyl radical by the surfactant monomers. In some cases, side processes can also be involved and affect the degradation kinetics at sub-CMC concentrations. Here, SDS degradation was for instance suspected to produce additional reactive species able to accelerate the degradation kinetics of Phe. Above the CMC, partitioning of the pollutant into surfactant micelles induces the decrease of the reaction rate constant, shielding it from the hydroxyl radical attack. Values of these second-order reaction rate constants were determined for pollutant molecules trapped into micelles and shown to be a factor of about three times lower compared to those obtained in premicellar condition, in good agreement with previous studies. Estimated K_{mw} values are in good agreement with K_{ow} values for both tested pollutants, with little impact of the nature of the surfactant for CBZ.

Organic pollutants are known to be enriched in surfactant-rich environments such as the SML, atmospheric aerosols and fog droplets or even in the case of evaporative waters. This work appears to provide a suitable foundation for evaluating the impact of such surfactant-rich environments on the half-life and fate of these pollutants and an interesting indication of how reaction kinetics may be modified in various organized surfactant structures. Future studies may wish to explore the influence of surfactants on direct photolysis and degradation processes initiated by other reactive species, such as the triplet state of organic matter or singlet oxygen. Another emerging question is whether the reaction pathways and the structure of the degradation products formed in these environments are modified.

Data availability

Raw data files can be obtained contacting the corresponding author.

Conflicts of interest

There are no conflicts to declare.

Acknowledgments

The authors gratefully acknowledge the financial support of the French National Research Agency (ANR) through the project PhotoSeaSMiL.

References

- 1 J. Socorro, A. Durand, B. Temime-Roussel, S. Gligorovski, H. Wortham and E. Quivet, The persistence of pesticides in atmospheric particulate phase: An emerging air quality issue, *Sci. Rep.*, 2016, **6**, 33456.
- 2 J. Socorro, P. S. J. Lakey, L. Han, T. Berkemeier, G. Lammel, C. Zetzsch, U. Pöschl and M. Shiraiwa, Heterogeneous OH Oxidation, Shielding Effects, and Implications for the Atmospheric Fate of Terbutylazine and Other Pesticides, *Environ. Sci. Technol.*, 2017, **51**, 13749–13754.

- 3 M. Shrivastava, S. Lou, A. Zelenyuk, R. C. Easter, R. A. Corley, B. D. Thrall, P. J. Rasch, J. D. Fast, S. L. Massey Simonich, H. Shen and S. Tao, Global long-range transport and lung cancer risk from polycyclic aromatic hydrocarbons shielded by coatings of organic aerosol, *Proc. Natl. Acad. Sci.*, 2017, **114**, 1246–1251.
- 4 C. Mattei, H. Wortham and E. Quivet, Heterogeneous atmospheric degradation of pesticides by ozone: Influence of relative humidity and particle type, *Sci. Total Environ.*, 2018, **625**, 1544–1553.
- 5 B. T. Mmereki, D. J. Donaldson, J. B. Gilman, T. L. Eliason and V. Vaida, Kinetics and products of the reaction of gas-phase ozone with anthracene adsorbed at the air–aqueous interface, *Atmos. Environ.*, 2004, **38**, 6091–6103.
- 6 E. A. Henderson and D. J. Donaldson, Influence of Organic Coatings on Pyrene Ozonolysis at the Air–Aqueous Interface, *J. Phys. Chem. A*, 2012, **116**, 423–429.
- 7 C. Zhou, J. Chen, H. Xie, Y. Zhang, Y. Li, Y. Wang, Q. Xie and S. Zhang, Modeling photodegradation kinetics of organic micropollutants in water bodies: A case of the Yellow River estuary, *J. Hazard. Mater.*, 2018, **349**, 60–67.
- 8 M. Bodrato and D. Vione, APEX (Aqueous Photochemistry of Environmentally occurring Xenobiotics): a free software tool to predict the kinetics of photochemical processes in surface waters, *Environ. Sci. Process. Impacts*, 2014, **16**, 732–740.
- 9 L. S. Albi, Z. Guo, G. Chen and C. Yoshimura, Potential effect of atmospheric condition on incident light and photo-production of reactive intermediates in freshwater systems, *Environ. Adv.*, 2023, **11**, 100346.
- 10 A. Rosso and D. Vione, Pollutant Photodegradation Affected by Evaporative Water Concentration in a Climate Change Scenario, *Molecules*, 2024, **29**, 2655.
- 11 J. W. Park and P. R. Jaffe, Partitioning of three nonionic organic compounds between adsorbed surfactants, micelles, and water, *Environ. Sci. Technol.*, 1993, **27**, 2559–2565.
- 12 J. C. Mata-Sandoval, J. Karns and A. Torrents, Effect of Rhamnolipids Produced by *Pseudomonas aeruginosa* UG2 on the Solubilization of Pesticides, *Environ. Sci. Technol.*, 2000, **34**, 4923–4930.
- 13 C. N. Mulligan, Environmental applications for biosurfactants, *Environ. Pollut.*, 2005, **133**, 183–198.
- 14 J. A. Marqusee and K. A. Dill, Solute partitioning into chain molecule interphases: Monolayers, bilayer membranes, and micelles, *J. Chem. Phys.*, 1986, **85**, 434–444.
- 15 V. P. Arkhipov, R. V. Arkhipov, E. V. Petrova and A. Filippov, Micellar and solubilizing properties of rhamnolipids, *Magn. Reson. Chem.*, 2023, **61**, 345–355.
- 16 K. T. Valsaraj and L. J. Thibodeaux, Relationships between micelle-water and octanol-water partition constants for hydrophobic organics of environmental interest, *Water Res.*, 1989, **23**, 183–189.
- 17 L. Zhu and S. Feng, Synergistic solubilization of polycyclic aromatic hydrocarbons by mixed anionic–nonionic surfactants, *Chemosphere*, 2003, **53**, 459–467.
- 18 B. Samiey, C.-H. Cheng and J. Wu, Effects of Surfactants on the Rate of Chemical Reactions, *J. Chem.*, 2014, **2014**, e908476.
- 19 G. La Sorella, G. Strukul and A. Scarso, Recent advances in catalysis in micellar media, *Green Chem.*, 2015, **17**, 644–683.
- 20 D. Yordanova, E. Ritter, T. Gerlach, J. H. Jensen, I. Smirnova and S. Jakobtorweihen, Solute Partitioning in Micelles: Combining Molecular Dynamics Simulations, COSMOmic, and Experiments, *J. Phys. Chem. B*, 2017, **121**, 5794–5809.
- 21 S. M. Thomas and S. K. Harrison, Surfactant-Altered Rates of Chlorimuron and Metsulfuron Photolysis in Sunlight, *Weed Sci.*, 1990, **38**, 602–606.

- 22 K. Kalyanasundaram, *Photochemistry in Microheterogeneous Systems*, Elsevier, 2012. View Article Online
DOI: 10.1039/D5EM00203F
- 23 C. A. Bunton and G. Savelli, in *Advances in Physical Organic Chemistry*, eds. V. Gold and D. Bethell, Academic Press, 1986, vol. 22, pp. 213–309.
- 24 O. Wurl, E. Wurl, L. Miller, K. Johnson and S. Vagle, Formation and global distribution of sea-surface microlayers, *Biogeosciences*, 2011, **8**, 121–135.
- 25 B. Sabbaghzadeh, R. C. Upstill-Goddard, R. Beale, R. Pereira and P. D. Nightingale, The Atlantic Ocean surface microlayer from 50°N to 50°S is ubiquitously enriched in surfactants at wind speeds up to 13 m s⁻¹, *Geophys. Res. Lett.*, 2017, **44**, 2017GL072988.
- 26 B. Čosović, in *The Mediterranean Sea*, Springer, Berlin, Heidelberg, 2005, pp. 269–296.
- 27 O. Wurl, L. Miller, R. Röttgers and S. Vagle, The distribution and fate of surface-active substances in the sea-surface microlayer and water column, *Mar. Chem.*, 2009, **115**, 1–9.
- 28 M. Cunliffe, A. Engel, S. Frka, B. Gašparović, C. Guitart, J. C. Murrell, M. Salter, C. Stolle, R. Upstill-Goddard and O. Wurl, Sea surface microlayers: A unified physicochemical and biological perspective of the air–ocean interface, *Prog. Oceanogr.*, 2013, **109**, 104–116.
- 29 P. Schmitt-Kopplin, G. Liger-Belair, B. P. Koch, R. Flerus, G. Kattner, M. Harir, B. Kanawati, M. Lucio, D. Tziotis, N. Hertkorn and I. Gebefügi, Dissolved organic matter in sea spray: a transfer study from marine surface water to aerosols, *Biogeosciences*, 2012, **9**, 1571–1582.
- 30 Y. Song, J. Li, N. T. Tsona, L. Liu and L. Du, Enrichment of short-chain organic acids transferred to submicron sea spray aerosols, *Sci. Total Environ.*, 2022, **851**, 158122.
- 31 N. Radoman, Department of Materials and Environmental Chemistry, Stockholm University, 2022.
- 32 G. de Leeuw, E. L. Andreas, M. D. Anguelova, C. W. Fairall, E. R. Lewis, C. O'Dowd, M. Schulz and S. E. Schwartz, Production flux of sea spray aerosol, *Rev. Geophys.*, DOI:10.1029/2010RG000349.
- 33 C. Guigue, M. Tedetti, S. Giorgi and M. Goutx, Occurrence and distribution of hydrocarbons in the surface microlayer and subsurface water from the urban coastal marine area off Marseilles, Northwestern Mediterranean Sea, *Mar. Pollut. Bull.*, 2011, **62**, 2741–2752.
- 34 O. Wurl and J. P. Obbard, A review of pollutants in the sea-surface microlayer (SML): a unique habitat for marine organisms, *Mar. Pollut. Bull.*, 2004, **48**, 1016–1030.
- 35 A. Cincinelli, A. M. Stortini, L. Checchini, T. Martellini, M. D. Bubba and L. Lepri, Enrichment of organic pollutants in the sea surface microlayer (SML) at Terra Nova Bay, Antarctica: influence of SML on superficial snow composition, *J. Environ. Monit.*, 2005, **7**, 1305–1312.
- 36 C. Guitart, N. García-Flor, J. C. Miquel, S. W. Fowler and J. Albaigés, Effect of the accumulation of polycyclic aromatic hydrocarbons in the sea surface microlayer on their coastal air–sea exchanges, *J. Mar. Syst.*, 2010, **79**, 210–217.
- 37 G. Casas, A. Martínez-Varela, J. L. Roscales, M. Vila-Costa, J. Dachs and B. Jiménez, Enrichment of perfluoroalkyl substances in the sea-surface microlayer and sea-spray aerosols in the Southern Ocean, *Environ. Pollut.*, 2020, **267**, 115512.
- 38 A. Cincinelli, A. M. Stortini, M. Perugini, L. Checchini and L. Lepri, Organic pollutants in sea-surface microlayer and aerosol in the coastal environment of Leghorn—(Tyrrhenian Sea), *Mar. Chem.*, 2001, **76**, 77–98.
- 39 R. Uning, M. T. Latif, K. L. Yu, S. Y. Cheng, F. Ahamad, M. F. Khan, E. A. Bedurus and S. Suratman, Surfactants in the Sea Surface Microlayer, Underlying Water and Atmospheric Particles of Tropical Coastal Ecosystems, *Water. Air. Soil Pollut.*, 2018, **229**, 305.

- 40 S. Shaharom, M. T. Latif, M. F. Khan, S. N. M. Yusof, N. A. Sulong, N. B. A. Wahid, R. Uning and S. Suratman, Surfactants in the sea surface microlayer, subsurface water and fine marine aerosols in different background coastal areas, *Environ. Sci. Pollut. Res.*, 2018, **25**, 27074–27089. View Article Online
DOI: 10.1039/D5EM00203F
- 41 F. Pawlak, K. Koziol, M. Frankowski, Ł. Nowicki, C. Marlin, A. M. Sulej-Suchomska and Ż. Polkowska, Sea spray as a secondary source of chlorinated persistent organic pollutants? - Conclusions from a comparison of seven fresh snowfall events in 2019 and 2021, *Sci. Total Environ.*, 2023, **891**, 164357.
- 42 J. H. Johansson, M. E. Salter, J. C. A. Navarro, C. Leck, E. D. Nilsson and I. T. Cousins, Global transport of perfluoroalkyl acids via sea spray aerosol, *Environ. Sci. Process. Impacts*, 2019, **21**, 635–649.
- 43 B. T. Mmereki and D. J. Donaldson, Laser induced fluorescence of pyrene at an organic coated air–water interface, *Phys. Chem. Chem. Phys.*, 2002, **4**, 4186–4191.
- 44 B. T. Mmereki, S. R. Chaudhuri and D. J. Donaldson, Enhanced Uptake of PAHs by Organic-Coated Aqueous Surfaces, *J. Phys. Chem. A*, 2003, **107**, 2264–2269.
- 45 E. M. McLay, C. A. Nour, Y. Y. Huang, Z. M. Golay, P. Wong-Wah-Chung, S. Rossignol and D. J. Donaldson, Experimental determination of the partitioning of representative organic pollutants to the air–water interface, *Environ. Sci. Process. Impacts*, 2024, **26**, 510–518.
- 46 Y.-J. An and E. R. Carraway, PAH degradation by UV/H₂O₂ in perfluorinated surfactant solutions, *Water Res.*, 2002, **36**, 309–314.
- 47 T. Scheytt, P. Mersmann, R. Lindstädt and T. Heberer, 1-Octanol/Water Partition Coefficients of 5 Pharmaceuticals from Human Medical Care: Carbamazepine, Clofibric Acid, Diclofenac, Ibuprofen, and Propyphenazone, *Water. Air. Soil Pollut.*, 2005, **165**, 3–11.
- 48 A. Pyca, M. Babuska and M. Zachariasz, A comparison of theoretical methods of calculation of partition coefficients for selected drugs, *Acta Pol. Pharm. Ń Drug Res.*, 2006, **63**, 159–167.
- 49 S. Lee, S.-I. Kang, J.-L. Lim, Y. J. Huh, K.-S. Kim and J. Cho, Evaluating controllability of pharmaceuticals and metabolites in biologically engineered processes, using corresponding octanol–water distribution coefficient, *Ecol. Eng.*, 2011, **37**, 1595–1600.
- 50 P. G.-J. de Maagd, D. Th. E. M. ten Hulscher, H. van den Heuvel, A. Opperhuizen and D. T. H. M. Sijm, Physicochemical properties of polycyclic aromatic hydrocarbons: Aqueous solubilities, n-octanol/water partition coefficients, and Henry's law constants, *Environ. Toxicol. Chem.*, 1998, **17**, 251–257.
- 51 F. Alves de Lima Ribeiro and M. M. C. Ferreira, QSPR models of boiling point, octanol–water partition coefficient and retention time index of polycyclic aromatic hydrocarbons, *J. Mol. Struct. THEOCHEM*, 2003, **663**, 109–126.
- 52 A. Monod, L. Poulain, S. Grubert, D. Voisin and H. Wortham, Kinetics of OH-initiated oxidation of oxygenated organic compounds in the aqueous phase: new rate constants, structure–activity relationships and atmospheric implications, *Atmos. Environ.*, 2005, **39**, 7667–7688.
- 53 T. Charbouillot, M. Brigante, G. Mailhot, P. R. Maddigapu, C. Minero and D. Vione, Performance and selectivity of the terephthalic acid probe for OH as a function of temperature, pH and composition of atmospherically relevant aqueous media, *J. Photochem. Photobiol. Chem.*, 2011, **222**, 70–76.
- 54 S. Page, W. Arnold and K. McNeill, Terephthalate as a probe for photochemically generated hydroxyl radical, *J. Environ. Monit. JEM*, 2010, **12**, 1658–65.
- 55 A. Cid, A. Acuña, M. Alonso-Ferrer, G. Astray, L. García-Río, J. Simal-Gándara and J. C. Mejuto, in *Microemulsion - a Chemical Nanoreactor*, IntechOpen, 2019.
- 56 F. M. Menger and C. E. Portnoy, Chemistry of reactions proceeding inside molecular aggregates, *J. Am. Chem. Soc.*, 1967, **89**, 4698–4703.

- 57 M. Pérez-Rodríguez, G. Prieto, C. Rega, L. M. Varela, F. Sarmiento and V. Mosquera, A Comparative Study of the Determination of the Critical Micelle Concentration by Conductivity and Dielectric Constant Measurements, *Langmuir*, 1998, **14**, 4422–4426.
- 58 I. M. Umlong and K. Ismail, Micellization behaviour of sodium dodecyl sulfate in different electrolyte media, *Colloids Surf. Physicochem. Eng. Asp.*, 2007, **299**, 8–14.
- 59 G. Basu Ray, I. Chakraborty and S. P. Moulik, Pyrene absorption can be a convenient method for probing critical micellar concentration (cmc) and indexing micellar polarity, *J. Colloid Interface Sci.*, 2006, **294**, 248–254.
- 60 M. Layek, S. Kundu, P. Karmakar, S. M. Rahaman, T. Mandal, A. Patra, A. Nandy, M. Chakravarty, P. Sar and B. Saha, Optimizing homologous alcohol oxidation: elucidating the impact of surfactant-alcohol hydrophobic interaction and micellar surface charge, *Res. Chem. Intermed.*, 2025, **51**, 311–330.
- 61 Z. Li, Y. Zhang, J. Lin, W. Wang and S. Li, High-Yield Di-Rhamnolipid Production by *Pseudomonas aeruginosa* YM4 and its Potential Application in MEOR, *Molecules*, 2019, **24**, 1433.
- 62 Y. Zhang, T. L. Placek, R. Jahan, P. Alexandridis and M. Tsianou, Rhamnolipid Micellization and Adsorption Properties, *Int. J. Mol. Sci.*, 2022, **23**, 11090.
- 63 I. E. Kłosowska-Chomiczewska, K. Mędrzycka, E. Hallmann, E. Karpenko, T. Pokynbroda, A. Macierzanka and C. Jungnickel, Rhamnolipid CMC prediction, *J. Colloid Interface Sci.*, 2017, **488**, 10–19.
- 64 G. V. Buxton, C. L. Greenstock, W. P. Helman and A. B. Ross, Critical Review of rate constants for reactions of hydrated electrons, hydrogen atoms and hydroxyl radicals ($\cdot\text{OH}/\cdot\text{O}^-$ in Aqueous Solution), *J. Phys. Chem. Ref. Data*, 1988, **17**, 513–886.
- 65 R. Xiao, J. Ma, Z. Luo, W. Zeng, Z. Wei, R. Spinney, W. Hu and D. D. Dionysiou, Experimental and theoretical insight into hydroxyl and sulfate radicals-mediated degradation of carbamazepine, *Environ. Pollut.*, 2020, **257**, 113498.
- 66 E. De Laurentiis, S. Chiron, S. Kouras-Hadef, C. Richard, M. Minella, V. Maurino, C. Minero and D. Vione, Photochemical Fate of Carbamazepine in Surface Freshwaters: Laboratory Measures and Modeling, *Environ. Sci. Technol.*, 2012, **46**, 8164–8173.
- 67 D. Vogna, R. Marotta, R. Andreozzi, A. Napolitano and M. d'Ischia, Kinetic and chemical assessment of the UV/H₂O₂ treatment of antiepileptic drug carbamazepine, *Chemosphere*, 2004, **54**, 497–505.
- 68 M. Srabovic, M. Poljakovic and E. Pehlic, Micellar Solubilization of Carbamazepine, *J. Sci. Res. Rep.*, 2014, 3106–3116.
- 69 J. R. Crison, N. D. Weiner and G. L. Amidon, Dissolution media for in vitro testing of water-insoluble drugs: Effect of surfactant purity and electrolyte on in vitro dissolution of carbamazepine in aqueous solutions of sodium lauryl sulfate, *J. Pharm. Sci.*, 1997, **86**, 384–388.
- 70 G. Duplâtre, M. F. Ferreira Marques and M. da Graça Miguel, Size of Sodium Dodecyl Sulfate Micelles in Aqueous Solutions as Studied by Positron Annihilation Lifetime Spectroscopy, *J. Phys. Chem.*, 1996, **100**, 16608–16612.
- 71 F. J. Beltrán, G. Ovejero and J. Rivas, Oxidation of Polynuclear Aromatic Hydrocarbons in Water. 3. UV Radiation Combined with Hydrogen Peroxide, *Ind. Eng. Chem. Res.*, 1996, **35**, 883–890.
- 72 M. E. Lindsey and M. A. Tarr, Inhibited hydroxyl radical degradation of aromatic hydrocarbons in the presence of dissolved fulvic acid, *Water Res.*, 2000, **34**, 2385–2389.
- 73 W. R. Haag and C. C. D. Yao, Rate constants for reaction of hydroxyl radicals with several drinking water contaminants, *Environ. Sci. Technol.*, 1992, **26**, 1005–1013.

- 74 A. Lebrón-Paler, J. E. Pemberton, B. A. Becker, W. H. Otto, C. K. Larive and R. M. Maier, Determination of the Acid Dissociation Constant of the Biosurfactant Monorhamnolipid in Aqueous Solution by Potentiometric and Spectroscopic Methods, *Anal. Chem.*, 2006, **78**, 7649–7658. Article Online
DOI: 10.1039/DT5EM00203F
- 75 J. T. Champion, J. C. Gilkey, H. Lamparski, J. Retterer and R. M. Miller, Electron Microscopy of Rhamnolipid (Biosurfactant) Morphology: Effects of pH, Cadmium, and Octadecane, *J. Colloid Interface Sci.*, 1995, **170**, 569–574.
- 76 R. J. Eismín, E. Munusamy, L. L. Kegel, D. E. Hogan, R. M. Maier, S. D. Schwartz and J. E. Pemberton, Evolution of Aggregate Structure in Solutions of Anionic Monorhamnolipids: Experimental and Computational Results, *Langmuir*, 2017, **33**, 7412–7424.
- 77 S. I. M. Ng, K. H. Ng, P. W. F. Yeung, R. Xu, P.-K. So, Y. Huang, J. Z. Yu, C. K. K. Choi, Y.-L. S. Tse and M. N. Chan, Chemical transformation of a long-chain alkyl organosulfate via heterogeneous OH oxidation: a case study of sodium dodecyl sulfate, *Environ. Sci. Atmospheres*, 2022, **2**, 1060–1075.
- 78 L. Wojnárovits and E. Takács, Rate constants of sulfate radical anion reactions with organic molecules: A review, *Chemosphere*, 2019, **220**, 1014–1032.
- 79 Y. Tao, O. Monfort, M. Brigante, H. Zhang and G. Mailhot, Phenanthrene decomposition in soil washing effluents using UVB activation of hydrogen peroxide and peroxydisulfate, *Chemosphere*, 2021, **263**, 127996.
- 80 R. Matta, S. Tlili, S. Chiron and S. Barbat, Removal of carbamazepine from urban wastewater by sulfate radical oxidation, *Environ. Chem. Lett.*, 2011, **9**, 347–353.
- 81 P. Neta, V. Madhavan, H. Zemel and R. W. Fessenden, Rate constants and mechanism of reaction of sulfate radical anion with aromatic compounds, *J. Am. Chem. Soc.*, 1977, **99**, 163–164.
- 82 Y.-J. An and E. R. Carraway, PAH degradation by UV/H₂O₂ in perfluorinated surfactant solutions, *Water Res.*, 2002, **36**, 309–314.
- 83 K. M. Bansal, L. K. Patterson, E. J. Fendler and J. H. Fendler, Reactions of hydrated electrons, hydrogen atoms and hydroxyl radicals in micellar systems, *Int. J. Radiat. Phys. Chem.*, 1971, **3**, 321–331.
- 84 J. A. Faust and J. P. D. Abbatt, Organic Surfactants Protect Dissolved Aerosol Components against Heterogeneous Oxidation, *J. Phys. Chem. A*, 2019, **123**, 2114–2124.
- 85 C. T. Jafvert, L. V. H. Patricia and J. K. Heath, Solubilization of non-polar compounds by non-ionic surfactant micelles, *Water Res.*, 1994, **28**, 1009–1017.

View Article Online
DOI: 10.1039/D5EM00203F

Data availability statement

Raw data files can be obtained contacting the corresponding author.

