## Environmental Science Processes & Impacts



## CORRECTION

View Article Online
View Journal | View Issue



Cite this: Environ. Sci.: Processes Impacts, 2025, 27, 513

## Correction: Responses of CO<sub>2</sub> and CH<sub>4</sub> in the alpine wetlands of the Tibetan Plateau to warming and nitrogen and phosphorus additions

Wenbao Zhang,<sup>a</sup> Huijuan Xin,<sup>a</sup> Zongxing Li,\*<sup>bc</sup> Qiao Cui,<sup>d</sup> Bin Xu,<sup>e</sup> Biao Tang,<sup>a</sup> Yaning Wang,<sup>c</sup> Chong Xu<sup>a</sup> and Jian Xue<sup>d</sup>

DOI: 10.1039/d5em90001h

rsc.li/espi

Correction for 'Responses of  $CO_2$  and  $CH_4$  in the alpine wetlands of the Tibetan Plateau to warming and nitrogen and phosphorus additions' by Wenbao Zhang et al., Environ. Sci.: Processes Impacts, 2024, 26, 1516–1525, https://doi.org/10.1039/D4EM00174E.

The authors regret that there were errors in the Experimental design Section 2.2.1.

The corrected text and updated Table 1 in section 2.2.1 are shown below:

Nitrogen and phosphorus addition treatments were applied at each temperature level, utilizing urea  $(CO(NH_2)_2)$  for nitrogen fertilization, with three gradients of nitrogen addition: N1 (5 g N per m<sup>2</sup> per year), N2 (10 g N per m<sup>2</sup> per year), and N3 (15 g N per m<sup>2</sup> per year). Calcium dihydrogen phosphate  $(Ca(H_2PO_4)_2)$  was employed for phosphorus fertilization, with three gradients of phosphorus addition: P1 (5 g P per m<sup>2</sup> per year), P2 (10 g P per m<sup>2</sup> per year), and P3 (15 g P per m<sup>2</sup> per year). Additionally, a combined nitrogen and phosphorus treatment, N2P2 (10 g N per m<sup>2</sup> per year, 10 g P per m<sup>2</sup> per year), and a control (CK) with no nutrient additions were implemented, with the NW and no nutrient addition treatment serving as controls.

Table 1 Sample conditions<sup>a</sup>

	NW		W1		W2		W3	
	N	P	N	P	N	P	N	P
CK	0	0	0	0	0	0	0	0
N2P2	10	10	10	10	10	10	10	10
P3	0	15	0	15	0	15	0	15
P2	0	10	0	10	0	10	0	10
P1	0	5	0	5	0	5	0	5
N3	15	0	15	0	15	0	15	0
N2	10	0	10	0	10	0	10	0
N1	5	0	5	0	5	0	5	0

<sup>&</sup>lt;sup>a</sup> In the table, N and P represent the annual additions of nitrogen (g N per m per year) and phosphorus (g P per m<sup>2</sup> per year) per plot, respectively.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

aSchool of Environment and Municipal Engineering, Lanzhou Jiao Tong University, Lanzhou 730070, Gansu, China. E-mail: zwenbao@163.com

<sup>&</sup>lt;sup>b</sup>Observation and Research Station of Eco-Hydrology and National Park by Stable Isotope Tracing in Qilian Mountains, Key Laboratory of Ecological Safety and Sustainable Development in Arid Lands, Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, Lanzhou 730000, China

<sup>&</sup>lt;sup>e</sup>College of Geography and Environmental Science, Northwest Normal University, Lanzhou 730070, China

<sup>&</sup>lt;sup>d</sup>Northwest Institute of Eco-Environment and Resources, Chinese Academy of Sciences, Lanzhou 730000, China

<sup>&</sup>lt;sup>e</sup>College of Energy and Power Engineering, Lanzhou University of Technology, Lanzhou 730050, China