

# Sustainable Food Technology

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: A. Sreedharan, K. V. Sunooj, V. Sathish kumar, N. Sahil, M. Navaf, S. Sabu, A. Sasidharan and V. Venugopal, *Sustainable Food Technol.*, 2025, DOI: 10.1039/D5FB00236B.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

## Sustainability spotlight

View Article Online  
DOI: 10.1039/D5FB00236B

The global food system faces growing challenges, resource depletion, climate change, and food insecurity, making sustainable transformation urgent. Industries' intensive exploitation of marine resources contributes significantly to overfishing, habitat damage, and emissions. A shift toward ecosystem-based fisheries, sustainable aquaculture, and circular strategies like seafood byproduct valorization is essential. Valorising byproducts like shells, skins, and bones reduces waste and pollution while producing high-value compounds like collagen, omega-3s, and chitin. These materials have diverse applications in food, pharma, and cosmetic industries, boosting local economies and reducing environmental burdens. Integrating such approaches fosters circular bioeconomy practices and supports Sustainable Development Goals, particularly SDG 12 (responsible consumption) and SDG 13 (climate action).



# Valorization of side streams to enhance seafood sustainability

View Article Online  
DOI: 10.1039/D5FB00236B

Received May 2025

Accepted -

DOI: 10.1039/x0xx00000x

Sreedharan Advaita<sup>a</sup>, Kappat Valiyapeediyekkal Sunooj<sup>\*a</sup>, Sathish kumar vellaisamy<sup>a</sup>, Sahil Negi<sup>a</sup>, Muahmmmed Navaf<sup>a</sup>, Sarasan Sabu<sup>b</sup>, Abhilash Sasidharan<sup>c</sup>, Vazhiyil Venugopal<sup>d</sup>

## Corresponding author

Kappat Valiyapeediyekkal Sunooj

[sunooj4u@pondiuni.ac.in](mailto:sunooj4u@pondiuni.ac.in)

<sup>a</sup> Department of Food Science and Technology, Pondicherry University, Puducherry, 605014, India.

<sup>b</sup> School of industrial fisheries, cochin university of science and technology, kochi, 682506, India.

<sup>c</sup> Department of Fish Processing Technology, Kerala University of Fisheries and Ocean Studies, Kochi, 682506, India.

<sup>d</sup> Retired scientist, Food Technology Division, Bhabha Atomic Research Center, Mumbai, India 400085

## Abstract

The global seafood industry produces many byproducts, often called seafood side streams, encompassing fish heads, skins, bones, viscera, and shells. Traditionally viewed as waste, these byproducts possess considerable potential for valorization, supporting sustainability, circular economy principles, and fostering progress toward the Sustainable Development Goals (SDGs) of the United Nations. This review explores the composition, bioactive potential, and innovative valorization strategies for seafood side streams, focusing on green processing including enzymatic hydrolysis, fermentation, and recovery of bioactive compounds. Diverse applications of the recovered compounds in food, pharmaceuticals, cosmetics, packaging, and other fields highlight economic and environmental advantages of using seafood side streams. Nonetheless, technical limitations, financial feasibility, regulatory hurdles, and consumer



acceptance remain significant obstacles. Policy recommendations, advancements in bioprocessing research, and integration of emerging technologies such as AI and blockchain for traceability are crucial for addressing these challenges. Emphasizing the value of cross-disciplinary collaboration, this review advocates the potential of zero-waste conversion of seafood side streams into valuable ingredients for a healthy blue economy and sustainable seafood industry.

### Sustainability spotlight

The global food system faces growing challenges, resource depletion, climate change, and food insecurity, making sustainable transformation urgent. Industries' intensive exploitation of marine resources contributes significantly to overfishing, habitat damage, and emissions. A shift toward ecosystem-based fisheries, sustainable aquaculture, and circular strategies like seafood byproduct valorization is essential. Valorizing byproducts like shells, skins, and bones reduces waste and pollution while producing high-value compounds like collagen, omega-3s, and chitin. These materials have diverse applications in food, pharma, and cosmetic industries, boosting local economies and reducing environmental burdens. Integrating such approaches fosters circular bioeconomy practices and supports Sustainable Development Goals, particularly SDG 12 (responsible consumption) and SDG 13 (climate action).

#### 1. Introduction

Seafood industries generate a wide range of byproducts during processing and post-harvest handling, commonly including heads, shells, skins, scales, bones, fins, viscera, roe, and frames, which account for 30–70% of the total biomass, depending on species and processing methods<sup>1</sup>. Although traditionally discarded, these off-cuts are abundant in valuable bioactive compounds and can be repurposed into nutraceuticals, animal feed, and eco-friendly packaging materials<sup>2</sup>. The proper utilization of seafood side streams integrates within the principles of circular economy, enhancing sustainability by reducing waste and increasing resource efficiency. As world food demand is projected to rise by 50–60% by 2050, existing food production systems face mounting pressures due to resource depletion, climate change, pollution, and biodiversity loss<sup>3,4</sup>. Additionally, food waste contributes to inefficiencies by placing undue strain by imposing excessive demands on land, water, and energy<sup>5</sup>. Greenhouse gas emissions from food systems have increased by approximately 25% since 2010, cropland use has increased by about 6% since 2000, and blue water consumption continues to rise due to intensified agricultural and aquaculture practices (FAO, 2023; IPCC, 2021)<sup>6</sup>.



Sustainable food systems are progressively recognized as essential for ensuring worldwide food security. Production systems must evolve to achieve sustainability by incorporating decarbonization strategies that mitigate climate change while enhancing resilience<sup>4</sup>. The seafood industry presents significant opportunities for reducing environmental impact by adopting sustainable aquaculture and fisheries management practices. Investments in green technologies and diverse aquaculture systems can significantly reduce the sector's carbon footprint. Moreover, green innovation has enhanced sustainable performance, bridging environmentally friendly human resource initiatives and improved sustainability outcomes for small and medium enterprises<sup>7</sup>. The marine food sector's role in food security, economic stability, and sustainable fisheries management is crucial to prevent overexploitation and ensure the long-term availability of seafood<sup>8</sup>. Unsustainable fishing methods threaten marine ecosystems, but implementing responsible fishing practices can help maintain biodiversity and ecosystem health. Certification programs such as the Marine Stewardship Council (MSC) provide critical oversight to promote environmentally accountable fishing<sup>9</sup>. Beyond environmental concerns, sustainability in seafood also involves social and economic aspects, including fair labor practices, equitable distribution of resources, and the well-being of fishing communities. A socio-ecological framework is integral for promoting sustainable seafood practices worldwide, accounting for cultural and regional differences<sup>10</sup>. Recent developments have significantly advanced the creation of regional and national ocean accounts. Collaborations like the Global Ocean Accounts Partnership play a crucial role by offering technical expertise and facilitating the sharing of knowledge across nations. Ocean accounts are built upon internationally recognized frameworks such as the System of National Accounts (SNA) and the System of Environmental-Economic Accounting (SEEA). These accounts provide a way to move "beyond GDP" by capturing the state and value of natural capital essential to the global economy. By integrating diverse data streams, these accounts support the creation of key indicators that help monitor progress towards a sustainable blue economy and fulfill global commitments like the SDGs<sup>11</sup>.

The seafood industry produces significant byproducts that, when managed sustainably, can be repurposed into valuable products like animal feed, biodegradable packaging, etc. Consumer engagement is a critical driver of sustainable seafood demand. Enhancing labelling transparency enables consumers to make informed choices, thereby fostering market incentives for responsible fisheries and aquaculture operations<sup>12</sup>. However, the global seafood trade remains complex, often disconnecting consumers and the sustainability of the fishing and aquaculture practices behind their



seafood choices. Establishing stringent traceability mechanisms and international sustainability standards ensures that seafood products meet environmental and ethical criteria<sup>8</sup>. DOI: 10.1039/D5FB00236B

The UN SDGs introduced in 2015, serve as a guiding framework for fostering sustainability in various sectors, including seafood. Among these, SDG 12 (Responsible Consumption and Production, Circular Economy and Waste Reduction), and SDG 14 (Life Below Water- Sustainable Fisheries and Ecosystem Preservation) are related to seafood production and marine conservation<sup>4</sup>. SDG Target 12.3 aims to halve per capita global food waste at the retail and consumer levels and reduce food losses along production and supply chains by 2030<sup>13</sup>. The SDG is also oriented to promote efficient resource utilization in seafood production by advocating for sustainable fishing techniques, bycatch reduction, and waste minimization. Environmentally responsible aquaculture practices further support this objective<sup>14</sup>. The SDG 14 calls for improved regulatory frameworks and sustainable fisheries management in order to preserve marine habitats, biodiversity, and mitigate overfishing<sup>15</sup>. To achieve these goals, policy reforms, ecosystem-based management strategies, and innovative governance mechanisms must be prioritized<sup>16</sup>.

Wild-capture fisheries have plateaued in the past 30 years, putting pressure on sustainable seafood production. At the same time, conventional processing operations result in discards as high as 50% of the processed finfish and shellfish, which include heads, skins, bones, viscera, and shells. If not properly utilized, these discards not only deny valuable nutrients to consumers, but also are responsible for environmental pollution and global warming due to putrefaction and release of GHG emissions. Strategies are therefore essential for climate-smart seafood production, food waste reduction, dietary adaptations to combat the current dual emergencies of climate change, together with associated setbacks, and sustainable fisheries. Therefore, a radical transformation of food systems is imperative to reach the UN SDGs and to fulfil the Paris Climate Agreement to limit global warming to well below 2 °C and make efforts to limit it to 1.5 °C. This is urgent for global food security and environmental sustainability<sup>17</sup>. Sustainable seafood production, encompassing fish, shellfish, and seaweed, must be managed to enable species regeneration, habitat restoration, and ecosystem resilience<sup>18</sup>. Since the agriculture and food sectors significantly contribute to greenhouse gas emissions, minimizing food loss and optimizing byproduct utilization are essential<sup>19</sup>. According to the FAO's 2024 report, world fisheries and aquaculture production reached a record of 223.2 million tonnes in 2022, comprising 185.4 million tonnes of aquatic animals and 37.8 million tonnes of algae. However, approximately 27% of landed fish is lost or wasted between landing and consumption, highlighting significant inefficiencies in the seafood supply chain<sup>20</sup>. This substantial loss underscores the need for improved handling, processing, and distribution practices



to enhance sustainability and food security in the global seafood sector. Currently, seafood accounts for only 17% of total edible meat production, but projections indicate that by 2050, the sector could expand by 21- 44 million tonnes, fulfilling 12-25% of the anticipated global meat demand<sup>18</sup>. However, the issue of food waste remains substantial, with nearly 1/3 of all food produced, approximately 1.6 billion tonnes annually, being lost, including 35% of fish and marine products. Addressing this inefficiency is crucial for strengthening global food security<sup>21</sup>.

This review explores the potential of seafood side streams as sources of nutrients and their usefulness, focusing on sustainable utilization strategies such as enzymatic hydrolysis, fermentation, and extraction of bioactive compounds. The valorization of these byproducts presents opportunities for different applications in the food processing, healthcare, and personal care industries, contributing to food security, health, and environmental sustainability. However, several challenges hinder widespread adoption, including technical limitations, cost constraints, regulatory barriers, and consumer scepticism. This review also attempts to address these challenges and provide future research directions, emphasizing advancements in bioprocessing, policy interventions, and the application of innovative technologies such as AI and blockchain to improve traceability and proficiency. By addressing these aspects, this review highlights the role of seafood side-stream valorization in promoting a circular bioeconomy and improving sustainability in the seafood sector.

## 2. Valorisation Strategies for Seafood Side Streams

A resilient food system cannot be achieved without minimizing losses of valuable resources; hence, it is imperative to adopt integrated approaches that reduce food loss and waste<sup>22</sup>. Maximizing food utilization requires coordinated efforts worldwide and within local communities. Using seafood byproducts is essential for enhancing sustainability and economic viability in the fisheries and aquaculture sectors. The key to transformative change lies in adopting new technologies and innovative solutions, including e-commerce for marketing and mobile food processing, improved workflows, and effective practices for managing food quality and minimizing waste. Through implementing circular economy principles, the sector may convert low-value waste into high-value products, thereby enhancing food security and promoting environmental conservation<sup>23</sup>. Recent advancements in process optimization, green extraction methods, and enzyme-assisted technologies have improved the efficiency and sustainability of recovering well-established compounds such as collagen, gelatin, omega-3 fatty acids, and chitin from seafood by-products<sup>24</sup>. Moreover, hydrothermal and enzymatic procedures have surfaced as effective techniques for transforming





seafood waste into biofuels, bioplastics, and useful components<sup>25</sup>. Fermentation and microbial processing augment the extraction of bioactive substances, facilitating sustainable resource utilization<sup>26</sup>. Considering the increasing apprehensions regarding environmental sustainability and food security, incorporating innovative valorization technologies for seafood side streams offers a feasible approach to minimize waste while producing economic and ecological advantages<sup>27</sup>.

### 2.1. Seafood side streams: composition and Bioactive potential

Fishery byproducts refer to the side-streams generated during the processing of seafood. These include off-cuts of aquatic food that are not typically consumed directly, such as head, skin, tail, viscera, shells, and other cut-offs. These side streams are rich in components such as enzymes, proteins, healthy fatty acids (MUFA & PUFA), gelatin, and collagen<sup>25</sup>. Seafood side streams can be classified according to the type of seafood and the particular parts of the organism, and are presented in **Table 1**. Seafood offcuts are high in nutritional and bioactive components, offering opportunities to develop valuable products with various uses in the food processing, healthcare, and personal care industries. The concentration and composition of these components are determined through various analytical methods, and their bioactive potential presents promising avenues for creating functional and health-promoting products. Seafood side streams, such as fish processing byproducts, contain moderate to high protein and lipid levels. For instance, sprat trimmings, marinated herring, and mackerel in tomato sauce had protein levels ranging from 28% to 32% of dry matter and lipid levels ranging from 34% to 43%. Additionally, sprat trimmings have approximately 29% ash and 1.5% phosphorus, indicating the presence of essential minerals that add value to these side streams<sup>28</sup>. Seafood byproducts such as fish viscera, skin, and heads contain notable levels of lipids, particularly MUFAs (e.g., oleic acid) and PUFAs (e.g., EPA and DHA). The fatty acid profile varies by species and tissue type, with oily fish wastes (e.g., from mackerel, sardines) being especially rich in omega-3 PUFAs, while lean fish may yield lower lipid fractions<sup>1</sup>. Seafood offcuts are rich in bioactive ingredients and fractions contain proteins, peptides, collagen, fish oils, gelatin, enzymes, chitin, and minerals. These bioactive components have been shown to possess various health-promoting properties such as antimicrobial, antioxidative, antihypertensive, and antihyperglycemic activities. Furthermore, the bioactive components from seafood side streams have potential applications in nutrient-rich foods, special feeds, nutraceuticals, drugs, and cosmetic products<sup>29</sup>. The concentration and composition of seafood side stream components are determined through various analytical methods, including the assessment of protein, lipid, ash, and phosphorous levels in the side streams<sup>28</sup>. The proteins found in marine food byproducts are rich in nutritional value and possess functional properties. These proteins contain bioactive peptides that





offer various health benefits, such as anti-inflammatory, anti-coagulant, anti-cancer, and hypocholesterolemic effects<sup>24</sup>. Fish fats are also an exceptional source of omega-3 (PUFAs), with valuable therapeutic properties. The bioactive potential of seafood side stream components is evident in their diverse benefits on health, including cardioprotective, neuroprotective, antioxidant, anti-inflammatory, anti-cancer, antiobesity, anticoagulant, antimicrobial, and immunomodulatory activities<sup>30</sup>. Furthermore, the bioactive components from seafood side streams have the potential to be developed into functional ingredients for use in functional foods and nutraceuticals, contributing to the maintenance of health and wellness<sup>31</sup>. Bioactive components from seafood side streams are incorporated into foods to enhance their nutritional value and health benefits. Many bioactives are used for their therapeutic properties in drug formulations and skincare products. In addition to proteins and lipids, seafood byproducts particularly shells and exoskeletons of crustaceans like shrimp and crabs are rich in minerals. Calcium is a major component, often present as calcium carbonate, accounting for up to 20–40% of dry shell weight. These mineral-rich residues have potential applications in nutraceuticals, bone tissue scaffolding, and as natural calcium supplements. Other essential minerals such as magnesium, phosphorus, and trace metals (e.g., zinc, and as natural calcium supplements. Other essential minerals such as magnesium, phosphorus, and trace metals (e.g., zinc, copper) are also found, although their concentrations vary by species and processing methods<sup>32</sup>.

**Table 1 Classification and composition of Seafood off-cuts**

Source / Species Group	Residue Type	Protein (%)	Lipid (%)	Carbohydrate (%)	Ash (%)	Moisture (%)	Reference
Shrimp (Crustacean)	Shell	25–40	2–5	20–30 (chitin)	15–25	8–15	33
Tuna (Teleost)	Head	18–25	10–15	<1	5–10	50–55	34
Salmon (Teleost)	Skin	20–30	15–25	<1	5–7	40–50	35
Skate (Cartilaginous)	Cartilage/Bone	15–22	2–5	1–2	25–35	35–40	36
Fish (Mixed species)	Mixed Waste	57.9 ± 5.3	19.1 ± 6.1	1.2 ± 1.2	21.8 ± 3.5	70–75	37



REVIEW		Sustainable food technology						
Shrimp	Waste	94.6 ±	4.1 ±	8.6 ± 0.1	1.3 ±	65 - 75	33	
		0.2	0.1		0.02		<a href="#">View Article Online</a> DOI: 10.1039/D5FB00236B	
Lobster	Waste	28.6 ±	2.2 ±	22.8 ± 1.5	33.7	65 - 72	38	
		0.5	0.2		± 2.2			
Mackerel	Head	12.3	17.2	1.17	3.74	55-60	39	
	Frame	14.2	10.4	0.31	3.48	50-55		
	Fins/Skin/	12.2	20.8	0.00	1.36	55-60		
	Gut Mix							
Salmon (Viscera Meal)	Meal	63.6	11.8	3.32	12.0	10.45	40	
Anchovy (Whole meal)	Meal	73.6	10.3	1.52	12.0	6.53	40	
Sprat (Whole meal)	Meal	70.1	9.9	3.77	12.0	7.15	40	
Cod/Hake (Bone Powder)	Bone	~13.2	~7.0	~1.8	49.9	28.2	41	

2.2. Extraction methods and Technological approaches for Byproduct Valorization

The role of seafood side streams in circular food systems is paramount, with their utilization and waste management practices being more critical than those of other food sources. The seafood offcuts are indispensable in worldwide food security and human nutrition, yet substantial waste generation occurs throughout the seafood supply chain. The seafood industry discards millions of waste each year, predominantly consisting of crab, shrimp, and lobster shells, which contribute to environmental problems linked to COD and BOD<sup>42</sup>. Fish waste, animal-derived non-lignocellulosic biomass, lacks lignin, cellulose, and hemicellulose, distinguishing it from traditional lignocellulosic feedstocks. Instead, it is composed of carbohydrates, proteins, and fats. The biochemical composition of various seafood wastes has been documented; however, a comprehensive characterization across different fish species remains limited. Studies suggest that the carbohydrate content in fish waste varies between 8% and 15%, whereas the protein content ranges from 30% - 50%, depending on the type of waste. Comparatively, meat and shrimp wastes contain 20–25% and 30–45% protein, respectively. Furthermore, the fatty acid composition of fish waste can range from 10% to 60%<sup>33</sup>. Unlike lignocellulosic biomass, seafood waste is rich in proteins and fats, making it

a valuable substrate for protein hydrolysates, bioactive compounds, and bioenergy applications<sup>43</sup>.

**Table 2** presents valorization of seafood by-products, their sources, composition, and bioactivities. A thermochemical process called Hydrothermal carbonization (HTC), particularly suited for wet biomass (>50% moisture content), offers an efficient approach for converting fish waste into hydrochar and bio-oil<sup>43</sup>. Unlike pyrolysis, which requires an energy-intensive drying step, HTC directly utilizes wet biomass, thereby reducing energy demands<sup>44</sup>. This process involves submerging the biomass in subcritical water conditions, facilitating the conversion of organic components into bio-oil and carbonaceous solids (Hydrochar), while producing minimal gaseous byproducts<sup>43</sup>. However, the application of HTC to seafood waste is limited due to its poor carbohydrate profile, as seafood residues are predominantly protein-rich (e.g., fish oils, proteins, peptides, collagen, gelatin, enzymes, chitin, and minerals). Recent research has explored optimizing HTC for such feedstocks, with studies on microwave-assisted hydrothermal carbonization (MHTC) showing promise in producing hydrochar comparable to that from municipal and sewage wastes<sup>44</sup>. Innovations in reactor design and process optimization have further enhanced the efficiency of HTC for fish waste treatment<sup>45</sup>. While this approach provides a sustainable waste management solution and generates valuable carbonaceous materials and energy resources<sup>44</sup>. Traditional disposal techniques like landfills and ocean dumping have been hugely criticized due to environmental risks, its industrial-scale adoption remains limited<sup>46</sup>. Scientific findings reveal that fish side-streams harbor essential bioactive ingredients, including fish oils, proteins, peptides, collagen, gelatin, enzymes, chitin, and minerals, offering significant potential for use in functional foods, high-performance feeds, nutraceuticals, pharmaceuticals, and cosmetic formulation. Some of the waste is converted into low-value products like animal feed for animals and aquaculture<sup>29</sup>. Given the high protein content of seafood residues, which contrasts with the carbohydrate-rich feedstocks typically suited for HTC, industrial-scale application of this technique for seafood waste is not yet widespread. Instead, valorization methods such as enzymatic hydrolysis and extraction are more commonly employed to leverage the protein-rich nature of these byproducts for higher-value applications<sup>45</sup>.

Several strategies and innovative technologies can be employed to improve the efficiency of seafood side-stream valorization. pH-Shift Technology uses acid or base solubilization and isoelectric precipitation to recover proteins from fish-side streams. Compared to conventional methods, it has shown significant reductions in environmental impacts such as carbon footprint, acidification, and water use. Its key benefits include a substantial reduction in environmental impacts and the potential for further enhancement with renewable energy integration<sup>47</sup>.



Conventional protein recovery methods such as thermal coagulation and mechanical pressing often suffer from lower yields and higher environmental footprints. The term 'acidification' in this context refers to environmental acidification from disposal of untreated seafood waste, not from the pH-shift processing itself<sup>48</sup>. Membrane concentration helps to recover molecules from side streams, while flocculation aids in recovering proteins and phosphates, minimizing the organic load of effluents. Green extraction techniques such as microwave- and ultrasound-assisted extraction, supercritical fluid extraction, and subcritical water extraction are effective for extracting bioactive compounds from seafood by-products<sup>49</sup>. Implementing circular economy principles to reduce waste and enhance resource reuse can result in greater production efficiency and reduced demand for natural resources, improving sustainability and economic viability, thereby contributing to global food security and environmental conservation<sup>50</sup>. Several studies have explored the techno-economic feasibility of zero-waste processing of seafood side streams. Zero-waste biorefinery concepts are based on the full valorization of seafood side streams. Their proper utilization aligns with circular economy principles by reducing waste and increasing sustainability<sup>2</sup>. The proper utilization of seafood side streams integrates within the principles of circular economy, enhancing sustainability by reducing waste and increasing. By integrating these innovative technologies, optimization strategies, and circular economy principles, the efficiency of seafood side-stream valorization can be significantly improved, leading to enhanced sustainability and economic benefits.

The seafood processing side streams through green chemistry present a transformative and eco-sustainable solution to the seafood industry's environmental challenges. Traditionally, these discards, which can constitute up to 70% of the raw material, are disposed of through methods such as landfilling, ocean dumping, or incineration, all of which contribute significantly to environmental degradation. Landfilling releases potent greenhouse gases like methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O), ammonia (NH<sub>3</sub>), and hydrogen sulfide (H<sub>2</sub>S), which not only accelerate global warming but also contaminate soil and water systems. On the other hand, incineration emits CO<sub>2</sub>, sulfur dioxide (SO<sub>2</sub>), and nitrogen oxides, further polluting the atmosphere and harming biodiversity. Ocean dumping leads to the depletion of oxygen levels in marine environments, disrupting aquatic ecosystems<sup>51</sup>.

Techniques such as enzymatic hydrolysis, microbial fermentation, anaerobic digestion, and green solvents allow for efficient biotransformation of seafood waste under low-energy conditions. These processes minimize hazardous emissions and reduce reliance on fossil fuel-intensive practices. Integrating these green valorization methods into biorefinery platforms enhances



environmental stewardship by achieving near-total utilization of waste biomass, thus embracing a circular economy model. This approach ensures that materials are continually recycled into the production chain, eliminating the "take-make-dispose" model and contributing to resource conservation. According to life cycle analysis (LCA) studies referenced in the paper, products recovered from seafood discards through green bioprocesses exhibit 88–94% lower greenhouse gas emissions than conventional products. For instance, managing protein-rich waste via valorization can prevent the release of up to 750 kg of CO<sub>2</sub>-equivalents per kilogram of wasted protein<sup>52</sup>. Moreover, implementing cleaner production strategies and decarbonization pathways can reduce up to 35% of total GHG emissions in seafood supply chains, as observed in practical examples from Western Australia. Valorization also contributes to the achievement of key Sustainable Development Goals (SDGs), particularly SDG 12 (responsible consumption and production), SDG 13 (climate action), and SDG 14 (life below water). Ultimately, adopting green chemistry for seafood waste valorization reduces the seafood industry's ecological footprint and promotes economic viability and social responsibility, facilitating the emergence of a resilient, low-carbon, and sustainable blue economy<sup>44</sup>.

**Table 2 -Valorization of Seafood By-products: sources, composition, bioactivities**

COMPONENT	CONCENTRATION/ COMPOSITION	BIOACTIVE POTENTIAL	SOURCES	REFER ENCES
Protein	Rich in fish protein hydrolysates (FPH), bioactive peptides, collagen, and gelatin. Shellfish- 65% Muscles- 17-22% Body parts 8-35%	Antioxidant, antihypertensive, anticoagulant, immunomodulatory, antimicrobial	Fish muscle, skin, viscera, shellfish	25,26
Fatty Acids	Present in fish oils and other lipid fractions Shellfish- 7-19% EPA - 24.7 DHA-28.3% PUFA- 7-19% Omega-3-10.95%	Cardioprotective, anti-inflammatory, neuroprotective, anticancer	Fish heads, viscera, shellfish	29



REVIEW

Sustainable food technology

minerals/Ash	Contains phosphorus, magnesium, sodium, calcium other minerals Inorganic compounds- 60-70% Fishbone- 70% Shellfish – 21% Crab- 28.5%	Bone health, mineral supplements	Fish bones, shellfish, crab shells	24,30 <small>New Article Online DOI: 10.1039/D5FB00236B</small>
Carotenoids	Present in fish skin and other pigmented tissues Astaxanthin, canthaxanthin, and zeaxanthin 32%	Antioxidant, anti-inflammatory, anticancer	Crustacean shells (primary), fish skin (secondary) Shrimp shells, fish skin	24,30
Enzymes	Chitinase, alkaline phosphatase, Proteases, transglutaminase, lipases hyaluronidase, and acetyl glycosaminidase	Used in food processing, pharmaceuticals	Fish viscera, shellfish	24
Polysaccharides (Chitin, Chitosan and glycosaminoglycans (GAG))	Proteoglycans, glycolipids, glycoconjugates Glycosaminoglycans (GAGs, sGAGs), hyaluronic acid (HA), chondroitin sulfate (CS), heparan sulfate, dermatan sulfate, and keratan sulfate Extracted from shells and other crustacean by-products 15-20% lobster shell- 70% Dry crab shell- 67-72 % squid skeletal pen- 41%	Anti-inflammatory, joint health, skin health Anticoagulant, antiallergic, antidiabetic, antihypertensive, and anti-obesity properties, Antimicrobial, antioxidant, anti-inflammatory, wound healing	Fish cartilage, shrimp shells	29


Open Access Article. Published on 30 September 2025. Downloaded on 10/6/2025 6:44:56 AM.  
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.





Vitamins Vitamin D (cholecalciferol), B12, General health, Fish liver, fish skin  
 A antioxidant properties

24  
 DOI: 10.1039/D5FB00236B



**Figure 1 Bioactive compounds extraction from seafood byproducts**






Figure 2 Fermentation methods for different seafood side streams

Biotechnological approaches in enzymatic hydrolysis and fermentation play a crucial role in utilizing seafood side streams, offering the possibility of retrieving valuable nutrients, minimizing waste, and producing value-added products<sup>53</sup>. In the enzymatic hydrolysis process, enzymes break down proteins in seafood side streams into smaller peptides and amino acids. This eco-friendly and efficient method produces valuable products such as fish protein hydrolysates (FPH), which have applications in food, nutraceuticals, and animal feed. **Table 3** depicts the enzymatic hydrolysis of various seafood side-streams. Fermentation utilizes microbes to convert seafood offcuts into value-added products. This process can produce biofuels, enzymes, and animal feed and is particularly effective in recovering nutritional components such as chitin, chitosan, and fish protein hydrolysates<sup>26</sup>. Since conventional techniques rely on large quantities of organic solvents, consume high amounts of energy, and are time-intensive, they are expected to be gradually replaced by more sustainable extraction approaches. These eco-friendly approaches improve selectivity and offer broader application potential<sup>54</sup>. **Figure 2** shows various fermentation methods used for seafood side streams involving enzymes.

Table 3 Enzymatic hydrolysis of various seafood side streams.



Substrate	Enzyme(s) Used	Hydrolysis Conditions	Yield (%)	Bioactivity Measured	Reference
Tuna waste	Alcalase	pH 8.0, 55°C, 2 h	65	Antioxidant, ACE-inhibitory	51
Salmon skin	Papain, Trypsin	pH 7.5, 50°C, 4 h	60	Collagen peptides, anti-aging	55
Shrimp head	Protamex	pH 7.0, 50°C, 2 h	58	Calcium-binding peptides	56
Cod frame	Flavourzyme	pH 6.5, 50°C, 3 h	62	Antimicrobial activity	57
Mackerel viscera	Neutrase	pH 7.0, 50°C, 1.5 h	70	DPPH radical scavenging capacity	39

View Article Online

DOI: 10.1039/D5FB00236B

### 2.3. Purpose and applications of extracted compounds

Seafood side streams, often considered waste, are increasingly recognized for potential applications across different industries, including food, pharmaceuticals, and cosmetics<sup>42</sup>. These peptides also enhance food ingredient solubility, water holding capacity, and gel formation<sup>58</sup>. Proteins and hydrolysates derived from seafood byproducts are used to develop protein-enriched foods. These proteins have functional properties like emulsification, solubility, and gelling capacities, making them suitable for use in protein-enriched foods, emulsified meat products, and surimi-based formulations. Due to their intense umami profile, peptides derived from enzymatic hydrolysis also serve as flavor enhancers, such as fish solubles used in broths and soups<sup>59</sup>. Fish soluble from side streams can be used as broth and flavor enhancers due to their high flavor intensity<sup>60</sup>. PUFA-rich oils, particularly those containing EPA and DHA, are recovered from fish viscera and heads. These omega-3 fatty acids are essential for cardiovascular and cognitive health and are used in fortified food products, pharmaceutical formulations, and infant nutrition. For instance, Hofseth BioCare (Norway) produces ProGo™, a clinically validated fish protein hydrolysate from salmon backbones and trimmings, which is used in nutritional supplements. Thai Union has commercialized functional tuna essence derived from side streams of tuna processing<sup>61</sup>. Collagen and gelatin, sourced from fish skin, bones and crustacean shells, have growing applications in biomedical engineering, such as scaffold materials for tissue repair and drug delivery systems. In cosmetics, they are integrated into anti-aging creams and serums for their ability to



REVIEW

improve skin hydration, elasticity, and overall appearance<sup>62</sup>. Marine bioactive compounds are utilized for their health benefits, including anti-tumor, anti-microbial, and anti-inflammatory effects, making them valuable in pharmaceuticals<sup>63</sup>. Carotenoids such as astaxanthin and fucoxanthin, derived from crustacean shells and marine algae, possess potent antioxidant and anti-inflammatory properties. These pigments are applied in functional foods, dietary supplements, and cosmetic products, especially those targeting skin health and UV protection. Chitin and chitosan, extracted from shrimp and crab shells, are used in cosmetics for their moisturizing and film-forming properties<sup>64</sup>. Chitosan is used as a drug delivery vehicle and wound healing agent in pharmaceuticals due to its biocompatibility and ability to stimulate tissue regeneration. These compounds also have antimicrobial effects, making them suitable for skin care products<sup>65</sup>. Collagen and peptides from seafood byproducts are incorporated into anti-aging creams and lotions due to their ability to improve skin elasticity and hydration<sup>66</sup>. **Table 4** demonstrates selected case studies of industrial seafood side stream valorization.

**Table 4 Selected Case Studies of Industrial Seafood Side Stream Valorization**

Company Name	Side Stream Used	Valorization Method	Product Developed	Application Area	Country	References
Hofseth BioCare ASA	Salmon heads and bones	Enzymatic hydrolysis	ProGo™ (fish protein hydrolysate)	Clinical nutrition	Norway	61
Seagarden AS	Cod skin and heads	Collagen extraction	Marine collagen powder	Cosmetics, nutraceuticals	Norway	67
Thai Union Group	Tuna cooking water	Filtration and concentration	SEALECT Tuna Essence	Functional beverages	Thailand	68
Ocean Harvest Tech	Seaweed and fish waste	Fermentation	Functional animal feed	Aquaculture	Ireland	69
Primex Iceland	Shrimp shells	Chitosan extraction	High-purity chitosan	Pharmaceuticals, cosmetics, agriculture	Iceland	70
BlueBioChain Project	Various marine by-products	Multiple (R&D initiative)	Enzymes, bioactive peptides, omega-3 oils	R&D, biotechnology	EU-wide	71
Biomega Group	Salmon side streams	Enzymatic hydrolysis	Bioactive peptides and marine oils	Functional foods, pet nutrition	Norway	72



Company Name	Side Stream Used	Valorization Method	Product Developed	Application Area	Country	View Article Online DOI: 10.1039/D5FB00236B
						References
Sofina Foods	Fish skins and bones	Collagen and gelatin recovery	Collagen, gelatin	Food, pharmaceuticals	Canada	73


## 2.4. Economic feasibility and scale-up potential

Valorization of fish byproducts can lead to significant economic benefits by reducing waste management costs and generating additional revenue from high-value products such as fishmeal, oil, gelatin, collagen, and fish protein hydrolysates<sup>74</sup>. The global market for marine collagen was valued at approximately USD 700 million in 2022 and is projected to reach USD 1.2 billion by 2030, growing at a compound annual growth rate (CAGR) of 7.3%. Similarly, the global chitosan market, largely derived from crustacean shells, is expected to exceed USD 3.3 billion by 2027, driven by demand in pharmaceuticals, cosmetics, and biodegradable packaging. The fish protein hydrolysates market is also growing steadily due to increased demand in clinical nutrition and functional food sectors, particularly in Europe and Southeast Asia, which are leading the pace<sup>60</sup>. Establishing valorization processes often requires significant capital investment in infrastructure and technology, which can deter many businesses<sup>75</sup>. Ensuring that the valorization processes are economically viable at an industrial scale is a major challenge. Variability in feedstock can lead to fluctuations in product yield and quality, affecting the overall cost-effectiveness<sup>76</sup>. Valorized products must compete with conventional products that are often cheaper due to established production processes and economies of scale<sup>77</sup>. European nations and the United States are actively exploring the circular economy to advance sustainable valorization, aiming to generate valuable products from waste while maintaining environmental balance<sup>78</sup>. Consumers' preferences for sustainability influence firms' incentives to coordinate the introduction of sustainable product variants, which can lead to multiple equilibria and the idea of sustainable practices "first-mover disadvantage" as an explanation for an agreement between competitors<sup>79</sup>. Embracing an international sustainability standard (ISS) can improve firms' sustainability performance and market value, but beyond a certain point, firms' markets decline, creating a "penalty zone" that acts as an important barrier to further sustainability agenda through additional ISS adoption<sup>80</sup>. Challenges in integrating management tools and the vital role of collaboration between academia and industry call for enhanced support in technology commercialization and methods that promote



industrial sustainability<sup>81</sup>. Developing integrated value chains that connect various stakeholders can improve the efficiency and economic viability of the valorization process<sup>82</sup>.

DOI: 10.1039/D5FB00236B



**Figure 3 Current market distribution of key bioactive compounds recovered from seafood side streams** <sup>56,83–85</sup>

### 3. Integration with Sustainable Development Goals (SDGs)

#### 3.1.SDG 2: Zero Hunger – Contribution to food security and nutrition

By 2030, SDG 2 seeks to purge hunger, strengthen sustainable food access, improve nutrition, and encourage sustainable agricultural practices. This goal is interconnected with other SDGs, requiring a holistic framework to discuss the complex challenges of food security and nutrition<sup>86</sup>. Various agricultural interventions, such as extension services, input subsidies, and value chain enhancements, have been implemented to improve food security. These interventions often include pro-poor features and community engagement. Approximately 67% of evaluated interventions positively impacted food security, 23% showed no measurable impact, and 10% had negative outcomes. The success of these interventions often depends on their design and implementation<sup>87</sup>. Effective pathways to achieve food security include agroecosystem diversification, ecological management, and place-based adaptive solutions. These practices ensure sustainability and resilience in food systems<sup>88</sup>. PUFA-rich oils, particularly those containing EPA and DHA, are vital for cognitive development and cardiovascular health. The inclusion of omega-3 fatty acids in therapeutic foods or nutritional supplements can address micronutrient deficiencies in vulnerable populations, especially among pregnant women and children. Gelatin and collagen, recovered from fish skin and crustacean shells, offer additional amino acid sources and may be incorporated into dietary supplements to support joint health and skin integrity, especially among the elderly and malnourished individuals. Policies promoting healthy and sustainable diets are essential for





achieving nutrition security, including undernutrition and overnutrition, as seen in the European Union's efforts to combat childhood obesity<sup>89</sup>. Achieving zero hunger must be balanced with climate change goals. Pivotal changes in agricultural practices, gender equality, and education are necessary to minimize trade-offs<sup>90</sup>. Addressing these issues through social protection programs and economic development is crucial for long-term success<sup>91</sup>.

### 3.2. SDG 3: Good Health and Wellness – Bioactive compounds and health benefits

Enforcing good health and wellness for all ages is the core of SDG 3, which focuses on maternal and child health, combating diseases, and providing access to essential healthcare services. The goal is interconnected with other SDGs, emphasizing the need for a holistic approach to health and well-being<sup>92</sup>. Bioactive peptides obtained from the enzymatic hydrolysis of fish proteins have shown promising effects in reducing blood pressure, inhibiting free radical activity, and modulating immune responses. They can be incorporated into functional foods, nutraceuticals, and clinical nutrition products targeting cardiovascular diseases, inflammation, and immune disorders<sup>96</sup>. These compounds offer numerous health benefits, such as anti-inflammatory, antioxidant, anticancer, antidiabetic, and antibacterial properties. They are essential in mitigating and controlling chronic diseases like cardiovascular diseases, diabetes, cancer, and neurodegenerative disorders<sup>93</sup>. Fish oils are routinely formulated into omega-3 capsules, fortified foods, and infant formulas, contributing to the prevention of heart disease, Alzheimer's, and cognitive decline. The positive impact of bioactive compounds ensures healthy lives and overall well-being. Developing functional foods and nutraceuticals from these compounds can enhance public health and provide therapeutic advantages<sup>94</sup>. Collagen and gelatin, extracted from marine sources like fish skin and crustacean shells, are widely used in joint and bone health supplements, tissue engineering, and wound healing applications. Collagen peptides have also demonstrated roles in improving skin elasticity, muscle recovery, and gut barrier function. Chitin and chitosan, derived from shrimp and crab shells, possess antimicrobial, anti-obesity, and cholesterol-lowering effects. Due to their biocompatibility and bioactivity, these compounds are employed in dietary supplements, drug delivery systems, and wound dressings. Marine carotenoids like astaxanthin and fucoxanthin exhibit strong antioxidant and anti-inflammatory activities. These are associated with reduced risk of oxidative stress-related diseases, such as cancer and metabolic disorders, and are formulated into nutraceuticals and health beverages. Opportunities for addressing health-related aspects of SDG 3 through bioactive compounds include developing novel delivery systems, such as nano-formulations, to overcome physiological barriers and improve effectiveness in treating cardiovascular diseases<sup>95</sup>.

### 3.3. SDG 6: Clean Water and Sanitation – Reduction of seafood processing waste impact



SDG 6 aims to promote the sustainable harnessing and management of water resources while ensuring sanitation for all. A significant aspect of this goal involves addressing the impact of seafood valorization on water quality and sanitation. The industry is a major consumer of water and energy used in various processes such as cleaning, freezing, and refrigeration. This high consumption results in significant wastewater generation, which poses a challenge for sustainable management<sup>96</sup>. Effluents from marine processing plants can lead to water quality degradation in nearby water bodies, which includes increased biochemical oxygen demand (BOD) levels, chemical oxygen demand (COD), and nutrients like nitrogen and phosphorus<sup>97</sup>. In regions like the Southwest coast of India, seafood processing waste has been linked to hydrogeochemical alterations in shallow aquifers, making the water unsuitable for drinking and posing risks to agricultural use<sup>96</sup>. Implementing advanced wastewater treatment technologies, including biological, chemical, and physical methods, can significantly reduce the environmental footprint of seafood processing plants<sup>96</sup>. Techniques such as bioremediation and absorbing materials can help to treat effluent waters and recover valuable nutrients.

#### 3.4. SDG 9: Industry, Innovation, and Infrastructure – Advancements in valorization technologies

SDG 9 works toward developing resilient infrastructure, ensuring inclusive industrial growth, and fostering a culture of innovation. Advancements in valorization technologies contribute significantly to achieve these objectives by reforming waste and byproducts into valuable resources, thus supporting sustainable industrial practices. Technologies like gasification and anaerobic digestion are being adapted for seafood waste valorization due to their ability to process high-moisture organic residues. Integrated frameworks, supported by AI and sensor-based controls, can optimize feedstock utilization and energy recovery from seafood processing side streams. The development of frameworks like DPSIR (Driver, Pressure, State, Impact, and Response) and the use of interval-valued intuitionistic fuzzy number-based methods can help to address the uncertainties and optimize the valorization process<sup>98</sup>. WaSeaBi Project focuses on valorizing seafood byproducts into sellable products like protein-based food ingredients and bioactive peptides. Technologies such as pH shift, enzymatic hydrolysis, and membrane concentration are employed to extract valuable components<sup>23</sup>. The project NEPTUNUS aims to support the sustainability of the seafood sector by defining eco-innovation approaches and methodologies for eco-labeling products, aligning with the UN SDGs<sup>99</sup>. The use of digital transformation, including Information and Communication Technology (ICT), Internet of Things (IoT), AI, and blockchain, can support the expansion needs of the fisheries industry, contributing to SDG 9<sup>100</sup>. Using new technologies such as Industry 4.0 innovations, offer a pioneering strategy to improve the blue



economy and food sustainability globally<sup>27</sup>. Artificial Intelligence (AI) can significantly enhance seafood valorization by optimizing enzyme loading, fermentation time, and yield prediction using machine learning models trained on historical process data. Blockchain technology ensures transparent traceability of valorized products, documenting each stage from raw material origin to end-use application. Coupled with IoT sensors, these tools can monitor pH, temperature, or contamination in real-time, ensuring consistent quality in hydrolysates, peptides, and oils derived from seafood residues. AI-enabled image recognition can also aid in real-time sorting and segregation of seafood waste streams, improving efficiency and reducing cross-contamination<sup>101</sup>.

### 3.5. SDG 12: Responsible Consumption and Production – Circular economy and waste reduction

The circular economy framework is integral to achieve SDG 12, which focuses on responsible consumption and production. A circular economy aims at practices such as recycling, reusing, and reducing waste generation. For instance, fish gelatin and chitosan extracted from seafood waste can be used to create biodegradable food packaging materials, reducing reliance on synthetic plastics<sup>102</sup>. Biological methods, such as microbial fermentation and algal biotechnology, offer sustainable ways to convert seafood waste into valuable products while maintaining their bioactivity. These processes can produce biofuels and other high-value chemical feedstock, promoting a circular bioeconomy within the seafood sector<sup>103</sup>. The Fish Waste-based Eco-Industrial Park (FWEIP) model aims to convert linear waste management into a circular system, favoring biofuel production and other sustainable practices<sup>104</sup>. Circular economy practices in seafood processing can mitigate environmental pollution, reduce emissions of greenhouse gases, and support the sustainability of aquatic resources<sup>105</sup>.

### 3.6. SDG 14: Life Under Water – Sustainable fisheries and ecosystem preservation

Sustainable fisheries face several setbacks, including overfishing, forbidden, unreported, and unregulated (IUU) fishing, and the readiness of stakeholders to adopt sustainable practices. Valorizing seafood byproducts into bioactive compounds, nutraceuticals, cosmetics, and feed ingredients reduces environmental impact, promotes a circular bioeconomy, minimizes overharvesting, and supports biodiversity conservation<sup>102</sup>. Moreover, it alleviates the pressure on marine habitats and ecosystems by preventing the illegal dumping of discards and limiting eutrophication and habitat degradation. Effective governance and stakeholder engagement are crucial for achieving sustainable fisheries<sup>106</sup>. Regional fisheries management organizations play a significant role in managing international fisheries and can contribute to ending overfishing and protecting marine areas. However, there is a need for better cooperation with other maritime organizations and more comprehensive management measures<sup>27</sup>. Establishing Marine Protected



Areas (MPAs) can protect critical habitats and biodiversity. Fully and partially protected areas contribute significantly to achieve SDG 14 targets<sup>107</sup>. Achieving SDG 14 can bring significant economic benefits, particularly for small-scale fisheries and developing countries that depend on marine resources for revenue and sustainable economic growth<sup>108</sup>. Using marine resources through sustainability can contribute to economic growth, livelihoods, and community sustainability, aligning with the broader economic goals outlined in the SDGs<sup>109</sup>. Implementing Ecosystem-Based Fisheries Management (EBFM) can help to balance the needs of fisheries with the health of marine ecosystems, ensuring long-term sustainability. **Table 5** points out key actions to restore the seafood system.

**Table 5 Key actions to restore the seafood system**

Key actions	Suggestions
Increase product efficiency	Innovations and growth
Increase biodiversity	Provide technical and financial support for regenerative practices with added environmental benefits.
Explore new food sources	Close collaboration with business partners and consumers within specific regions guarantees natural ingredient volumes and quality.
Reduce waste	Re-cycle waste and up-cycle byproducts
Reduced energy system	Encourage net-zero emissions, allow carbon sequestration and efficient water usage.

Source: Adapted from Knorr and Augustine<sup>110</sup>

**4. Future Perspectives and Research Directions**

The future of seafood side stream valorization lies in developing and integrating innovative technologies, addressing environmental and economic challenges, and overcoming technological and market barriers. Future research should focus on scaling up the pH-shift method technology, membrane concentration, flocculation technologies, and integrating renewable energy sources to enhance sustainability<sup>47</sup>. Biorefineries integrate various technologies such as enzymatic treatments, microbial fermentation, and extraction by green solvents like ionic liquids and deep eutectic solvents, among others<sup>111</sup>. Studies have highlighted the importance of reducing chemical consumption and improving energy efficiency to optimize the environmental performance of valorization technologies<sup>23</sup>. A five-tier model suggests applications for seafood side-streams in medicine, food manufacturing, animal feed, fertilizers, and energy fuel, maximizing their potential and reducing waste<sup>42</sup>. Incorporating renewable energy sources can further enhance the



sustainability of valorization processes<sup>23</sup>. Combining different treatment methods based on the composition and properties of side-streams can improve environmental performance<sup>112</sup>. Increasing awareness and acceptance among stakeholders is essential for successfully implementing valorization strategies<sup>50</sup>. Continued research and development, along with stakeholder engagement, are key to achieve a sustainable and circular seafood industry.

#### 4.1. Technical and processing limitations to sustainable valorisation

Sustainable valorization, the technique of converting waste into value-added products, faces several technical and processing challenges that hinder its widespread adoption and efficiency. The inconsistency in the composition of waste materials, such as waste generated from food, affects the efficiency and yield of valorization processes. This variability can lead to fluctuations in product quality and process performance<sup>75</sup>. Effective valorization requires precise separation and classification of waste materials, which is technically challenging, especially for mixed waste streams like plastic films and food waste<sup>113</sup>. Hydrothermal carbonization and microbial electrosynthesis are promising novel technologies, but they are still in the developmental stage and face scalability issues<sup>114</sup>. While pyrolysis may require significant energy inputs, anaerobic digestion is a net energy-producing process due to the generation of biogas, offering a sustainable solution for seafood waste management<sup>115</sup>. The regeneration of green sorbents used in pollutant mitigation from marine-derived wastes presents technical challenges, particularly in maintaining their sorption capacity and surface functionality over multiple cycles<sup>57</sup>. Significant investment in infrastructure and equipment is necessary to implement valorization technologies at an industrial scale, which includes the need for specialized facilities and machinery, which can be a major financial barrier<sup>116</sup>. The complexity of supply chains for sustainable chemical technologies adds another layer of difficulty, requiring coordination among various stakeholders to ensure smooth operation and integration<sup>117</sup>. **Table 6** represents the category-wise challenges in the sustainable valorization.

**Table 6 Challenges in sustainable valorization**

Category	Changes	Reference
Regulatory	Lack of clear regulations, over-regulation.	118
Technological	Path-dependence lock-ins, infrastructure requirements.	119
Economic	Financial constraints, and high operational costs.	120
Social	Stakeholder involvement, market acceptance.	121

#### 4.2. Regulatory and policy constraints

**Table 7** depicts policy recommendations for sustainable valorization. The seafood industry yearly generates metric tons of offsets, overwhelmingly from shrimp, crab, and lobster shells, leading to



environmental impacts due to COD and BOD issues<sup>42</sup>. Implementing stringent traceability regulations like those in the EU and the US can create conflicts between extraterritorial fisheries management and local market access. These regulations require detailed information on when, where, what, who, and how seafood is produced, which can complicate the valorization process<sup>122</sup>. The EU's Common Fisheries Policy reform mandates that fishers land all catches, including unwanted ones, instead of discarding them at sea. It can create a need for efficient management of low-value marine biomass, which can be challenging due to the lack of infrastructure and economic incentives<sup>123</sup>. The Marine Stewardship Council (MSC) certification has been shown to have implications for companies' value creation processes, with a disparity in economic outcomes of MSC-certified businesses over those that were not, indicating the economic benefits of sustainable fishing practices<sup>9</sup>. **Table 8** suggests protocols to protect the oceans and marine life.

**Table 7 Policy recommendations for sustainable valorization**

Policy recommendations for sustainable valorization	Actions to be taken	References
Implement Sustainable Fishing Practices	Enforce the Common Fisheries Policy (CFP) to ensure sustainable fishing through conservation measures, fleet management, and by-catch strategies. Adopt individual quota systems based on total allowable catch and maximum sustainable yield.	124
Enhance Traceability and Transparency	Implement policies to minimize illegal, unreported, and unregulated (IUU) fishing. Improve catch documentation and supply chain transparency.	125
Promote Certification and Eco-labeling	Encourage the use of third-party certifications and eco-labels like the Marine Stewardship Council (MSC) and Seafood Watch. Develop government-operated certification programs based on established principles.	124
Foster Consumer Awareness and Education	Launch media campaigns and partnerships with advocacy groups to raise consumer awareness about sustainable seafood. Develop school meal programs and hospitality menus featuring sustainable seafood.	125





Support Sustainable Aquaculture	Promote sustainable aquaculture practices, including the use of eco-friendly feed and minimizing environmental impacts. Expand sustainable coastal mariculture to enhance food security and economic resilience.	
Valorize Seafood Processing Waste	Implement bioconversion and biorefinery processes to convert seafood waste into products like biofuels, omega-3 fatty acids, and nutraceuticals. Develop circular bioeconomies within the seafood processing sector.	103
Integration with Food Security Goals	Apply Marine Spatial Planning (MSP) and EBM to protect marine ecosystems and biodiversity. Mainstream aquatic biodiversity in sectoral policies and planning.	126
Develop Sustainable Supply Chains	Transform supply chains to ensure sustainability from production to consumption. Support policies and regulations that facilitate sustainable commoditization of seafood.	127
Minimize Discards and Optimize By-catch Utilization	Implement guidelines to minimize discards and valorize unavoidable by-catch through efficient management networks. Develop protocols for the optimal valorization of discarded species.	128
Address Social Sustainability	Ensure fair wages and working conditions in the seafood industry. Endorse local seafood as a fresh, sustainable, and ethical choice.	129
Provide Economic Incentives	Introduce tax rebates and green loans for companies investing in seafood valorization infrastructure.	126
Strengthen Certification Systems	Expand adoption of MSC (Marine Stewardship Council) and eco-labeling schemes for valorized products.	127
Harmonize Waste Utilization Standards	Align national seafood waste reuse standards with international FAO and EU circular economy frameworks.	128

**Table 8 Protocols to protect the oceans and marine life**

Protocols	Advantages
-----------	------------



Preventing overfishing, catch restrictions, seasonal restrictions, and bycatch reductions	Ensure the fisheries population remains at a sustainable level.	View Article Online DOI: 10.1039/D5FB00236B
Establishing marine reserves	Provide havens for marine life, allowing the ecosystem to thrive without human interference. Helps preserve biodiversity, supports fish stock recovery, and the resilience of marine habitats.	
Promoting aquaculture	Allows a sustainable source of alternate seafood with eco-friendly practices, disease management, and feed sustainability	
Educating fishermen	Provides awareness of the benefits of sustainable practices and provides training on alternative livelihoods	
Educating consumers	It provides awareness of the importance of choosing sustainably sourced seafood. Can drive demand for responsible fishing practices and foster sustainability in the seafood industry	
Supporting policy and regulation	Essential for enforcing sustainable fishing practices and protecting marine ecosystems	
Research and innovation	Developments in innovative practices for sustainable protocols are highly beneficial.	

Source: Adapted from Sustainable Fisheries: 6 Powerful Ways to Protect Our Oceans and Marine Life, <https://ecochatters.com/sustainable-fisheries-ways-to-protect-marine-life/>

### 4.3.Consumer perception and acceptance

Research indicates that taste, freshness, and source attributes often influence seafood-related behavior. To shift consumer tendency toward green alternatives, coastal communities promote manufacturing and source attributes of local seafood, such as being wild-caught, ecologically sustainable, and harvested locally. However, consumers may lack confidence in distinguishing these attributes when purchasing seafood, indicating a gap between the importance and confidence in seafood attributes<sup>130</sup>. Customer impressions and retail choices are highly situational and vary significantly across different cultures and countries. To address consumer scepticism, transparent labeling with traceable QR codes, displaying the origin and transformation journey of seafood side streams into value-added products, can enhance consumer confidence. Public awareness campaigns featuring nutritionists, sustainability influencers, and testimonials have proven effective in shifting public perceptions. Additionally, integrating byproducts derived from seafood side-stream into



recognizable formats such as soups, bars, or dietary supplements can improve acceptance, lower psychological resistance, and improve market adoption<sup>131</sup>. This variability makes it challenging to develop universally accepted sustainable seafood byproducts. The limited availability of clear and comprehensible data on the sustainability and benefits of seafood byproducts is a major barrier to social acceptance<sup>132</sup>. Effective communication strategies are essential to overcome this barrier. Adding new production attributes, such as "farmed in marine waters," increases the difficulty of consumer choices, which can further hinder acceptance<sup>130</sup>. Improving marketing and consumer education about the benefits and safety of sustainable seafood products can serve as a link between importance and confidence. Developing targeted communication strategies that consider cultural and contextual differences can help improve consumer perceptions and acceptance of sustainable seafood products<sup>132</sup>.

#### 4.4. Industry-academia collaborations for innovation

Industry-academia collaborations contribute greatly to the innovation and valorization of seafood side streams, transforming what was once considered waste into valuable products. Projects like WaSeaBi, funded by the EU, focus on developing innovative technologies to efficiently valorize seafood off-cuts into market-based products such as high-protein food ingredients, bioactive peptides, and mineral supplements<sup>23</sup>. Effective communication, trust, and adequate funding are essential for successful university-maritime industry collaborations (UMICs). These elements help to build strong partnerships and support the development of new technologies and solutions. Collaborations can lead to significant environmental benefits by reducing waste and improving sustainability. For example, the WaSeaBi project demonstrated that minimizing chemical consumption and enhancing energy efficiency is key to evaluate valorization technologies' ecological proficiency. Economically, these collaborations can enhance gear efficiency, create new fishery opportunities, and sustain fishing operations<sup>42</sup>. Decision-making tools developed within these collaborations help industry stakeholders select appropriate valorization strategies by considering technical, legal, economic, and environmental aspects. Successful collaborations, such as the development of the CTD (Conductivity, Temperature, Depth) tag by SMRU (Sea Mammal Research Unit) and Valeport, highlight the mutual benefits for academia and industry. These partnerships can lead to rapid technological advancements and improved product performance<sup>133</sup>. Despite the benefits, collaborations often face challenges, such as differing agendas and goals. Addressing these challenges requires clear communication, mutual respect, and trust. Guidelines and best practices have been developed to promote sustainable and respectful collaborative research.



#### 4.5. Prospects of integrating digital technologies

View Article Online  
DOI: 10.1039/D5FB00236B

Artificial Intelligence (AI) can significantly enhance the efficiency and scalability of seafood byproduct valorization. AI can optimize the extraction parameters for bioactive compounds from seafood byproducts, such as enzyme/substrate ratio, pH, time, temperature, and performance of extraction instruments. This optimization can lead to higher yields of valuable compounds like polyunsaturated fatty acids, amino acids, peptides, and collagen<sup>134</sup>. Machine learning algorithms can streamline these processes by predicting optimal conditions and adjusting parameters in real-time. AI technologies, including machine learning and computer vision, can monitor the quality of extracted bioactive compounds. These systems can analyze large volumes of data to ensure consistent product quality and detect anomalies or deviations from optimal conditions. AI-driven monitoring systems can provide real-time alerts, allowing for timely interventions to maintain high standards<sup>135</sup>. Predictive analytics powered by AI can forecast the yield of bioactive compounds based on historical data and current process conditions. This capability helps in planning and optimizing production schedules, ensuring efficient resource utilization and minimizing waste<sup>136</sup>. AI models can achieve high accuracy in predicting yields, as demonstrated in bioethanol production optimization. AI can classify and manage waste streams effectively, identifying valuable components within seafood byproducts and directing them towards appropriate valorization processes. This classification can reduce environmental impact and enhance the sustainability of seafood processing. AI-driven systems can also optimize waste management practices, ensuring that valuable bioactive compounds are not lost<sup>49</sup>.

Blockchain's immutable ledger allows for real-time data sharing, which enhances traceability and transparency in the seafood supply chain. This capability helps to verify the origins of seafood products, which ensures compliance with sustainability standards, and combat illegal, unreported, and unregulated (IUU) fishing. By reducing the need for intermediaries, blockchain can lower transaction costs and increase profit margins for fishermen and other stakeholders in the supply chain<sup>137</sup>. The integration of blockchain with other technologies like AI and IoT can improve sustainability practices by enabling better resource allocation, ecosystem modeling, and intelligent monitoring systems<sup>138</sup>. Regulatory uncertainty and the need for reforms in regulatory compliance are major factors affecting the adoption of blockchain technology in the seafood industry<sup>139</sup>. Incorporating De-Fi solutions can provide fishermen with better access to capital and global markets, enhancing the economic viability of blockchain projects in the seafood industry. Using automation and IoT in data collection can improve the quality and trust in data, further enhancing the benefits of blockchain technology. Blockchain can support sustainable practices by enabling



better waste management techniques, innovative packaging solutions, and energy-efficient methods<sup>140</sup>.

DOI: 10.1039/D5FB00236B

## 5. Conclusion

This review highlights the promising aspects of seafood side streams in promoting sustainability through valorization strategies that convert waste into high-value-added products. Bioactive compounds derived from aquatic food side streams offer diverse applications in the food, pharmaceutical, nutraceutical, and cosmetic industries. Emerging technologies such as enzymatic hydrolysis and microbial fermentation have demonstrated promising results in extracting valuable compounds while minimizing environmental impact. However, technological limitations, regulatory hurdles, and market acceptance challenges should be addressed to meet the requirements of sustainable development goals, particularly SDG 12.

Policies should support circular economy principles, green processing technologies, and transparent supply chains to enhance sustainable seafood byproduct utilization. Strengthening anti-IUU (Illegal, Unreported, and Unregulated) fishing policies, promoting certification programs, and encouraging sustainable aquaculture practices are crucial. Industry stakeholders, including seafood processors and biotechnology firms, must collaborate to invest in research, infrastructure, and innovation for effective seafood side-stream valorization. Government incentives and funding for sustainable initiatives will further drive adoption.

Achieving sustainable seafood valorization requires a multidisciplinary effort involving biotechnologists, policymakers, environmental scientists, and industry leaders. Further research should emphasize developing cost-effective, scalable, and energy-efficient technologies. Artificial intelligence and blockchain integration can improve traceability and market transparency. Additionally, consumer awareness and education are pivotal in fostering demand for sustainable seafood-derived products. By embracing an interdisciplinary and collaborative approach, the seafood industry can transform towards a circular bioeconomy, aligning with SDGs and ensuring long-term environmental and economic benefits.

## Author contributions

**Sreedharan Advaita** – design of methodology, acquisition of data, analysis, and interpretation, and writing the original draft, **Kappat Valiyapeediyekkal Sunooj** – design of methodology, acquisition of data, review, editing of the draft, supervision and project administration, **Sathish kumar vellaisamy** -



review, **Sahil Negi** – review, **Muhammed Navaf** – review and editing of the draft, **Abhilash Sasidharan** – review and editing of the draft, **Vazhiyil Venugopal** – idea conceptualization, review and editing of the draft.

DOI: 10.1039/D5FB00236B

## Conflicts of interest

There are no conflicts to declare.

## Data availability

No primary research results, software, or code have been included, and no new data were generated or analysed as part of this review.

## Acknowledgements

This review work did not receive any specific grant from public, commercial, or not-for-profit funding agencies. The authors acknowledge the DST-FIST LEVEL B (2023-2028) Department of Food Science and Technology, Pondicherry University for providing facilities.

## References

- 1 F. Ozogul, M. Cagalj, V. Šimat, Y. Ozogul, J. Tkaczewska, A. Hassoun, A. A. Kaddour, E. Kuley, N. B. Rathod and G. G. Phadke, Recent developments in valorisation of bioactive ingredients in discard/seafood processing by-products, *Trends Food Sci Technol*, 2021, **116**, 559–582.
- 2 M. E. Lam, The Ethics and Sustainability of Capture Fisheries and Aquaculture, *J Agric Environ Ethics*, 2016, **29**, 35–65.
- 3 D. Knorr and M. A. Augustin, Food systems restoration, *Sustainable Food Technology*, 2024, **2**, 1365–1390.
- 4 Food and Agriculture Organization of the United Nations Rome, *The future of food and agriculture: alternative pathways to 2050*, FAO Rome, Italy, 2018, vol. 60.
- 5 M. H. Wong, D. Purchase and N. Dickinson, *Impacts, Management, and Recycling of Food Waste: Global Emerging Issues*, 2023.
- 6 I. F. Policy Research Institute, *Global food policy report 2023: Rethinking food crisis responses*, 2023.
- 7 T. M. Garlock, F. Asche, J. L. Anderson, H. Eggert, T. M. Anderson, B. Che, C. A. Chávez, J. Chu, N. Chukwuone, M. M. Dey, B. Tibesigwa and R. Tveteras, Environmental, economic, and social sustainability in aquaculture: the aquaculture performance indicators, *Nat Commun*, DOI:10.1038/s41467-024-49556-8.





- 8 K. M. Blincow, A. C. Haynie and B. X. Semmens, Disparities between sustainability of country-level seafood production and consumption, *PLoS One*, DOI:10.1371/journal.pone.0313823. DOI: 10.1039/D5FB00236B
- 9 L. Miret-Pastor, T. Peiró-Signes, M.-D.-V. Segarra-Oña and P. Herrera-Racionero, Empirical analysis of sustainable fisheries and the relation to economic performance enhancement: The case of the Spanish fishing industry, *Mar Policy*, 2014, **46**, 105–110.
- 10 R. Hilborn, E. A. Fulton, B. S. Green, K. Hartmann, S. R. Tracey and R. A. Watson, When is a fishery sustainable?, *Canadian Journal of Fisheries and Aquatic Sciences*, 2015, **72**, 1433–1441.
- 11 E. Lockerbie, T. G. Loureiro, A. J. C. Schramm, J. Gacutan, I. Yulianto, A. Rosdiana and F. A. K. Kurniawan, in *Handbook of Sustainable Blue Economy*, Springer Nature Switzerland, Cham, 2024, pp. 1–33.
- 12 T. Vella, L. Roberson, C. Kuempel and C. Klein, Quantifying the accessibility of sustainable seafood in south-eastern Queensland, Australia, *Mar Freshw Res*, 2023, **74**, 1340–1354.
- 13 S. Ardra and M. K. Barua, Halving food waste generation by 2030: The challenges and strategies of monitoring UN sustainable development goal target 12.3, *J Clean Prod*, 2022, **380**, 135042.
- 14 J. Cropotova, K. Kvangarsnes, G. H. Aas, S. Tappi and T. Rustad, *Protein from seafood*, 2023.
- 15 A. Charles, *Sustainability and resilience*, 2023.
- 16 L. R. Luttenberger, I. Ančić and A. Luttenberger, The challenges for croatian fisheries within current regulatory environment, *Pomorstvo*, 2021, **35**, 170–178.
- 17 N. Stern and A. Valero, Innovation, growth and the transition to net-zero emissions, *Res Policy*, 2021, **50**, 104293.
- 18 C. Costello, L. Cao, S. Gelcich, M. Á. Cisneros-Mata, C. M. Free, H. E. Froehlich, C. D. Golden, G. Ishimura, J. Maier, I. Macadam-Somer, T. Mangin, M. C. Melnychuk, M. Miyahara, C. L. de Moor, R. Naylor, L. Nøstbakken, E. Ojea, E. O'Reilly, A. M. Parma, A. J. Plantinga, S. H. Thilsted and J. Lubchenco, The future of food from the sea, *Nature*, 2020, **588**, 95–100.
- 19 M. Crippa, E. Solazzo, D. Guizzardi, F. Monforti-Ferrario, F. N. Tubiello and A. Leip, Food systems are responsible for a third of global anthropogenic GHG emissions, *Nat Food*, 2021, **2**, 198–209.
- 20 *FAO. 2024. The State of Food and Agriculture 2024 – Value-driven transformation of agrifood systems. Rome.*, FAO, 2024.
- 21 *The State of World Fisheries and Aquaculture 2020.*, Food and Agriculture Organization of the United Nations, 2018.
- 22 R. Carvalho, M. R. Lucas and A. Marta-Costa, Food Waste Reduction: A Systematic Literature Review on Integrating Policies, Consumer Behavior, and Innovation, *Sustainability*, 2025, **17**, 3236.
- 23 E. Cadena, O. Kocak, J. Dewulf, B. Iñarra, C. Bald, M. Gutierrez, D. San Martin, J. Ibarruri, A.-D. M. Sørensen, G. Hyldig, G. Bruggeman and C. Jacobsen, Valorisation of Seafood Side-Streams through the Design of New Holistic Value Chains: WaSeaBi Project, *Sustainability (Switzerland)*, DOI:10.3390/su16051846.



## REVIEW

## Sustainable food technology

- 24 V. Venugopal, A. Sasidharan and T. Rustad, Green Chemistry to Valorize Seafood Side Streams: An Ecofriendly Roadmap toward Sustainability, *J Agric Food Chem*, 2023, **71**, 17494–17509. DOI: 10.1039/D5FB00236B
- 25 S. A. Siddiqui, H. Schulte, D. Pleissner, S. Schönfelder, K. Kvangarsnes, E. Dauksas, T. Rustad, J. Cropotova, V. Heinz and S. Smetana, Transformation of Seafood Side-Streams and Residuals into Valuable Products, *Foods*, DOI:10.3390/foods12020422.
- 26 S. A. Siddiqui, D. Lakshmikanth, C. Pradhan, Z. Farajinejad, R. Castro-Muñoz and A. Sasidharan, Implementing fermentation technology for comprehensive valorisation of seafood processing by-products: A critical review on recovering valuable nutrients and enhancing utilisation, *Crit Rev Food Sci Nutr*, 2025, **65**, 964–991.
- 27 A. Hassoun, J. Cropotova, H. Trollman, S. Jagtap, G. Garcia-Garcia, C. Parra-López, N. Nirmal, F. Özogul, Z. Bhat, A. Aït-Kaddour, A. Aït-Kaddour and G. Bono, Use of industry 4.0 technologies to reduce and valorize seafood waste and by-products: A narrative review on current knowledge, *Curr Res Food Sci*, DOI:10.1016/j.crfs.2023.100505.
- 28 N. Warwas, M. Langeland, J. A. C. Roques, M. Montjouridès, J. Smeets, H. Sundh, E. Jönsson and K. Sundell, Fish processing side streams are promising ingredients in diets for rainbow trout (*Oncorhynchus mykiss*) — Effects on growth physiology, appetite, and intestinal health, *J Fish Biol*, 2025, **106**, 75–92.
- 29 A.-L. Välimaa, S. Mäkinen, P. Mattila, P. Marnila, A. Pihlanto, M. Mäki and J. Hiidenhovi, Fish and fish side streams are valuable sources of high-value components, *Food Quality and Safety*, 2019, **3**, 209–226.
- 30 N. M. Stephen, T. Maradagi, Y. P. Kavalappa, H. Sharma and G. Ponesakki, *Seafood nutraceuticals: Health benefits and functional properties*, 2021.
- 31 M. Hayes and J. Valverde, *Nutraceuticals derived from marine sources: Extraction and application of bioactive rates, proteins and peptides as functional ingredients*, 2014.
- 32 K. Mohan, T. Muralisankar, R. Jayakumar and C. Rajeevgandhi, A study on structural comparisons of  $\alpha$ -chitin extracted from marine crustacean shell waste, *Carbohydrate Polymer Technologies and Applications*, 2021, **2**, 100037.
- 33 J. M. J. Rodríguez-Jiménez, L. M. Anaya-Esparza, E. Martínez-Montaño, E. Montalvo-González and M. L. García-Magaña, Bioactive peptides from by-products of shrimp processing: A review, *Int Food Res J*, 2024, **31**, 530–550.
- 34 A. Sasidharan, T. Rustad and G. M. Cusimano, Tuna sidestream valorization: a circular blue bioeconomy approach, *Environmental Science and Pollution Research*, 2023, **31**, 62230–62248.
- 35 B. Zieniuk and A. Fabiszewska, Fish and fish wastes as source of bioactive compounds and raw material for energy industry, *Zywnosc Nauka Technologia Jakosc/Food Science Technology Quality*, 2018, **114**, 5–16.
- 36 S. Patil, N. K. Rajendran, U. Bhardwaj and L. Bareja, Size-Dependent Biochemical Properties of Fish Skin: A Focus on Collagen and Protein Content, *Nat Eng Sci*, 2025, **10**, 457–467.
- 37 Kirti and S. S. Khora, *Marine fish-derived proteins and peptides as potential antioxidants*, 2022.



- 38 M. Veeraiyan, J. Somasundaram, S. Poorni, R. Sujith, G. J. Thirukonda and D. Thomas, **Lobster Shells: A Sustainable Source of Biogenic Materials and Their Extractions and Applications**, *UTTAR PRADESH JOURNAL OF ZOOLOGY*, 2024, **45**, 644–651. DOI: 10.1039/D5FB00236B
- 39 X. Wang, H. Yu, R. Xing, X. Chen, S. Liu and P. Li, Optimization of antioxidative peptides from mackerel (*Pneumatophorus japonicus*) viscera, *PeerJ*, 2018, **6**, e4373.
- 40 B. Bayraklı and S. Yıldız, Comparative Analysis of Nutritional Values of Fishmeals Produced From Whole Anchovy and Sprat and Farmed Salmon Viscera in the Black Sea Region, *Acta Natura et Scientia*, 2024, **5**, 150–159.
- 41 M. B. Jensen, J. Jakobsen, C. Jacobsen, J. J. Sloth, J. Ibarruri, C. Bald, B. Iñarra, N. Bøknæs and A.-D. M. Sørensen, Content and Bioaccessibility of Minerals and Proteins in Fish-Bone Containing Side-Streams from Seafood Industries, *Mar Drugs*, DOI:10.3390/md22040162.
- 42 R. Anbarasan, B. K. Tiwari and R. Mahendran, *Upcycling of seafood side streams for circularity*, 2024, vol. 108.
- 43 J. A. Libra, K. S. Ro, C. Kammann, A. Funke, N. D. Berge, Y. Neubauer, M.-M. Titirici, C. Fühner, O. Bens, J. Kern and K.-H. Emmerich, Hydrothermal carbonization of biomass residuals: a comparative review of the chemistry, processes and applications of wet and dry pyrolysis, *Biofuels*, 2011, **2**, 71–106.
- 44 S. Kannan, Y. Gariepy and G. S. V. Raghavan, Optimization and characterization of hydrochar produced from microwave hydrothermal carbonization of fish waste, *Waste Management*, 2017, **65**, 159–168.
- 45 J. G. Lynam, M. T. Reza, W. Yan, V. R. Vásquez and C. J. Coronella, Hydrothermal carbonization of various lignocellulosic biomass, *Biomass Convers Biorefin*, 2015, **5**, 173–181.
- 46 R. A. Nazeer, I. Joshi, S. Mahankali, A. Mazumdar, B. Sridharan and S. J. Sankar, *Is Marine Waste a Boon or Bane? An Insight on Its Source, Production, Disposal Consequences, and Utilization*, 2023.
- 47 E. Cadena, O. Kocak, J. Dewulf, I. Undeland and M. Abdollahi, Environmental Sustainability Assessment of pH-Shift Technology for Recovering Proteins from Diverse Fish Solid Side Streams, *Sustainability (Switzerland)*, DOI:10.3390/su17010323.
- 48 V. Venugopal, Valorization of Seafood Processing Discards: Bioconversion and Bio-Refinery Approaches, *Front Sustain Food Syst*, DOI:10.3389/fsufs.2021.611835.
- 49 F. Ling Wen Xia, S. Supri, H. Djamaludin, R. Nurdiani, L. Leong Seng, K. Wee Yin and K. Rovina, Turning waste into value: Extraction and effective valorization strategies of seafood by-products, *Waste Management Bulletin*, 2024, **2**, 84–100.
- 50 R. Cooney, D. B. de Sousa, A. Fernández-Ríos, S. Mellett, N. Rowan, A. P. Morse, M. Hayes, J. Laso, L. Regueiro, A. H. Wan, A. H. Wan and E. Clifford, A circular economy framework for seafood waste valorisation to meet challenges and opportunities for intensive production and sustainability, *J Clean Prod*, DOI:10.1016/j.jclepro.2023.136283.
- 51 V. Venugopal and S. Kim, Mitigation of Seafood-Related Environmental Pollution: A Green Chemistry Perspective, *Green and Low-Carbon Economy*, DOI:10.47852/bonviewGLCE52023459.



## REVIEW

## Sustainable food technology

- 52 M. Peydayesh, M. Bagnani, W. L. Soon and R. Mezzenga, Turning Food Protein Waste into Sustainable Technologies, *Chem Rev*, 2023, **123**, 2112–2154. Available Online  
DOI: 10.1039/D5FB00236B
- 53 S. A. Ismail and A. A. Hassan, Utilization of seafood byproducts: biological activities and biotechnological applications, *Egypt J Chem*, 2022, **65**, 1531–1546.
- 54 L. C. Freitas, J. R. Barbosa, A. L. C. da Costa, F. W. F. Bezerra, R. H. H. Pinto and R. N. de Carvalho Junior, From waste to sustainable industry: How can agro-industrial wastes help in the development of new products?, *Resour Conserv Recycl*, 2021, **169**, 105466.
- 55 Y. Zhao, F. Wang, Z. Zhang, Y. Pan, J. Ou and H. Liu, Research progress and prospect of artificial seafood | 人造海鲜的研究进展与展望, *Journal of Fisheries of China*, 2021, **45**, 306–316.
- 56 T. Silva, J. Moreira-Silva, A. Marques, A. Domingues, Y. Bayon and R. Reis, Marine Origin Collagens and Its Potential Applications, *Mar Drugs*, 2014, **12**, 5881–5901.
- 57 K. Y. Cheong, S. H. Kong, S. Y. Foong, P. N. Yuh Yek, T.-B. Nguyen, N. L. Ma, C.-D. Dong and S. S. Lam, Valorization of marine-derived wastes as green sorbents, *Curr Opin Green Sustain Chem*, DOI:10.1016/j.cogsc.2024.100922.
- 58 I. Ucak, M. Afreen, D. Montesano, C. Carrillo, I. Tomasevic, J. Simal-Gandara and F. J. Barba, Functional and Bioactive Properties of Peptides Derived from Marine Side Streams, *Mar Drugs*, DOI:10.3390/MD19020071.
- 59 S. Smaoui, E. Hosseini, Z. T. Tsegay, T. D'Amore and T. Varzakas, Sustainable valorisation of seafood-derived proteins: Current approaches for recovery and applications in biomedical and food systems, *Food Biosci*, DOI:10.1016/j.fbio.2024.105450.
- 60 T. Aspevik, T. A. Samuelsen, M. Øvrur Gaarder and Å. Oterhals, Sensory Properties and Chemical Composition of Fish Solubles Obtained from Upcycling of Fish Filleting Side Streams, *Journal of Aquatic Food Product Technology*, 2023, **32**, 336–348.
- 61 <https://www.dkshdiscover.com/en/promo/fbi-my-hofseth?utm>.
- 62 S.-K. Kim, *Seafood processing by-products: Trends and applications*, 2013, vol. 9781461495.
- 63 M. Atef and S. Mahdi Ojagh, Health benefits and food applications of bioactive compounds from fish byproducts: A review, *J Funct Foods*, 2017, **35**, 673–681.
- 64 C. Espinales, M. Romero-Peña, G. Calderón, K. Vergara, P. J. Cáceres and P. Castillo, Collagen, protein hydrolysates and chitin from by-products of fish and shellfish: An overview, *Heliyon*, DOI:10.1016/j.heliyon.2023.e14937.
- 65 T. S. Trung, N. Van Hoa, P. T. D. Phuong, N. C. Minh, N. T. H. Mai, H. N. Cuong, P. T. Loc and W. F. Stevens, *Recovery of bioactive components from seafood by-products toward zero-waste processing*, 2020.
- 66 Y. Zou, M. Heyndrickx, J. Debode, K. Raes, D. de Pascale, P. Behan, M. Giltrap, C. O'Connor, R. G. Solstad, K. Lian, I. Undeland and J. Robbens, Valorisation of crustacean and bivalve processing side streams for



industrial fast time-to-market products: A review from the European Union regulation perspective, *Front Mar Sci*, DOI:10.3389/fmars.2023.1068151. Frontiers in Marine Science Online  
DOI: 10.1039/D5FB00236B

- 67 <https://seagarden-norway.com/>.
- 68 <https://investor.thaiunion.com/>.
- 69 <https://oceanharvesttechnology.com/>.
- 70 <https://www.primex.is/>.
- 71 <https://jpi-oceans.eu/en/bluebiochain>.
- 72 [https://seafood.media/fis/companies/details.asp?l=e&company\\_id=159443](https://seafood.media/fis/companies/details.asp?l=e&company_id=159443).
- 73 <https://www.sofinafoods.com/>.
- 74 C. Lopes, L. T. Antelo, A. Franco-Uría, A. A. Alonso and R. Pérez-Martín, Valorisation of fish by-products against waste management treatments - Comparison of environmental impacts, *Waste Management*, 2015, **46**, 103–112.
- 75 U. Danya, M. Kamaraj, P. S. Babu and J. Aravind, Bibliometric analysis and review of food waste management via sustainable approaches, *Global Journal of Environmental Science and Management*, 2024, **10**, 2127–2144.
- 76 E. Capanoglu and F. A. Tomás-Barberán, Introduction to Novel Approaches in the Valorization of Agricultural Wastes and Their Applications, *J Agric Food Chem*, 2022, **70**, 6785–6786.
- 77 J. C. Warner, *Green Chemistry and Innovation*, 2015, vol. 1205.
- 78 C. Vichoray, B. Narkhede and A. Deogaonkar, *Role of supply chain in sustainable valorization*, 2023.
- 79 R. Inderst, E. S. Sartzetakis and A. Xepapadeas, Firm Competition and Cooperation with Norm-Based Preferences for Sustainability, *Journal of Industrial Economics*, 2023, **71**, 1038–1071.
- 80 N. Darnall, K. Iatridis, E. Kesidou and A. Snelson-Powell, Penalty Zones in International Sustainability Standards: Where Improved Sustainability Doesn't Pay, *Journal of Management Studies*, 2024, **61**, 2373–2405.
- 81 C. Farrukh and M. Holgado, in *Towards Sustainable Technologies and Innovation - Proceedings of the 27th Annual Conference of the International Association for Management of Technology, IAMOT 2018*, 2018.
- 82 T. B. Ribeiro, G. B. Voss, M. C. Coelho and M. E. Pintado, *Food waste and by-product valorization as an integrated approach with zero waste: Future challenges*, 2022.
- 83 J. Synowiecki and N. A. Al-Khateeb, Production, Properties, and Some New Applications of Chitin and Its Derivatives, *Crit Rev Food Sci Nutr*, 2003, **43**, 145–171.
- 84 F. Shahidi and P. Ambigaipalan, Omega-3 Polyunsaturated Fatty Acids and Their Health Benefits, *Annu Rev Food Sci Technol*, 2018, **9**, 345–381.





## REVIEW

## Sustainable food technology

- 85 J. Dumay, M. Allery, C. Donnay-Moreno, G. Barnathan, P. Jaouen, M. E. Carbonneau and J. P. Bergé, Optimization of hydrolysis of sardine (*Sardina pilchardus*) heads with Protamex: enhancement of lipid and phospholipid extraction, *J Sci Food Agric*, 2009, **89**, 1599–1606. Article Online  
DOI: 10.1039/D5FB00236B
- 86 S. Sharma, J. Sahni and R. Joy, *Linking the Path to Zero Hunger: Analysing Sustainable Development Goals Within the Context of Global Sustainability*, 2025, vol. 557 LNCE.
- 87 L. Bizikova, S. Jungcurt, K. McDougal and S. Tyler, How can agricultural interventions enhance contribution to food security and SDG 2.1?, *Glob Food Sec*, DOI:10.1016/j.gfs.2020.100450.
- 88 J. Blesh, L. Hoey, A. D. Jones, H. Friedmann and I. Perfecto, Development pathways toward “zero hunger”, *World Dev*, 2019, **118**, 1–14.
- 89 G. Grosso, A. Mateo, N. Rangelov, T. Buzeti and C. Birt, Nutrition in the context of the Sustainable Development Goals, *Eur J Public Health*, 2020, **30**, I19–I23.
- 90 C. Radcliffe and J. Singh, *Achieving Food and Nutrition Security and Climate Change: Clash of the Titans or Alignment of the Stars?*, 2020.
- 91 K.-D. A, A. RS, S. J and M. Steiner-Asiedu, Food And Nutrition Security Situation In Ghana: Nutrition Implications For National Development, *African Journal of Food, Agriculture, Nutrition and Development*, 2021, **21**, 18005–18018.
- 92 J.-F. Guégan, G. Suzán, S. Kati-Coulibaly, D. N. Bonpamgue and J.-P. Moatti, Sustainable development goal #3, “health and well-being”, and the need for more integrative thinking, *Veterinaria Mexico OA*, DOI:10.21753/vmoa.5.2.443.
- 93 A. M. Martín Ortega and M. R. Segura Campos, *Bioactive Compounds as Therapeutic Alternatives*, 2019.
- 94 G. Dable-Tupas, V. Tulika, V. Jain, K. Maheshwari, D. D. Brakad, P. N. Naresh and S. Suruthimeenakshi, *Bioactive compounds of nutrigenomic importance*, 2022.
- 95 R. K. Sindhu, A. Goyal, E. A. Yapar and S. Cavalu, Bioactive compounds and nanodelivery perspectives for treatment of cardiovascular diseases, *Applied Sciences (Switzerland)*, DOI:10.3390/app112211031.
- 96 S. Murali, V. S. Krishnan, P. R. Amulya, P. V. Alfiya, D. S. A. Delfiya and M. P. Samuel, Energy and water consumption pattern in seafood processing industries and its optimization methodologies, *Clean Eng Technol*, DOI:10.1016/j.clet.2021.100242.
- 97 V. Vidya, G. Prasad and A. M. Sheela, Assessment of threats to a Ramsar site from seafood processing operation effluents, *Lakes and Reservoirs: Science, Policy and Management for Sustainable Use*, 2020, **25**, 196–213.
- 98 M. A. Moktadir and J. Ren, Promoting sustainable management of hazardous waste-to-wealth practices: An innovative integrated DPSIR and decision-making framework, *J Environ Manage*, DOI:10.1016/j.jenvman.2023.118470.
- 99 J. Laso, I. Ruiz-Salmón, M. Margallo, P. Villanueva-Rey, L. Poceiro, P. Quinteiro, A. C. Dias, C. Almeida, A. Marques, E. Entrena-Barbero, S. Mellett and R. Aldaco, Achieving Sustainability of the Seafood Sector in





- the European Atlantic Area by Addressing Eco-Social Challenges: The NEPTUNUS Project, *Sustainability* (Switzerland), DOI:10.3390/su14053054. Online  
DOI: 10.1039/D5FB00236B
- 100 N. J. Rowan, The role of digital technologies in supporting and improving fishery and aquaculture across the supply chain – Quo Vadis?, *Aquac Fish*, 2023, **8**, 365–374.
  - 101 J. Grecuccio, E. Giusto, F. Fiori and M. Rebaudengo, Combining Blockchain and IoT: Food-Chain Traceability and Beyond, *Energies (Basel)*, 2020, **13**, 3820.
  - 102 Z. Zhao, Y. Li and Z. Du, Seafood Waste-Based Materials for Sustainable Food Packing: From Waste to Wealth, *Sustainability (Switzerland)*, DOI:10.3390/su142416579.
  - 103 S. Singh, T. Negi, N. A. Sagar, Y. Kumar, A. Tarafdar, R. Sirohi, R. Sindhu and A. Pandey, Sustainable processes for treatment and management of seafood solid waste, *Science of the Total Environment*, DOI:10.1016/j.scitotenv.2022.152951.
  - 104 E. T. T. Tan, L. S. Chua and I. M. Yusoff, *Promoting sustainable growth and renewable energy through food waste valorization*, 2024.
  - 105 F. O. Areche, J. A. J. Dominguez, J. T. Huaman, R. J. M. Yapias, T. J. C. Rivera, L. D. M. Perales, J. D. H. T. Quispe, C. Y. H. Zea, C. R. C. Galarza, A. R. Rodríguez, A. R. Rodríguez and D. Y. Areche-Mansilla, Maximizing the Potential of Marine Resources: a Sustainable Approach to High-Value Product Development from Seafood by-Products and Waste, *Current Research in Nutrition and Food Science*, 2024, **12**, 1074–1092.
  - 106 N. Zulfainarni, D. Indrawan and N. I. Khumaera, in *IOP Conference Series: Earth and Environmental Science*, 2020, vol. 420.
  - 107 J. M. Reimer, R. Devillers and J. Claudet, Benefits and gaps in area-based management tools for the ocean Sustainable Development Goal, *Nat Sustain*, 2021, **4**, 349–357.
  - 108 I. Okafor-Yarwood, Illegal, unreported and unregulated fishing, and the complexities of the sustainable development goals (SDGs) for countries in the Gulf of Guinea, *Mar Policy*, 2019, **99**, 414–422.
  - 109 A. Said and R. Chuenpagdee, Aligning the sustainable development goals to the small-scale fisheries guidelines: A case for EU fisheries governance, *Mar Policy*, DOI:10.1016/j.marpol.2019.103599.
  - 110 D. Knorr and M. A. Augustin, The future of foods, *Sustainable Food Technology*, 2024, **2**, 253–265.
  - 111 W. Dong, J. Tang, J. Cropotova, D.-W. Sun and B. K. Tiwari, Green technologies for bio-refinery in marine crustacean shell valorisation from chitin perspective, *Trends Food Sci Technol*, DOI:10.1016/j.tifs.2024.104580.
  - 112 R. N. Mankaa and M. Traverso, Regional management options for floating marine litter in coastal waters from a life cycle assessment perspective, *International Journal of Life Cycle Assessment*, 2023, **28**, 1705–1722.
  - 113 A. Nzihou, Toward the valorization of waste and biomass, *Waste Biomass Valorization*, 2010, **1**, 3–7.



## REVIEW

## Sustainable food technology

- 114 A. S. T. Tan, H. Uthayakumar, L. S. Yeo, K. G. H. Kong, S. L. Y. Lo, V. Andiappan, A. C. M. Loy, S. Y. Teng and B. S. How, Shapley-Shubik Agents Within Superstructure-Based Recycling Model: Circular Economy Approaches for Fish Waste Eco-Industrial Park, *Process Integration and Optimization for Sustainability*, 2024, **8**, 487–501. DOI:10.1039/D5FB00236B
- 115 Y. Hu, X. Yu, J. Ren, Z. Zeng and Q. Qian, Waste tire valorization: Advanced technologies, process simulation, system optimization, and sustainability, *Science of the Total Environment*, DOI:10.1016/j.scitotenv.2024.173561.
- 116 H.-M. Stander and J. L. Broadhurst, Understanding the opportunities, barriers, and enablers for the commercialization and transfer of technologies for mine waste valorization: A case study of coal processing wastes in South Africa, *Resources*, DOI:10.3390/resources10040035.
- 117 T. Fennelly, *Navigating Supply Chains*, 2021.
- 118 R. Nicastro, M. Papale, G. M. Fusco, A. Capone, B. Morrone and P. Carillo, Legal Barriers in Sustainable Agriculture: Valorization of Agri-Food Waste and Pesticide Use Reduction, *Sustainability (Switzerland)*, DOI:10.3390/su16198677.
- 119 E. Ntostoglou, V. Martin, D. Khatiwada and F. Urban, Path-dependencies in the transition to sustainable biowaste valorization: Lessons from a socio-technical analysis of Sweden and Greece, *Waste Management*, 2025, **192**, 47–57.
- 120 A. Khan, A. A. Shah, B. A. Alotaibi and I. Khan, Prioritizing Resource and Policy Barriers to Post-Disaster Recovery in Pakistan: A Multi-Expert Fermatean Fuzzy AHP Analysis, *Sustainability (Switzerland)*, DOI:10.3390/su162310414.
- 121 N. E. I. Boumali, F. Mamine, E. Montaigne and F. Arbouche, Drivers and Barriers for the Valorization of the Apricot Pit, *International Journal of Fruit Science*, 2021, **21**, 158–179.
- 122 J. He, IMPORTED SEAFOOD TRACEABILITY REGULATIONS: A MISHAP FOR THE WTO'S DISREGARD FOR NON-PRODUCT RELATED PROCESSES AND PRODUCTION METHODS?, *Asian Journal of WTO and International Health Law and Policy*, 2020, **15**, 169–207.
- 123 L. T. Antelo, G. M. De Hijas-Liste, A. Franco-Uría, A. A. Alonso and R. I. Pérez-Martín, Optimisation of processing routes for a marine biorefinery, *J Clean Prod*, 2015, **104**, 489–501.
- 124 R. Gormley, Ethics and sustainability in the seafood chain, *Food Science and Technology*, 2010, **24**, 14–17.
- 125 T. Hirokawa and B. S. Thompson, The influence of new sustainable fisheries policies on seafood company practices and consumer awareness in Japan, *Mar Policy*, DOI:10.1016/j.marpol.2023.105819.
- 126 Z. Kyriazi, L. R. de Almeida, A. Marhadour, C. Kelly, W. Flannery, A. Murillas-Maza, R. Kalaydjian, D. Farrell, L. M. Carr, D. Norton, D. Norton and S. Hynes, Conceptualising Marine Biodiversity Mainstreaming as an Enabler of Regional Sustainable Blue Growth: The Case of the European Atlantic Area, *Sustainability (Switzerland)*, DOI:10.3390/su152416762.
- 127 B. Belton, T. Reardon and D. Zilberman, Sustainable commoditization of seafood, *Nat Sustain*, 2020, **3**, 677–684.



- 128 T. Ordóñez-Del Pazo, L. T. Antelo, A. Franco-Uría, R. I. Pérez-Martín, C. G. Sotelo and A. A. Alonso, Fish discards management in selected Spanish and Portuguese métiers: Identification and potential valorisation, *Trends Food Sci Technol*, 2014, **36**, 29–43. DOI: 10.1039/D5FB00236B
- 129 L. McClenachan, S. T. M. Dissanayake and X. Chen, Fair trade fish: consumer support for broader seafood sustainability, *Fish and Fisheries*, 2016, **17**, 825–838.
- 130 L. W. Jodice and W. C. Norman, Comparing importance and confidence for production and source attributes of seafood among residents and tourists in South Carolina and Florida coastal communities, *Appetite*, DOI:10.1016/j.appet.2019.104510.
- 131 S. G. Lewis and M. Boyle, The Expanding Role of Traceability in Seafood: Tools and Key Initiatives, *J Food Sci*, DOI:10.1111/1750-3841.13743.
- 132 J. Petereit, C. Hoerterer and G. Krause, Country-specific food culture and scientific knowledge transfer events – Do they influence the purchasing behaviour of seafood products?, *Aquaculture*, DOI:10.1016/j.aquaculture.2022.738590.
- 133 M. R. Baker, R. Alverson, R. Christiansen, K. Criddle, D. Eilertsen, R. J. Foy, J. Gauvin, S. E. Goodman, L. Habegger, B. P. Harris, K. D. E. Stokesbury and D. R. Sykes, Mechanisms and models for industry engagement in collaborative research in commercial fisheries, *Front Mar Sci*, DOI:10.3389/fmars.2023.1077944.
- 134 Z. T. Tsegay, S. Agriopoulou, M. Chaari, S. Smaoui and T. Varzakas, Statistical Tools to Optimize the Recovery of Bioactive Compounds from Marine Byproducts, *Mar Drugs*, 2024, **22**, 182.
- 135 A. Ettalibi, A. Elouadi and A. Mansour, AI and Computer Vision-based Real-time Quality Control: A Review of Industrial Applications, *Procedia Comput Sci*, 2024, **231**, 212–220.
- 136 A. Casas, D. Rodríguez-Llorente, G. Rodríguez-Llorente, J. García and M. Larriba, Machine learning screening tools for the prediction of extraction yields of pharmaceutical compounds from wastewaters, *Journal of Water Process Engineering*, 2024, **62**, 105379.
- 137 R. A. I. Pratiwi, L. A. Fani and F. Kusasi, in *BIO Web of Conferences*, 2024, vol. 134.
- 138 S. Zulkarnain, Development of Blockchain-Based System for Tracking Fish Species' Taxonomy and Conservation Status, *Journal of Fish Taxonomy*, 2024, **31**, 22–31.
- 139 F. Tolentino-Zondervan, P. T. A. Ngoc and J. L. Roskam, Use cases and future prospects of blockchain applications in global fishery and aquaculture value chains, *Aquaculture*, DOI:10.1016/j.aquaculture.2022.739158.
- 140 A. Subash, H. N. Ramanathan and M. Šostar, From catch to consumer: enhancing seafood processing management with Industry 4.0 innovations, *Discover Food*, DOI:10.1007/s44187-024-00115-6.



**Data availability**

No primary research results, software, or code have been included, and no new data were generated or analysed as part of this review.

