

Sustainable Food Technology

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: M. J. Lawrence, V. Hitlamani, A. Nougou Bissou, A. A. Indamdar, M. R. Mouangue and N.Y. Njintang, *Sustainable Food Technol.*, 2025, DOI: 10.1039/D5FB00326A.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.


Sustainability Spotlight Statement

View Article Online
DOI: 10.1039/D5FB00326A

Sub-Saharan Africa faces major food security challenges due to post-harvest losses (PHL) in roots, tubers, and plantains (RTPs), which are vital calorie sources. Traditional artisanal processing of RTPs into semolina such as gari, attiéké, and tapioca offers a sustainable solution by reducing PHL, extending shelf life, and increasing market value. The process involves peeling, grating, fermenting, squeezing, and heat treatment, with fermentation playing a key role in safety and flavor. This method not only minimizes waste but also empowers women-led rural enterprises and improves local nutrition. To scale this approach, supportive policies, infrastructure investment, and digital traceability (like IoT and blockchain) are essential. By leveraging traditional knowledge and technology, SSA can enhance food security, boost rural economies, and build a more resilient food system advancing key Sustainable Development Goals (SDGs) like Zero Hunger, Gender Equality, and Responsible Consumption.



Artisanal production of semolina: an agro-industrial prospect for the reduction of post-harvest losses of roots, tubers, and plantains in Sub-Saharan Africa.

Mundéné-Timothée Junior Lawrence^{1,2}, Veeranna Hitlamani², Nougba Bissoué Achille¹, Aashitosh A Inamdar ², Mouangue Ruben³ & Njintang Yanou Nicolas⁴.

¹Advanced Teacher's Training College for Technical Education, Department of Chemical Engineering, University of Douala, Cameroon;

²Flour Milling, Baking, & Confectionery Technology Department, CSIR-Central Food Technological Research Institute, Mysuru, India;

³National Higher Polytechnic School of Douala, University of Douala, Cameroon;

⁴National School of Agro-Industrial Science, Department of Food Sciences and Nutrition, University of Ngaoundere, Cameroon;

 **Corresponding Author** (aainamdar@cftri.res.in)

Aashitosh, A, Inamdar, Senior Principal Scientist, Department of Flour Milling, Baking, & Confectionery Technology, Central Food Technological Research Institute, (Council of Scientific and Industrial Research), Mysuru, 500 020, India.



Abstract

The region of sub-Saharan Africa faces major food and nutrition security challenges because of post-harvest losses (PHL), which affect its essential roots, tubers, and plantains (RTPs) as calorie sources. The combination of inefficient supply chains, inadequate infrastructure, equipment, and these losses creates serious economic and nutritional consequences.

The evaluation examines the scalability of traditional food crop processing into semolina in Sub-Saharan Africa (SSA) to fight PHL while creating economic opportunities for rural communities. The inclusion criteria required studies demonstrating artisanal RTP processing into semolina in sub-Saharan Africa while showing the traditional food crops associated with PHL origins.

The research data demonstrated that different production methods exist across various regional and cultural groups. The studies reveal that RTP semolina production involves five fundamental steps: peeling, grating, fermentation and squeezing, and finally heat treatment through roasting or steaming. The research demonstrates that fermentation is a crucial process that shapes final product characteristics and operating conditions by reducing toxic cyanogenic compounds and creating distinctive semolina flavours. The artisanal processing method decreases post-harvest losses through its ability to lengthen product shelf life and create higher market value from processed products. The socio-economic advantages of artisanal production include greater earnings for predominantly women producers and better availability of nutritious food in local communities. Implementing integrated policies is essential for establishing a sustainable food system to support artisanal semolina production and consumption of m'bahou, attiéké, gari, and tapioca. Using systems such as the Internet of Things (IoT) or blockchain to ensure the traceability of traditional food crops from farm to consumer could also help reduce PHL significantly.

Keywords: Agro-industry; Attiéké; Gari; M'bahou; Post-harvest losses; Semolina; Tapioca.



1.0.Introduction

View Article Online
DOI: 10.1039/D5FB00326A

Sub-Saharan Africa is at a crossroads characterized by increasing urbanization and rapidly changing dietary habits¹. As urban centers expand and populations seek to diversify their diets, RTPs remain fundamental pillars of food security and local economies in the region^{2,3}. Rich in energy and essential nutrients, these crops form the basis of the diets of millions of people and play a crucial role in traditional and emerging farming systems⁴⁻⁷.

However, despite their vital importance, SSA's RTPs sector is seriously affected by massive PHL⁶. These losses, which can reach alarming percentages, estimated at 30-40% of total production in some locations throughout the value chain from farm to consumer, are multifactorial^{6,8,9}. They are mainly due to biological factors (rotting, sprouting), technical aspects (often rudimentary farming practices, inadequate harvesting and handling technologies, poor storage conditions), and socio-economic factors (limited access to markets, storage infrastructure, and processing technologies)¹⁰⁻¹³. The socio-economic consequences of these losses are devastating: reduced income for farmers, reduced food availability, increased vulnerability of populations, waste of precious resources, and damage to regional food security^{13,14}. PHLs do not contribute to the sustainable development of emerging countries as defined by the United Nations, jeopardizing global efforts to eradicate hunger by 2030 while promoting sustainable agriculture¹⁵.

Faced with this critical problem, a reorientation of the management system is required to explore innovative and sustainable agro-industrial solutions to add value to RTPs while drastically reducing post-harvest losses. In this context, the production of semolina from the processing of these crops is emerging as a promising avenue¹⁶. Locally produced from roots (such as cassava), tubers (such as yams), or plantains, semolina (such as *attiéké*, *gari*, *tapioca*, and *m'bahou*) is easy to prepare, nutritious, and meets the logistical and economic constraints of local and international producers and consumers⁶. Semolina, a dry and stable product, offers the advantage of long-term preservation, diversification of culinary uses, and creation of added value within local value chains, thereby strengthening culinary and cultural sovereignty and food security¹⁷.

To ensure the reliability of this work, only studies published in French or English between 2000 and 2024 that dealt with semolina production processes (*attiéké*, *gari*, *tapioca*, or *m'bahou*), the PHL of RTPs, the challenges of agro-industrialization, and the development of sustainable agriculture in Sub-Saharan Africa were included. Research on inaccessible or concerned with other cereals, fruits, and vegetables was excluded. This approach enabled a focus on recent,



peer-reviewed bibliographic data, while reflecting the current realities of this part of the continent. The results will guide all stakeholders, including policymakers, researchers, and practitioners.

This study proposes to explore the causes of post-harvest losses and the potential of local semolina production as a viable agro-industrial alternative to reduce losses of traditional food crops in Sub-Saharan Africa. It aims to synthesise available scientific and technical knowledge, identify challenges and opportunities, and contribute to the development of effective strategies for adding value to these important crops that play a central role in food security and local economies. Considering all this data, this review aims to inform the various stakeholders of the key role that semolina production could play in the sustainable transformation of the traditional food crop sector in Sub-Saharan Africa, by combining it with innovative traceability systems.

2.0.Post-harvest losses of roots, tubers, and plantains in sub-Saharan Africa.

According to the FAO report “Agricultural Production Statistics 2000-2021”⁵, the world production of roots and tubers was estimated at 800 million tonnes in 2021. Africa and Asia were the primary producers, with about 340 million tonnes, accounting for almost 80% of world production. Potato was the most produced tuber in this category, with 376 million tonnes, followed by cassava with 315 million tonnes. Cassava (*Manihot esculenta*) production increased significantly by 79% compared to 2000, reflecting its growing importance as a staple food, particularly in sub-Saharan Africa. Significant roots and tubers such as sweet potato, yam, and taro contributed to world production, albeit in smaller quantities. Sweet potato production fell by 36% between 2000 and 2021, while yam and taro production increased by 90% and 17%, respectively. Despite these production levels, the region faces significant post-harvest losses threatening millions of smallholder farmers' food security and economic stability.

RTP's post-harvest losses constitute a significant challenge in SSA. Contrary to estimates of RTP production in many producing countries, there is a lack of actual, quantified data on post-harvest losses¹⁸. According to the FAO in its May report, “Reducing post-harvest losses for improved food and nutrition security in IGAD member states,” losses in SSA are estimated to be between 40 and 50 percent for RTP and occur mainly during transport, storage, and processing^{19,20}. They result from many factors, including rotting, pest infestation, poor storage conditions, lack of infrastructure, excess supply, lack of expertise about effective post-harvest management, and climatic conditions (**Figure 1**)¹⁰. Beyond their agricultural impact, these



losses have economic, social, and environmental dimensions. They affect food availability and the economic stability of smallholder farmers.




Figure 1. Steps involved in post-harvest losses from producer to consumer.

2.1. Post-Harvest Loss Causes

The PHL of RTPs, which can vary by crop and region in SSA, results from various factors that can be classified into biological, technical, and socio-economic.

2.1.1. Biological factors

Biological factors are one of the leading causes of PHL. They include rotting, sprouting, parasitic infestation, and pathogen contamination.

2.1.1.1. Rot

Microorganisms irreversibly damage food during microbial attacks, rendering it unacceptable due to rotting or producing toxic substances ¹¹. Root and tuber rots are often caused by fungi (such as *Ceratocystis fimbriata* (black rot), *Aspergillus niger* (dry rot), *Fusarium solani* (root and stem rot), *Plenodomus destruens* (foot rot), *Rhizopus stolonifer* (soft rot), *Lasiodiplodia theobromae* (stem end rot and dieback), and *Penicillium* spp. (blue mould)) ^{21,22}. Those losses due to yam dry rot, exacerbated by nematodes, can reach alarming levels ^{21,23}. Tubers like sweet potato and cassava are particularly susceptible to fungal and bacterial attack, especially under



humid and hot storage conditions²⁴. Mycotoxins, such as aflatoxin (a liver carcinogen), are also a major risk to PTRs, resulting in significant economic losses¹¹.

2.1.1.2. Physiological activities

Several physiological mechanisms controlled by respiration can occur in RTPs during storage, causing changes in the sensory and nutritional properties of the food. Natural respiration, which promotes the processes of ripening, senescence, wilting, and germination in RTPs, is responsible for a significant proportion of weight loss through the heat it generates. These changes can increase the susceptibility of the food to mechanical damage or infection by pathogens, and reduce nutritional value and consumer acceptance¹¹. Premature germination of tubers, such as potatoes and yams, leads to quality deterioration and weight loss due to the mobilization of nutrient reserves²⁵. Sprouting is often favored by inappropriate temperatures and humidity during storage²⁶. Losses due to sprouting can be significant, especially if tubers are not stored under optimal conditions. According to Adeniyi and Ayandiji¹⁸, a significant proportion of PHL of plantain in the Cote d’Ivoire and Nigeria, estimated at 35%, is due to the rapid ripening of this fruit, and most products made from it can only be stored for a few days. The bio-deterioration of sweet potato is responsible for about 32.5% of post-harvest losses of this tuber²¹. Degebas¹⁹ and Liu et al.²⁶ also report that maturity at harvest and tuber stress influence the degree of sprouting and reduce natural dormancy. Considering storage conditions, increased respiration (use of sugar converted from starch) during sprout growth, and moisture loss, damaged and diseased tubers sprout earlier than healthy tubers.

2.1.1.3. Pest infestation

Insect pests such as weevils, moths, rodents, and birds cause significant damage to RTP stocks. Their consumption of feed results in direct losses, not to mention the contamination caused by hair, feathers, excrement, and the unpleasant odors of urine and pheromones¹¹. Nematodes, in particular, are responsible for yam dry rot, which exacerbates post-harvest losses^{21,23}. Insect pests such as *Euzopherodes vapidella* contribute significantly to yam deterioration²⁷.

2.1.2. Technical factors

Open Access Article. Published on 19 September 2025. Downloaded on 10/7/2025 1:43:32 AM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.


Sustainable Food Technology Accepted Manuscript

Technical factors are crucial in PHL, primarily due to poor storage conditions and inadequate infrastructure, and suboptimal handling practices.

2.1.2.1. Poor storage conditions and inadequate infrastructure

Traditional storage structures commonly used in sub-Saharan Africa fail to adequately protect crops from pests and environmental hazards, compounded by critical deficits in infrastructure. Variations in temperature and humidity during storage accelerate losses, particularly in humid tropics, where inadequate facilities at harvest trigger rapid deterioration of produce²¹. Nwankwo and Chiekezie²⁸ documented cassava losses reaching 35% in Nigeria due to deficient storage systems, while Degebasa¹⁹ highlighted physical damage to potato tubers from improper bag stacking and insufficient ventilation during storage, leading to bruising. These conditions exacerbate physiological stress in tubers, increasing water loss and quality decline²⁹. Furthermore, the absence of essential infrastructure, such as modern storage silos and rural roads, severely limits farmers' capacity to preserve and market harvests³⁰. This is evidenced by Chiekezie et al.³¹ reporting 266,116 kg of cassava post-harvest losses, predominantly during harvesting (55%) and processing (33%) phases. These findings align with Adeniyi and Ayandiji¹⁸, who attributed 46.45% of plantain losses to transportation gaps and 33.60% to marketing constraints.

2.1.2.2. Suboptimal handling practices

Farmers often lack training in proper handling and storage practices. Kuyu et al.³², show that inappropriate harvesting techniques damage tubers, accelerating their deterioration. In addition, mechanical damage during harvesting and transport creates entry points for pathogens, increasing the risk of rot¹⁹. Kouadio et al.³³ Also, improper handling during storage and transport, combined with inadequate temperature control, exacerbates yam losses. Similarly, Sugri and Johnson³⁴, indicate that post-harvest losses for plantain in various African states are often due to inadequate harvesting, transport, and storage practices.

2.1.3. Socio-economic factors

Socioeconomic factors, such as limited market access and low-value addition, exacerbate post-harvest losses.



2.1.3.1.Limited access to markets

The remoteness of production areas from urban markets, combined with poor transport infrastructure, leads to significant losses. Farmers are often forced to sell their produce at low prices to avoid degradation, reducing their income and ability to invest in better conservation practices³⁵. In Nigeria, for example, these constraints can reach 50-60%³¹. Lack of market access and high temperatures contribute to increased losses, especially in countries such as Ghana^{36,37}.

2.1.3.2.Low product value

RTPs are often undervalued in the market due to their high perishability, which limits incentives to invest in preservation and processing technologies. The low value of RTP products, such as flour, crisps, or drinks, is a significant challenge for the agri-food sector in Africa, which can be supported by the analysis of several studies on the characteristics, preservation practices, and market demand for these products, which are indicators of commercial outlets.

The low value added of tubers can be attributed to problems related to product quality and preservation. Research has shown that the storage time of untreated tubers can lead to significant weight losses of up to 43%, while properly treated tubers lose only 17% of their weight³⁸. It is also important to note that the quality of tubers such as cassava is influenced by various factors, including the variety grown and ecological conditions^{39,40}. Therefore, it is necessary to favor adapted varieties and develop effective preservation techniques to improve cassava's value.

According to the data collected, there is a growing demand for alternatives to imported wheat flour. However, national production of tubers such as cassava and sweet potato is still insufficient to meet the needs of the industrial market⁴¹. Despite their high nutritional potential, tuber-based products have failed to penetrate the market due to increasing wheat imports and an ill-defined substitution strategy⁴¹. In addition, efforts must be made to process these products, particularly by producing better-quality bread flour^{41,42}.

The value of products such as plantain also appears to be under-exploited. The processing of plantain into products such as chips, juice, and semolina could benefit from research into methods to prevent browning and recover extraction residues⁴³. In addition, the use of



innovative approaches, such as traditional storage in refrigerated pots, could help to preserve these products better, thereby improving their quality and competitiveness in the market ⁴⁴.

2.1.3.3. Poverty and lack of finance

According to Anyoha et al.¹⁰, the leading cause of post-harvest losses in cassava in Imo State, Nigeria, is a lack of finance. Small-scale producers, who make up the majority of farmers in SSA, often lack the financial resources to invest in modern storage infrastructure or processing technologies ⁴⁵. This forces them to resort to inefficient, traditional practices, perpetuating loss cycles.

2.2. Impact of post-harvest losses

The impact of PHL on roots and tubers in SSA is seen in two main areas: food security and farmers' incomes.

2.2.1. Social Impact

Post-harvest losses of RTPs in SSA directly impact food security, mainly by reducing the availability of staple foods. Roots and tubers such as cassava, yams, sweet potatoes, and plantains are staple foods for much of the sub-Saharan population. However, their high perishability and poor storage and transport conditions result in significant losses.

2.2.1.1. Threats to food security

Post-harvest losses (PHL) in Roots, Tubers, and Plantains (RTPs), reaching up to 50% of production or 20% according to Rwubitse et al.⁴⁶, severely reduce food availability in sub-Saharan Africa (SSA), where over 250 million people faced undernourishment in 2019⁴⁷. Staple crops like cassava, vulnerable to deterioration within 1 to 3 days post-harvest⁴⁸, exemplify this crisis. These losses not only diminish food quantity but also degrade nutritional quality, exacerbating malnutrition as pests and poor storage contaminate calorie- and nutrient-dense RTPs essential for rural populations^{48,49}. Consequently, urban and peri-urban households increasingly rely on imported processed foods to meet demand for convenient diets⁵⁰, while rural communities become dependent on volatile global markets. This import dependency



exposes populations to price shocks and supply-chain disruptions, undermining local food sovereignty. As Thomas et al.⁵¹ emphasize, strengthening post-harvest systems is critical to enhancing food access, stabilizing farmer livelihoods, and securing sustainable food systems across SSA.

2.2.1.2.Impact on Sustainable Development Goals (SDGs)

Cultivating plantains, sweet potatoes, cassava, yams, taro, and potatoes is critical to achieving the Sustainable Development Goals (SDGs), especially Goal 2, which aims to end hunger and promote sustainable agriculture ²⁰ Root and tuber crops are recognized for their resilience and adaptability to various climates. They also make significant nutritional contributions, which enhance food security in vulnerable regions ⁵².

The production of these crops directly supports food security by providing calorie-dense foods essential for large populations, especially in developing regions. Sweet potatoes and yams, for example, have been extensively promoted due to their high yields and nutritional value. These crops have the potential to reduce hunger levels if widely adopted by households ^{53,54}. Furthermore, studies highlight the importance of sweet potatoes for subsistence and for bolstering local economies, especially when smallholder farmers cultivate them ^{54,55}. The genetic diversity of these crops allows them to adapt better to climate change, thus contributing to the resilience of agricultural systems under varying conditions ^{56,57}.

Additionally, cultivating these tuberous crops within agroecological frameworks is associated with benefits aligned with the SDGs. These practices improve soil quality, reduce reliance on chemical fertilizers, and promote environmental sustainability and enhanced agricultural productivity ^{55,58}. For instance, integrating sustainable practices into sweet potato and cassava cultivation can mitigate environmental degradation and foster healthier soil, supporting more sustainable agricultural methods ^{58,59}.

Moreover, these crops' nutritional benefits provide a way to combat micronutrient deficiencies, a challenge highlighted in SDG 2. Incorporating biofortified varieties, such as orange-fleshed sweet potatoes, has been recognized as an effective strategy to address issues like vitamin A deficiency ^{60,61}. These crops are useful beyond direct food consumption because they can also



be used to produce biofuels and other industrial products, which aligns with SDG 7, promoting affordable and clean energy ⁶².

Furthermore, intercropping these root and tuber crops with other food varieties enhances agricultural diversity and resilience against pests and diseases. This improves overall food systems and contributes to the socioeconomic stability of rural communities ^{54,63,64}. These integrative approaches are essential for achieving multiple SDGs because they ensure that agricultural practices are sustainable, inclusive, and equitable ⁶⁵.

Emphasizing the cultivation of root and tuber crops, such as sweet potatoes, cassava, yams, and taro, is crucial for advancing the goals of the 2030 Agenda for Sustainable Development. They address immediate food security needs, promote sustainable agricultural practices, and improve nutritional outcomes. Thus, they provide resilience against economic and environmental challenges while supporting overall socioeconomic development ^{54,63,66}. Strategically integrating these crops into agricultural policies and programs is vital for achieving Sustainable Development Goal 2.

2.2.2. Economic impact

Post-harvest losses significantly impact farmers' incomes, especially for smallholder farmers who depend on crop sales for their primary source of income. This issue is particularly acute in the cultivation of root and tuber crops, such as taro, sweet potatoes, cassava, and yams. For example, viral diseases can devastate taro production, resulting in yield losses ranging from 20% to 60%. This significantly compromises farmers' financial viability and necessitates the implementation of effective management strategies to mitigate these losses ^{67,68}. Additionally, studies suggest that inadequate handling and storage practices after harvest can result in losses, further exacerbating economic challenges for small-scale farming operations ^{10,69}.

The economic implications of post-harvest losses are evident in the analysis of specific crops, such as cassava. Cassava is vulnerable to excess supply, improper processing, and inadequate storage infrastructure, resulting in substantial waste and decreased income potential for farmers ¹⁰. Statistical data suggests that post-harvest losses for cassava can exceed 30%, further illustrating the pressing economic challenges faced by these communities ⁷⁰.



Environmental factors such as high humidity and poor infrastructure further exacerbate the situation by accelerating the deterioration of storage quality for root and tuber crops. For example, poor storage conditions accelerate the physiological deterioration of taro, resulting in rapid quality loss and increased post-harvest waste^{71,72}. Addressing these storage challenges is crucial. Adopting better preservation methods, such as treatments that delay enzymatic browning, could markedly improve the shelf life of crops and enhance farmers' economic returns^{71,72}.

2.2.2.1.Reduced income

Post-harvest physiological deterioration caused by physical damage during harvest reduces root and tuber quality, leading to price drops that directly lower farmer incomes. Global economic losses from this waste amount to \$940 billion annually¹⁰, as observed in plantain systems, where pre-harvest phytosanitary issues create economic gaps for dependent farmers⁷³, and cassava value chains where extreme perishability limits market access and income opportunities⁴⁸.

This income loss perpetuates a cycle of poverty: with 66% of West African employment tied to the food economy⁵⁰, farmers' inability to invest in improved practices or storage tech reduces long-term productivity. Consequently, diminished reinvestment capacity deepens rural poverty and food insecurity⁴⁶, trapping communities in financial vulnerability. This self-reinforcing dynamic disproportionately affects smallholders, who lose 15-35% of potential earnings from PHL, while weakening regional food systems' resilience.

2.2.2.2.Market instability

Market instability triggered by post-harvest losses (PHL) manifests through volatile price mechanisms and disrupted supply-demand equilibria. When PHL reduces available crop volumes, particularly perishable staples like cassava or plantain, local markets experience *supply shocks* that inflate consumer prices while depressing farmgate prices. This asymmetry erodes trader margins and discourages long-term contracts, fragmenting value chains⁵¹. Consequently, risk-averse investors avoid capital injection into agricultural cooperatives, processing facilities, or storage infrastructure⁷⁴. The resulting underinvestment stifles productivity gains, entrenches subsistence farming models, and perpetuates rural economic stagnation. Critically, this cycle undermines SDG targets for poverty reduction (Goal 1) and



sustainable agri-food systems (Goal 2), as confirmed by Thomas et al.'s analysis of West African economies⁵¹.

3.0.A Technical analysis of unit operations and their impact on the properties: Overview of semolina made from cassava and plantain

Semolina is a granular product obtained by coarsely grinding starchy materials (cereals, roots, tubers, or fruits). It is intermediate in size between flour and whole grains, characterized by its grainy texture and ability to absorb liquids. Traditionally produced from durum wheat, it can also be made from tropical resources such as cassava, plantain, or yam, serving as a base for various food preparations like couscous and traditional dishes consumed mainly in Africa and South America^{75–77}. It is a form of valorization aimed at reducing post-harvest losses thanks to its stability and versatility.

Gari, *tapioca*, and *attiéké* are three staple foods widely consumed in many parts of the world, particularly in Africa and South America, all of which are derived from cassava root. Cassava is a tuber exceptionally rich in starch, making it a primary source of energy for millions of people⁷⁸. Although the raw material is the same, complex and varied processing methods result in finished products with distinct physical, organoleptic, and nutritional properties. Unlike these three products, *m'bahou* is a traditional Ivorian dish that uses another major starch, plantain, as its main ingredient⁶. **Figure 2** shows the artisanal production process for certain semolinas derived from cassava and plantain (*attiéké*, *gari*, *tapioca*, and *m'bahou*). These processes vary from region to region and directly influence all the characteristics of the semolina produced (**Table 1**). Artisanal methods, although mostly manual, can be optimized to improve product quality and safety while preserving their cultural identity. Understanding these processes is therefore crucial for the development and improvement of the quality of traditional semolina in Sub-Saharan Africa.

Table 1. Impacts of unit operations on semolina properties.

Process steps	Impact on properties			
	Nutritional	Functional	Rheological	Sensory





Peeling and washing	Reduction of anti-nutrients and contaminants.	Increased contact surface for subsequent steps.	N/A	<div>View Article Online DOI: 10.1039/D5FB00326A</div> N/A
Grating	N/A	Fine grating		
		Increases water absorption capacity and swelling power.	Improves dispersion and texture.	Produces a smoother texture.
		Coarse grating		
		Reduces water absorption capacity and swelling power.	More granular texture.	Slightly coarse texture.
Fermentation	Reduction of cyanogenic glycosides (cyanide).	Improves dispersibility and solubility.	Fermentation too long	
			Reduces maximum viscosity.	Risk of excessive acidity.
	Improves nutrient bioavailability.	Fermentation too short		Develops a slightly acidic flavour.
		Risk of poor dough formation.	Produces firmer semolina.	
Squeezing	Concentrates nutrients, but may result in a minor loss of soluble nutrients.	Reduces moisture, which improves product stability.	N/A	N/A
Roasting	Extends shelf life by reducing moisture.	Low temperature		
		Preserves the integrity of starch.	Produces firmer semolina.	Produces a more attractive colour and flavour.
	High temperature			

	Loss of certain nutrients (ex., carotenoids).	Reduces water absorption capacity.	Reduces maximum viscosity.	Risk of excessive browning and burnt flavour.
Stockage	Stored incorrectly		N/A	N/A
	Risk of oxidation and loss of nutrients	degrades functional properties		

416






Figure 2. Schematic diagram of the production process of some types of semolina made from the roots (cassava) and plantain.

3.1. *Attiéké*: Cassava couscous, a fermented product

View Article Online
DOI: 10.1039/D5FB00326A

Attiéké is a fermented cassava couscous, an iconic dish in Ivorian cuisine⁷⁹. It is distinguished by its grainy texture and slightly sour taste, which are the direct result of its manufacturing process, particularly fermentation. Consumed initially in the south of Côte d'Ivoire, the product has grown in popularity and is now produced and consumed throughout the country and in other parts of West Africa^{80,81}. Women have traditionally dominated the production process of *Attiéké* and require specific technical skills. However, the quality of *Attiéké* can vary considerably depending on the cassava variety used, processing techniques, and fermentation inoculants^{81,82}.

3.1.1. Nutritional value of *Attiéké*

Attiéké is an energy-rich food, but its caloric value is significantly lower than that of gari, ranging from 161.95 to 215.26 kcal per 100 g⁸³. This lower energy density is attributed to its higher moisture content, which ranges from 50 to 55% of the product^{83,84}. It is mainly composed of carbohydrates (36.6 g to 47.11 g per 100 g)⁸³, making it a sustainable source of energy. Like other cassava products, *Attiéké* is low in protein (0.77 to 1.74 g) and fat (0.15 to 3.60 g)^{83,84}. It is a source of fiber (approximately 1.5 g), which aids digestion and promotes a feeling of satiety⁸⁴. Minerals are present, with potassium (47.22 mg per 100 mg of ash) being the most abundant, followed by calcium (6.84 mg), phosphorus (6.20 mg), and magnesium (4.38 mg)⁸⁴. This profile indicates that *Attiéké* is typically consumed as a complementary component of a meal, paired with protein-rich dishes to achieve a balanced diet. The final nutritional quality of the product is not solely a function of the raw cassava but is significantly influenced by the production processes.

3.1.2. Unit operations of *Attiéké* production and their impacts

The production of *Attiéké* is a complex process involving a series of distinct unit operations. Each of these steps plays a critical role, not only in transforming the raw cassava into the final semolina but also in shaping its fundamental properties. The following sections detail each unit operation and its specific, integrated impacts on the nutritional, functional, rheological, and sensory characteristics of the finished product.



3.1.2.1. Peeling and Washing

This foundational step is the initial point of processing for cassava roots. It involves the physical removal of the outer skin and a subsequent cleaning with water. This operation is critical for food safety. It removes impurities and, most importantly, eliminates toxic compounds such as cyanogens that are present in the cassava skin^{85,86}. This reduction of anti-nutrients and contaminants is a prerequisite for producing a safe food product^{84,87}. Peeling and washing also serve a functional purpose by increasing the surface area of the cassava root that is exposed. This is a crucial preparation step for subsequent operations, as a larger surface area facilitates more effective grinding and fermentation.

3.1.2.2. Grinding

Following the initial preparation, the cassava roots are ground into a fine paste⁸⁰. The fineness of this grinding process has a significant and direct influence on the properties of the semolina. Grinding is essential for breaking down the cellular structures of the cassava, a necessary action to improve starch availability for the subsequent fermentation process⁸⁸. The particle size resulting from grinding is a primary determinant of the final product's texture. A fine grating process increases the cassava's water absorption capacity and swelling power, which in turn leads to a smoother, more dispersible texture in the final semolina⁸⁹. Conversely, a coarser grating reduces the water absorption and swelling power, resulting in a more granular and slightly coarse texture⁸⁹. This direct relationship between particle size and texture allows producers to control the final mouthfeel of the *Attiéké*.

3.1.2.3. Fermentation

This is a crucial, multi-hour process that typically lasts between 6 and 48 hours^{81,85,88}. It is initiated by a traditional cassava-based starter known as "*magnan*," which contains a variety of microbial strains, notably yeast-like *Candida tropicalis* and lactic acid bacteria⁹⁰. Fermentation is the principal detoxification mechanism, responsible for significantly reducing toxic cyanogenic glycosides and hydrocyanic acid⁹⁰⁻⁹². It also improves the bioavailability of nutrients, such as vitamins and amino acids^{93,94}. The duration and control of fermentation have a profound influence on the final product's quality. An optimally controlled fermentation results in a light, granular texture and the development of the characteristic, slightly acidic

Open Access Article. Published on 19 September 2025. Downloaded on 10/7/2025 1:43:32 AM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.



Sustainable Food Technology Accepted Manuscript

flavor and aroma^{95,96}. However, fermentation that is too short poses a risk of incomplete dough formation and leaves high levels of residual cyanide^{97,98}. Conversely, an excessively long fermentation period can lead to an overly acidic taste and a reduction in the maximum viscosity of the cooked product^{99,100}. The variability often seen in traditional starters can lead to inconsistent product quality, highlighting the importance of standardized fermentation conditions to ensure uniform sensory and safety attributes in the final product.

3.1.2.4.Squeezing

The fermented cassava pulp is subjected to a pressing operation to remove excess water^{88,101}. The removal of water during this stage concentrates the remaining nutrients in the pulp^{101,102}. However, it is important to acknowledge that there can be a minor loss of soluble nutrients in the expelled water, which represents a trade-off inherent in the process. By reducing the moisture content, squeezing significantly improves the product's stability and facilitates subsequent processing steps such as sieving and drying^{80,103}. This step is also fundamental in achieving the granular texture that is characteristic of *Attiéké*^{88,102}.

3.1.2.5.Sieving and Granulation

This operation involves passing the squeezed pulp through a sieve to separate fine particles and achieve a semolina with a uniform size¹⁰⁴. The primary function of this step is to produce a semolina of uniform granulometry. This directly impacts the final texture of the *Attiéké*, which must be granular and light to be acceptable to consumers^{86,104}. The sieving and granulation process ensures the consistency of this key textural property.

3.1.2.6.Drying

The semolina's moisture content is reduced to between 12% and 15% through drying, which is typically done in the sun or with mechanical dryers^{105,106}. Drying is a crucial preservation step. The reduction in moisture is essential for prolonging the product's shelf life and inhibiting the growth of undesirable microorganisms, thereby ensuring its microbiological safety⁹⁰.

3.1.2.7.Steaming





518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551

Steaming is the final essential heating step in *Attiéké* production. It gives the semolina its final texture and is critical for both safety and digestibility⁸⁸. This operation is vital for eliminating any remaining traces of cyanide residues that may have survived the fermentation process¹⁰⁷. Steaming also improves the digestibility of the product and aids in the retention of key nutrients, such as carotenoids, in varieties of biofortified cassava¹⁰⁶. The steaming process is fundamental to developing the final texture, which is a key sensory attribute that determines consumer acceptance.

3.2. *Gari*: A highly energetic cassava semolina

Gari, also known as “saviour” or tapioca in Cameroon, is a staple food in sub-Saharan Africa. It is a cassava semolina characterized by a fine or coarse texture, whose color can vary from white to yellow depending on the production methods¹⁰⁸. It is obtained through a transformation process that includes grating, fermentation, pressing, and roasting the cassava pulp⁷⁸. Each of these stages influences not only the food safety of the product but also its nutritional, functional, and rheological properties.

3.2.1. Nutritional value of *Gari*

Gari is a staple food in sub-Saharan Africa, and its nutritional properties, like those of other cassava-based foods, are primarily shaped by its production process. It is primarily a source of energy and carbohydrates. It has a high energy value, with data ranging from 330 to 370 kcal per 100 g. This calorie density comes mainly from its high carbohydrate/starch content, which makes up most of its composition, with values ranging from 62.6 g to 84.4 g per 100 g of product^{108,109}. The proportion of sugars is very low, generally less than 0.5 g per 100 g. On the other hand, *Gari* is very low in protein and fat. Protein content is marginal, ranging from 1.96 g to 2.88 g, while fat is almost absent, with values ranging from 0.33 to 1.58 g^{108,110}. It is also a source of dietary fiber, with an average value between 1.24 and 3.46 g per 100 g¹⁰⁸, as reported in one source. However, another source mentions an exceptionally high value of 48 g, which is likely a measurement error. Micronutrients such as vitamin A (13.2–732 IU/100 g)¹⁰⁸, calcium (31.41–49.86 mg), and iron (23.0–62.4 mg) are also listed¹¹⁰. The final nutritional composition is a direct consequence of a series of unit operations, most notably fermentation and roasting.

3.2.2. Unit operations of Gari production and their impacts

3.2.2.1. Peeling and Washing

This initial step involves peeling the cassava roots and washing them to remove dirt, impurities, and the outer skin. This is a crucial preparatory stage that removes anti-nutrients and contaminants, thereby laying the groundwork for a safe final product ^{111,112}.

3.2.2.2. Grating

The peeled cassava is grated into a fine pulp. The fineness of this grating, whether done manually or mechanically, is a critical variable. The particle size of the grated pulp directly impacts the final texture and functional properties of the gari ^{109,113}. A fine grating increases the contact surface area, which improves the product's water absorption capacity and swelling power. A coarser grating, on the other hand, results in a more granular and gritty texture ^{113,114}.

3.2.2.3. Fermentation

This is a critical process that typically lasts for 2 to 3 days, during which the cassava pulp is detoxified by the action of microorganisms. Fermentation is the primary step for reducing toxic cyanogenic glycosides to safe levels. Optimal fermentation time not only ensures food safety but also improves the bioavailability of nutrients. The duration of fermentation has a direct effect on the cooked gari's texture and viscosity. An optimal fermentation period develops a desirable flavor and texture ^{115–117}, whereas excessive fermentation can lead to an overly acidic product and cause significant starch degradation, which reduces the viscosity of the cooked gari ^{99,100}.

3.2.2.4. Squeezing

The fermented cassava pulp is pressed to remove excess moisture. This dehydration step is essential for preventing product deterioration and improving its stability during storage ¹¹⁸. It also prepares the pulp for the subsequent roasting stage and plays a role in shaping the final texture.



3.2.2.5. Roasting (Garification)

This is the final stage of *Gari* production, where the dehydrated pulp is heated to high temperatures to form the characteristic crunchy granules. The high temperatures used during roasting can lead to a loss of certain nutrients, such as carotenoids, particularly in fortified cassava varieties^{113,119}. The roasting temperature and duration are the most significant factors in determining the final *gari*'s color, flavor, and texture. Well-controlled roasting produces *gari* with a slightly acidic flavor, an attractive color, and a uniform texture^{115–117}. Insufficient roasting results in a poor texture, while excessive roasting can lead to a burnt flavor and excessive browning^{113,119,120}. The temperature also affects the rheological properties; lower roasting temperatures tend to preserve the integrity of the starch, which results in a firmer final product^{112,113}.

3.2.3. Comparison of traditional and mechanized *Gari* production methods

The production of *Gari* can be broadly categorized into two methods: traditional and mechanized. Each approach presents distinct advantages and disadvantages that influence the final product's quality, safety, and efficiency (Table 2).

Traditional Methods (Figure 3): These methods, often manual, are characterized by high variability in product quality. A key feature of traditional production is the potential for prolonged fermentation times, which are highly effective at reducing cyanide content and enhancing the final flavor and texture. However, the roasting process in traditional methods is often less controlled, which can lead to a greater loss of nutrients, such as carotenoids^{119,121}.

Mechanized Methods (Figure 4): Mechanized production is more uniform and efficient, allowing for more precise control over process parameters¹²². This precision, particularly in the roasting stage, can improve nutrient retention and lead to a more consistent overall product quality^{111,123}. However, mechanized methods may not always guarantee the same level of optimal cyanide reduction as the longer, traditional fermentation processes.

This comparison reveals a fundamental trade-off: traditional methods, while less uniform, may offer superior benefits in terms of detoxification and flavor development due to extended fermentation, whereas mechanized methods provide greater consistency, efficiency, and better



control over parameters like roasting, which can enhance nutrient retention. An integrated approach that combines the best practices of both methods, such as using a standardized mechanical process with an optimized, biologically driven fermentation step, could represent a path toward producing a consistently high-quality and safe gari^{99,100}.

Particular attention should be paid to the fermentation and roasting stages, which are crucial for reducing anti-nutrients and improving gari's functional and sensory properties. Finally, promoting improved production techniques could increase food security and strengthen the economic empowerment of local communities, especially women, who play a central role in artisanal gari production¹²⁴.

Table 2. Comparison of Traditional and Mechanized Gari Production Methods.

Feature	Traditional Gari Production	Mechanized Gari Production
Product Uniformity	High variability	High uniformity
Efficiency	Lower (manual)	Higher (automated)
Cyanide Reduction	Highly effective due to long fermentation	It may not always be optimal
Nutrient Retention	Potential loss of nutrients from prolonged roasting	Improved retention with precise roasting control
Final Flavor & Texture	Superior flavor from long fermentation	More consistent, but flavor may differ




Figure 3. The artisanal gari production process.¹²⁵

(1) Cassava tubers after harvesting, (2) tubers are peeled and washed, (3) mechanical grinding of the tubers, (4) collection of the ground tubers, (5) palm oil and soy bean added, (6) placed in buckets, sealed with lids, and left to ferment for 48 h, (7) and (8) mixture is placed in a large bag, (9) the dewatering process, (10) sieving to remove large particles, (11) area where mixture is fried/garified, (12) frying/garification, (13) fine sieving (14) comparison between traditional gari (left) and gari fortified with soybean and palm oil.



Figure 4. Mechanised gari production process diagram.¹²⁶

3.3. *Tapioca*: A pure and versatile cassava starch

Tapioca, also known as sago in India, is a product distinct from *Gari* and *Attiéké*. It is an almost pure starch extracted from cassava root and sold in the form of flour, flakes, or pearls. As a starch, its nutritional composition is heavily focused on carbohydrates and has a limited nutritional profile.

3.3.1. Nutritional profile

The energy value of dry *Tapioca* is similar to that of *Gari*, with values of around 342-358 kcal per 100 g¹²⁷. It is almost exclusively composed of carbohydrates (80.9 g to 89 g per 100 g)^{127,128}. This concentration of carbohydrates is its main characteristic. In contrast, *Tapioca* contains very little protein (1.03 g to 1.1 g), fat (0.24 g to 1.6 g), and dietary fiber (1.8 g to 2.7 g)^{127,128}. Its micronutrient density is also low, although some sources mention the presence of iron, potassium, and magnesium¹²⁸. *Tapioca* contains no vitamins A, C, E, or B12. Its production involves complex processes directly influencing its final characteristics, such as texture, viscosity, and nutritional properties.

3.3.2. Unit operations of *Tapioca* production and their impacts

3.3.2.1. Starch Extraction

This multi-stage process is the initial and most critical set of operations for *Tapioca* production. It involves the preparation of the cassava roots and the physical separation of the starch from other components.

This process begins with washing and peeling the cassava roots to remove impurities and the toxic, cyanogenic skin⁸⁵. The peeled roots are then grated to facilitate starch extraction. The resulting mixture is then washed and filtered, and the starch is separated by sedimentation¹²⁹. The quality of the raw cassava and the fineness of the grating directly influence the starch yield and the quality of the final product¹³⁰. Modern mechanical grating methods are capable of producing a higher-quality starch with less fibrous residue¹²⁹. This initial phase is also essential for safety, as the washing and peeling steps remove the majority of the toxic compounds. The quality of the water used during sedimentation can also affect the physicochemical properties of the starch, demonstrating that seemingly minor details can have a significant impact on the final product.

3.3.2.2. Heat Treatment (Gelatinization)

Once the starch has been extracted, it undergoes a heat treatment process that alters its functional and sensory properties. Starch pearls are immersed in boiling water, causing the granules to absorb water, swell, and become translucent. This gelatinization process is critical for achieving the desired soft, elastic texture of finished products like tapioca pearls^{131,132}. It also significantly affects the viscosity of the tapioca, which is crucial for its use as a thickening agent in sauces, soups, and desserts¹³⁴. Moisture heat treatment can be used to modify the starch structure, which lowers the gelatinization temperature, demonstrating how process parameters can be adjusted to achieve specific functional outcomes.¹³² However, there is a limit to this process; excessive heat treatment can reduce the starch's swelling capacity, which would negatively impact the final texture¹³³. This underscores the need for precise process control to achieve the desired result.

3.3.2.3. Drying



The gelatinized starch is then dried to form the *Tapioca* granules. Drying is a key step that improves both the texture and the shelf life of the *Tapioca* granules^{136,1}. Advanced techniques, such as microwave drying, can offer further benefits by also inducing a form of pre-gelatinization, which increases the starch's binding capacity and results in more uniform granules¹³⁷. This shows how technological innovation can combine multiple process benefits into a single, efficient operation.

3.4. *M'bahou*: A traditional dish made from overripe plantains

M'bahou differs fundamentally from the other foods studied, as it is made from plantains rather than cassava. It is a traditional dish eaten in eastern and central Côte d'Ivoire (culinary heritage of the Agni and Baoulé ethnic groups^{3,6}), and its production is notably focused on utilizing “senescent” plantains, a product that is often rejected in the market due to its perishability⁶.

3.4.1. Specific nutritional composition: a different profile from cassava products

The nutritional profile of *M'bahou* is exceptionally rich, particularly in micronutrients. A study⁶ on dried *M'bahou* reveals that it has a low moisture content (7.13%) and is an excellent source of carbohydrates (87.64%) and starch (61.15%).⁴ Its energy value is approximately 364.87 kcal per 100 g. What sets it apart from cassava products is its richness in micronutrients. *M'bahou* contains significant amounts of vitamins, including vitamin A (134.67 µg), vitamin B1 (49.57 µg), and vitamin B2 (564.87 µg) per 100 g. Its mineral composition is also remarkable, with high levels of potassium (120.32 mg), calcium (85.50 mg), sodium (55.71 mg), and magnesium (46.23 mg). The dish is also a source of phytochemicals, such as polyphenols, phytates, tannins, and flavonoids. The most significant aspect of its nutritional profile, however, lies in its potential for enhancement through fortification. The replacement of a portion of the cassava powder with legume powders, such as soya or cowpea, can substantially increase the protein and lipid content, thereby improving the overall nutritional balance and reducing the glycemic index of the food^{3,6}.

3.4.2. Unit operations of *M'bahou* production and their impacts

3.4.2.1. Preparation of ingredients



The production of *M'bahou* begins with the preparation of its two primary ingredients, senescent plantain and cassava. The plantains are washed, peeled, and mashed ⁶, while the cassava roots are peeled, grated, dried, and ground into a powder. A key process variable in this stage is the optional pre-fermentation of the cassava flour for 2 to 10 hours ³.

3.4.2.2. Mixing, Granulation, and Drying

The plantain puree is mixed with the cassava flour (fermented or not) and salt to form a paste. This paste is then sieved and rolled into granules, which are pre-cooked (20 min) via steaming and finally dried at 50°C to reduce moisture ⁶. The fermentation of the cassava, mentioned in the previous step, plays a significant role in limiting the agglomeration of the granules after cooking by reducing the cohesiveness and stickiness of the final product ³. This is a desirable outcome for texture. However, this same fermentation process also reduces the elasticity and chewability of the food, which represents a processing trade-off ³. The drying process, which reduces the moisture content to a low 7.13%, is critical for the long-term preservation of *m'bahou*. The low lipid content (0.24%) and acidic pH (5.55) of the food also contribute to its stability and resistance to rancidity and microbial growth ⁶.

3.4.2.3. Fortification

To improve the nutritional quality of the dish, 10% to 20% of the cassava powder can be replaced with protein-rich legume powders like soya or cowpea ³. The addition of these legumes significantly increases the protein and lipid content of the *M'bahou*, thereby improving its nutritional balance. Fortification does alter the sensory attributes, such as color and flavor, but these fortified formulations remain generally acceptable to consumers ³. The addition of the legumes can also partially counteract the reduction in elasticity caused by cassava fermentation, thereby increasing the hardness and elasticity of the final product while maintaining its low stickiness ³. This demonstrates a sophisticated interplay of ingredients and processing that can be used to achieve a desired textural profile.

3.4.3. Role of *M'bahou* in food waste reduction and food security

The production of *M'bahou* is unique in its direct contribution to food waste reduction and food security. The dish makes use of senescent plantains, a product that is often rejected by the



market due to its high perishability and advanced stage of ripeness³. By transforming this otherwise unmarketable crop into a valuable food product, the *M'bahou* production process not only reduces post-harvest losses but also plays a crucial role in the local economy. This practice provides a low-cost source of nutrition for communities, thereby contributing to food security and income diversification in both rural and urban areas. The optimization of the production process, particularly through fortification and improved techniques, makes the food more attractive to consumers, further strengthening its positive socio-economic impact.

3.5.Comparative Analysis and Broader Implications

3.5.1. Nutritional profiles

To facilitate comparison, the table below summarizes the nutritional profiles of the semolina foods studied, presenting average values or ranges per 100 g of product (**Table 3**). Analysis of the profiles reveals significant commonalities and important specificities. All of these foods are primarily sources of energy and carbohydrates, with low protein and fat content, confirming their role as staple foods in many diets. *Gari*, *Tapioca*, and *Attiéké*, all derived from cassava, are also naturally gluten-free, making them suitable for people with intolerances.

Table 3. summarizes the nutritional profiles of the semolina of cassava and plantain-based.

Product category	Raw material	Energy value (kcal)	Total carbohydrates (g)	Protein (g)	Fat (g)	Dietary fibers (g)
<i>Attiéké</i>	Cassava	161.5 - 215	36.6 – 47.11	0.77 – 1.74	0.15 – 3.60	2
<i>Gari</i>	Cassava	330 - 370	62.6 – 84.4	1.96 – 2.88	0.33 – 1.58	1.24 – 3.46
<i>Tapioca</i>	Cassava	342 - 358	80.9 - 89	1.03 – 1.1	0.24 – 1.6	1.8 – 2.7
<i>M'bahou</i>	Plantain	365	87.6	3.07	0.24	2.87

However, there are notable differences. *Tapioca* stands out as the purest food, being an almost exclusive source of starch, with the most limited nutritional profile in terms of micronutrients.



Gari, with its lower moisture content, is the most calorie-dense of the three cassava products. *Attiéké*, due to its fermentation process and higher water content, is the least energy-dense and provides fiber and minerals.

The most striking exception is *M'bahou*. Its raw material, plantain, gives it a micronutrient (vitamins A, B1, B2) and mineral (potassium, calcium) profile that is significantly richer than that of cassava products. These characteristic highlights that the raw material, beyond the processing methods, is the determining factor in the final nutritional richness.

3.5.2. Synthesis of Unit Operations and Their Critical Impacts Across Products

The analysis of *Attiéké*, *Gari*, *Tapioca*, and *M'bahou* production reveals several shared principles while highlighting unique differences. The initial steps of peeling and washing are universally critical for food safety, as they remove toxic cyanogenic compounds and other contaminants across all products. The fineness of grating is a consistent determinant of final product texture and functional properties, influencing attributes like water absorption and dispersion in *Attiéké* and *Gari*, and starch yield in *Tapioca*.

However, the most critical unit operations differ between the products. For *Attiéké* and *Gari*, fermentation is arguably the single most important step. It is a dual-purpose process that is vital for both food safety (detoxifying the cassava) and for defining key quality attributes (flavor, aroma, and texture). In *Tapioca* production, the transformative step is the heat treatment of gelatinization, which modifies the starch's structure to create the desired functional properties of swelling capacity and viscosity. *M'bahou* unique character is shaped by a combination of the use of a traditional, perishable ingredient (senescent plantain) and the innovative application of fortification to enhance its nutritional and textural properties. The final heating step also differs significantly; *Attiéké* is steamed, *Gari* is roasted (garified), and *Tapioca* is gelatinized and then dried, with each method imparting distinct characteristics to the final product.

3.5.3. An Overview of Food Safety and Quality Management

The reports repeatedly emphasize that food safety is critically important in producing all cassava-based semolinas. The presence of cyanogenic compounds in raw cassava requires a



series of processing steps to ensure the final product is safe. The analysis indicates that peeling and washing are the initial defenses, followed by more advanced detoxification methods like fermentation and final heating (steaming for *Attiéké*, roasting for *Gari*). Nonetheless, microbiological risks still exist, especially due to the use of traditional inoculums and potentially contaminated water. To reduce these risks and maintain consistent quality, implementing comprehensive food safety management systems such as HACCP is crucial. These systems offer a structured approach to monitor and control critical points throughout the entire production process^{85,145,146}.

3.5.4. Socio-economic significance and future outlook of cassava semolinas

The production of these cassava-based foods holds significant socio-economic importance in West Africa. *Attiéké*, *Gari*, and *M'bahou* production not only provides a source of nutrition but also serves as a source of income, thereby contributing to poverty reduction and food security. The unique role of *M'bahou* in adding value to otherwise unmarketable plantains is a particularly strong example of how these traditional processes can reduce food waste and strengthen the local economy.

However, the artisanal production of these foods faces several challenges, including a lack of technical training for producers and a significant variability in quality, which is often a result of using different cassava varieties and inconsistent processing methods^{81,92,146}. The future of these products lies in a strategic combination of traditional knowledge and technological innovation. By optimizing fermentation techniques and using improved cassava varieties, the organoleptic and nutritional quality of the products can be significantly enhanced. Furthermore, the professionalization of the production process through training and the implementation of food safety standards can lead to a more consistent, higher-quality product, which would ultimately meet consumer preferences and strengthen the economic empowerment of the local communities that produce them.

4.0. Potential of semolina processing as a sustainable solution to post-harvest losses of roots, tubers, and plantains in sub-Saharan Africa.

In the wake of artisanal processes for producing semolina from RTPs, it is essential to explore the potential of this processing as a sustainable solution to post-harvest losses in sub-Saharan



Africa. Although nutritious and an essential part of the staple diet, RTPs are highly perishable products that suffer significant post-harvest losses due to rapid deterioration and inadequate storage conditions. Processing these products into semolina using artisanal and semi-industrial techniques offers a unique opportunity to reduce these losses while providing economic, social, and environmental benefits. Analyzing these aspects shows how semolina processing can contribute to the region's food security and sustainable development.

4.1.Better preservation

Processing RTPs into semolina significantly extends their shelf life, thereby reducing post-harvest losses, which can be as high as 30% in some regions ^{18,147}. RTPs are particularly susceptible to postharvest spoilage. Cassava, for example, begins to decompose within 24 to 48 hours of harvest, resulting in significant losses if no preservation measures are taken ¹⁴⁸. Processing into semolina, especially using drying techniques, significantly extends the shelf life of these products ⁴⁹. This preservation method is essential to improve the availability of products in local and regional markets while reducing losses due to physical and biological deterioration. Complementary preservation techniques, such as vacuum packaging or natural preservatives, could be explored to optimize semolina preservation.

4.2.Diversification of food and feed uses

Semolina made from RTPs offers excellent versatility, allowing for diversifying food products for human and animal consumption. Cassava semolina is a staple food in many African cultures, providing a nutritious and affordable alternative ⁴⁰. It can also produce functional foods and animal feeds, contributing to food security and dietary diversification ^{149,150}. Semolina could also be fortified with nutrients (such as vitamins and minerals) to address nutritional deficiencies prevalent in the region, while being used in animal feed to reduce costs for livestock farmers. This diversification opens up new economic and nutritional opportunities while meeting the needs of local communities.

4.3.Reducing food waste

Processing into semolina reduces food waste using products that would otherwise be lost after harvest. By extending shelf life and creating new products from surplus or damaged produce,

Open Access Article. Published on 19 September 2025. Downloaded on 10/7/2025 1:43:32 AM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.



Sustainable Food Technology Accepted Manuscript

this approach minimizes losses and adds value to available resources^{151,152}. For example, food waste recovery initiatives turn losses into valuable products, thereby reducing waste and contributing to more sustainable management of agricultural resources. Awareness campaigns and training could be provided to encourage farmers to adopt these processing techniques, while partnerships with local businesses could facilitate the collection and processing of surpluses.

4.4.Recovery of by-products

Processing RTPs into semolina generates by-products that can be used in other sectors, such as animal feed, composting, or even the production of bioplastics^{74,153}. Processing residues can be used to feed livestock or to enrich soil, promoting a circular economy and sustainable agricultural practices¹⁵⁴. By-products could also produce energy (biogas) or biodegradable materials, contributing to a greener, more sustainable economy. Further research could explore these opportunities to maximize by-product recovery. In addition, the recovery of agricultural by-products could lead to new business models focused on sustainability and resource efficiency¹⁵⁵. By-product recovery reduces waste and maximizes resource use while creating new economic opportunities.

4.5.Creating jobs and improving incomes for farmers

Processing into semolina creates significant economic opportunities, mainly by creating jobs in the processing, marketing, and distribution sectors^{8,50}. This activity is particularly beneficial for women and youth, who find this sector a stable source of income and an opportunity for empowerment¹⁵⁶. In rural areas with limited employment opportunities, these activities improve farmers' incomes and contribute to community development and economic resilience¹⁵⁷. By selling processed rather than raw products, farmers can increase their incomes and access broader markets, contributing to the economic development of rural communities. In addition, ongoing education and training programs could be introduced to help farmers and local workers master new farming practices, technologies, and techniques for processing and marketing semolina, helping them adapt to market demands and improve their livelihoods¹⁹. In addition, agricultural cooperatives could be promoted to facilitate access to markets and increase the incomes of small producers.



5.0.Challenges in processing roots, tubers, and plantains into semolina

The processing of RTPs into semolina is critical to sub-Saharan Africa's agricultural and economic development. However, this process faces several challenges that hinder its widespread adoption, including access to technology, training of local actors, access to markets, and competitiveness, as well as obstacles to the widespread adoption of these practices on an industrial scale. Each aspect deserves special attention to promote the processing and add value to local products.

5.1.Access to processing technologies

Access to processing technologies is challenging when producing semolina from RTPs in SSA. Modern technologies are essential to improving the productivity and quality of processed products. However, their adoption remains limited due to insufficient infrastructure and adequate finance, especially in remote areas. Agbe and Atake¹⁵⁸ note that agricultural credit could enable farmers to acquire modern equipment, but the majority of smallholder farmers often suffer from a lack of adequate finance ¹⁵⁹. This lack of technology prevents local processors from adopting modern techniques that could improve yields and product consistency. In addition, Forsythe et al. ¹⁶⁰ note that plant breeding programs (RTBs) do not always consider the needs of processors and consumers, which hinders the adoption of improved varieties. Similarly, Jideani et al.¹⁶¹, explain that the rise of industrialization has marginalized traditional food processing methods, reflecting the impact of inadequate technology on local practices. To overcome these barriers, it is crucial to increase investment in infrastructure and promote public policies that encourage technological innovation in the agricultural sector.

5.2.Training local stakeholders

Training local stakeholders is key to efficiently processing RTPs in Semolina. Many farmers and processors lack the knowledge to use available technologies and adopt optimal practices. Temple and Bon suggest that appropriate training programs could improve the productivity and quality of processed products ¹⁶². Forsythe et al. also emphasize the importance of understanding community contexts when implementing new practices, as local knowledge contributes significantly to the successful adoption of innovations ¹⁶⁰. By integrating traditional



knowledge with modern techniques, as Kouakou et al. suggest, communities could increase the efficiency of processing ¹⁶³. The ability to process RTP into semolina requires not only technical skills but also an understanding of market preferences and nutritional implications, as Adeyeye's review of food processing shows ¹⁶⁴. However, the lack of educational infrastructure and financial support for agricultural training remains a significant constraint, requiring increased investment in education and local capacity building.

5.3. Market Access and Competitiveness

Access to markets is a major challenge hampering the competitiveness of semolinas that process RTPs. Processed products must meet strict quality standards to compete with imports, but systemic problems such as inadequate transport infrastructure, logistical challenges, lack of financial capital to efficiently scale up production, and regulatory barriers limit access to national and international markets ¹⁶⁵. Nzossié and Temple⁴¹, show that local producers struggle to compete with imported wheat flour, which is often subsidized, putting pressure on local prices. Economic Partnership Agreements (EPAs), with their tariff exemptions, put local products at a disadvantage compared to imported products that meet strict international food standards ¹⁶¹. Furthermore, Royen et al.¹⁶⁶ points out that economic and logistical constraints in SSA limit opportunities for local producers. Developing appropriate market strategies, strengthening transport infrastructure, and supporting local producers with favorable trade policies are essential to improving competitiveness.

5.4. Large-scale adoption barriers

Socio-economic and cultural barriers hamper the adoption of large-scale RTP processing technologies. The high costs associated with modernization often deter smallholder farmers from investing in improved technologies ¹⁶⁰. These barriers are closely linked to cultural perceptions and historical practices, as illustrated by the gender dynamics in food processing identified by Okoye et al. ¹⁶⁷. Gender disparities in agricultural roles and decision-making can complicate the path to broader adoption of advanced processing methods. ¹⁶⁸, also point out that climate instability and food security issues discourage long-term investment. Given the socio-economic and cultural contexts, it is clear that a multifaceted approach is needed to address the challenges associated with processing RTP. Adeyeye argues that innovative food processing technologies are key to improving food security in Africa; their success is highly dependent on local engagement and investment strategies tailored to farmers' needs ¹⁶⁴.



Overcoming these barriers requires targeted studies of community needs and public policies that support adopting innovative technologies while integrating local realities and traditional practices.

6.0.Prospective

Developing integrated traceability systems accessible to local stakeholders is crucial to enable the adoption of technological solutions (IoT, blockchain) for transparent product tracking across the value chain. Standardizing traceability standards, aligned with existing certifications, would enhance access to international markets for traditional sub-Saharan African food crops while raising awareness among local actors (producers, distributors, processors, traders) about traceability’s role in reducing losses and boosting consumer trust. A practical example would be the use of devices connected (tablets, smartphones, etc.) to smart sensors via predictive algorithms to monitor real-time weather conditions (temperature, humidity, rainfall), soil irrigation, and the presence of any contaminants during cultivation, harvesting, storage, and transport. Partnerships with entities possessing advanced technologies, such as Central Africa’s Rader Group, could support implementation. Additionally, assessing the technical and economic feasibility of predictive algorithms would help anticipate spoilage risks and optimize post-harvest logistics in rural and peri-urban settings, accounting for energy and financial constraints.

To maximize the impact of this approach, the following measures are essential. Strengthen public policies to support local value chains. Governments should promote regulatory frameworks and financial incentives that favor local processing and encourage investment in this sector. They should also promote affordable and sustainable technologies like solar cookers or semi-automatic dryers. These innovations would help reduce the environmental footprint while improving the efficiency of artisanal processing. Strengthen the technical skills of producers and processors through training and extension programs tailored to the specific needs of rural communities, thereby making a significant contribution to optimizing the quality and profitability of finished products. Encourage public-private partnerships to facilitate access to finance and appropriate infrastructure. Such collaborations could help mobilize the resources needed to modernize processing facilities while preserving artisanal and cultural aspects. Existing studies highlight these products’ cultural significance and the challenges of modernizing traditional methods. Future research should refine mechanical processing,



fermentation control, and optimization techniques to enhance safety without compromising traditional qualities. Further opportunities include expanding geographical research coverage, standardizing quality metrics, and developing safety innovations that align with indigenous practices.

7.0. Conclusion

In conclusion, the artisanal processing of RTP into semolina is an essential strategy for addressing the challenges of post-harvest losses, food security, and socio-economic development in SSA. The scientific data collected and analyzed in this review show that this practice can significantly reduce post-harvest losses by improving product preservation and economic value. It is also a viable alternative for strengthening food systems' resilience, improving rural communities' livelihoods, and responding to climatic and economic challenges. It represents a holistic approach that transforms a persistent problem into an opportunity for sustainable and inclusive development.

The associated socio-economic benefits are manifold: this activity could generate additional income, particularly for rural women, strengthen local food security, and help reduce dependence on food imports. What is more, small-scale semolina production is based on technologies that are accessible and adapted to local resources, making it easier to adopt in rural areas.

However, several challenges remain: limited access to modern equipment and adequate infrastructure, and the actors' low level of technical training continues to hamper the widespread adoption of these practices. Regional disparities are also observable, with tremendous success in areas benefiting from institutional support programs and proven technology transfer initiatives. The analysis revealed limited research coverage outside West Africa, inconsistent measurement methods across studies, and minimal data on m'bahou production.

Finally, the local semolina agro-industry represents an important opportunity to transform food systems in sub-Saharan Africa. More research is needed to explore further the collaborations between artisanal practices and modern industrial solutions and to assess their long-term impact on community food resilience. While challenges remain, this approach offers immense potential to ensure future generations' food security and economic prosperity.



Acknowledgments

The work presented in this paper was carried out under the **CSIR-TWAS Sandwich Postgraduate Research Fellowship (22/FF/CSIR-TWAS/2024)** awarded to the first author, and CSIR-TWAS is gratefully acknowledged. The authors thank the CSIR-Central Food Technological Research Institute, Mysuru, India, for providing all the necessary facilities to conduct this research. The authors would also like to thank Pr. Nguimbou Richard Marcel and Dr. Bouelet Ntsama Isabelle Sandrine, teacher-researchers at ENSAI, University of Ngaoundere, and ENSET, University of Douala, respectively, for their guidance during the local editing of the manuscript.

Mundéné-Timothée Junior Lawrence: Writing – original draft, Visualization, Validation, Investigation, Graphics, Conceptualization. **Veeranna Hitlamani:** Writing – review & editing, Visualization, Validation, Investigation, Graphics. **Nouga Bissoue Achille:** Writing – review & editing, Supervising. **Aashitosh A Inamdar:** Writing – review & editing, Visualization, Supervision, Resources, Conceptualization. **Mouangue Ruben:** Writing – review & editing, Supervising. **Njintang Yanou Nicolas:** Writing – review & editing, Supervising



References

- 1 Bricas N and Walser M. L'évolution des habitudes alimentaires. *Une écologie de l'alimentation* :91 (2021).
- 2 Bricas N, Tchamda C, and Martin P. Les villes d'Afrique de l'Ouest et du Centre sont-elles si dépendantes des importations alimentaires? *Cahier Agricultures* **25**:1–10 (2016).
- 3 N'Goran CA, Petit J, N'Guessan AA, Gonnety JT, and Scher J. Improvement of the production process and the sensory and nutritional quality of m'bahou, a traditional plantain semolina, enriched with soy or cowpea. *International Journal of Food Science and Technology* **58**:4171–4185 (2023).
- 4 Diallo Y, Talla Gueye M, Sakho M, Gbaguidi Darboux P, Kane A, Barthelemy J-P, and Lognay G. Importance nutritionnelle du manioc et perspectives pour l'alimentation de base au Sénégal (synthèse bibliographique). *Biotechnol Agron Soc Environ* (2013).
- 5 FAO. Agricultural production statistics 2000–2021. Rome, Italy: Food and Agriculture Organization of the United Nations; p. 17 2022Report No.: 60.
- 6 N'Guessan AA, Sinh J-N, N'Gbo KML, Disseka KW, Djina Y, N'Goran CA, and Gonnety JT. Physicochemical and sensory characteristics of m'bahou: A traditional dish consumed in Côte d'Ivoire. *GSC Biological and Pharmaceutical Sciences* **18**:182–192 (2022).
- 7 Vernier P, N'Zué B, and Zakhia-Rozis N. Le manioc, entre culture alimentaire et filière agro-industrielle. éditions Quae; 2018.
- 8 Chiapo AC and Akahoua SN. Determinants of post-harvest losses of plantain in the Sassandra-Marahou district, Cte dlvoire. *J Dev Agric Econ* **16**:26–33 (2024).
- 9 Mundéné-Timothée JL, Nguimbou RM, Bissoue AN, Ntsama ISB, Tamo SPB, Agamou JAA, Ngane RAN, Mouangue R, and Yanou NN. Cooking practices, consumption and sensory perception of Ntuba ekōn: A traditional dish consumed in Cameroon. *Journal of Food Stability* **7**:21–37 (2024).
- 10 Anyoha N-PO, Udemba C, Ogbonnaya A, and Okoroma E. Causes Of Cassava Post-Harvest Losses Among Farmers In Imo State, Nigeria. *Journal of Agricultural Extension* **27**:73–79 (2023).
- 11 Atanda SA, Agoda S, and Ikotun I. The concepts and problems of post-harvest food losses in perishable crops. *African Journal of Food Science* **5**:603–6013 (2011).
- 12 Lepengue A, Mouaragadja I, Dick E, Mbatchi B, and Ake S. Amélioration de la durée de conservation des bananes plantain aux températures ambiantes. *International Journal of Biological and Chemical Sciences* **4** (2010).
- 13 Muroyiwa B, Shokopa L, Likoetla P, and Rantlo M. Integration of post-harvest management in agricultural policy and strategies to minimise post harvest losses in Lesotho. *JDAE Academic Journals*; **12**:84–94 (2020).
- 14 Benoit-Cattin M and Dorin B. Disponible alimentaire et productivité agricole en Afrique subsaharienne. *Cahiers Agricultures* **21**:337–347 (1) (2012).





1114 15 Leisher C, Robinson N, Brown M, Kujirakwinja D, Schmitz MC, Wieland M, and Wilkie D. Ranking the direct threats to biodiversity in sub-Saharan Africa. *Biodivers Conserv* **31**:1329–1343 (2022).
1115

1116 16 Janin P and Dury S. The new frontiers of food security. A prospective review. *Cahiers Agricultures* **21**:285–292 (2012).
1117

1118 17 Mundéné-Timotheé JL, Nougá Bissoué A, Nguimbou RM, Bissim SM, Bouelet Ntsama IS, Bouopda Tamo SP, Fokam L, Mouangue R, and Njintang Yanou N. Plantain flour: production processes, technological characteristics, and its potential use in traditional African dishes - a review. *J Sci Food Agric* (2024).
1119
1120
1121

1122 18 Adeniyi OR and Ayandiji A. Economic Analysis of Post Harvest Losses in Plantain (and Banana): A Case Study of South Western Nigeria. *British Journal of Applied Science & Technology* **4**:4456–4467 (2014).
1123
1124

1125 19 Degebase AC. Prospects and Challenges of Postharvest Storage and Losses of Potato (*Solanum tuberosum* L.) in Central highlands of Ethiopia: A Review. *Journal of Natural Sciences Research* **10**:36 (2020).
1126
1127

1128 20 FAO. Réduction des pertes après récolte pour une meilleure sécurité alimentaire et nutritionnelle dans les États membres de l'IGAD. Rome, Italy: Bureau sous-régional de la FAO pour l'Afrique orientale; p. 11 2022 MayReport No.: TCP/SFE/3702.
1129
1130

1131 21 Dania VO. Bioefficacy of *Trichoderma* species against important fungal pathogens causing post-harvest rot in sweet potato (*Ipomoea batatas* (L.) Lam): Bioefficacy of *Trichoderma* metabolites of sweetpotato. *Journal of the Bangladesh Agricultural University* **17**:446–453 (2019).
1132
1133

1134 22 Rees D, Kapinga R, Mtunda K, Chilosa D, Mbilinyi L, Rwiza E, Kilima M, Kiozya H, Amour R, Ndondi T, Chottah M, Mayona D, Tomlins K, Aked J, Carey E, and Oirschot QV. Sweetpotato post-harvest assessment: experiences from East Africa. Chapter 5. Extending root shelf-life during marketing by cultivar selection. *The University of Greenwich* Chatham, UK: Natural Resources Institute; p. 51–66 2003.
1135
1136
1137
1138

1139 23 Pirzada T, Affokpon A, Guenther RH, Mathew R, Agate S, Blevins A, Byrd MV, Sit TL, Koenning SR, Davis EL, Pal L, Opperman CH, and Khan SA. Plant-biomass-based hybrid seed wraps mitigate yield and post-harvest losses among smallholder farmers in sub-Saharan Africa. *Nat Food* Nature Publishing Group; **4**:148–159 (2023).
1140
1141
1142

1143 24 Ikechi-Nwogu CG and Nworuka N. Isolation And Identification of Common Fungal Pathogens Invading Sweet Potatoes (*Ipomoea batatas*) Sold in Choba Market, Port Harcourt, Nigeria. *Journal of Applied Sciences and Environmental Management* **27**:43–49 (2023).
1144
1145

1146 25 Sonnewald S and Sonnewald U. Regulation of potato tuber sprouting. *Planta* **239**:27–38 (2014).

1147 26 Liu T, Wu Q, Zhou S, Xia J, Yin W, Deng L, Song B, and He T. Molecular Insights into the Accelerated Sprouting of and Apical Dominance Release in Potato Tubers Subjected to Post-Harvest Heat Stress. *International Journal of Molecular Sciences* Multidisciplinary Digital Publishing Institute; **25**:1699 (2024).
1148
1149
1150

1151 27 Essis B, Soro S, Samaké A, Hala KA, Dibi K, Kouamé KT, Ehounou A, N'Zué B, and Amani Michel K. Inventory of major post-harvest deterioration agents of the national yam genetic resources collection at the cnra food crops research station in Bouaké. *International Journal of Agriculture and Environmental Research* **10**:810–828 (2024).
1152
1153
1154

- 1155 28 Nwankwo EC and Chiekezie NR. Agricultural Productivity and Postharvest Loss Among Cassava
1156 Farmers, In Anambra State, Nigeria. *Global Journal of Agricultural Research* **12** (2024).
- 1157 29 Upadhyay KP, Paudel N, Aryal S, Simkhada R, Bhusal B, Thapa B, Subedi G, and Gautam IP. Post-
1158 Harvest Losses of Potato Genotypes at Farmers' Storage Conditions. *Sustainability in Food and*
1159 *Agriculture* **2**:51–56 (2021).
- 1160 30 Mohamed A, Liet Z, Goru D, and David J. Losses after Harvesting and Management. *E3S Web of*
1161 *Conferences* **477** (2024).
- 1162 31 Chiekezie JC, Chiekezie NR, Nwankwo EC, and Ozor MU. Assessment of the extent of post-
1163 harvest losses along the cassava value chain in Anambra state. *Journal of Advance Research in*
1164 *Food, Agriculture and Environmental Science (ISSN 2208-2417)* **9**:1–6 (2023).
- 1165 32 Kuyu CG, Tola YB, and Abdi GG. Study on post-harvest quantitative and qualitative losses of
1166 potato tubers from two different road access districts of Jimma zone, South West Ethiopia.
1167 *Heliyon* **5**:e02272 (2019).
- 1168 33 Kouadio Y, Kouassi K, Kouassi K, Ouattara D, N'dri Y, and Amani N. Practices and damages
1169 encountered by wholesale traders during 'Kponan' yams (*Dioscorea cayenensis-rotundata*)
1170 storage in Abidjan, Côte d'Ivoire. *International Journal of Biological and Chemical Sciences*
1171 **16**:1570–1579 (2022).
- 1172 34 Sugri I and Johnson P. Effect of two storage methods on the keeping and sensory qualities of
1173 four plantain varieties. *African Journal of Food, Agriculture, Nutrition and Development* **9** (2009).
- 1174 35 Kaminski J and Christiaensen L. Post-harvest loss in sub-Saharan Africa—what do farmers say?
1175 *Global Food Security* **3**:149–158 (2014).
- 1176 36 Kitinoja L and AlHassan HY. Identification of Appropriate Postharvest Technologies for Small
1177 Scale Horticultural Farmers and Marketers in Sub-Saharan Africa and South Asia - Part 1.
1178 Postharvest Losses and Quality Assessments. *Acta Horti* :31–40 (2012).
- 1179 37 Naziri D, Quaye W, Siwoku B, Wanlapatit S, Tu Viet Phu, and Bennett B. The diversity of
1180 postharvest losses in cassava value chains in selected developing countries. *Journal of*
1181 *Agriculture and Rural Development in the Tropics and Subtropics* Kassel University Press;
1182 **115**:111–123 (2014).
- 1183 38 Issa AH, Doumma A, and Bello MT. Inventaire des variétés, des méthodes locales de stockage et
1184 de protection contre les ravageurs de la patate douce (*Ipomea batatas* L.) dans la bande Ouest
1185 du Niger. *International Journal of Biological and Chemical Sciences* **9**:1962–1971 (2015).
- 1186 39 Djinadou AKA, Olodo NI, and Adjanohoun A. Evaluation du comportement de variétés
1187 améliorées de manioc riches en bêta-carotène au Sud du Bénin. *International Journal of*
1188 *Biological and Chemical Sciences* **12**:703–715 (2018).
- 1189 40 Kouadio KKH, Ettien DJB, Bakayoko S, Soro D, and Girardin O. Variabilité physico-morphologique
1190 des racines tubéreuses de manioc (*Manihot esculenta* CRANTZ) cultivées sur ferralsol en zone de
1191 forêt d'Afrique de l'Ouest. *Journal of Applied Biosciences* **82**:7316–7325 (2014).
- 1192 41 Nzossié EJF and Temple L. Politique d'import-substitution au blé et compétitivité des farines
1193 panifiables à base de manioc, banane plantain et patate douce au Cameroun. *Cah Agric EDP*
1194 *Sciences*; **32**:25 (2023).





1195 42 Archimède H, Bastianelli D, Boval M, Tran G, and Sauvant D. Ressources tropicales : disponibilité
1196 et valeur alimentaire. *INRAE Productions Animales* **24**:23–40 (2011). Article Online
DOI: 10.1039/D5FB00326A

1197 43 Yeo MA, Koffi EK, Chatigre OK, and Elleingand EF. Prévention du brunissement lors de
1198 l'extraction de jus de banane plantain et valorisation des résidus issus de cette extraction |
1199 Request PDF. *Int J Biol Chem Sci* **9**:1809–1821 (2015).

1200 44 Kouakou K.A K, M. B K, L. C S, and -Anin Atchibri L O. Storing Plantain Using A Traditional Cold
1201 Jar. *IJSE* **2**:114–120 (2022).

1202 45 Ji CY, Jin R, Xu Z, Kim HS, Lee C-J, Kang L, Kim S-E, Lee H-U, Lee JS, Kang CH, Chi YH, Lee SY, Xie Y,
1203 Li H, Ma D, and Kwak S-S. Overexpression of Arabidopsis P3B increases heat and low
1204 temperature stress tolerance in transgenic sweetpotato. *BMC Plant Biology* **17**:139 (2017).

1205 46 Rwubatsé B, Akubor PI, and Mugabo E. Traditional Drying Techniques for Fruits and Vegetables
1206 Losses Alleviation in Sub-Saharan Africa. *IOSR Journal of Environmental Science, Toxicology and*
1207 *Food Technology* **8**:52–56 (2014).

1208 47 Kimani S. The Role of Agricultural Innovation in Enhancing Food Security in Sub-Saharan Africa.
1209 *International Journal of Developing Country Studies* **6**:58–73 (2024).

1210 48 Vutula N. The Reduction of Post-harvest Losses is Crucial for a Successful Cassava Value Chain
1211 and Food Security in Africa. *The Open Agriculture Journal* Bentham Science Publishers; **18**:1–11
1212 (2024).

1213 49 Onzo A, Biaou JT, Loko LY, Tamo M, and Dansi A. Vulnérabilité des cossettes issues de quelques
1214 cultivars d'igname à l'attaque de *Dinoderus porcellus* Lesne (Coleoptera: Bostrichidae) en
1215 conditions de laboratoire. *International Journal of Biological and Chemical Sciences* **8**:2494–2507
1216 (2014).

1217 50 Allen T, Heinrigs P, and Heo I. Agriculture, alimentation et emploi en Afrique de l'Ouest. *Notes*
1218 *ouest-africaines* **14**:36 (2018).

1219 51 Thomas A, Alpha A, Barczak A, and Zakhia-Rozis N (eds.). Durabilité des systèmes pour la
1220 sécurité alimentaire: Combiner les approches locales et globales. Quæ. Versailles: éditions Quæ;
1221 2024.

1222 52 Dury S, Vall É, and Imbernon J. Production agricole et sécurité alimentaire en Afrique de l'Ouest.
1223 *Cah Agric* EDP Sciences; **26**:61001 (2017).

1224 53 Mabhaudhi T, Mpandeli S, Nhamo L, Chimonyo VGP, Nhemachena C, Senzanje A, Naidoo D, and
1225 Modi AT. Prospects for Improving Irrigated Agriculture in Southern Africa: Linking Water, Energy
1226 and Food. *Water* Multidisciplinary Digital Publishing Institute; **10**:1881 (2018).

1227 54 Afzal N, Afionis S, Stringer LC, Favretto N, Sakai M, and Sakai P. Benefits and Trade-Offs of
1228 Smallholder Sweet Potato Cultivation as a Pathway toward Achieving the Sustainable
1229 Development Goals. *Sustainability* Multidisciplinary Digital Publishing Institute; **13**:552 (2021).

1230 55 Sakai P, Afionis S, Favretto N, Stringer LC, Ward C, Sakai M, Weirich Neto PH, Rocha CH, Alberti
1231 Gomes J, Souza NM de, and Afzal N. Understanding the Implications of Alternative Bioenergy
1232 Crops to Support Smallholder Farmers in Brazil. *Sustainability* Multidisciplinary Digital Publishing
1233 Institute; **12**:2146 (2020).

- 1234 56 Yıldız BC, Demir EF, and Hancı F. Some Bioactive Components of Sweet Potato and Taro: A Comparative Study Based On Plant Organs. *J Inst Sci and Tech Igdir University*; **13**:2315–2324 (2023). [View Article Online](#)
DOI: 10.1039/D5FB00326A
- 1237 57 Hamidov A, Helming K, Bellocchi G, Bojar W, Dalgaard T, Ghaley BB, Hoffmann C, Holman I, Holzkämper A, Krzeminska D, Kværnø SH, Lehtonen H, Niedrist G, Øygarden L, Reidsma P, Roggero PP, Rusu T, Santos C, Seddaiu G, Skarbøvik E, Ventrella D, Żarski J, and Schönhart M. Impacts of climate change adaptation options on soil functions: A review of European case-studies. *Land Degrad Dev* **29**:2378–2389 (2018).
- 1242 58 Altieri MA, Funes-Monzote FR, and Petersen P. Agroecologically efficient agricultural systems for smallholder farmers: contributions to food sovereignty. *Agron Sustain Dev* **32**:1–13 (2012).
- 1244 59 Lal P, Tiwari RK, Behera B, Yadav MR, Sharma E, Altaf MA, Jena R, Ahmad A, Dey A, Kumar A, Singh B, Lal MK, and Kumar R. Exploring potato seed research: a bibliometric approach towards sustainable food security. *Front Sustain Food Syst* **7** (2023).
- 1247 60 Zhang J, Huang L, Tang X, Pan M, Ye X, Li X, and Ma H. High β -sitosterol-*D*-glucoside content in sweet potato varieties and its anti-breast cancer potential through multiple metastasis-associated signals. *Food Science and Human Wellness* **13**:2779–2789 (2024).
- 1250 61 Lim S, Xu J, Kim J, Chen T-Y, Su X, Standard J, Carey E, Griffin J, Herndon B, Katz B, Tomich J, and Wang W. Role of anthocyanin-enriched purple-fleshed sweet potato p40 in colorectal cancer prevention. *Molecular Nutrition & Food Research* **57**:1908–1917 (2013).
- 1253 62 Carvalho DG, Trierweiler LF, and Trierweiler JO. Production of sweet potato distilled beverage: Economic evaluation via enzymatic and acid hydrolysis. *Biofuels, Bioproducts and Biorefining* **18**:1880–1891 (2024).
- 1256 63 Hagenimana T. Food safety priority, a critical gap and a window for effective food, nutrition security and sustainable development in Rwanda: A contextual analysis. *Journal of Regulatory Science* **10**:1–7 (2022).
- 1259 64 Hinz R, Sulser TB, Huefner R, Mason-D'Croz D, Dunston S, Nautiyal S, Ringler C, Schuengel J, Tikhile P, Wimmer F, and Schaldach R. Agricultural Development and Land Use Change in India: A Scenario Analysis of Trade-Offs Between UN Sustainable Development Goals (SDGs). *Earth's Future* **8**:e2019EF001287 (2020).
- 1263 65 Calicioglu O, Flammini A, Bracco S, Bellù L, and Sims R. The Future Challenges of Food and Agriculture: An Integrated Analysis of Trends and Solutions. *Sustainability Multidisciplinary Digital Publishing Institute*; **11**:222 (2019).
- 1266 66 Olaniran AF, Okonkwo CE, Iranloye YM, Morakinyo OO, Taiwo AE, Erinle OC, Bamidele OP, Ojo OA, Malomo AA, and Osemwegie OO. Fermented Gluten-Free Multi-Grain Cereal Paste Development: The Role of the Orange-Fleshed Sweet Potato (OFSP) as a Dietary Supplement. *Nutr Metab Insights* **16**:11786388231155007 (2023).
- 1270 67 Muruu D, Kinyua J, Kepue M, Kerubo L, Njaci I, and Mware B. Diversity of viruses and viroids infecting Taro in Kenya based on small RNA sequencing and pcr detection. *American Journal of Agriculture AJPO JOURNALS*; **4**:15–33 (2022).



Open Access Article. Published on 19 September 2025. Downloaded on 10/7/2025 1:43:32 AM.
This article is licensed under a Creative Commons Attribution 3.0 Unported Licence.



1273 68 Kidanemariam DB, Sukal AC, Abraham AD, Stomeo F, Dale JL, James AP, and Harding RM. Identification and molecular characterization of Taro bacilliform virus and Taro bacilliform CH
1274 virus from East Africa. *Plant Pathol* **67**:1977–1986 (2018).
1275

1276 69 Nkwain KT, Odiaka EC, Ikwuba AA, and Nkwi GE. Analysis of post-harvest losses of banana and
1277 the economic wellbeing of farmers in Boyo Division, North West Region of Cameroon. *Dutse*
1278 *Journal of Pure and Applied Sciences* **7**:undefined-undefined (2022).

1279 70 Djè KM and Eba KP. Influence of water cooking on the macronutrients loss rate of yam tuber
1280 parts during post-harvest storage. *Research Journal of Life Sciences, Bioinformatics,*
1281 *Pharmaceutical and Chemical Sciences* **04**:122–133 (2018).

1282 71 Wang B, Huang Y, Zhang Z, Xiao Y, and Xie J. Ferulic Acid Treatment Maintains the Quality of
1283 Fresh-Cut Taro (*Colocasia esculenta*) During Cold Storage. *Front Nutr* **9**:884844 (2022).

1284 72 Diaguna R, Santosa E, and Budiman C. The Effect of the Taro Corm Type and Storage on
1285 Morphophysiological Deterioration and Early Growth in the Field. *International Journal of*
1286 *Agronomy* **2023**:3280339 (2023).

1287 73 Pierrot J, Achard R, Temple L, Abadie C, and Fogain R. Déterminants de la production de plantain
1288 dans le sud-ouest du Cameroun : intérêt d'un observatoire. *Fruits* **57**:75–86 (2002).

1289 74 Kouadio Kouakou KA, Kone MB, and Ocho-Anin Atchibri L. Inventaire des méthodes de stockage
1290 des bananes plantain dans la localité de Man. *JCBPS* **11**:283–291 (2021).

1291 75 Huffstetler E. Learn All About Semolina, the Essential Flour Used To Make Pasta. The Spruce
1292 Eats. 2024. Available: <https://www.thespruceeats.com/easy-semolina-flour-substitute-4142772>
1293 [17 March 2025].

1294 76 Posner ES. MILLING | Characteristics of Milled Products. In: Caballero B, ed. *Encyclopedia of*
1295 *Food Sciences and Nutrition (Second Edition)* Oxford: Academic Press; p. 3997–4005 2003.

1296 77 Zamarripa M. Semolina: Nutrition, Benefits, Uses, and Downsides. Healthline. 2023. Available:
1297 <https://www.healthline.com/nutrition/semolina> [17 March 2025].

1298 78 Awoyale W, Alamu E, Chijioke U, Tran T, Tchuenta HN, Ndjouenkeu R, Ngoualem F, and Maziya-
1299 Dixon B. A review of cassava semolina (gari and eba) end-user preferences and implications for
1300 varietal trait evaluation. *International Journal of Food Science & Technology* **56** (2020).

1301 79 Flibert G, Siourime SN, Hamidou C, Donatien K, Cissé H, Ulrich MNJ, Tankoano A, Hagretou S-L,
1302 and Aly S. Lactic Acid Bacteria and Yeasts Associated with Cassava Fermentation to attiéké in
1303 Burkina Faso and Their Technological Properties. *American Journal of Food Science and*
1304 *Technology* Science and Education Publishing; **9**:173–184 (2021).

1305 80 Abel BZBI, Jean-Paul BKM, Alfred KK, Hermann CW, Celah KA, Nevry RK-, and Marcellin DK.
1306 Technical Sheet of the Preparation of Traditional Cassava Starters Used for Attieke Production in
1307 Côte d'Ivoire. *Biotechnology Journal International* :11–20 (2020).

1308 81 Assanvo JB, Agbo GN, Beez JB, Monsan V, and Farah Z. Sensory Profiling and Hedonic Evaluation
1309 of Attiéké from Local and Improved Cassava Varieties. *Food and Nutrition Sciences* Scientific
1310 Research Publishing; **9**:1472–1497 (2018).

- 1311 82 Toka MD, Djéni TN, and Dje MK. Improved Process of Cassava Processing into “Attiéké” a
 1312 Traditional Food Product of Côte D’Ivoire. *International Journal of Food Engineering* De Gruyter;
 1313 **4** (2008). View Article Online
DOI: 10.1039/D5FB00326A
- 1314 83 Flibert G, Donatien K, Hagrétou S-L, and Aly S. Hygienic Quality and Nutritional Value of Attiéké
 1315 from Local and Imported Cassava Dough Produced with Different Traditional Starters in Burkina
 1316 Faso. *Food and Nutrition Sciences* Scientific Research Publishing; **7**:555–565 (2016).
- 1317 84 Yéboué KH, Amoikon KE, Kouamé KG, and Kati-Coulibaly S. Valeur nutritive et propriétés
 1318 organoleptiques de l’attiéké, de l’attoukpou et du placali, trois mets à base de manioc,
 1319 couramment consommés en Côte d’Ivoire. *Journal of Applied Biosciences* **113**:11184–11191
 1320 (2017).
- 1321 85 Djeni NT, Bouatenin KMJ-P, Assohoun NMC, Toka DM, Menan EH, Dousset X, and Dje KM.
 1322 Biochemical and microbial characterization of cassava inocula from the three main attieke
 1323 production zones in Côte d’Ivoire. *Food Control* **50**:133–140 (2015).
- 1324 86 Djeni NT, N’Guessan KF, Toka DM, Kouame KA, and Dje KM. Quality of attieke (a fermented
 1325 cassava product) from the three main processing zones in Côte d’Ivoire. *Food Research*
 1326 *International* **44**:410–416 (2011).
- 1327 87 Mohidin SRNSP, Moshawih S, Hermansyah A, Asmuni MI, Shafqat N, and Ming LC. Cassava
 1328 (Manihot esculenta Crantz): A Systematic Review for the Pharmacological Activities, Traditional
 1329 Uses, Nutritional Values, and Phytochemistry. *J Evid Based Complementary Altern Med* SAGE
 1330 Publications Inc STM; **28**:1–26 (2023).
- 1331 88 Assamoi AA, Regina Krabi E, Fafadzi Ehon A, Amani N’guessan G, Sébastien Niamké L, and
 1332 Thonart P. Isolation and screening of Weissella strains for their potential use as starter during
 1333 attiéké production. *Biotechnol Agron Soc Environ* **20**:355–362 (2016).
- 1334 89 Illoh C, Agu HO, Okolo CA, and Jideani VA. Physicochemical and consumer acceptability of
 1335 different particle-sized cassava strips. *AJFS Academic Journals*; **16**:71–80 (2022).
- 1336 90 Konan MK, Assamoi AA, and Germain KT. Diversity of yeast strains isolated from the “magnan” a
 1337 traditional ferment used for the production of “attiéké” a cassava food in Côte d’Ivoire. *Journal*
 1338 *of Applied Biosciences* **136**:13861–13867 (2019).
- 1339 91 Assanvo JB, Agbo GN, Behi YEN, Coulin P, and Farah Z. Microflora of traditional starter made
 1340 from cassava for “attiéké” production in Dabou (Côte d’Ivoire). *Food Control* **17**:37–41 (2006).
- 1341 92 Diby NAS, Deffan KP, Adinsi L, Bechoff A, Kanon AL, Bouniol A, Yapi YE, Deuscher Z, Bugaud C,
 1342 N’Zué B, and Ebah Djedji CB. Use of sensory and physico-chemical parameters to understand
 1343 consumer perception of attiéké, a fermented cassava product. *Journal of the Science of Food*
 1344 *and Agriculture* **104**:4596–4605 (2024).
- 1345 93 Oloya B, Adaku C, Andama M, Oloya B, Adaku C, and Andama M. The Cyanogenic Potential of
 1346 Certain Cassava Varieties in Uganda and Their Fermentation-Based Detoxification. *Cassava -*
 1347 *Recent Updates on Food, Feed, and Industry* IntechOpen; 2023.
- 1348 94 Knez E, Kadac-Czapska K, and Grembecka M. Effect of Fermentation on the Nutritional Quality of
 1349 the Selected Vegetables and Legumes and Their Health Effects. *Life Multidisciplinary Digital*
 1350 *Publishing Institute*; **13**:655 (2023).





1351 95 Daouda N, Achille TF, Abodjo KC, Charlemagne N, and Georges AN. Influence of Traditional View Article Online
DOI: 10.1039/D5FB00326A
1352 Inoculum and Fermentation Time on the Organoleptic Quality of "Attiéké". *Food and Nutrition*
1353 *Sciences Scientific Research Publishing*; **3**:1335–1339 (2012).

1354 96 Thierry APM, Kouakoua YE, Abouo NV, Coulibaly A, and Amani NG. Optimisation of the "attiéké"
1355 Fermentation Process: Evaluation of the Efficiency of Various Starter Ferments. *J Food Nutr Sci*
1356 *Science Publishing Group*; **12**:261–269 (2024).

1357 97 Samson SO, O.m A, and F.k O. Fermentation: A Means of Treating and Improving the Nutrition
1358 Content of Cassava (*Manihot esculenta* C.) Peels and Reducing Its Cyanide Content. *Genomics*
1359 *and Applied Biology* **8** (2017).

1360 98 Oloya B, Adaku C, Ntambi E, and Andama M. Detoxification of Nyar-Udota Cassava Variety in
1361 Zombo District by Fermentation. *International Journal of Nutrition and Food Sciences* **6**:118–121
1362 (2017).

1363 99 Ismaila A. R, Alakali J. S, and Atume T. G. Effect of Local Processing Techniques on the Nutrients
1364 and Anti-Nutrients Content of Bitter Cassava (*Manihot Esculenta* Crantz). *American Journal of*
1365 *Food Science and Technology Science and Education Publishing*; **6**:92–97 (2018).

1366 100 Ojo O and Deane R. Effects of cassava processing methods on antinutritional components and
1367 health status of children. *Journal of the Science of Food and Agriculture* **82**:252–257 (2002).

1368 101 Lambebo T and Deme T. Evaluation of Nutritional Potential and Effect of Processing on
1369 Improving Nutrient Content of Cassava (*Mannihot esculenta* crantz) Root and Leaves. *bioRxiv*; p.
1370 2022.02.04.479097 2022.

1371 102 Felber C, Azouma YO, and Reppich M. Evaluation of analytical methods for the determination of
1372 the physicochemical properties of fermented, granulated, and roasted cassava pulp - gari. *Food*
1373 *Science & Nutrition* **5**:46–53 (2017).

1374 103 Rosario MM. Industrialization, production, and processing of cassava (*Manihot Esculenta* Crantz)
1375 for the extraction of flour and potential commercial usage. *ijaeb* **8**:1–9 (2023).

1376 104 Fronza P, Costa ALR, Franca AS, and Oliveira LS de. Extraction and Characterization of Starch
1377 from Cassava Peels. *Starch - Stärke* **75**:2100245 (2023).

1378 105 Aghogho CI, Eleblu SJY, Bakare MA, Kayondo SI, Asante I, Parkes EY, Kulakow PA, Offei S, and
1379 Rabbi IY. Genetic variability and genotype by environment interaction of two major cassava
1380 processed products in multi-environments. *Frontiers in Plant Science* *Frontiers Media*; **13**:1–17
1381 (2022).

1382 106 Alamu EO, Sangodoyin OM, Diallo TA, Kolawole PO, Olajide JO, Jekayinfa SO, Abass A, Tran T,
1383 Awoyale W, Parkes E, and Maziya-Dixon B. Development of an Improved Steamer for Optimum
1384 Retention of Carotenoids in Attiéké Produced from Biofortified Cassava (*Manihot esculenta*
1385 *Crantz*) Roots. *Journal of Food Processing and Preservation* **2023**:1–13 (2023).

1386 107 Yao K, Koffi R, and Aboua F. Composition of dehydrated attiéké powder. *Tropical Science*
1387 **46**:224–226 (2006).

1388 108 Ngoualem Kégah F and Ndjouenkeu R. Gari, a Cassava (*Manihot esculenta* Crantz) Derived
1389 Product: Review on Its Quality and Their Determinants. *Journal of Food Quality* **2023**:7238309
1390 (2023).

- 1391 109 Kawalawu WKC, Asiedu R, Maziya-Dixon B, Abass A, Edet M, and Awoyale W. Evaluation of the
 1392 chemical composition and functional properties of *gari* from Liberia. *Croat j food sci technol*
 1393 (Online) **11**:157–167 (2019). View Article Online
DOI: 10.1039/D5FB00326A
- 1394 110 Olaoye O, Lawrence IG, Cornelius G, and Ihenetu ME. Evaluation of quality attributes of cassava
 1395 product (*gari*) produced at varying length of fermentation. 2015.
- 1396 111 Dahdouh L, Escobar A, Rondet E, Ricci J, Flidel G, Adinsi L, Dufour D, Cuq B, and Delalonde M.
 1397 Role of dewatering and roasting parameters in the quality of handmade *gari*. *International*
 1398 *Journal of Food Science & Technology* **56**:1298–1310 (2021).
- 1399 112 Sobowale SS, Adebisi JA, and Adebo OA. Design, construction, and performance evaluation of a
 1400 *gari* roaster. *Journal of Food Process Engineering* **40**:e12493 (2017).
- 1401 113 Awoyale W, Alamu EO, Chijioke U, Tran T, Takam Tchunte HN, Ndjouenkeu R, Kegah N, and
 1402 Maziya-Dixon B. A review of cassava semolina (*gari* and *eba*) end-user preferences and
 1403 implications for varietal trait evaluation. *Int J Food Sci Technol* **56**:1206–1222 (2021).
- 1404 114 Adesokan M, Alamu EO, Fawole S, and Maziya-Dixon B. Prediction of functional characteristics
 1405 of *gari* (cassava flakes) using near-infrared reflectance spectrometry. *Front Chem* **11**:1156718
 1406 (2023).
- 1407 115 Adinsi L, Akissoé N, Escobar A, Prin L, Kougblenou N, Dufour D, Hounhouigan DJ, and Flidel G.
 1408 Sensory and physicochemical profiling of traditional and enriched *gari* in Benin. *Food Science &*
 1409 *Nutrition* **7**:3338–3348 (2019).
- 1410 116 Ezeocha CV, Ihesie LC, and Kanu AN. Comparative evaluation of toasting variables and the
 1411 quality of *gari* produced by different women in Ikwuano LGA, Abia State, Nigeria. *Journal of Food*
 1412 *Processing and Preservation* **43**:e14060 (2019).
- 1413 117 Ndjouenkeu R, Ngoualem Kegah F, Teeken B, Okoye B, Madu T, Olaosebikan OD, Chijioke U,
 1414 Bello A, Oluwaseun Osunbade A, Owoade D, Takam-Tchunte NH, Biaton Njeufa E, Nguidem
 1415 Chomdom IL, Forsythe L, Maziya-Dixon B, and Flidel G. From cassava to *gari*: mapping of quality
 1416 characteristics and end-user preferences in Cameroon and Nigeria. *International Journal of Food*
 1417 *Science & Technology* **56**:1223–1238 (2021).
- 1418 118 Sobowale SS, Awonorin SO, Shittu T, Ajisegiri E, Adebo O, and Olatidoye O. Modeling of the
 1419 Garification Process of Fermented Cassava Mash. *J Bioprocess Biotech* **07** (2017).
- 1420 119 Eyinla TE, Maziya-Dixon B, Alamu OE, and Sanusi RA. Retention of Pro-Vitamin A Content in
 1421 Products from New Biofortified Cassava Varieties. *Foods* **8**:177 (2019).
- 1422 120 Abass AB, Awoyale W, Ogundapo A, Oluwasoga O, Nwaoliwe G, Oyelekan J, and Olarinde LO.
 1423 Adoption of improved cassava varieties by processors is linked to processing characteristics and
 1424 products biophysical attributes. *Journal of Food Processing and Preservation* **46**:e16350 (2022).
- 1425 121 Bechoff A, Tomlins KI, Chijioke U, Ilona P, Westby A, and Boy E. Physical losses could partially
 1426 explain modest carotenoid retention in dried food products from biofortified cassava. *PLOS ONE*
 1427 Public Library of Science; **13**:e0194402 (2018).
- 1428 122 Saediman H, Amini A, Basiru R, and Nafiu LO. Profitability and Value Addition in Cassava
 1429 Processing in Buton District of Southeast Sulawesi Province, Indonesia. *Journal of Sustainable*
 1430 *Development* **8**:p226 (2015).





123 Adegbite SA, Asiru WB, Salami MO, Nwaeche CF, Ebun KK, and Ogunbiyi AA. Design and Development of Power Driven Gari Fryer. *Journal of Engineering Research and Reports* :1–15 (2019).

124 Taiwo KA. Utilization Potentials of Cassava in Nigeria: The Domestic and Industrial Products. *Food Reviews International* Taylor & Francis; **22**:29–42 (2006).

125 Edward V, Egounlety, Huch, zyl P van, Singh S, Nesengani, Haakuria V, and Franz C. Isolation and screening of microorganisms from a gari fermentation process for starter culture development. *African Journal of Biotechnology* **11**:12865–12877 (2012).

126 Doing Holdings - Henan Jinrui Food Engineering Co L. What is the gari production process? Cassava processing machine. 2019. Available: [//www.cassavaprocessingplant.com/faq/gari_production_process_997.html](http://www.cassavaprocessingplant.com/faq/gari_production_process_997.html) [5 March 2025].

127 Godwin Tejiri U, Julius I, Abraham Omo E, Momoh, Shemahu M-, Bernard Ojeizuanbi A, and Odion Paul O. Improving the Dietary Fiber Content of Tapioca Using Corn Fiber as a Fiber Supplement. *International Journal of Innovative Science and Research Technology (IJISRT)* :850–857 (2024).

128 Abimbola .A N and Chioma .V A. Nutritional Composition and Sensory Evaluation of Tapioca Fortified with Soy-Coconut Flour. *Food Science and Quality Management* **92**:36 (2019).

129 Moorthy S, SANKARAKUTTY S, Ambrose K, and Anish R. Tropical tuber starches: structural and functional characteristics. *Starch in Food*; 2021.

130 Adebowale AR, Sanni L, Awonorin S, Daniel I, and Kuye A. Effect of cassava varieties on the sorption isotherm of tapioca grits. *International Journal of Food Science & Technology* **42**:448–452 (2007).

131 Bulathgama U, Gunasekara D, Wickramasinghe I, and Somendrika D. Development of Commercial Tapioca Pearls used in Bubble Tea by Microwave Heat–Moisture Treatment in Cassava Starch Modification. *European Journal of Engineering and Technology Research* **5**:103–106 (2020).

132 Lin C-L, Lin J-H, Lin J-J, and Chang Y-H. Properties of High-Swelling Native Starch Treated by Heat–Moisture Treatment with Different Holding Times and Iterations. *Molecules* Multidisciplinary Digital Publishing Institute; **25**:5528 (2020).

133 Herceg Z, Batur V, Režek Jambrak A, Vukušić T, Gmajnički I, and Špoljarić I. The Effect of Tribomechanical Micronization and Activation on Rheological, Thermophysical, and Some Physical Properties of Tapioca Starch. *International Journal of Carbohydrate Chemistry* **2013**:657951 (2013).

134 Martin Torrejon V, Song H, Wu B, Luo G, and Song J. Effect of Starch Type and Pre-Treatment on the Properties of Gelatin–Starch Foams Produced by Mechanical Foaming. *Polymers* Multidisciplinary Digital Publishing Institute; **15**:1775 (2023).

135 Apeji YE, Olayemi OJ, Anyebe SN, Oparaech C, Orugun OA, Olowosulu AK, and Oyi AR. Impact of binder as a formulation variable on the material and tableting properties of developed co-processed excipients. *SN Appl Sci* **1**:561 (2019).

- 1470 136 Fu Y-C, Dai L, and Yang BB. Microwave finish drying of (tapioca) starch pearls. *International* View Article Online
1471 *Journal of Food Science & Technology* **40**:119–132 (2005). DOI: 10.1039/D5FB00326A
- 1472 137 Krishnakumar T, SANKARAKUTTY S, Saravanan R, Giri N, Chintha P, and Bansode V. Studies on
1473 the development of cassava based reconstituted dry starch sago with modified starch as binder
1474 and characterization of its physico-functional properties. *Journal of Environmental Biology*
1475 **41**:29–34 (2020).
- 1476 138 Chauynaron N, Bhuiyan M, Kanto U, and Iji P. Variation in Nutrient Composition of Cassava Pulp
1477 and its Effects on in vitro Digestibility. *Asian Journal of Poultry Science* **9**:203–212 (2015).
- 1478 139 Samuel F, Otegbayo B, and Alalade T. Nutrient and Anti-Nutrient Content of Soy-Enriched
1479 Tapioca. *Food and Nutrition Sciences* **336105**:784–789 (2012).
- 1480 140 Adebawale O and Ajibode OO. Fortification of cassava starch with coconut residue: effects on
1481 flours' functional properties and products' (Tapioca meals) nutritional and sensory qualities.
1482 *Natural Resources for Human Health* **2**:200–207 (2022).
- 1483 141 Ollé Resa CP, Jagus RJ, and Gerschenson LN. Do fillers improve the physicochemical properties of
1484 antimicrobial tapioca starch edible films? *Journal of Food Safety* **41**:e12880 (2021).
- 1485 142 Manchun S, Nunthanid J, Limmatvapirat S, and Sriamornsak P. Effect of Ultrasonic Treatment on
1486 Physical Properties of Tapioca Starch. *Advanced Materials Research* Trans Tech Publications Ltd;
1487 **506**:294–297 (2012).
- 1488 143 Cayot N. Preliminary tests on a flavoured model system: elaboration process and rheological
1489 characterization of a custard dessert. *Flavour and Fragrance Journal* **21**:25–29 (2006).
- 1490 144 Adeboye A, Adefunke B, Adebo O, Okafor D, and Tajudeen B. Physicochemical, functional and
1491 sensory properties of tapioca with almond seed (*Terminalia catappa*) flour blends. *African*
1492 *Journal of Food Science* **13**:182–190 (2019).
- 1493 145 Edith K-KN, Benjamin YN, Julien CK, and Ibrahim K. Microbiological Quality of Attiéké (Steamed
1494 Cassava Semolina) Sold in Côte d'Ivoire. *Journal of Applied Biotechnology* **8**:14 (2020).
- 1495 146 Kouamé A k., Djéni T n., N'Guessan F k., and Dje M k. Postprocessing microflora of commercial
1496 attiéke (a fermented cassava product) produced in the south of Côte d'Ivoire. *Letters in Applied*
1497 *Microbiology* **56**:44–50 (2013).
- 1498 147 Brennan CS. Regenerative Food Innovation: The Role of Agro-Food Chain By-Products and Plant
1499 Origin Food to Obtain High-Value-Added Foods. *Foods Multidisciplinary Digital Publishing*
1500 *Institute*; **13**:427 (2024).
- 1501 148 Gnagne G-BAA, Koffi EK, Assanvo JB, and Soro S. Influences de la congélation et du séchage de
1502 l'attiéké sur ses caractéristiques physico-chimiques et organoleptiques. *International Journal of*
1503 *Biological and Chemical Sciences* **10**:808–819 (2016).
- 1504 149 Adeyi A, Adeyi O, Oke O, Ogunsola A, Ajayi O, Otolorin J, Areghan S, Owoloja A, Isola B, and
1505 Akinyemi O. Convective drying of unripe plantain: a comparative response surface methodology
1506 and genetic algorithm optimization study, certainty and sensitivity analysis. *JGhIE* **23**:1–9 (2023).





1507 150 Melini V, Melini F, Luziatelli F, and Ruzzi M. Functional Ingredients from Agri-Food Waste: Effect
1508 of Inclusion Thereof on Phenolic Compound Content and Bioaccessibility in Bakery Products.
1509 *Antioxidants* Multidisciplinary Digital Publishing Institute; **9**:1216 (2020). View Article Online
DOI: 10.1039/D5FB00326A

1510 151 Donner M, Verniquet A, Broeze J, Kayser K, and De Vries H. Critical success and risk factors for
1511 circular business models valorising agricultural waste and by-products. *Resources, Conservation*
1512 *and Recycling* **165**:105236 (2021).

1513 152 Kover A, Kraljić D, Marinaro R, and Rene ER. Processes for the valorization of food and
1514 agricultural wastes to value-added products: recent practices and perspectives. *Syst Microbiol*
1515 *and Biomanuf* **2**:50–66 (2022).

1516 153 Verdini F, Calcio Gaudino E, Grillo G, Tabasso S, and Cravotto G. Cellulose Recovery from Agri-
1517 Food Residues by Effective Cavitation Treatments. *Applied Sciences* Multidisciplinary Digital
1518 Publishing Institute; **11**:4693 (2021).

1519 154 Dhiman S, Kaur ,Pardeep, Narang ,Jasjeet, Mukherjee ,Gunjan, Thakur ,Babita, Kaur
1520 ,Sukhminderjit, and Tripathi M and. Fungal bioprocessing for circular bioeconomy: Exploring
1521 lignocellulosic waste valorization. *Mycology* Taylor & Francis; **15**:538–563 (2024).

1522 155 Rusu AV, Schwarze A-K, and Alvarez Penedo B. FUNGUSCHAIN EU Project: Extracting Value from
1523 the Agricultural Offcuts of Commercial Mushroom Farming. *Bulletin of University of Agricultural*
1524 *Sciences and Veterinary Medicine Cluj-Napoca Food Science and Technology* **77**:104–106 (2020).

1525 156 Mbang MO. Commerce et Autonomisation des Femmes en Zone CEMAC. *European Scientific*
1526 *Journal, ESJ* **19**:232–232 (2023).

1527 157 Hina M, Chauhan C, Sharma R, and Dhir A. Circular economy business models as pillars of
1528 sustainability: Where are we now, and where are we heading? *Business Strategy and the*
1529 *Environment* **32**:6182–6209 (2023).

1530 158 Agbe YFT and Atake E-H. Les déterminants de la productivité agricole dans les pays de l’Afrique
1531 subsaharienne. *La Revue Internationale des Économistes de Langue Française* **8**:84–106 (2023).

1532 159 Totouom A. Les obstacles au développement industriel de l’Afrique. *ae HEC Montréal*; **94**:363–
1533 387 (2018).

1534 160 Forsythe L, Tufan H, Bouniol A, Kleih U, and Fliedel G. An interdisciplinary and participatory
1535 methodology to improve user acceptability of root, tuber and banana varieties. *International*
1536 *Journal of Food Science and Technology* **56**:1115–1123 (2021).

1537 161 Jideani AIO, Onipe OO, and Ramashia SE. Classification of African Native Plant Foods Based on
1538 Their Processing Levels. *Front Nutr* Frontiers; **9** (2022).

1539 162 Temple L and Bon H de. L’agriculture biologique : controverses et enjeux globaux de
1540 développement en Afrique. *Cah Agric* EDP Sciences; **29**:3 (2020).

1541 163 Kouakou ARS, Kouassi KN, Kouadio YH, and Kouame AF. Identification of the processes and the
1542 uses of Tanga: Dried ripe plantain (*Musa paradisiaca*) pulp produced in Côte d’Ivoire. *World*
1543 *Journal of Advanced Research and Reviews* World Journal of Advanced Research and Reviews;
1544 **21**:399–409 (2024).

- 1545 164 Adeyeye SAO. The role of food processing and appropriate storage technologies in ensuring food security and food availability in Africa. *Nutrition & Food Science* Emerald Publishing Limited; **47**:122–139 (2017). [View Article Online](#)
DOI: 10.1039/D5FB00326A
- 1548 165 Konte MA, Cisse M, Birba BJ-J, and Behanzin TDT. Les déterminants du développement des
1549 archés boursiers en Afrique Subsaharienne. *La Revue Internationale des Économistes de Langue*
1550 *Française* **2**:7–24 (2017).
- 1551 166 Royen KV, Lachat C, Holdsworth M, Smit K, Kinabo J, Roberfroid D, Nago E, Orach CG, and
1552 Kolsteren P. How Can the Operating Environment for Nutrition Research Be Improved in Sub-
1553 Saharan Africa? The Views of African Researchers. *PLOS ONE* Public Library of Science; **8**:e66355
1554 (2013).
- 1555 167 Okoye B, Ofoeze M, Ejechi M, Onwuka S, Nwafor S, Onyemauwa N, Ukeje B, Eluagu C,
1556 Obidiegwu J, Olaosebikan O, and Madu T. Prioritizing preferred traits in the yam value chain in
1557 Nigeria: a gender situation analysis. *Front Sociol* Frontiers; **8** (2023).
- 1558 168 Dandonougbo Y and Agbodji EA. Variabilité climatique en Afrique de l'Ouest : incidence de l'aide
1559 publique au développement sur la sécurité alimentaire. *La Revue Internationale des Économistes*
1560 *de Langue Française* **5**:220–234 (2020).
- 1561
- 1562



Data availability

[View Article Online](#)
DOI: 10.1039/D5FB00326A

The data that support the findings of this study are available from the authors upon reasonable request.

