

# Sustainable Food Technology

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: Z. Wang, Y. Zhou, L. Chen, T. Chen, Y. Wang, S. Ding, Y. Du, J. Hu and D. Wu, *Sustainable Food Technol.*, 2025, DOI: 10.1039/D5FB00446B.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

## Sustainability Spotlight Statement

1. First identification of *Ulva lactuca* lectin allergenicity *via* cellular/murine models.
2. UL lectin triggers IgE (45 ng/mL) & IgG1 (2888 ng/mL) surge and splenomegaly in mice.
3. UL lectin was regarded as hidden algal allergen requiring food safety assessment.



## Molecular Properties and Allergenicity of Lectin from *Ulva lactuca*

Zhicong Wang<sup>a, b</sup>, Yingying Zhou<sup>a, b</sup>, Lihang Chen<sup>a, b</sup>, Tao Chen<sup>a, b</sup>, Yuying Wang<sup>a, b</sup>,  
Shanshan Ding<sup>a, b</sup>, Yinan Du<sup>a, b</sup>, Jiangning Hu<sup>a, b</sup>\*, Di Wu<sup>a, b</sup>\*

<sup>a</sup> SKL of Marine Food Processing & Safety Control, National Engineering Research Center of Seafood, Collaborative Innovation Center of Seafood Deep Processing, Dalian Polytechnic University, Dalian, Liaoning, 116034, China

<sup>b</sup> School of Food Science and Technology, Dalian Polytechnic University, Dalian, 116034, China

Corresponding author: Prof. Jiangning Hu, Dr. Di Wu, SKL of Marine Food Processing & Safety Control, National Engineering Research Center of Seafood, Collaborative Innovation Center of Seafood Deep Processing, School of Food Science and Technology, Dalian Polytechnic University, Dalian 116034, China, Tel: +86-411-86332275, Fax: +86-411-86323262, E-mail: hujiangning2005@hotmail.com, m13039998695@163.com





1     **ABSTRACT**

View Article Online  
DOI: 10.1039/D5FB00446B

2           *Ulva lactuca*, the type species of the genus algae, holds ecological significance in  
3 coastal ecosystems and has attracted considerable interest in food and medical  
4 applications due to its bioactive constituents. Proteins or peptides from algae that  
5 exhibit high abundance, stability, or significant homology to specific allergens were  
6 identified as antigens by immune cells. In response, the immune system mounts a  
7 variable reaction to each protein, ultimately leading to cellular degranulation and the  
8 manifestation of a range of diseases. Lectin, derived and purified from *Ulva lactuca* (*U.*  
9 *lactuca*), displaying distinct bands at molecular weights of 107.55, 75, 67.6, 35, 32.02,  
10 28.75, 23.51, 18.7, and 10.7 kDa. Subsequent bioinformatic alignment against the PDB  
11 database revealed significant homology between *U. lactuca* lectin and documented  
12 allergenic lectins. *In vitro* evaluation using rat basophilic leukemia mast cells (RBL-  
13 2H3) demonstrated that *U. lactuca* lectin significantly promoted mast cell degranulation,  
14 whereas histamine (His) release rate was reached at  $45.28 \pm 2.40$  ng mL<sup>-1</sup>. *In vivo*  
15 experiments revealed that *U. lactuca* lectin induced splenomegaly, and promoted  
16 substantial elevation of His and mast cell proteases in mice. Moreover, *U. lactuca* lectin  
17 samples exhibited significantly upregulated levels of allergen-specific Immunoglobulin  
18 E (IgE), Immunoglobulin G1 (IgG1) inducing a Th2 immune response. These findings  
19 provide foundational evidence for the allergenic potential of *U. lactuca* lectin and  
20 contribute to safety evaluations of marine-derived proteins.

21     **Keywords:** *U. lactuca* lectin, allergenicity, mouse model

22

## 1. Introduction

Green algae have been recognized as one of the most significant allergenic sources in aquatic products. *U. lactuca*, a macroalga widely distributed in intertidal zones from temperate to tropical regions globally, was extensively studied due to its rapid growth, high biomass, and rich content of proteins, polysaccharides, minerals, and bioactive compounds. In recent decades, it has been regarded as a promising functional food ingredient and sustainable bioresource.<sup>1</sup> In Asian countries, *U. lactuca* was traditionally consumed as food for centuries. Its applications were also explored in bioenergy production, pharmaceuticals, and cosmetics.<sup>1</sup> The presence of umami amino acids in *U. lactuca* was identified, which contributed to its distinct marine flavor. This characteristic enabled its utilization in developing unique seafood-flavored products.<sup>2</sup> Furthermore, polyphenolic compounds extracted from *U. lactuca* were demonstrated to exhibit antioxidant and antimicrobial properties. These bioactive substances were proposed as natural food additives for meat preservation.<sup>2</sup> However, with the increasing industrial-scale application of *U. lactuca* and its derivatives, their potential allergenic risks have become a critical concern for both academia and the food industry.

The lectin from *U. lactuca* was characterized as carbohydrate-binding proteins with biological functions in cell signaling and pathogen defense through specific recognition of glycoconjugates.<sup>3</sup> Lectins were first isolated from castor beans (*Ricinus communis*) in 1888, designated as Ricin.<sup>4</sup> The development of affinity chromatography in the 1970s significantly advanced lectin research. However, Concanavalin A, crystallized from jack beans (*Canavalia ensiformis*) in 1919, was recognized as the first lectin to be systematically studied by Sumner and Howell.<sup>4,5</sup> Immunomodulatory, antiviral, and antitumor activities of lectins were documented, though their interaction



mechanisms with mammalian immune systems remained incompletely elucidated. Notably, specific lectins (e.g., peanut agglutinin, soybean lectin) were confirmed to trigger IgE-mediated allergies by binding to intestinal epithelial receptors, potentially inducing anaphylaxis.<sup>7</sup> Compared to known allergens in other algae, allergic reactions related to lectins were on the rise. In China, lectins accounted for 36.59% of plant food allergy cases in 2017, attributed to their high abundance (2.4-5% of total proteins) in edible plants.<sup>8,9</sup> Gastrointestinal symptoms including nausea, vomiting, and abdominal distension were observed in both children and adults. Current research predominantly focused on legume lectins, while systematic risk assessment of *U. lactuca* lectin allergenic potential remained unexplored. Thus, comprehensive evaluation of *U. lactuca* lectin allergenicity was urgently required to ensure the safety of algal-based food products.

Food allergy has been acknowledged as a worldwide food safety issue, and its prevalence has significantly increased over the past two decades. IgE-mediated type I hypersensitivity was established as the primary mechanism. Following gastrointestinal processing of allergens, epitopes were recognized by antigen-presenting cells.<sup>10</sup> Th2-type immune responses were subsequently activated. This cascade induced the production of allergen-specific IgE antibodies by B cells.<sup>11</sup> These antibodies were bound to mast cells and basophils via FcεRI receptors. Re-exposure to homologous allergens triggered cellular degranulation. Inflammatory mediators including His were released. Multisystem symptoms ranging from mild pruritus to anaphylactic shock were observed.<sup>12</sup> Lectin proteins were recognized as potential hidden allergens due to their gastrointestinal stability and capacity to enhance antigen presentation. The allergenic potential of *U. lactuca* lectin was therefore considered critical for ensuring the safety of Ulva-derived food products.



Internationally recognized frameworks for assessing protein allergenicity had been established by regulatory bodies including the European Food Safety Authority, Codex Alimentarius Commission, and World Health Organization. These guidelines encompassed animal models, cellular immunological assays, serological analyses, and bioinformatics approaches.<sup>13, 14</sup> Previous research efforts were predominantly directed toward developing cellular and animal experimental systems.<sup>15</sup> In this study, lectin was isolated from *U. lactuca* using ammonium sulfate precipitation. RBL-2H3 cell model was established to evaluate His release following *U. lactuca* lectin stimulation. Subsequently, BALB/c mice were sensitized with *U. lactuca* lectin. Clinical allergy symptoms, serum-specific antibody levels, and cellular degranulation responses were systematically measured. The allergenic potential of *U. lactuca* lectin was comprehensively investigated through this multidimensional experimental strategy. These findings were intended to support the safe utilization of algal resources and contribute theoretical insights for identifying marine-derived allergens.

## 2. Materials and Methods

### 2.1. Materials and Chemicals

*Ulva lactuca* were purchased from Daqin Island, Changdao, Yantai, Shandong Province. BCA Protein Concentration Detection Kit, Sodium dodecyl sulfate, Tris, Toluidine blue stain solution and DEAE-Sepharose Fast Flow were procured from Solarbio Science & Technology Co., Ltd. (Beijing, China). Coomassie Brilliant Blue R-250, glycerol, glycine, acrylamide, N, N'-methylene diacrylamide, TEMED, Ammonium persulfate, bromophenol blue,  $\beta$ -Mercaptoethanol, MTT, cholera toxin



Open Access Article. Published on 23 September 2025. Downloaded on 10/4/2025 7:03:43 AM.  
This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.



(CT) and albumin(OVA) were purchased from Sigma-Aldrich (St. Louis, MO, USA). H<sub>2</sub>SO<sub>4</sub>, NaCl, CH<sub>3</sub>COOH and NaOH were procured from Tianjin Damao Chemical Reagent Co., Ltd. (Tianjin, China). (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, HCl were bought from Tianjin Kemiou Chemical Reagent Co., Ltd. (Tianjin, China). Advanced DMEM and Trypsin-EDTA were supplied by Gibco (Waltham, MA, USA). FBS was purchased from Procell Co., Ltd. (Wuhan, China). Penicillin-Streptomycin Solution was procured from Neyotime Co., Ltd. (Shanghai, China). RBL-2H3 cells were purchased from the Cell Resource Center (Shanghai, China). ELISA kits of His , IgE, IgG1 and Mouse mast cell protease 1 (mMCP-1) were bought from Jianglai Co., Ltd. (Shanghai, China).

**2.2. Extraction and Purification of *U. lactuca* Lectin**

A mixer grinder equipment (Dade, DFY-1000C, China) was applied to produce *U. lactuca* powder. *U. lactuca* powder (1:15, w/v) was dissolved in 10 mM PBS with 0.15 mol L<sup>-1</sup> NaCl (pH 7.0). The mixture was centrifuged at 9000 rpm for 20 min at 4 °C by hybrid refrigerated centrifuge (Techcomp, CAX-571, China), after extracted for 16 h at 4 °C. (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> with 20% saturation were added into the supernatant at 4 °C, the mixture was stirred at magnetic stirrer (YUHUA, 98-2, China) overnight (4 °C). After centrifugation at 11000 rpm for 20 min at 4 °C, the precipitate was discarded and the supernatant was brought to 60% saturation with (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> and stored for 12 h. The precipitation was dissolved in distilled water, and was dialyzed for 36 h against the same buffer to eliminate residual (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>. Then, the solution was lyophilized and stored in a dryer, which was named as lectin (L). The target proteins were purified using ion exchange chromatography (GE, AKTA avant 25, China) in a DEAE-Sepharose Fast Flow, with a 39 mL column (Henghuibio, HC-0220-02A, China) with the eluting buffer of NaCl (0-1 mol L<sup>-1</sup>) in the Tris-HCl (50 mM, pH 8.0).<sup>16</sup> Then, the penetrating peak



(A) and elution peak (B) of L were obtained. All purification procedures were conducted at 4 °C, with subsequent lectin aliquots stored at -20 °C pending downstream assays.

View Article Online  
DOI: 10.1039/D5FB00446B

### 2.3. SDS and Native Polyacrylamide Gel Electrophoresis

Native-PAGE and SDS-PAGE analyses were performed using the following methods with some modifications.<sup>17</sup> In Native-PAGE, samples were prepared in a buffer containing 20% glycerol and 25 mM Tris-HCl (pH 8.0). And then, 10 µL of each sample was loaded onto Native-PAGE gels. In SDS-PAGE, samples were dissolved to 4 mg mL<sup>-1</sup> with the sample buffer which contains 25 mM Tris-HCl (pH 8.0), SDS, glycerinum and β-mercaptoethanol. After boiling for 10 min, the samples (10 µL) were centrifuged and loaded onto a stacking gel (3%) and separating gel (12.5%). The electrophoresis was performed under constant flow conditions of 10 mA. After electrophoresis, the gel was fixed with a solution of 100 mM ammonium acetate dissolved in methyl alcohol: acetic acid (5:1, v/v) for 2 h. After fixing, gels were stained with Coomassie Brilliant Blue R-250 and destained with 10% acetic acid solution for 12 h, and washed by water. Imaging was performed using a GS-800 densitometer (DNR BioImaging Systems Ltd., Israel).<sup>18</sup>

### 2.4. RBL-2H3 Cell Model Experiment

#### 2.4.1. RBL-2H3 Cell Viability Assay

The cell viability test was measured in accordance to MTT cytotoxicity assay utilizing RBL-2H3 cells (Shanghai cell bank of Chinese Academy of Sciences). Medium for cells culture was Minimum Essential Medium (Hyclone, Logan, UT, USA)



with addition of 1% penicillin-streptomycin (Hyclone, Logan, UT, USA) and 15% fetal bovine serum (PAN-Biotech, UK). Typically, RBL-2H3 cells (200  $\mu$ L) were suspended in cell culture medium and seeded into 96-well plates at a density of  $5 \times 10^4$  cells per mL, being incubated for 24 h at 37  $^{\circ}$ C in 5% CO<sub>2</sub> to obtain a monolayer of cells.<sup>19</sup> After 24 h of incubation, 200  $\mu$ L of L, A and B in different concentrations (0, 50, 100, 200, 300, 400, 500, 700 and 1000  $\mu$ g/mL) were added for 24 h of stimulation. The sample solution was removed after incubation and the cells were further cultured with of MTT (5 mg mL<sup>-1</sup>, 15  $\mu$ L) solution for 4 h. Finally, the culture medium was substituted with DMSO (150  $\mu$ L) to dissolve the formazan and the absorbance of the DMSO solution at 490 nm was tested by a micro plate reader (Infinite<sup>TM</sup> M200, TECANE, Switzerland).

**2.4.2. Degranulation and Morphological Observation of RBL-2H3 Cells**

RBL-2H3 cells (1 mL) were suspended in cell culture medium and seeded into 12-well plates at a density of  $5 \times 10^4$  cells mL<sup>-1</sup>, being incubated for 24 h at 37  $^{\circ}$ C in 5% CO<sub>2</sub> to obtain a monolayer of cells.<sup>19</sup> After 24 h of incubation, 200  $\mu$ L of A, B and lectin (300  $\mu$ g mL<sup>-1</sup>) were added for 24 h of stimulation. At the end of excitation, the supernatants were collected and centrifuged (1000 rpm, 20min). Then the supernatant was collected, and His content were determined using ELISA kits,<sup>20</sup> which were purchased from Jianglai Co., Ltd. (Shanghai, China). At the end of the excitation, the cells were observed under a microscope and photographed after staining with toluidine blue stain solution.



## 2.5. Allergenicity Assessment

### 2.5.1. Mice Allergy Model

Female BALB/c mice (three to four weeks) were purchased from Changsheng Biotechnology Co., Ltd (Benxi, Liaoning, China). All mice were housed in an air-conditioned room ( $22 \pm 2$  °C, with a 12 h/12 h dark/light cycle,  $50 \pm 15\%$  relative humidity) with free access to water and sample-free chow. All animal experimentation was approved by the Ethics Committee of Dalian Polytechnic University.

### 2.5.2. Sensitization and Challenge

All BALB/c mice were distributed to 8 groups ( $n = 10$  for each group) as follows: saline control group (Control), mice gavaged with CT (CT), mice gavaged with CT + OVA (OVA), mice gavaged with CT + high concentration of B (BH), mice gavaged with CT + low concentration of B (BL), mice gavaged with CT + high concentration of L (LH), mice gavaged with CT + low concentration of L (LL), and mice gavaged with CT + *U. lactuca* powder (UL). CT as an adjuvant can stimulate the immune system and effectively enhance the immune response of the antigen.<sup>21</sup> To mitigate potential confounding effects of CT on immune responses, a dedicated control group receiving CT-only gavage was established in this study. For the stage of basic sensitization, mice in each group were given different doses of corresponding samples by intragastric administration. OVA, BL, and LL groups were gavaged 200  $\mu\text{L}$  sample with a concentration of 2  $\text{mg mL}^{-1}$ . BH, LH, and UL groups were gavaged 200  $\mu\text{L}$  sample with a concentration of 20  $\text{mg mL}^{-1}$ . The mice were sensitized weekly by intraperitoneal injection with samples plus CT (5  $\mu\text{g}$ ) per mice, and other two groups were gavaged with saline and CT, respectively. For the stage of challenge, OVA and sample groups



mice were given 5 times the dose of OVA and samples respectively sacrificed on Day 36.<sup>22</sup> All the experimental design of animal were shown in **Figure 3 a**. In this study, all animal operations were conducted by following the National Institutes of Guide for the Care and Use of Laboratory Animals, and approved by Dalian Polytechnic University Ethical Committee for Animal Experimentation (approval number: DLPU2024077).

### 2.5.3. Serum Collection

Blood samples were collected by intraocular canthal vein puncture at 30, 33, 35 and 36 days. Serum samples were centrifuged at 4000 rpm for 20 min at 4 °C. All serum samples were kept at -80 °C until analysis

### 2.5.4. Analysis of Spleen Index

On day 36, the target organs (spleen) were removed and weighed. The spleen index was calculated according to the formula :<sup>23</sup>

$$\text{Spleen index} = \frac{\text{Weight of spleen (g)}}{\text{Body weight (g)}}$$

### 2.5.5. Measurement of Vascular Permeability

The levels of albumin in the peritoneal fluid of BALB/c mice were determined 40 min after challenge.<sup>24</sup> Mice were intraperitoneally injected with 1 mL of PBS. After 1 min of abdominal massage, the abdominal massage-derived fluids were processed through refrigerated centrifugation (4 °C, 500×g, 6 min). The supernatant was stored at -80 °C. Albumin concentration in the supernatant was quantified by a BCA kit. The BCA working fluid and standard solution were prepared according to the requirements of the kit. The sample or standard solution was mixed with the BCA working fluid. The sample or standard solution was placed at 37 °C for 30 min and cooled to room



205 temperature. The absorbance of each sample was measured at 562 nm. The protein  
206 concentration was calculated according to the standard curve.

#### 207 **2.5.6. ELISA Assays for Specific IgE, IgG1, His and mMCP-1 Levels**

208 The serum from each mouse was collected, and IgE, IgG1, mMCP-1 and His  
209 concentrations were detected by ELISA assay using a Mouse IgE ELISA Kit, Mouse  
210 IgG<sub>1</sub> ELISA Kit, Mouse HIS ELISA Kit and mMCP-1 ELISA Kit according to the  
211 manufacturer's instructions. Absorbance was measured at 450 nm using a microplate  
212 reader, and serum concentrations were calculated based on standard curves.

#### 213 **2.5.7. Histopathology of Lung, Spleen, Jejunum and Duodenum**

214 Resected lung, spleen, jejunum and duodenum tissues were removed after  
215 euthanasia and the jejunum was washed with normal saline. Then, the lung, spleen  
216 jejunum and duodenum tissues were fixed in 4% paraformaldehyde and embedded in  
217 paraffin. Serial paraffin sections with 4-6  $\mu$ m thickness were made and prepared for  
218 staining of Hematoxylin and Eosin (H&E) to observe the histopathological.<sup>25</sup> To  
219 measure the mast cell degranulation, 5 spleen sections in each group were stained with  
220 toluidine blue (TB) and the number of intact and degranulated mast cells in each section  
221 was counted under a microscope in 5 randomly chosen fields.<sup>26</sup>

#### 222 **2.6. Statistical Analysis**

223 The mean  $\pm$  standard deviations (SD) of three sets of data from three independent  
224 experiments and technical replicates were calculated. Data were assessed using one-  
225 way ANOVA and the post hoc test HSD Tukey with the SPSS software (SPSS 16.0 Inc,  
226 Chicago, USA). The significance level was set at  $p < 0.05$ .



## 3. Results and Discussions

### 3.1 Preparation of *U. lactuca* Lectin

The *U. lactuca* L was extracted by salting out with NaCl and  $\text{NH}_4\text{SO}_4$  and purified by DEAE-Sepharose Fast Flow chromatography. The subunits of L were composed at 250.46, 180, 141.65, 67.6, 35, 28.72, and 18.7 kDa in Native-PAGE (**Figure 1 b**). During the process of ion exchange chromatography, with the ascending of the concentration of NaCl, peaks were collected. One single elution peak B and A1, A2 were observed during the elution process (**Figure 1 a**) and subjected to SDS-PAGE. From the results of the SDS-PAGE (**Figure 1 b**), the molecular weights of B were at 107.55, 75, 67.6, 35, 32.02, 28.75, 23.51 and 18.7 kDa. Lectins are carbohydrate-binding proteins that recognize and selectively bind to specific sugar structures. Developments in bioinformatics reveal that the plant's proteome includes a range of putative amino acid sequences encoding lectins with diverse carbohydrate-recognition domains.<sup>27</sup> As reported by the molecular weight of lectins 1VIW (*Phaseolus vulgaris*), 7KDU (*Ricinus communis*) and 7KBI (*Ricinus communis*) were distributed at 74.87, 74.6 and 74.98 kDa (**Figure 1 c**).<sup>3, 28, 29</sup> Lectins 2WGC (*Triticum aestivum*), 4AML (*Triticum aestivum*), 1LOC (*Lathyrus ochrus*), 6VGF (*Arachis hypogaea*) and 5AVA (*Phaseolus vulgaris*) were distributed at 35.54, 35.66, 105.01, 106.08 and 251.14 kDa.<sup>30-32</sup> Molecular structures of these lectins, which had the similar molecular weight in *U. lactuca* L were downloaded from the Protein Data Bank (<https://www.rcsb.org>). Allergenic protein might be contained in *U. lactuca* L and purified lectin (B).



### 3.2 Experimental Allergenicity Assessment of RBL-2H3 cells

To explore the allergenicity of the *U. lactuca* L at the cellular level, we evaluated the effects of L, A and B on cell degranulation with RBL-2H3 cell model.<sup>33</sup> Selection of the appropriate drug concentration for subsequent experiments was first made by MTT experiments, the concentration of 300  $\mu\text{g/mL}$  was identified as the optimal concentration for cell stimulation.<sup>34</sup> As shown in **Figure 2 e**, the His content of RBL-2H3 were determined. His release rate of RBL-2H3 cells was significantly increased after induction by B and L, reaching at  $42.40 \pm 1.47 \text{ ng mL}^{-1}$  and  $45.28 \pm 2.40 \text{ ng mL}^{-1}$ , respectively. Mast cells and basophils were activated through allergen stimulation, triggering exocytotic release of granular contents. His was identified as a key biomarker of cellular degranulation. Therefore, His levels in RBL-2H3 cell culture supernatants were quantified.<sup>35</sup> The results show that B and L were more sensitizing. This result can also be seen in the cellular morphology of the **Figure 2 e-h**. Effects of control (**Figure 2 e**), lectin (**Figure 2 f**), A (**Figure 2 g**), and B (**Figure 2 h**) on RBL-2H3 cells were observed and photographed by microscope. After being stimulated for 24 hours, compared with the control group, the cell morphology changed and became more rounded in lectin (**Figure 2 f**) and B (**Figure 2 h**), whereas there was no significant difference between control and A (**Figure 2 g**).

### 3.3 Changes in Splenic Index in BALB/c Mice

The experimental protocol for mice was shown in the **Figure 3 a**. During this study, all groups of mice were kept under the same conditions and allowed to drink and eat freely. The hair of all mice remained shiny without shedding, no obvious fighting or biting behavior was observed, and the general condition was good. Furthermore, there



271 were no deaths or significant toxicological symptoms throughout the whole experiment.  
272 Spleen is the largest and most important immune tissue in the body, and after antigenic  
273 stimulation causes an immune response in which macrophages and lymphocytes in the  
274 spleen phagocytose the exogenous material, which in turn leads to swelling of the  
275 splenic tissue.<sup>36</sup> As shown in **Figure 3 b**, mice treated with OVA, BL, LL, and LH  
276 showed a larger size of spleen than control. During allergic responses, the immune  
277 system was activated, and the spleen, as a critical immune organ, was observed to  
278 exhibit substantial proliferation of immune cells including T lymphocytes, B  
279 lymphocytes, and dendritic cells. Concurrently, a variety of inflammatory cytokines  
280 such as interleukin-4 and interleukin-12 were induced. These mediators were found to  
281 not only participate in immune regulation but also contribute to structural and functional  
282 alterations in splenic tissues. Subsequent cellular activation and hyperplasia were  
283 triggered in the spleen, ultimately resulting in a measurable increase in splenic  
284 volume.<sup>37</sup> As demonstrated in the **Figure 3 c**, the spleen indices in OVA, BL, BH, LL,  
285 LH, and UL groups were significantly higher than control and CT groups ( $p < 0.05$ ),  
286 whereas the LH group exhibiting the highest index of  $0.74 \pm 0.05\%$ . A notable increase  
287 in splenic index was also detected in the CT group relative to control group ( $p < 0.05$ ),  
288 which might be attributed to CT-induced inflammatory responses leading to structural  
289 alterations in splenic architecture.<sup>38</sup> This result suggested that lectins (L and B) in *U.*  
290 *lactuca* induce significant increase in splenic index in mice, causing swelling of  
291 immune tissues. Comparative analysis of images of the spleen revealed significant  
292 splenomegaly in experimental groups relative to the control and CT groups. This  
293 morphological deviation was pathologically associated with hypersensitivity  
294 manifestations, providing histopathological confirmation of allergic reactivity in the  
295 sample-administered groups.





### 3.4 Changes in Vascular Permeability in BALB/c Mice

When an allergic reaction occurs, mast cells and basophils release mediators such as His, which can lead to inflammatory responses such as vasodilation and increased permeability.<sup>39</sup> Increased vascular permeability is one of the hallmarks of inflammation, and measuring its changes can reflect the severity of inflammation and help determine the intensity of an allergic reaction.<sup>40</sup> Therefore, vascular permeability in mice was quantified by determining the level of peritoneal albumin in each group of mice. As shown in **Figure 3 d**, highest intra-abdominal albumin concentrations in peritoneal fluid were detected in both OVA ( $2.21 \pm 0.15 \text{ mg mL}^{-1}$ ) and UL ( $2.04 \pm 0.29 \text{ mg mL}^{-1}$ ) ( $p < 0.05$ ). There was no significant difference in abdominal albumin content between BL, BH, LH and CT. The results of the vascular permeability experiments described above indicated that there was a significant vascular leakage response in L-treated and B-treated mice, causing them to develop an allergic reaction.

### 3.5 Analysis of Specific Antibody Levels in the Serum of BALB/c Mice

Most food allergic reactions were mediated by IgE,<sup>41</sup> so an increase in allergen-specific IgE is a distinctive feature of the occurrence of allergic reactions. It is typical for serum antibody levels to be elevated in the presence of food allergy.<sup>42</sup> As shown in **Figure 4**, mice in the control group did not exhibit allergic reactions. The levels of various specific antibodies in the mouse serum were then detected. Serum levels of specific IgE antibodies in mice were shown in **Figure 4 a**, serum-specific IgE antibody level in the adjuvant (CT) group was maintained at a lower concentration of  $21.65 \pm 3.58 \text{ ng mL}^{-1}$ , indicating that CT adjuvant was specifically employed to enhance the immune response to allergens without inducing allergic responses itself. Compared



with CT groups, serum-specific IgE levels ( $p < 0.05$ ) were significantly elevated in BL (33.88  $\pm$  5.38 ng mL<sup>-1</sup>), LL (32.59  $\pm$  6.23 ng mL<sup>-1</sup>), LH (31.24  $\pm$  2.65 ng mL<sup>-1</sup>), and UL (45.02  $\pm$  7.43 ng mL<sup>-1</sup>) groups ( $p < 0.05$ ). The results showed that UL, LL and LH induced B cells to produce specific IgE antibodies. Serum-specific IgG1 antibodies can also induce an allergic reaction in the organism.<sup>43</sup> The serum-specific IgG1 antibody levels in each group of mice were presented in the **Figure 4 b**, compared with control and CT groups, LH group showed the highest concentration (2888.82  $\pm$  549.43 ng mL<sup>-1</sup>) in serum-specific IgG1 antibody levels. The serum specific IgE and IgG1 antibody levels in mice revealed that *U. lactuca* powder and lectins in *U. lactuca* triggered a Th2 type of immune response and sensitized the mice.<sup>44</sup>

### 3.6 Changes in Serum His and mMCP-1 levels in BALB/c Mice

His was a key allergic mediator released during mast cell degranulation, and changes in its levels also reflect the degree of sensitization of the organism.<sup>41</sup> mMCP-1 was stored and secreted by intestinal mucosal mast cells, which was predominantly expressed in mucosal mast cells.<sup>41</sup> Researchers studied mast cell degranulation also causes elevated levels of mMCP-1 and increased permeability of intestinal epithelial cells, leading to food allergic diseases.<sup>45</sup> As demonstrated in **Figure 4 c-d**, there were no significant differences between control and CT groups in serum His and serum mMCP-1 levels. Compared with the BALB/c mice challenged with OVA (100.18  $\pm$  13.32 ng mL<sup>-1</sup>), there were no significant differences in BH (84.72  $\pm$  12.54 ng mL<sup>-1</sup>), LH (98.55  $\pm$  11.59 ng mL<sup>-1</sup>) and UL (95.20  $\pm$  8.90 ng mL<sup>-1</sup>) groups ( $p > 0.05$ ) in His. Compared with control and CT groups, mMCP-1 concentrations were significantly elevated in the OVA, BH, LL, LH, and UL groups, with the highest level (6.77  $\pm$  2.32 ng mL<sup>-1</sup>) being recorded in the BH group ( $p < 0.05$ ). The above results indicate that



lectins in *U. lactuca* induced an allergic reaction in mice, resulting in the rupture of mast cells in the spleen and the release of large amounts of inflammatory mediators. His and mMCP-1, which were transported to various tissues and organs through the bloodstream, causing severe allergic reactions in mice. As a result, His and mMCP-1 were detected in large quantities in the serum of *U. lactuca* lectins-sensitized mice, and their levels were significantly higher than those of non-sensitized mice (control).

### 3.7 Histopathological Changes in BALB/c Mice

A systematic histopathological evaluation of multiple organ systems was conducted to comparatively assess the pathophysiological effects induced by differential dietary allergen exposure in murine models. Histopathological evaluation examined by H&E staining of jejunal sections across experimental groups (**Figure 5 a**) revealed preserved intestinal architecture in control and CT groups, characterized by intact mucosal integrity, well-organized villous alignment, and maintenance of crypt-villus axis proportions. As shown in **Figure 5 a**, distinct pathological alterations were observed in the OVA, BL, BH, LL, LH, and UL experimental cohorts, where jejunal villus architecture exhibited marked disruption, including villous fragmentation and goblet cell hyperplasia. As shown in **Figure 5 b**, the intestinal villi of duodenum sections in OVA group and sample groups were damaged and broken, and the intestinal barrier was damaged. In contrast, well-organized alignment of intestinal villi with preserved structural integrity was observed in both control and CT groups. The duodenal architecture and histological features were maintained without detectable pathological damage or necrotic changes. Consistent histopathological alterations in the gastrointestinal tract are commonly observed in conjunction with IgE-mediated food hypersensitivity reactions.<sup>46, 47</sup> This leads to the conclusion that lectins in *U. lactuca*



caused an allergic reaction in the intestine. Normal alveolar septal architecture without significant histopathological alterations was observed in pulmonary tissues of control and CT groups. In contrast, OVA-sensitized mice exhibited marked bronchial epithelial hyperplasia and alveolar septal thickening. Although less pronounced than in the OVA group, similar pathological features were detected in BL, BH, LL, LH, and UL groups (**Figure S1**). This study revealed that alveolar septal thickening might contribute to airway obstruction, potentially exacerbating allergic responses in murine models.<sup>48</sup>

### 3.8 Changes in the spleen of BALB/c mice

Splenic tissues from allergic mice demonstrated marked white pulp hyperplasia, accompanied by dilated and congested red pulp with extensive immune cell infiltration, widened marginal zones containing densely packed lymphocytes, and vascular dilation with blood stasis compared to normal controls.<sup>49</sup> As shown in **Figure 6 a**, splenic architecture was examined using H&E staining. Well-demarcated white pulp compartments with preserved periarteriolar lymphoid sheaths were observed in both control and CT groups, accompanied by loosely arranged red pulp exhibiting intact sinusoidal networks. Distinct demarcation between the marginal zone and red pulp was maintained across all specimens. Distinct white pulp hyperplasia with compromised interface integrity between white and red pulp compartments was observed in OVA-sensitized, BL, BH, LL, LH, and UL groups. The experimental findings demonstrated that dietary lectins exposure was associated with hypersensitivity reactions in murine models, which subsequently triggered immune activation characterized by augmented lymphoproliferative responses.



Mast cells, which are ubiquitously distributed throughout somatic tissues, were demonstrated to play pivotal roles in allergic pathogenesis.<sup>50,51</sup> Allergic mice exhibited a significantly increased splenic mast cell population with prominent degranulation, whereas normal controls showed sparse distribution of morphologically intact mast cells retaining cytoplasmic granules. As illustrated in **Figure 6 b**, TB staining was systematically performed on splenic tissue sections to evaluate morphological characteristics and degranulation activity of mast cells within this primary immune organ. Mast cells infiltrations were significant observed in splenic tissues of OVA and experimental groups, with evident membrane disintegration and perivascular dispersion of cytoplasmic granules, indicative of advanced degranulation processes. Elevated mast cell populations were observed in the BL, BH, and UL groups compared to the LL and LH groups, with concurrent degranulation activity detected. In contrast, minimal mast cell presence was detected in control and CT groups. In addition, the morphometric analysis of degranulated mast cells in the spleen was presented in **Figure S2**. The Control and CT groups showed a relatively low level of mast cells degranulation, with an average of approximately 6.67% and 11.11%. Compared to mice challenged with normal saline and CT, mice stimulated with OVA, B and L showed a notable increase in the percentage of degranulated mast cells ( $p < 0.05$ ), with an average exceeding 60.71%. The results showed that B and L could induce the degranulation of mast cells in the spleen of BALB/c mice. Thus, the *U. lactuca* lectin has potential high allergenic.

#### 4. Conclusions

Within the broader context of simultaneous social development and escalating allergic disease burdens, pronounced limitations and asymmetries were recognized in



Open Access Article. Published on 23 September 2025. Downloaded on 10/4/2025 7:03:43 AM.  
This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.



existing allergen recognition frameworks. Researches have long been predominantly confined to terrestrial plant-derived allergens (e.g., peanuts and soybeans), whereas insufficient attention has been paid to potential allergens present in marine flora, particularly in macroalgae species that are utilized as food or food additives. The allergenic potential of these components, which may trigger human immune responses, has not been thoroughly investigated, and related research remained in its infancy, constituting a critical knowledge gap urgently requiring further investigation. Therefore, the allergenic potential of lectin, a significant allergen from *U. lactuca* (a common edible macroalga), was investigated and was considered essential for a comprehensive understanding of seaweed allergy. Previous researches have confirmed that lectins in the diet could interfere with nutrient digestion and absorption, and high dietary levels of lectins could seriously threaten the growth and health of consuming animals. Compelling evidence was provided in this study demonstrating that the lectin purified from the green alga (*U. lactuca*) represents a novel marine-derived allergen with significant functional properties. Systematic investigation revealed that its immunomodulatory characteristics underlie the allergenic potential. These findings offered critical insights for fundamental research in allergology and the applied utilization of marine resources.

This study systematically investigated the allergenic potential of *U. lactuca* lectin, establishing it as a novel marine-derived allergen with distinct immunomodulatory properties. Experimental validation demonstrated that highly purified *U. lactuca* lectin significantly induced mast cell degranulation and His release *in vitro*, mechanistically linked to the activation of Th2-polarized immune responses. *U. lactuca* lectin elevated serum levels of allergen-specific IgE and IgG1 antibodies, His, and mMCP-1 in

437 BALB/c mice. These findings revealed that *U. lactuca* lectin induced type I hypersensitivity through Th2-biased immune activation and mast cell degranulation. 438  
439 This study emphasized the necessary to evaluate allergenic risks during the development of marine-algae resources. The study provided critical experimental 440  
441 evidence for safety assessments of *U. lactuca* in food and pharmaceutical applications, while established a method for identifying and characterizing marine-lectin allergens. 442  
443 Future research should focus on developing desensitization strategies to enable the safe utilization of *U. lactuca* lectin in food research development. 444

## 445 **Conflicts of interest**

446 There are no conflicts of interest to declare.

## 447 **ACKNOWLEDGEMENTS**

448 This study was financially supported by the National Key Research and Development Program of China (2023YFF1104001), the National Natural Science 449  
450 Foundation of China (32302004), Youth Science and Technology Star Program of Dalian (2023RQ092), and the Special Fund of Basic Scientific Research Expenses of 451  
452 Undergraduate Universities in Liaoning Province (LJ212510152042).

## 453 **Abbreviations**

454 His: histamine

455 IgE: immunoglobulin E

456 IgG1: immunoglobulin G1



457 RBL-2H3: rat basophilic leukemia mast cells

458 CT: cholera toxin

459 OVA: albumin

460 mMCP-1: mouse mast cell protease 1

461 L: lectin

462 A: target protein A, the penetrating peak of ion exchange chromatography in Figure 1a

463 B: target protein B, the elution peak of ion exchange chromatography in Figure 1a

464 BL: low concentration of B

465 BH: high concentration of B

466 LL: low concentration of L

467 LH: high concentration of L

468 UL: *Ulva lactuca* powder

469 H&E: hematoxylin and Eosin

470 TB: toluidine blue

## 471 5. REFERENCES

472 [1] Lahaye, M., Robic, A. Structure and functional properties of ulvan, a polys  
473 accharide from green seaweeds[J]. *Biomacromolecules*, 2007, 8(6), 1765-17  
474 74.

475 [2] Qi, H. M., Sun, et al. Antioxidant activity of high sulfate content derivativ  
476 e of ulvan in hyperlipidemic rats[J]. *International Journal of Biological M*  
477 *acromolecules: Structure, Function and Interactions*, 2015, 7(6), 326-329.





- 478 [3] Elumalai, P., Rubeena, A. S., Arockiaraj, J., et al. The Role of Lectins in Fish  
 479 infish: A Review[J]. *Reviews in Fisheries Science & Aquaculture*, 2019, 2  
 480 7(2), 152-169.
- 481 [4] Tsaneva M, Damme, E. J. M. V. 130 years of Plant Lectin Research[J]. *Glycoconjugate Journal*, 2020, 37, 1594074.
- 483 [5] Damme, E. V. V. Purification of GNA-Related Lectins from Natural Sources[J]. *Methods in Molecular Biology*, 2020, 2132, 413-419.
- 485 [6] Hopper, Jonathan, T. S., Stephen, et al. The Tetrameric Plant Lectin BanLe  
 486 c Neutralizes HIV through Bidentate Binding to Specific Viral Glycans[J].  
 487 2017, 5(25), 773-782.
- 488 [7] Geijtenbeek, T. B. H., Gringhuis, S. I. C-type lectin receptors in the control  
 489 of T helper cell differentiation[J]. *Nature Reviews Immunology*, 2016, 16,  
 490 433--448.
- 491 [8] Damme, E. J. M. V., Peumans, W, J., Barre, A, et al. Plant lectins: a composite  
 492 of several distinct families of structurally and evolutionary related proteins  
 493 with diverse biological roles[J]. *Critical Reviews in Plant Sciences*,  
 494 1998, 17(6), 575-692.
- 495 [9] Van, Damme, E. J. M., Lannoo, N., Peumans, W. J. Plant lectins [M]. *Advances in botanical research*. Academic Press, 2008, 48, 107-209.
- 497 [10] Sampson, H, A., O'Mahony, L., Burks, A. W., et al. Mechanisms of food  
 498 allergy[J]. *The Journal of Allergy and Clinical Immunology*, 2018, 141(1),  
 499 11-19.



- [11] Gowthaman, U., Chen, J. S., Zhang, B., et al. Identification of a T follicular helper cell subset that drives anaphylactic IgE[J]. *Science*, 2019, 365(6456), eaaw6433.
- [12] Sicherer, S. H., Sampson, H. A. Food Allergy: A review and update on epidemiology, pathogenesis, diagnosis, prevention and management[J]. *Journal of Allergy and Clinical Immunology*, 2017, 141(1), 41-58.
- [13] Herman, R. A., Ladics, G. S. Allergenic sensitization versus elicitation risk criteria for novel food proteins[J]. *Regulatory Toxicology & Pharmacology*, 2018, 94, 283-285.
- [14] Kedar, O., Golberg, A., Obolski, U., Confino-Cohen, R. Allergic to bureaucracy? Regulatory allergenicity assessments of novel food: Motivations, challenges, compromises, and possibilities[J]. *Comprehensive Reviews in Food Science and Food Safety*, 2024, 23, e13300.
- [15] Han, K., Xie, F., Phoo, M. T., et al. Inulin-gel-based oral immunotherapy remodels the small intestinal microbiome and suppresses food allergy[J]. *Nature Materials*, 2024, 23(10), 30.
- [16] Li, H., Yuan, D., Wang, C., Wang, Y., Zhang, J., Li, Z., et al. Zwitterionic peptide pendant as a homing agent to achieve lymphatic targeting antigen-specific immunotherapy of allergenic protein[J]. *Chemical Engineering Journal*, 2024, 485, 149315.
- [17] Zang, J. C., Chen, H., Zhang, X. R., Zhang, C. X., Guo, J., Du, M., Zhao G. H. Disulfide-mediated conversion of 8-mer bowl-like protein architect



- 522 ure into three different nanocages[J]. *Nature Communications*, 2019, 10(1), View Article Online  
DOI: 10.1039/D5FB00446B
- 523 778.
- 524 [18] Li, Y. J., Wang, Z. C., Liu, H. X., Chen, T., Du, Y. N., Hu, J. N., Wu,  
525 D. Controlled ionic-induced self-assembly of phycocyanin: Structure, bindi  
526 ng mechanism, and molecular dynamics simulation[J]. *Food Chemistry*, 202  
527 5, 477, 143603, 1-9.
- 528 [19] Liu, Y., Lin, S., Liu, K., Wang, S., Li, W., & Sun, N. Insights into sens  
529 itizing and eliciting capacity of gastric and gastrointestinal digestion produc  
530 ts of shrimp (*penaeus vannamei*) proteins in balb/c mice[J]. *Food Science  
531 and Human Wellness*, 2024, 13(1), 10.
- 532 [20] Wang, Y., Zhang, K., Chen, W. M., Mao, J. H., Shao, Y. H., & Tu, Z.  
533 C., et al. Allergenicity assessment of  $\beta$ -lactoglobulin ferulic acid-glucose c  
534 onjugates[J]. *Food Chemistry*, 2024, 460, 140605.
- 535 [21] Scott, C. L., Bain, C. C., Wright, P. B., Sichien, D., Kotarsky, K., Persso  
536 n, E. K., Luda, K., Guilliams, M., Lambrecht, B. N., Agace, W. W., Milli  
537 ng, S. WF., & Mowat, A. M. Ccr2(+)cd103(-) intestinal dendritic cells de  
538 velop from dc-committed precursors and induce interleukin-17 production b  
539 y t cells[J]. *Mucosal Immunology*, 2015, 8, 327-339.
- 540 [22] Liu, Y., Lin, S. Y., Liu, K. X., Wang, S., Liu, Q. Z., Sun., N., Explorat  
541 ion of digestion-resistant immunodominant epitopes in shrimp (*Penaeus van  
542 namei*) allergens[J]. *Food Chemistry*, 2024, 438, 137920, 1-14.



- [23] Xue, L., Li, Y., Li, T., Pan, H., & Wang, L. Phosphorylation and enzymatic hydrolysis with alcalase and papain effectively reduce allergic reactions to gliadins in normal mice[J]. *Journal of Agricultural and Food Chemistry*, 2019, 67(22), 6313-6323.
- [24] Jiang, T., He, F., Han, S., et al. Characterization of cAMP as an anti-allergic functional factor in Chinese jujube (*Ziziphus jujuba* Mill.)[J]. *Journal of Functional Foods*, 2019, 60, 103414.
- [25] Yu, B., Bi, D., Yao, L., Li, T., Gu, L., Xu, H., et al. The inhibitory activity of alginate against allergic reactions in an ovalbumin-induced mouse model[J]. *Food & Function*, 2020, 11, 2704-2713.
- [26] Luo, C., Chen, G., Ahmed, I., Sun, L., Li, W., & Pavase, T. R., et al. Immunostimulatory and allergenic properties of emulsified and non-emulsified digestion products of parvalbumin (*scophthalmus maximus*) in RBL-2H3 cells and balb/c mouse models[J]. *Food & Function*, 2021, 12, 5351-5360.
- [27] Naithani, S., Komath, S. S., Nonomura, A., et al. Plant lectins and their many roles: Carbohydrate-binding and beyond[J]. *Journal of Plant Physiology*, 2021, 266, 153531.
- [28] Bourne, Y., Ayoub, A., Rougé, P., Cambillau, C. Interaction of a legume lectin with two components of the bacterial cell wall. a crystallographic study[J]. *Journal of Biological Chemistry*, 1994, 269(13), 9429.



- [29] Christine, S. W. 2.2 Å resolution structure analysis of two refined N-acetylneuraminyllactose-Wheat germ agglutinin isolectin complexes[J]. *Journal of Molecular Biology*, 1990, 4(215), 635-651
- [30] Cagnoni, A. J., Primo, E. D., Klinke, S., Cano, M. E., Giordano, W., Mariño, K. V., Kovensky, J., Goldbaum, F. A., Uhrig, M. L., Otero, L. H. C. Crystal structures of peanut lectin in the presence of synthetic  $\beta$ -N- and  $\beta$ -S-galactosides disclose evidence for the recognition of different glycomimetic ligands[J]. *Acta Crystallographica Section D*, 2020, 76, 1080-1091
- [31] David, S., Caroline, M., Johannes, G. B., Sonja, S., Kay, D., Heiko, M. M., Wolfram, W., Valentin, W. Structural Basis of Multivalent Binding to Wheat Germ Agglutinin[J]. *Journal of the American Chemical*, 2010, 132, 25
- [32] Michael, J. R., Amanda, Y. P., Simona, K., Anne, G. M., Claire, R. P., Simon, A. D., Kirsten, S., David, J. V., Nicholas, J. M. Structural Analysis of Toxin-Neutralizing, Single-Domain Antibodies that Bridge Ricin's A-B Subunit Interface[J]. *Journal of Molecular Biology*, 2021, 15(433) 167086
- [33] Hung, C. F. Anti-allergic and anti-inflammatory effects of neferine on RB L-2H3 cells[J]. *International Journal of Molecular Sciences*, 2021, 22(20), 10994.
- [34] Yang, M., Xu, Z., Wu, D., Dong, Y., Wang, Z. Y., Du, M. Characterizations and the Mechanism Underlying Osteogenic Activity of Peptides from Enzymatic Hydrolysates of *Stichopus japonicus*[J]. *Journal of Agricultural and Food Chemistry*, 2021, 69(51), 15611-15623.



- [35] Sahid, M., Kiyoi, T. Mast Cell Activation Markers for in Vitro Study[J]. *Journal of Immunoassay & Immunochemistry*, 2020, 41(4): 778-816.
- [36] Morafo, V., Srivastava, K., Huang, C. K., Kleiner, G., Lee, S. Y., Sampson, H. A., et al. Genetic susceptibility to food allergy is linked to differential th2-th1 responses in c3h/hej and balb/c mice[J]. *Journal of Allergy and Clinical Immunology*, 2003, 111(5), 1122-1128.
- [37] Lewis, M. S., Williams, A., Eisenbarth, C. S. Structure and function of the immune system in the spleen[J]. *Science Immunology*, 2019, 4(33), eaau6085.
- [38] Benedetti, R., Lev, P., Massouh, E., et al. Oral administration of one dose of cholera toxin induces a systemic immune response prior to a mucosal immune response by a direct presentation in the spleen[J]. *Immunology Letters*, 1998, 60(2-3), 149-156.
- [39] Sun N, Wang J, Zhou C, et al. Cell-based immunological assay: complementary applications in evaluating the allergenicity of foods with FAO/WHO guidelines[J]. *Food Research International*, 2014, 62, 735-745.
- [40] Medzhitov, R. Origin and physiological roles of inflammation[J]. *Nature*, 2008, 454(7203), 428-435.
- [41] Bai, J., Hui, J., Lu, Q., Yang, A., Yuan, J., & Gao, J., et al. Effect of transglutaminase cross-linking on the allergenicity of tofu based on a balb/c mouse model[J]. *Food & Function*, 2020, 11, 1-10.



- [42] Liu, K., Lin, S., Liu, Y., Wang, S., Liu, Q., Sun, K., Sun, N. Mechanism of the Reduced Allergenicity of Shrimp (*Macrobrachium Nipponense*) by Combined Thermal/Pressure Processing: Insight into Variations in Protein Structure, Gastrointestinal Digestion and Immunodominant Linear Epitopes[J]. *Food Chemistry*, 2023, 405, 134829.
- [43] Zhang, T. T., Hu, Z. Y., Cheng, Y. W., Xu, H. X., Velickovic, T. C., He, K., Sun, F., He, Z. D., Liu, Z. G., Wu, X. L. Changes in allergenicity of ovalbumin in vitro and in vivo on conjugation with quercetin[J]. *Journal of Agricultural and Food Chemistry*, 2020, 13(68), 4027–4035.
- [44] Kanagaratham, C., Ansari, Y. S. E., Lewis, O. L., et al. IgE and IgG Antibodies as Regulators of Mast Cell and Basophil Functions in Food Allergy[J]. *Frontiers in Immunology*, 2020, 11, 603050.
- [45] Rupa, P., & Mine, Y., Comparison of glycated ovalbumin–monosaccharides in the attenuation of ovalbumin-induced allergic response in a balb/c mouse model[J]. *Journal of Agricultural and Food Chemistry*, 2019, 29(67), 1–11.
- [46] Roth-Walter, F., Berin M. C., et al. Pasteurization of milk proteins promotes allergic sensitization by enhancing uptake through Peyer's patches[J]. *Allergy*, 2008, 63(7), 882–890.
- [47] He, S., Zhao, J., Zhang, Y., et al. Effects of Low-pH Treatment on the Allergenicity Reduction of Black Turtle Bean (*Phaseolus vulgaris* L.) Lectin and Its Mechanism[J]. *Journal of Agricultural and Food Chemistry*, 2021, 69(4), 1379–1390.



- 630 [48] Kim, D. E., Lee, Y., Kim, M. G., et al. Bilirubin nanoparticles ameliorate  
631 allergic lung inflammation in a mouse model of asthma[J]. *Biomaterials*,  
632 2017, 140, 37-44.
- 633 [49] Rajaura, S., Babu, R., Bhardwaj, N., et al. Aflatoxin B1 administration ca  
634 uses inflammation and apoptosis in the lungs and spleen[J]. *Toxicon*, 2024,  
635 238(000), 7.
- 636 [50] Kumar, S., Sharma, A., Gupta, R., et al. Allergenicity assessment of Buch  
637 anania lanzan protein extract in BALB/c mice[J]. *International Immunophar  
638 macology*, 2018, 63, 170-182.
- 639 [51] C. H. O. Mast cells in food allergy: Inducing immediate reactions and sh  
640 aping long-term immunity[J]. *The Journal of Allergy and Clinical Immunol  
641 ogy*, 2022, 151(1), 21-25.
- 642





643 **Figure Captions**View Article Online  
DOI: 10.1039/D5FB00446B

644 **Figure 1.** Elution of DEAE-Sepharose Fast Flow chromatography of *Ulva lactuca*  
645 lectin and electrophoresis of purified lectin. (a) Elution of DEAE-Sepharose Fast Flow  
646 chromatography of *Ulva lactuca* lectin. Crude protein sample was subjected to an anion  
647 exchange column pre-equilibrated with 50 mM Tris-HCl buffer (pH 8.0). The column  
648 was eluted with a linear gradient of NaCl (0-1.0 M) in the Tris-HCl at a flow rate of 2.0  
649 mL/min. The volume of each fraction was 5 mL. Protein was monitored by measuring  
650 absorbance at 280 nm. (b) Native-PAGE and SDS-PAGE of lectin and purified lectin.  
651 (c) Overview of lectins from Protein Data Bank.

652 **Figure 2.** Proliferation and histamine content on RBL-2H3 cells treated by lectin.  
653 Lectin (a), A (b), and B (c) toxicity for 24 on RBL-2H3 cells. (d) Histamine release rate  
654 of RBL-2H3 cells induced by lectin. Effects of control (e), lectin (f), A (g), and B (h)  
655 on RBL-2H3 cells were observed and photographed by microscope.

656 **Figure 3.** Protocol of food allergic model and hypersensitive reaction. (a) Illustration  
657 of mice allergenicity scheme. Images of the spleen (b), splenic index (c) and intra-  
658 abdominal albumin concentration in different groups.

659 **Figure 4.** Anaphylaxis in the BALB/c mice sensitized and challenged in different  
660 groups. Levels of IgE (a), IgG1 (b), histamine (c), and mMCP-1 (d) in serum from mice.  
661 The concentration of IgE, IgG1, histamine, and mMCPT-1 in each BALB/c mice was  
662 determined by a commercial ELISA kit (n = 5, data presented as mean  $\pm$  SD, different



663 letters above the bars represented significant differences, with a significance level of  
664 0.05).


665 **Figure 5.** Histopathology alterations of the jejunal (a) and duodenum (b) sections of  
666 the BALB/c mice administrated with different groups. Images were taken by a  
667 microscope at 20× magnification.

668 **Figure 6.** Hematoxylin-eosin (H&E, a) and toluidine blue (TB, b) staining of splenic.  
669 Images were taken by a microscope at 20× magnification. In the H&E staining, the  
670 white pulp with a clear boundary from the red pulp were indicated by white arrows in  
671 (a). In the TB staining, the white arrows denoted intact mast cells and the red arrows  
672 denoted degranulated mast cells in (b).







**Figure 1.** Elution of DEAE-Sephacrose Fast Flow chromatography of *Ulva lactuca* lectin and electrophoresis of purified lectin.



**Figure 2.** Proliferation and histamine content on RBL-2H3 cells treated by lectin.



**Figure 3.** Protocol of food allergic model and hypersensitive reaction.




**Figure 4.** Anaphylaxis in the BALB/c mice sensitized and challenged in different groups.






(a)




(b)



**Figure 5.** Histopathology alterations of the jejunal and duodenum sections of the BALB/c mice administrated with different groups.



(b)



**Figure 6.** Hematoxylin-eosin (H&E, a) and toluidine blue (TB, b) staining of splenic.



## Data Availability Statement

The authors confirmed that the data supporting the findings of this study are available within the article and its supplementary materials.

