


Sustainable Food Technology

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: I. K. Zubairua, N. Leksawasdi, S. Wangtueai, P. Boonchuay, S. L. Htike, F. Lao, J. M. Castagnini and Y. Phimolsiripol, *Sustainable Food Technol.*, 2025, DOI: 10.1039/D5FB00879D.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

Sustainability Spotlight Statement

This study introduces ultrasound-assisted extraction (UAE) as an environmentally friendly method that enhances the value of longan processing byproducts—usually thrown away—by turning them into phenolic-rich, bioaccessible extracts with potent antioxidant properties. By greatly decreasing solvent and energy use while improving extraction efficiency, the UAE method demonstrates sustainable progress in food processing. The research promotes a circular bioeconomy through resource recovery and waste reduction, directly supporting UN SDG 12 (Responsible Consumption and Production) and SDG 9 (Industry, Innovation and Infrastructure). Additionally, by encouraging eco-efficient conversion of agricultural waste into nutraceutical ingredients, it contributes to SDG 3 (Good Health and Well-being), emphasizing the environmental and societal advantages of green technologies in the food sector.



ARTICLE

View Article Online

DOI: 10.1039/D5FB00879D

Ultrasound-enhanced extraction as a green Intensification strategy for phenolic release and *in vitro* gastrointestinal bioaccessibility from longan byproducts

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Idris Kaida Zubairu,^a Noppol Leksawasdi,^{a,b} Suttee Wangtueai,^{a,b} Pinpanit Boonchuay,^c Su Lwin Htike,^{a,b} Fei Lao,^d Juan Manuel Castagnini^e and Yuthana Phimolsiripol^{a,b*}

Longan byproducts from industry, often discarded as waste, contain bioactive compounds with potential health benefits. In this study, temperature, solvent ratio, and extraction time were optimized using Response Surface Methodology, comparing water bath and ultrasonic probe techniques. The optimized extracts were evaluated for percentage yield, total phenolic content (TPC), total flavonoid content (TFC), and antioxidant capacity. Ultrasonic-assisted extraction (UAE) achieved significantly higher yield (11.23 %), as well as higher phenolic content (219.11 mg GAE/100 g DW), and antioxidant activity (DPPH IC₅₀: 3.78) than the water bath method. The optimized extracts were subjected to simulated gastrointestinal conditions. UAE extracts showed superior intestinal retention with more than 120 mg GAE/ 100g DW of TPC compared with > 20 mg GAE/ 100g DW for the water bath extracts. Antioxidant activities were also higher, with UAE extracts retaining > 95 % activity, whereas water bath extracts showed only 55-75 %. The digested UAE extract was analyzed using high-performance liquid chromatography (HPLC) to evaluate the impact of *in vitro* digestion on polyphenols. The compounds corilagin, ellagic acid, gallic acid, epicatechin, procyanidin A2, quercetin, and pinostrobin showed significant increases in oral digestion. These findings demonstrate that UAE increased the extraction efficiency of phenolics from longan processing waste and preserved their activity during oral digestion and, to a lesser extent, in the gastric and intestinal stages, as confirmed by HPLC results. Encapsulation to improve compound stability and the evaluation of cell absorption and digestion are recommended to improve the understanding of bioaccessibility. This study supports the valorization of longan processing waste as a sustainable source of nutraceutical ingredients.

1. Introduction

Longan (*Dimocarpus longan* Lour.) is a popular fruit in temperate and sub-tropical regions and a significant economic crop in Southeast Asia, with Thailand being the second-largest producer after China. In 2023, 80-90 % of Thailand's exports to China were valued at US\$474 million. In 2025, Thailand's longan production reached 1.57 million tons, a 10.8 % increase from 2024.¹ The processing of longan yields a substantial quantity of byproducts, including pericarp and seeds, which account for approximately 30-40 % of the fresh fruit by weight.² These byproducts are highly valued in traditional Chinese medicine for their health-promoting properties, including anticancer, antiaging, and memory-enhancing effects.³ Given their bioactive potential, particularly in extracts, there is increasing interest in using these byproducts to deliver additional functional and health benefits while contributing to sustainability.

Extraction of bioactive compounds from longan processing byproducts has gained attention due to their health benefits and various applications, particularly the presence of compounds such as corilagin, ellagic acid, and gallic acid.^{4,5} These compounds help mitigate oxidative stress and related diseases.^{6,7} Notably, longan seeds exhibit higher antioxidant activity than lychee seeds (6122 vs. 334 mg GAE/100 g DW) under comparable extraction conditions.⁸ The growing demand for natural products, coupled with the need to valorize longan processing byproducts, has encouraged research into efficient extraction methods that maximize yield while preserving bioactive compounds.^{4,6}

The stability and bioavailability of bioactive compounds from longan processing byproducts determine their efficacy in functional foods, as the benefits depend on their survival through digestion and absorption. Studies have reported varying stability of bioactive compounds, with some retaining their bioactivity despite exposure to acidic pH, digestive enzymes, and bile salts.⁹ Additionally, the extraction method can impact the bioaccessibility of bioactive compounds during digestion.^{10,11} For instance, phenolic compounds in longan peel maintained antioxidant capacity throughout digestion.¹² However, ensuring bioavailability remains a challenge, as not all compounds survive digestion or achieve effective concentrations in the bloodstream. Nevertheless, *in vitro* models suggest a significant proportion remains accessible for uptake, indicating potential *in vivo* benefits.^{2,5}

- a. Faculty of Agro-industry, Chiang Mai University, 50100, Thailand
- b. Center of Excellence in Agro Bio-Circular-Green Industry, Chiang Mai University, Chiang Mai, 50100 Thailand
- c. Food Innovation and Packaging Center, Chiang Mai University, 50100, Thailand
- d. College of Food Science and Nutritional Engineering, China Agricultural University, Beijing, 100083, China
- e. Research group in Innovative Technologies for Sustainable Food (ALISOST), Department of Preventive Medicine and Public Health, Food Science, Toxicology and Forensic Medicine, Faculty of Pharmacy and Food Science, Universitat de València, Avda. Vicent Andrés Estellés, s/n, 46100 Burjassot, Valencia, Spain

See DOI: 10.1039/x0xx00000x



Despite advances in extraction techniques, several obstacles remain to achieving optimal extraction conditions. Conventional extraction technologies often yield low recovery.¹³ and compromise compound quality due to thermal degradation of heat-sensitive molecules, oxidation during prolonged processing, non-selective solvent extraction, and limited release of bioactive from rigid plant cell matrices.⁸ Moreover, studies on the optimization of water bath extraction and ultrasound-assisted extraction (UAE) remain limited. This study applied Response Surface Methodology (RSM) to optimize extraction conditions, aiming to enhance the recovery of bioactive compounds and antioxidants from longan processing byproducts. This work focuses on optimizing extraction conditions for longan processing byproducts using the UAE and water bath methods via RSM, while assessing various extraction factors. It further examines the relationships among optimal extraction conditions, stability, and gastrointestinal bioaccessibility of bioactive compounds in longan processing byproducts. These findings provide valuable insights into the potential application of longan processing byproducts in functional foods and nutraceuticals, contributing to sustainability and supporting the growing demand for natural, health-promoting ingredients.

2. Materials and Methods

2.1 Materials and sample preparation

Longan byproducts (LBP) were used in a mixed form, consisting of pericarps and (seeds and residual pulp) in approximately ratios of 63%, 37%, and on a dry weight basis, respectively, as described in supplementary Fig. 1, obtained from PM 80 Ltd. (Lamphun, Thailand). Chemicals and reagents, including DPPH, TPTZ, ABTS (2,2-diphenyl-1-picrylhydrazyl, 2,4,6-Tris(2-pyridyl)-s-triazine, 2,2'-azinobis (3-ethylbenzothiazoline-6-sulfonic acid), respectively), gallic acid, catechin, and Trolox, were all from Sigma (St. Louis, MO, USA). Ethanol (99.8 %) was sourced from a liquor distillery, Bangkok. Folin-Ciocalteu reagent, HPLC-grade standards, including procyanidin A2, gallic acid, epicatechin, ellagic acid, corilagin, quercetin, pinocembrin, pinostrobin, and the 5-fluorouracil ($\geq 99\%$ purity) were supplied by Merck KGaA (Darmstadt, Germany). Sample preparation followed a modified version of the method described by Tan *et al.*⁵ Briefly, the byproducts were milled using an herb grinder (Heavybao, vertical, HLB-JGY-AB, China) and passed through a 60-mesh sieve to produce a fine powder. The powdered

samples were transferred into polyethylene bags and sealed until further analysis. [View Article Online](#) DOI: 10.1039/D5FB00879D

DOI: 10.1039/D5FB00879D

2.2 Experimental methods

2.2.1 Extraction procedure using RSM

The experiment was conducted in seventeen runs, including three center points to estimate pure error and assess model adequacy (Table 1), following a Box-Behnken Design with three factors: ethanol-to-water ratio (X_1 : 60-80 % v/v; Y_1 : 40-80 %), temperature (X_2 & Y_2 : 40-80 °C), and time (X_3 : 60-240 min; Y_3 : 30-60 min). A solid-to-liquid ratio (1:10) was adopted from Bai *et al.*² and the extraction volume was 30 mL, consisting of equal volumes of food-grade ethanol (99.8 % or 95 %) and distilled water. The suspension was stirred at 150 rpm at 30 ± 1 °C, a centrifuge (Rotina 380R, Hettich Zentrifugen, Tuttlingen, Germany) at $8,644 \times g$ for 15 min at 25 °C, and the supernatant was filtered through Whatman No. 1 paper. The solvent was removed under reduced pressure at 40 °C using a rotary evaporator (BUCHI, Rotavapor® R-300, Thailand Ltd.), and the resulting crude extract was stored at -20 °C until further analysis. Optimization variables, with coded and actual values, are summarized in Table 1. Regression coefficients were obtained via multiple linear regression fitted to a second-order polynomial model (Eqs. 1 & 2), and the desirability function in (Minitab® Version 21.1.1, Minitab LLC, Chicago, USA) was applied to determine the optimal extraction conditions.

where X_1, Y_1 , X_2, Y_2 , and X_3, Y_3 , are the independent variables that affect the response Y/Z ; β_0 represents the intercept; β_i ($i = 1, 2, 3$) denotes the linear terms; β_{ii} ($i = 1, 2, 3$) corresponds to the quadratic terms and β_{ij} ($i = 1, 2, 3$; $j = 2, 3$) represents the cross-product terms.

2.2.2 Model validation

The predictive RSM equations were used to determine the optimal extraction conditions for bioactive compounds and antioxidants, based on time, solvent ratio, and temperature (Table 1). Following the development of the 2nd-order model and the multifactor analysis of variance, the desirability function was applied to identify the optimal extraction conditions.

Table 1: Experimental codes, ranges, and levels of independent variables in the response surface methodology experiment for water bath and UAE-assisted extraction. [View Article Online](#) DOI: 10.1039/D5FB00879D

Extraction methods	Symbol codes	Variables	Units	Levels		
				Low (-1)	Center (0)	High (+1)
Water bath method	X_1	Solvent ratio	%	60	70	80
	X_2	Temperature	°C	40	60	80
	X_3	Time	min	60	150	240
UAE method	Y_1	Solvent ratio	%	40	60	80
	Y_2	Temperature	°C	40	60	80
	Y_3	Time	min	30	45	60

2.3 Determination of bioactive compounds and antioxidants

2.3.1 Total phenolics and flavonoids contents

Total phenolic contents were measured following a modified protocol of Seling *et al.*,¹⁴ using Folin-Ciocalteu's phenol reagent and measured using a microplate reader (TECAN, Infinite 200 Pro, Mplex). Results were expressed as mg gallic acid equivalents per gram of dried weight. Total flavonoid content was determined with modifications according to Ayele *et al.*¹⁵ Absorbance was measured at 510 nm using a microplate reader (TECAN, Infinite 200 Pro, Mplex, Switzerland). Catechin was used as the standard, and results were expressed as mg catechin equivalents per gram of sample.

2.3.2 Determination of antioxidant properties (DPPH, ABTS, and FRAP)

The DPPH radical scavenging assay was carried out according to the method of Chaiwong *et al.*¹⁶ A DPPH solution (0.08 mg/mL) was prepared by dissolving 8 mg of DPPH in 70 % ethanol and adjusting the final volume to 100 mL in a volumetric flask. For the assay, 100 μ L of DPPH solution was mixed with 25 μ L of the extract at varying concentrations in microplate wells, and the mixture was incubated in the dark at 30 °C for 30 min. Absorbance was measured at 517 nm using the microplate reader. All measurements were performed in triplicate. The radical scavenging activity was calculated using Eq. (3).

% Inhibition

$$\frac{\text{Absorbance of control} - \text{Absorbance of sample}}{\text{Absorbance of control}} \times 100 \dots (3)$$

The ABTS radical scavenging activity of the extracts was determined using the method described by Chaiwong *et al.*¹⁶ with minor modifications. Briefly, the ABTS radical cation (ABTS⁺) was generated, and before analysis, the ABTS⁺ solution was diluted with deionized water to an absorbance of 0.70 ± 0.02 at 734 nm and equilibrated at 30 °C for 5 min. Absorbance was measured at 734 nm using the microplate reader. Results were expressed as mg Trolox equivalents (TE) per 100 g dry weight (DW).

The FRAP was determined using the method described by Chaiwong *et al.*¹⁶ Absorbance was measured at 595 nm using the microplate reader. A Trolox standard curve (0–100 μ g/mL) was used, and results were expressed as mg Trolox equivalents (TE) per 100 g dry weight (DW).

2.3.3 Simulated gastrointestinal digestion of the extract

Three digestion phases, including oral, gastric, and intestinal, were simulated *in vitro* following the revised protocol of Chailangka *et al.*,¹⁷ Simulated saliva was prepared by dissolving Na₂HPO₄ (2.4 g), KH₂PO₄ (0.2 g), and NaCl (8.0 g) in 1 L of distilled water, then adjusting the pH to 6.7 with 0.03 mol/L HCl. Then the mixture was mixed with α -amylase (200 U/mL, human saliva). Simulated gastric fluid consisted of pepsin (300 U/mL) in 0.03 mol/L HCl, while simulated intestinal fluid contained pancreatin (0.05 g) and bile extract (0.3 g) in 35 mL of 0.1 mol/L NaHCO₃. For digestion, 100 mg of the sample was homogenized in 5 mL of simulated saliva and incubated at 37 °C for 10 min using a shaking incubator (Daihan: Shaking Incubator WIS-20) at 100 rpm. The pH was then adjusted to 1.2 with 0.03 mol/L HCl, followed by the addition of 5 mL gastric fluid, and the mixture was incubated at 37 °C for 120 min using an incubator (Daihan: Shaking Incubator WIS-20) at 100 rpm. Subsequently, the pH was adjusted to 6.0 with 1 M NaOH, 5 mL of intestinal juice was added, the pH was readjusted to 7.0 with 1 M NaOH, and the mixture was mixed with 5 mL NaCl (120 mmol/L) and 5 mL KCl. Digestion proceeded for 120 min at 37 °C in the dark. After digestion, samples were centrifuged (6800 \times g, 30 min, 4 °C), and the supernatants were ultrafiltered using a 3-kDa cut-off membrane. Enzyme activity was stopped by adding 1:1 (v/v) methanol to the digested extract. The bioaccessibility was determined using Eq. 4

$$\text{Bioaccessibility} = \left(\frac{BD}{AD} \right) \times 100 \dots (4)$$

where AD = compound/antioxidant after digestion and BD = compound/antioxidant before digestion.

2.4 Identification of compounds by HPLC

Phenolic compounds were quantified by an HPLC (1220 Infinity II LC System, Agilent, Santa Clara, CA, USA) using a C18 column (250 \times 4.6 mm, 5 μ m) at 30 °C with acetonitrile and 0.1 % formic acid in water as the mobile phases under gradient elution (1.0 mL/min, 10–20 μ L injection). Detection was performed at 254–280 nm. Samples were diluted in 80 % methanol and filtered (0.45 μ m). LOD and LOQ were determined from the standard deviation of the response and the calibration slope,^{2,5} supplementary information comprehensive.

2.5 Statistical analysis

All determinations were performed in triplicate and analyzed using Analysis of Variance (ANOVA) with SPSS software (Version 23.0, Chicago, IL, USA). Duncan's multiple-range test was used to separate means, and differences were considered statistically significant at $p < 0.05$. Statistical modelling and optimization were conducted using Minitab statistical software (Minitab® Version 21.1.1, Minitab LLC, Chicago, USA).

3. Results and Discussion

3.1 Experimental responses for the water bath method

The extraction yield ranged from 5.30 % to 9.03 % (Table 2), indicating that solvent ratio, temperature, and time significantly ($p < 0.05$) influenced the efficiency of bioactive compound recovery. The highest yield (9.03 %) was achieved at 70 % solvent, 70 °C, and 150 min, confirming that a combination of moderate temperature and extraction time, with a balanced solvent polarity, is optimal for extracting phenolics and flavonoids. The result was within the range (8.1 – 15.5 %) of extractable yield reported by Sai-Ut *et al.*,⁸ for longan seed. The relatively lower value observed in this study may be due to compound degradation or limited solubilization under extreme conditions. In contrast, the yield was higher than the maximum value (8.5 %) reported by Fikry *et al.*,¹³ for the ultrasound-assisted extraction of longan seeds, indicating that extreme extraction parameters may cause degradation or reduced efficiency. As shown in Fig. 1 (a-c), the contour plots illustrate the interaction of solvent ratio, temperature, and extraction time on extraction yield, with an optimum region evident at the central points. In contrast, extreme solvent concentrations, prolonged extraction times, or elevated temperatures resulted in reduced yields. The regression equation (Table 4) for percentage yield, derived from the coded factors, indicated a good fit for the linear, quadratic, and interaction models.

TPC ranged from 42.89 to 79.56 mg GAE/100g DW, with the highest value (79.56 mg GAE/100 g DW) observed at 70% solvent, 70 °C, and 150 min, indicating that moderate extraction conditions promote phenolic recovery. Lower values observed at solvent concentrations of 60% and 80% suggest that both overly polar and non-polar systems reduce solubility and extraction efficiency.¹² Similarly, Natungnuy *et al.*,¹⁸ reported a TPC of 84.73 mg GAE/g, which was higher than that obtained in this study, likely due to the longer extraction duration (7 days). Mahindrakar and Rathod¹⁹ reported that Soxhlet extraction yielded 30.05 mg GAE/g of phenolics at 100 °C after 6 h, whereas batch extraction achieved a markedly higher TPC of 79.87 mg GAE/g at 50 °C within 105 min. Both values were lower than those observed in the present study, possibly due to differences in extraction time, temperature, and the material matrix. Likewise, Cao *et al.*,²⁰ reported a maximum TPC of 27.84 ± 0.33 mg


GAE/g DW from pawpaw leaves using RSM-optimized microwave-assisted extraction (14 min, 460 W, 77 °C, 30 mL/g, DW). This was comparable to the values obtained in this study. Fig. 1 (d-f) further illustrates these effects, with optimum conditions observed near the central points, with temperature–solvent interaction exerting the strongest influence. The regression models (linear, quadratic, and interaction) provided an adequate description of the TPC prediction, as summarized in Table 3.

Total flavonoid content ranged from 2.18 to 3.74 mg CE/100 g DW. The highest TFC (3.74 mg CE/100 g DW) was observed at 60 % solvent, 70 °C, 60 min, whereas some runs with higher TPC showed comparatively low TFC values (2.18 mg CE/100 g DW). This indicates that phenolics and flavonoids respond differently to extraction conditions, likely due to structural differences that influence their solubility and stability.¹⁴

The contour plots (Fig. 1g-i) highlight an optimum region around the central points, with the solvent–temperature interaction showing the most substantial effect. Prolonged extraction times promoted higher TFC recovery and the recovery of other compounds; however, they increased the risk of degradation²¹. Excessively long extraction durations further reduce efficiency, suggesting potential degradation of flavonoids.

For antioxidant activity, DPPH IC₅₀ values ranged widely, with the lowest IC₅₀ (4.63), indicating strong radical-scavenging activity. ABTS values ranged from 16.03 to 17.00 mg TE/100 g DW, which were lower than the value (94.04 mg TE/g) obtained by Natungnuy *et al.*,¹⁸ for longan seeds. In contrast, FRAP values ranged from 35.07 to 51.29 mg TE/100 g DW, with the highest value (51.29 mg TE/100 g DW) corresponding to the maximum TPC. Antioxidant activity showed a closer association with total phenolic content than with total flavonoid content, suggesting that phenolic compounds were the main contributors to the observed antioxidant capacity.²² These further state that natural plant extracts are typically abundant in antioxidants, making them valuable sources of bioactive compounds. This may be due to the broader variety of compounds included in TPC, which exhibit a wider range of antioxidant properties than those represented in TFC²³. As shown in Fig. 1 (j-l), the contour plots reveal that moderate solvent ratios, around 70 %, combined with lower temperatures, favor higher antioxidant activity (lower IC₅₀). In contrast, extreme solvent concentrations and prolonged extraction times increased IC₅₀ values, indicating reduced antioxidant potential, likely due to thermal or oxidative degradation of phenolics.²⁴ Additionally, Fig. 1 (m-r) describes the contour plot interaction effects for ABTS and FRAP, which exhibited different response patterns to the coded variables.

The regression model (Table 3) confirmed highly significant fits for all response parameters (Adj- $R^2 > 0.91$, $p < 0.005$), indicating that extraction conditions substantially influence both yield and antioxidant activity, as measured by the ABTS and FRAP assays.






Fig. 1: Contour plots of predicted models demonstrating the effects of temperature, solvent proportion, and time on yield (a-c); bioactive compounds (TPC (d-f) & TFC (g-i)); and antioxidants (DPPH (j-l), ABTS (m-o), & FRAP (p-r) of the water bath extraction method.

These findings highlight that moderate extraction conditions (70 % solvent, 70 °C, 150 min) consistently produced higher yields, TPC, and antioxidant activities, while extreme conditions either reduced extraction efficiency or promoted degradation of sensitive compounds, consistent with the findings of Tourabi *et al.*²⁵ Thus, balanced extraction parameters are critical for achieving both high yield and functional quality of extracts.

The desirability function, as first described by Derringer and Suich²⁶ approach in Minitab was employed to determine the optimal extraction conditions by simultaneously considering multiple responses. Responses (% Yield, TPC, TFC, DPPH, ABTS, and FRAP) were optimized simultaneously to maximize all responses while minimizing DPPH (antioxidant activity was expressed as IC50, with lower values indicating stronger radical-scavenging activity), targeting practical extraction efficiency. Each response was converted into a dimensionless desirability value (d_i) ranging from 0 to 1 (undesirable to entirely desirable). These values were aggregated using the geometric mean to calculate the overall

desirability of 0.85 for the water bath. The optimization analysis predicted optimal conditions of 69.09 % solvent ratio, 72.32 °C, and 149.09 min. Under these conditions, the overall desirability was 0.85, indicating a balance across all responses. The predicted values were experimentally validated, with percentage errors ranging from 0.65 to 7.55 %, confirming the adequacy of the optimization model.

3.2 Experimental responses for the UAE method

Yields ranged from 7.03 % to 12.43 % (Table 4), which is generally higher than those obtained with water bath extraction. The highest yield (12.43 %) was achieved at 60 % solvent, 60 °C, and 45 min, indicating that moderate conditions support the highest yield. Similarly, Fikry *et al.*¹³ reported that longan seed extraction reached its highest yield at 55 % ultrasonic power, 55 °C, and 25 min, which was 2.36 times higher than the minimum yield in their study. The consistent yields above 10 % across multiple UAE extraction conditions suggest that it is more effective at breaking cell walls and enhancing solvent penetration than water bath extraction.

Table 2. Experimental responses of the extraction yield, bioactive compounds (TPC and TFC), and antioxidant activity (DPPH, ABTS, and FRAP) of LBP under different extraction conditions using the water bath extraction method.

Runs	X ₁ (Solvent ratio, %)	X ₂ (Temperature, °C)	X ₃ (Time, min)	Extraction yield (%)	TPC (mg GAE/100g DW)	TFC (mg CE/100g DW)	DPPH IC ₅₀	ABTS (mg TE/100g DW)	FRAP (mg TE/100g DW)
1	60.00	60.00	150.00	7.49±0.64	42.89±0.74	2.77±0.07	5.82±0.03	16.34±0.45	35.07±0.06
2	80.00	60.00	150.00	6.35±0.32	48.32±0.89	2.58±0.03	6.19±0.07	16.67±0.87	41.08±0.77
3	60.00	80.00	150.00	7.56±0.33	46.70±0.29	3.20±0.05	4.64±0.02	16.58±0.35	42.54±0.46
4	80.00	80.00	150.00	6.65±0.67	46.37±0.87	2.86±0.12	4.63±0.00	16.46±0.78	39.46±0.28
5	60.00	70.00	60.00	5.35±0.65	43.87±0.28	3.74±0.11	7.37±0.04	16.03±0.05	35.96±0.67
6	80.00	70.00	60.00	6.13±0.78	59.42±0.29	2.56±0.34	6.78±0.33	16.12±0.57	36.78±0.75
7	60.00	70.00	240.00	7.03±0.23	63.67±0.22	2.18±0.17	5.20±0.08	16.42±0.63	36.68±0.88
8	80.00	70.00	240.00	5.30±0.76	56.89±0.84	2.23±0.19	5.77±0.56	16.04±0.97	35.26±0.64
9	70.00	60.00	60.00	5.61±0.97	43.53±0.18	3.06±0.23	8.47±0.23	16.23±0.35	36.74±0.45
10	70.00	80.00	60.00	7.08±0.46	55.42±0.88	3.65±0.09	6.09±0.11	16.82±0.74	39.50±0.63
11	70.00	60.00	240.00	7.20±0.43	67.87±0.26	2.54±0.32	5.10±0.09	16.87±0.76	38.43±0.67
12	70.00	80.00	240.00	5.78±0.02	57.11±0.11	2.66±0.37	5.89±0.45	16.32±0.36	36.79±0.98
13	70.00	70.00	150.00	9.03±0.67	74.53±0.46	3.28±0.06	5.79±0.21	17.00±0.57	47.88±0.57
14	70.00	70.00	150.00	8.76±0.56	79.56±0.38	3.41±0.02	5.32±0.32	17.00±0.89	51.29±0.88
15	70.00	70.00	150.00	8.69±0.83	70.24±0.29	3.13±0.36	5.89±0.46	17.00±0.64	48.21±0.44
Adj-R ²	-	-	96.420	-	92.360	91.110	92.440	94.350	91.410
p-value	-	-	-	0.000	0.002	0.003	0.002	0.001	0.003

Note: - GAE: Gallic acid equivalent; CE: Catechin equivalent; TE: Trolox equivalent; DW: Dry weight

Table 3: Regression equations in uncoded units for water bath extraction methods.

[View Article Online](#)
DOI: 10.1039/D5FB00879D

Responses	Water bath extraction method
Yield	$-95.8 + 1.622X_1 - 0.01140X_1^2 - 0.00675X_2^2 - 0.000214X_3^2 - 0.000697X_1X_3 - 0.000803X_2X_3$
TPC	$-1570 + 1.100X_3 - 0.1436X_1^2 - 0.1434X_2^2 - 0.000549X_3^2 - 0.00620X_1X_3 - 0.00629X_2X_3$
TFC	$-16.70 + 0.459X_1 + 0.148X_2 - 0.01078X_3 - 0.003604X_1^2 - 0.000029X_3^2 + 0.000342X_1X_3$
DPPH	$3.9 + 0.048X_2 - 0.1247X_3 + 0.000104X_3^2 + 0.000881X_2X_3$
ABTS	$-18.29 - 0.004475X_1^2 - 0.000049X_3^2 - 0.001125X_1X_2 - 0.000131X_1X_3 - 0.000317X_2X_3$
FRAP	$-579.5 - 0.05642X_1^2 - 0.03947X_2^2 - 0.000903X_3^2 - 0.02272X_1X_2$

In their UAE study of acerola residue, this mechanism yielded a higher recovery of phenolic compounds (approximately 1034 mg GAE/100 g) than conventional extraction (780 mg GAE/100 g), a trend consistent with our observations.^{28,29} This finding is comparable to those of Zhang *et al.*,³⁰ who reported TPC values of 22.09–132.47 mg GAE/100 g fresh weight in representative Chinese longan pulps. However, it was lower than the 57.8 ± 0.6 mg GAE/g DW reported by Bai *et al.*,² for longan pericarp. The variation may be attributed to the use of coupled technology, combining microwave and ultrasound, in their study, as well as differences in the characteristics of the plant material.

The TFC obtained by the UAE ranged from 11.37 to 29.00 mg CE/100 g DW, with the highest value observed at 60 % solvent, 60 °C, and 45 min. This condition correlated with high TPC runs, indicating that flavonoids respond particularly well to UAE extraction. Enhanced recovery of flavonoids and phenolic acid using UAE was reported by Jan and Gavahian²⁸ in mulberry extracts. The suitability of this temperature for maximizing TFC agrees with the findings of Vo *et al.*,³¹ who observed that raising the temperature to 60 °C increased TPC and TFC by 1.2 and 1.1-fold, respectively, in passion fruit peels. For antioxidant Activity, DPPH IC₅₀ values ranged from 3.45 to 7.97, indicating stronger radical-scavenging capacity under UAE conditions. The best activity (lowest IC₅₀, 3.45) was observed at the center point. Comparable results were reported by Sai-Ut *et al.*,⁸ in their optimization of lychee and longan seeds extraction. Similarly, Ayele *et al.*,¹⁵ obtained IC₅₀ values ranging from 3.5 to 6.4 from *Croton macrostachyus* root extracts. ABTS values (87.34–142.00 mg TE/100 g DW) and FRAP values (72.34–105.46 mg TE/100 g DW), as presented in Table 4, correlated well with the DPPH results, indicating consistency among different antioxidant assays and confirming the strong radical-scavenging potential of the extracts. Antioxidant assays aligned strongly with TPC and TFC, supporting that phenolics are the main contributors to antioxidant capacity.²⁴ High adjusted R² values (90.77–95.10) with significant p-values (<0.01) indicate that the model reliably explains the influence of solvent ratio, temperature, and time on extraction outcomes with the UAE method. These findings demonstrate that UAE markedly enhances the extraction efficiency of LBP by enabling

higher yields, greater recovery of phenolics and flavonoids, and enhanced antioxidant activity compared to conventional methods.^{13,27,32,33} Optimal responses were observed under moderate conditions (60 % solvent ratio, 60 °C, 45 min), where both TPC and TFC reached maximum levels and antioxidant capacity was highest. This finding is consistent with those of Fuangchoom *et al.*,³³ who reported similar results using microwave-assisted extraction. In contrast, extreme extraction conditions reduced efficiency or compromised the stability of bioactive compounds.³⁴ Therefore, optimized UAE conditions are essential for maximizing bioactive recovery and preserving the functional integrity of the extracts.

The regression model described by the linear, quadratic, and interactive equations (Table 5), showed strong fits for all responses (Adj-R² > 0.91, p < 0.005), confirming that extraction conditions significantly influenced yield and antioxidant activity. Moderate conditions (60 % solvent, 60 °C, 45 min) obtained superior results, while extreme conditions reduced efficiency or caused compound degradation, consistent with previous reports^{12,14,19}. Therefore, balanced parameters are crucial for maximizing yield and functional quality.

3.3 Validation of the model

The predictive accuracy of the RSM models was validated by the close agreement between predicted and experimental values for both extraction methods (Table 6). The percentage errors across responses were all less than 10 %, which is statistically acceptable for model reliability. This confirms that the models provided robust descriptions of extraction performance and can be used with confidence for process optimization.



Table 4. Experimental responses of the extraction yield, bioactive compounds (TPC and TFC), and antioxidant activity (DPPH, ABTS, and FRAP) of LBP under different extraction conditions using UAE methods.

Runs	Y_1 (Solvent ratio, %)	Y_2 (Temperature , °C)	Y_3 (Time, min)	Extraction yield (%)	TPC (mg GAE /100g DW)	TFC (mg CE /100g DW)	DPPH IC ₅₀	ABTS (mg TE /100 g DW)	FRAP (mg TE /100 g DW)
1	40.00	40.00	45.00	7.40±0.03	87.34±0.98	21.80±0.48	7.85±0.03	103.64±0.69	73.29±0.86
2	80.00	40.00	45.00	8.43±0.06	97.79±0.34	14.37±0.07	7.21±0.02	112.33±1.04	91.23±0.57
3	40.00	80.00	45.00	9.36±0.12	102.38±0.62	16.81±0.37	7.48±0.06	116.34±0.98	92.07±0.76
4	80.00	80.00	45.00	7.03±0.11	89.23±0.27	20.56±0.38	7.87±0.01	102.34±0.73	72.34±0.92
5	40.00	60.00	30.00	7.52±0.34	220.16±0.67	28.35±0.18	7.97±0.05	87.34±0.82	85.34±0.84
6	80.00	60.00	30.00	9.05±0.26	85.55±0.37	11.37±0.85	6.57±0.03	88.24±0.78	86.23±0.47
7	40.00	60.00	60.00	10.11±0.11	123.22±0.38	15.23±0.95	5.67±0.09	124.89±0.99	93.45±0.73
8	80.00	60.00	60.00	8.67±0.07	200.32±0.92	25.47±0.02	6.32±0.03	129.13±0.78	77.48±0.85
9	60.00	40.00	30.00	7.21±0.23	149.78±0.28	18.78±0.05	7.43±0.02	91.23±1.09	76.02±0.58
10	60.00	80.00	30.00	9.84±0.12	89.21±0.18	18.08±0.73	6.07±0.00	107.45±0.89	87.28±0.48
11	60.00	40.00	60.00	10.23±0.55	52.40±0.84	18.83±0.37	5.34±0.07	123.89±1.03	95.43±0.94
12	60.00	80.00	60.00	7.32±0.27	134.57±0.74	17.68±0.29	5.62±0.03	117.49±1.30	73.68±1.03
13	60.00	60.00	45.00	12.11±0.00	219.00±0.49	29.00±0.02	3.45±0.09	142.00±1.06	103.56±0.87
14	60.00	60.00	45.00	11.23±0.46	218.28±0.37	25.98±0.37	4.12±0.02	137.46±0.78	99.87±1.07
15	60.00	60.00	45.00	12.43±0.28	214.39±0.48	27.00±0.54	4.38±0.10	135.73±0.97	105.46±1.05
Adj-R ²	-	-	-	90.770	94.270	95.100	93.110	87.360	90.780
p-value	-	-	-	0.003	0.001	0.001	0.002	0.007	0.003

Note:- Temp.: Temperature; GAE: Gallic acid equivalent; CE: Catechin equivalent; TE: Trolox equivalent; DW: Dry weight.


ARTICLE

View Article Online

DOI: 10.1039/D5FB00879D

Table 5: Regression equations in uncoded units for UAE extraction methods.

Responses	Regression equations
Yield	$-61.47 - 0.004602Y_1^2 - 0.005069Y_2^2 - 0.00554Y_3^2 - 0.002100Y_1Y_2 - 0.002473Y_1Y_3 - 0.004617Y_2Y_3$
TPC	$-335 - 0.0903Y_1^2 - 0.2173Y_2^2 - 0.1058Y_3^2 + 0.1764Y_1Y_3 + 0.1190Y_2Y_3$
TFC	$3.5 + 0.119Y_3 - 0.00898Y_1^2 - 0.01338Y_2^2 - 0.01615Y_3^2 + 0.00698Y_1Y_2 + 0.02269Y_1Y_3$
DPPH	$54.98 - 0.4593Y_3 + 0.005171Y_1^2 + 0.003877Y_2^2 + 0.002581Y_3^2 + 0.001708Y_1Y_3$
ABTS	$-407.1 + 7.90Y_3 - 0.04044Y_1^2 - 0.03390Y_2^2 - 0.0659Y_3^2$
FRAP	$-346.3 - 0.02276Y_1^2 - 0.02907Y_2^2 - 0.03660Y_3^2 - 0.02354Y_1Y_2 - 0.02751Y_2Y_3$






Fig. 2: Contour plots of predicted models demonstrating the effects of temperature, solvent proportion, and time on yield (a-c); bioactive compounds (TPC (d-f) & TFC (g-i)); and antioxidants (DPPH (j-l), ABTS (m-o), & FRAP (p-r) of the UAE method.

Table 6: RSM predicted and experimental responses for the water bath and UAE-assisted extraction methods.View Article Online
DOI: 10.1039/D5FB00879D

Responses	Water bath method			UAE method		
	Predicted values	Experimented values	% error	Predicted values	Experimented values	% error
Yield (%)	8.83	8.21±0.02	7.55	11.95	11.23±0.32	6.03
TPC (mg GAE/100g DW)	73.80	70.86±0.37	4.15	215.30	219.11±1.45	1.77
TFC (mg CE/100g DW)	3.33	3.49±0.85	4.59	27.21	28.12±0.08	3.34
DPPH (IC ₅₀)	5.54	5.67±0.78	2.29	3.89	3.78±0.05	2.83
ABTS (mg TE/100g DW)	17.00	16.89±0.47	0.65	140.73	141.24±0.27	0.36
FRAP (mg TE/100g DW)	49.09	52.43±0.26	6.37	102.89	101.98±0.13	0.88

The optimal extraction conditions were determined using the desirability function approach in Minitab software. % Yield, TPC, TFC, ABTS, and FRAP were maximized, whereas DPPH was minimized (antioxidant activity was expressed as IC₅₀, where lower values indicate stronger radical scavenging activity) to achieve practical extraction efficiency. Each response was converted into a dimensionless desirability value (d_i) ranging from 0 (undesirable) to 1 (fully desirable). These values were combined using the geometric mean to calculate an overall desirability of 0.93 for the UAE method. The optimization analysis predicted the optimal conditions to be 59.39%, 59.79 °C, and 47.89 min. Under these conditions, the overall desirability was 0.93, indicating an excellent balance across all responses. The predicted values were experimentally validated, with percentage errors ranging from 0.36 to 6.03 %, confirming the adequacy of the UAE optimization model and indicating greater efficiency than the water bath-assisted method. The contour plot in Fig. 2 illustrates the interactive effects of the factors on all measured responses, whereas Table 5 summarizes the regression equations for each response along with their corresponding factors.

A comparative analysis of the two methods revealed that the UAE method outperformed the conventional water bath in extraction yield (%) and bioactive compound content. UAE achieved a higher yield of 11.23 %, compared with 8.21 % from the water bath-assisted method, a difference likely attributable to the ultrasonic cavitation,²⁷ which promotes solvent penetration and enhances mass transfer of the bioactive compounds.³⁵ Similarly, the UAE yielded TPC and TFC over 3- and 8-fold higher, respectively, highlighting its efficiency in releasing phenolic compounds that are less accessible during conventional thermal extraction.^{28,33}

Antioxidant activity assays further highlighted the advantages of UAE over conventional extraction. Ultrasound-assisted extracts exhibited lower DPPH IC₅₀ values and significantly ($p < 0.05$) higher ABTS and FRAP activities than those obtained with the water bath method. For instance, FRAP activity was 2 times higher under UAE conditions, reflecting a greater concentration of redox-active metabolites. These findings align with previous reports that UAE not only enhances

antioxidant extraction but also preserves thermolabile compounds that may degrade during prolonged heating.^{34,36} These results suggest that UAE is a greener, more efficient alternative to conventional extraction, yielding higher phenolic levels and stronger antioxidant activity in shorter extraction times with lower solvent use. This demonstrates potential for industrial applications in the production of functional food and nutraceuticals. The validated RSM models further support this applicability by enabling predictive control of process parameters.

3.4 Stability of compounds and bioactivity under *in vitro* gastrointestinal conditions

In vitro digestion (Fig. 4) revealed dynamic changes in phenolic content and antioxidant activity across the oral, gastric, and intestinal phases, indicating the influence of digestive conditions on the extract's stability. In general, UAE extracts consistently exhibited the highest bioactive compound and antioxidant activities at all stages, followed by water-bath extracts, while the untreated control remained the lowest. This can be connected to the original contents before digestion, as the values correspond precisely with the results after digestion. Previous studies have suggested that the extraction method influences the bioaccessibility of bioactive compounds during digestion.^{10,11} The *in vitro* digested extracts of the samples are presented in Fig. 3.

Total phenolic content increased during the gastric phase compared to the oral phase for water bath and control extracts, followed by a significant ($p < 0.05$) decline in the intestinal phase (Fig. 4). TPC in the UAE extract decreased steadily from the oral (280.80 mg GAE/100 g) to the gastric (236.57 mg) and intestinal phases (153.93 mg), unlike the control and water bath extracts, which increased during the gastric stage. This difference reflects the extraction mechanism: ultrasound disrupts cell walls and liberates free phenolics during processing, leaving fewer bound compounds to be released under acidic gastric conditions³⁷ observed from a study on black chokeberry extract. However, these free phenolics are also more susceptible to acid-induced degradation, accounting for the observed decline in the gastric phase.³⁸ In the intestinal phase, the

alkaline pH and bile salts accelerate oxidation and structural transformation. Phenolic acids such as gallic, caffeic, and chlorogenic acids are particularly unstable under these conditions.³⁹ Proanthocyanidins, another polyphenol found in LBP, also undergo rapid degradation under intestinal conditions, which explains the more pronounced TPC reduction in UAE extracts compared to the other treatments.⁴⁰ Interestingly, total flavonoid content (TFC) increased progressively across the digestion phases, particularly in UAE extracts (from 99.03 to 208.33 mg CE/100 g DW). This suggests that flavonoid glycosides undergo enzymatic or pH-induced hydrolysis, liberating aglycone forms that are more readily detected during intestinal digestion.^{39,40}



Fig. 3: *In vitro* gastrointestinal digested extracts and control (non-extract powder). The first three are controls (Ctrl) with oral, gastric, and intestinal from left to right-hand side. The three at the center are water bath (WBM) and UAE (UAEM) extracts, respectively, with oral, gastric, and intestinal extracts from left to right for both methods.

The antioxidant assays corroborated the phenolic trends, with DPPH radical scavenging (expressed as IC_{50}) generally improving during digestion. UAE extracts exhibited the strongest activity, with an IC_{50} of 2.43 at the intestinal phase. This could be due to the release of more minor, more active phenolic metabolites during digestion. ABTS and FRAP values increased progressively across the digestive phases, with UAE extracts reaching significantly higher levels ($p < 0.05$) in the intestinal phase (1313.60 mg TE/100 g DW and 538.44 mg TE/100 g DW, respectively). This trend suggests that the intestinal breakdown of complex phenolics generates metabolites with enhanced radical-scavenging and reducing power, as observed in black and green tea phenolic extracts.⁴¹ This agrees with prior reports indicating that *in vitro* digestion can convert polymeric phenolics into smaller, bioaccessible compounds with higher antioxidant potential.³⁸ The results for pH variations throughout the *in vitro* digestive process are shown in Table 7.

The superior stability and bioaccessibility of UAE extracts throughout digestion underscore the advantages of ultrasound-assisted extraction in producing phenolic-rich fractions that withstand gastrointestinal conditions. Compared with water bath extracts, the UAE extract not only exhibited higher initial phenolic concentrations but also retained greater activity after intestinal digestion. Similar observations have been reported by Iftikhar *et al.*,⁴² for phenolic compounds from rye bran, supporting the robustness of ultrasound-assisted techniques across diverse matrices. These findings

collectively suggest that the UAE facilitates the release of phenolic compounds with enhanced resilience to digestive degradation. The improved stability of UAE extracts may be attributed to multiple factors. First, ultrasound minimizes thermal exposure during processing, reducing the risk of phenolic oxidation or structural alteration compared with conventional heating.⁴³ Second, the mechanical cavitation generated during sonication enhances the solubilization of bound phenolics, thereby increasing their extractability and potential bioaccessibility.⁴⁴ These advantages suggest that the UAE is a green and effective strategy for extracting fruit byproducts into bioactive-rich ingredients with potential applications in functional foods and nutraceuticals. Fig. 5 shows the bioaccessibility of bioactive compounds and antioxidants.

3.5 HPLC analysis of the UAE extract after stimulated digestion

The HPLC chromatogram (Fig. 6) showed that the identified compounds were present in the digested UAE longan byproduct extract under the used conditions. Using established standard curves, the concentrations of these (gallic acid, epicatechin, quercetin, ellagic acid, procyanidin A2, corilagin, and pinostrobin) compounds were quantified at various stages of digestion (oral, gastric and intestinal), including the non-extract sample (Fig. 6A-D) within a retention time from 5 to 28 min and at different absorbances of 254, 270, and 280 nm. Ellagic acid shows the highest peak height across all samples, with a retention time of 11.753–11.776 min, indicating a higher content of the compound than the other available compounds at all stages of digestion. Moreover, quercetin and procyanidin A2 peaks are not present in all digestion stages except the orally digested extract. The intestinal digestion stage has shown only two peaks (ellagic acid and pinostrobin). Zhang *et al.*,³⁰ detected quercetin at 350 nm in longan pulp using a mobile phase of 0.4 % acetic acid and acetonitrile with a 45–50 min gradient flow (solution B 35–50 %). Unlike the analytical conditions used in this study, in which quercetin was not detected, likely due to differences in polarity and chromatographic elution strength.

Previous investigations have consistently identified ellagic acid, corilagin, and gallic acid as the major phenolic compounds in longan seeds and peels.^{2,5–7} Notably, Bai *et al.*,² identified only three phenolics (gallic acid, corilagin, and ellagic acid) within the range of 0.1–10 mg/g dry longan pericarp powder, which corresponds well with the core compounds found in this study, confirming their predominance in longan matrices. Bai *et al.*,² quantified ellagic acid (17.37 mg/kg), corilagin (5.25 mg/kg), and quercetin (3.12 mg/kg), in purified longan pericarp extracts, while Chindaluang and Sriwattana³² quantified substantially higher levels of corilagin (35.62 mg/g), ellagic acid (7.02 mg/g), and gallic acid (16.55 mg/g) in ultrasound-assisted extracts of longan seed. Similarly, Tang *et al.*,⁴⁵ reported ellagic acid contents of 0.18 mg/g in longan seed, nearly fivefold higher than that in the pericarp (0.03 mg/g), reinforcing the current finding that ellagic acid is the predominant phenolic. However, the observed quantitative differences can be attributed to variations in extraction solvents, chromatographic conditions, plant part analyzed, and the digestion conditions.






Fig. 4: A) Bioactive compounds and B) Antioxidant activity *in vitro* digestion. Note: TPC: Total phenolic compounds; total flavonoid compounds; GAE: gallic acid equivalent; CE: catechin equivalent.

Table 7: pH changes *in vitro* gastrointestinal digestion.

Treatment	Digestion	pH					
		0 min	10 min	30 min	60 min	90 min	120 min
Control (LBP powder)	Oral	6.70±0.00	5.92±0.01	ND	ND	ND	ND
	Gastric	1.20±0.00	ND	1.34±0.01	1.47±0.00	1.66±0.01	1.63±0.00
	Intestinal	7.00±0.00	ND	7.38±0.02	7.34±0.01	7.59±0.01	7.76±0.02
Water bath method	Oral	6.70±0.00	5.83±0.02	ND	ND	ND	ND
	Gastric	1.20±0.00	ND	1.48±0.02	1.52±0.00	1.60±0.01	1.59±0.00
	Intestinal	7.00±0.00	ND	7.16±0.02	7.66±0.02	7.51±0.00	7.63±0.01
UAE method	Oral	6.70±0.00	5.92±0.01	ND	ND	ND	ND
	Gastric	1.20±0.00	ND	1.43±0.01	1.37±0.01	1.46±0.00	1.54±0.00
	Intestinal	7.00±0.00	ND	7.23±0.02	7.35±0.02	7.57±0.02	7.55±0.02

Note: LBP: longan byproduct; ND: not determined

ARTICLE

View Article Online

DOI: 10.1039/D5FB00879D




Fig. 5: A) Bioaccessibility of bioactive compounds and B) antioxidants at stages of gastrointestinal digestion. Note: TPC: Total phenolic compounds; total flavonoid compounds; GAE: gallic acid equivalent; CE: catechin equivalent.






Fig. 6: HPLC profile of the stimulated digested extract and non-extract: A1) Oral digested extract (1: corilagin, 2: procyanidin, 3: quercetin, 4: pinostrobin) and A2) Oral digested extract (1: gallic acid 2: epicatechin, 3: ellagic acid); B1) Gastric digested extract (1: corilagin and 2: pinostrobin) and B2) Gastric digested extract (1: gallic acid 2: epicatechin, 3: ellagic acid); and C1) Intestinal digested extract (1: pinostrobin) and C2) Intestinal digested extract (1: ellagic acid); D1) Non-extract (1: pinostrobin); D2) Non-extract (1: gallic acid and 2: ellagic acid).

Table 8: Calibration curve parameters, detection limits, and linearity for standard compounds.

Compound	Slope	Intercept	Residual SD	LOD ($\mu\text{g}/\text{mg}$)	LOQ ($\mu\text{g}/\text{mg}$)	R^2
Gallic acid	13.1	377.09	332.9	84	254.09	0.9952
Ellagic acid	0.29	-3.73	14.71	169	512.82	0.9807
Epicatechin	3.96	-421.90	116.82	97	295.16	0.9935
Corilagin	8.04	-465.07	144.95	60	180.29	0.9976
Procyanidin A2	2.92	-339.30	26.14	30	89.45	0.9994
Quercetin	21.1	62.24	63.4	9.9	30.11	0.9997
Pinostrobin	6.9	472.61	149.09	71	216.1	0.981

Note: LOD=limit of detection; LOQ=limit of quantification; SD=standard deviation

Table 9: Concentration (μg/g) of the UAE simulated digested extract of LBP by HPLC.View Article Online
DOI: 10.1039/D5FB00879D

Compound name	Oral digested extract	Gastric digested extract	Intestinal digested extract	Non-extract powder
Gallic acid	45.94±2.38 ^a	nd	-	nd
Corilagin	164.08±1.04 ^a	138.34±2.58 ^b	-	-
Ellagic acid	885.12±0.72 ^a	119.22±1.15 ^{bc}	208.53±1.55 ^b	70.69±0.61 ^c
Epicatechin	219.78±2.89 ^b	227.14±2.11 ^a	-	-
Quercetin	nd	-	-	-
Procyanidin A2	241.18±7.38 ^a	-	-	-
Pinostrobin	nd	nd	nd	nd

Note: nd means not detected. Values with different superscripts within a row are significantly different ($p>0.05$).

Previous investigations have consistently identified ellagic acid, corilagin, and gallic acid as the major phenolic compounds in longan seeds and peels.^{2,5-7} Notably, Bai *et al.*² identified only three phenolics (gallic acid, corilagin, and ellagic acid) within the range of 0.1–10 mg/g dry longan pericarp powder, which corresponds well with the core compounds found in this study, confirming their predominance in longan matrices. Bai *et al.*² quantified ellagic acid (17.37 mg/kg), corilagin (5.25 mg/kg), and quercetin (3.12 mg/kg), in purified longan pericarp extracts, while Chindaluang and Sriwattana³² quantified substantially higher levels of corilagin (35.62 mg/g), ellagic acid (7.02 mg/g), and gallic acid (16.55 mg/g) in ultrasound-assisted extracts of longan seed. Similarly, Tang *et al.*⁴⁵ reported ellagic acid contents of 0.18 mg/g in longan seed, nearly fivefold higher than that in the pericarp (0.83 mg/g), reinforcing the current finding that ellagic acid is the predominant phenolic. However, the observed quantitative differences can be attributed to variations in extraction solvents, chromatographic conditions, plant part analyzed, and the digestion conditions.

The reduction in phenolic diversity and concentration observed across digestive stages aligns with the findings of Fang *et al.*,⁴⁶ who reported that 5-hydroxymethylfurfural (5-HMF) levels in dried longan increased during oral digestion but declined significantly ($p < 0.05$) after gastric and intestinal digestion. Moreover, 5-HMF can undergo biotransformation into sulfonylmethylfurfural, a genotoxic compound, during *in vitro* digestion,⁴⁶ highlighting the complexity of gastrointestinal transformations. Similarly, Bao *et al.*⁴⁷ found that although phenolic content and antioxidant activity decreased in digested tartary buckwheat, the remaining antioxidant potential was still considerable, suggesting that degradation or transformation products retain functional bioactivity. Furthermore, Kessy *et al.*⁴⁸ demonstrated that hydrolytic enzymes such as hydropectinase, β -glucosidase, and tannase promote the release and transformation of phenolics in litchi pericarps, processes likely analogous to those occurring in longan digestion.

The persistence of ellagic acid and corilagin throughout digestion may indicate greater structural stability and potential bioaccessibility than those of other phenolics. Corilagin, in particular, exhibits broad pharmacological activities, including antifungal effects against *Candida glabrata*, antihypertensive action, and cardiovascular protection.⁴⁵ Additionally, biotransformation during digestion can

generate novel metabolites with enhanced or distinct bioactivities,⁴⁹ implying that gastrointestinal modification of longan phenolics may not solely result in degradation but could also yield beneficial derivatives.

These findings suggest that ellagic acid and corilagin are not only key phenolics in LBP but also among the most stable under simulated digestive conditions. The observed compound-specific variability highlights the importance of gastrointestinal biotransformation in modulating the bioaccessibility and potential bioactivity of longan-derived phenolics.

4. Conclusions

This study optimized the extraction of phenolic-rich compounds from longan industrial byproducts by comparing conventional water-bath-assisted and ultrasound-assisted extraction methods. Response surface methodology using a Box–Behnken design revealed significant differences ($p < 0.05$), with UAE achieving the highest desirability ($D = 0.93$). Furthermore, simulated gastrointestinal digestion demonstrated that phenolics from the extract endured the stress and remained bioaccessible, with UAE extracts exhibiting the highest retention of antioxidant activity, suggesting greater potential for *in vivo* functionality. Although intestinal digestion reduced TPC, the concurrent increase in antioxidant capacity implies functional transformation of polyphenols rather than degradation. Finally, these results uncovered the potential of the UAE method as a green, efficient, and potentially scalable strategy for valorizing longan industrial byproducts into functional food and nutraceutical ingredients. The HPLC profile of the UAE extract showed that compounds were available at the initial stage of digestion, whereas most became unavailable. Future investigations should first focus on encapsulating the phenolic-rich extract to improve targeted bioavailability and clinically validate its health-promoting effects.

Acknowledgments

The authors thank the Center of Excellence in Agro Bio-Circular-Green Industry, Chiang Mai University, for technical support. In



Journal Name

ARTICLE

addition, thanks for the financial support from the "Fundamental Fund", "Reinventing University Program" and the "International Research Fellowship" Chiang Mai University. The first author is a PhD student in the Division of Food Science and Technology, Faculty of Agro-Industry, Chiang Mai University, under the CMU Presidential Scholarship.

Author contributions

Idris Kaida Zubairu: Investigation, Data curation, Conceptualization, Writing - original draft. Noppol Leksawasdi: Writing - review & editing. Sutee Wangtueai: Writing - review & editing. Pinpanit Boonchuay: Writing - review & editing. Fei Lao: Writing - review & editing. Juan Manuel Castagnini: Writing - review & editing. Su Lwin Htike: Writing - review & editing. Yuthana Phimolsiripol: Supervision, Conceptualization, Writing - review & editing.

Conflicts of interest

The authors declare no known competing financial or personal interests that could have influenced the work reported in this article.

Data availability

Data is available from the corresponding author upon reasonable request. The cited references in the supplementary information are provided in the reference list section.

References

- 1 T. Sawasdee, Factors behind the increased yield of six Thai fruits in 2025: Market linkages and value enhancement, <https://thailand.go.th/useful-information-detail/-6---68-->.
- 2 X. Bai, R. Pan, M. Li, X. Li and H. Zhang, *Molecules*, 2019, **24**, 619–619.
- 3 Y. Hu, X. Zhang, D. Li, C. Ma, L. Dong, Y. Luo, X. Hu and F. Chen, *Food Chemistry*, 2025, 145985.
- 4 Z. Ke, S. Tan and S. Shi, *Analytical Sciences*, 2023, **39**, 1405–1412.
- 5 S. Tan, Z. Ke, C. Zhou, Y. Luo, X. Ding, G. Luo, W. Li and S. Shi, *Molecules*, 2023, **28**, 2083–2083.
- 6 K. Rakariyatham, D. Zhou, N. Rakariyatham and F. Shahidi, *Journal of Functional Foods*, 2020, **67**, 103846.
- 7 I. K. Zubairu, K. Rakariyatham, S. Bai-Ngew, N. Leksawasdi, J. M. Regenstein, F. Lao, H. Hong, W.-S. Shin, K. J. Alzahrani and Y. Phimolsiripol, *Current Nutrition Reports*, 2025, **14**, 28.
- 8 S. Sai-Ut, P. Kingwascharapong, Md. A. R. Mazumder and S. Rawdkuen, *Foods*, 2023, **12**, 2827.
- 9 W. Qin, S. Ketnawa and Y. Ogawa, *Food Science and Human Wellness*, 2022, **11**, 669–675.
- 10 S. Nisa, Y. Bibi, S. Masood, A. Ali, S. Alam, M. Sabir, A. Qayyum, W. Ahmed, S. Alharthi, E. Y. Santali, S. A. Alharthy, W. M. Bawazir and M. N. Almarshjary, *Molecules*, 2022, **27**, 7932.
- 11 M. Rudzińska, A. Grygier, G. Knight and D. Kmiecik, *Foods*, 2024, **13**, 1814.
- 12 M. Xu, L. Ran, N. Chen, X. Fan, D. Ren and L. Yi, *Food Chemistry*, 2019, **297**, 124970.
- 13 M. Fikry, S. Jafari, K. A. Shiekh, I. Kijpatanasilp, S. Khongtongsang, E. Khojah, H. Aljumayi and K. Assataarakul, *Ultrasonics Sonochemistry*, 2024, **108**, 106949.
- 14 G. S. Seling, R. C. Rivero, C. V. Sisi, V. M. Busch and M. P. Buera, *Foods*, 2025, **14**, 2927.
- 15 D. T. Ayele, M. L. Akele and A. T. Melese, *BMC Chemistry*, 2022, **16**, 30.
- 16 N. Chaiwong, M. Gavahian, P. Chompoorat Tridttanakiat, P. Therdtthata, C. Moukamnerd, N. Leksawasdi and Y. Phimolsiripol, *Innovative Food Science & Emerging Technologies*, 2025, **104**, 104118.
- 17 A. Chailangka, N. Autsavapromporn, S. Karnjanapratum, N. Leksawasdi, J. M. Castagnini, F. J. Barba, A. Mousavi Khanegah and Y. Phimolsiripol, *Food Hydrocolloids*, 2024, **146**, 109288.
- 18 K. Natungnuy, P. P. Chareonsap and S. Poeaim, *International Journal of Agricultural Technology*, 2018, **14**, 1505–1514.
- 19 K. V. Mahindrakar and V. K. Rathod, *Chemical Engineering and Processing - Process Intensification*, 2020, **149**, 107841.
- 20 D. Cao, X. Qiao, Y. Guo and P. Liu, *Food Chemistry X*, 2024, **22**, 101500.
- 21 E. Brglez Mojzer, M. Knez Hrnčič, M. Škerget, Ž. Knez and U. Bren, *Molecules*, 2016, **21**, 901.
- 22 Y. M. Muflihah, G. Gollavelli and Y.-C. Ling, *Antioxidants*, 2021, **10**, 1530.
- 23 J. A. Domínguez-Avila, *Foods*, 2023, **12**, 4205–4205.
- 24 S. Aryal, M. K. Baniya, K. Danekhu, P. Kunwar, R. Gurung and N. Koirala, *Plants*, 2019, **8**, 96.
- 25 M. Tourabi, K. Faiz, R. Ezzouggi, B. Louasté, M. Merzouki, M. Dauelbait, M. Bourhia, K. S. Almaary, F. Siddique, B. Lyoussi and E. Derwich, *Bioresour. Bioprocess.*, 2025, **12**, 24.
- 26 G. Derringer and R. Suich, *Journal of Quality Technology*, 1980, **12**, 214–219.
- 27 A. K. Adam, T. Sadhu, I. K. Zubairu, S. S. Nassarawa and M. M. Nasiru, *Journal of Food Composition and Analysis*, 2025, **148**, 108300.
- 28 K.-C. Jan and M. Gavahian, *Food Bioscience*, 2025, **68**, 106686.
- 29 Y. R. R. S. Rezende, J. P. Nogueira, T. O. M. Silva, R. G. C. Barros, C. S. D. Oliveira, G. C. Cunha, N. C. Gualberto, M. Rajan and N. Narain, *Food Research International*, 2021, **140**, 109869.
- 30 R. Zhang, S. A. Khan, Y. Lin, D. Guo, X. Pan, L. Liu, Z. Wei, Y. Zhang, Y. Deng and M. Zhang, *International Journal of Food Properties*, 2018, **21**, 746–759.
- 31 Vo Phat T., Chieng Ha M. N., and Nguyen Quan D., *Chemical Engineering Transactions*, 2024, **108**, 13–18.
- 32 Y. Chindaluang and S. Sriwattana, *Chiang Mai University Journal of Natural Sciences*, DOI:10.12982/CMUJNS.2014.0047.
- 33 L. Shen, S. Pang, M. Zhong, Y. Sun, A. Qayum, Y. Liu, A. Rashid, B. Xu, Q. Liang, H. Ma and X. Ren, *Ultrasonics Sonochemistry*, 2023, **101**, 106646.
- 34 K. Kumar, S. Srivastav and V. S. Sharanagat, *Ultrasonics Sonochemistry*, 2021, **70**, 105325.
- 35 S. A. Siddiqui, A. Ali Redha, M. Salauddin, I. A. Harahap and H. P. V. Rupasinghe, *Critical Reviews in Analytical Chemistry*, 2025, **55**, 139–160.
- 36 X. Lin, J. Chen, G. Xiao, Y. Xu, D. Tang, J. Wu, J. Wen and W. Chen, *Food Science and Biotechnology*, 2016, **25**, 701–706.
- 37 S. Lachowicz-Wiśniewska, M. Świeca, I. Kapusta, A. Sip and I. Ochmian, *Sci Rep*, 2025, **15**, 28805.
- 38 R. B. Cuvas-Limon, P. Ferreira-Santos, M. Cruz, J. A. Teixeira, R. Belmares and C. Nobre, *Antioxidants*, 2022, **11**, 2479.
- 39 Y. Tang, W. Liu, J. Zhang, B. Juan, Y. Zhu, L. Zhu, Y. Zhao, M. Daglia, X. Xiao and Y. He, *Nutrients*, 2025, **17**, 2598.



Journal Name

ARTICLE

40 J. N. S. Souza, T. Tolosa, B. Teixeira, F. Moura, E. Silva and H. Rogez, *Molecules*, 2022, **28**, 66.

41 G. Annunziata, M. Maisto, C. Schisano, R. Ciampaglia, P. Daliu, V. Narciso, G. C. Tenore and E. Novellino, *Nutrients*, 2018, **10**, 1711.

42 M. Iftikhar, H. Zhang, A. Iftikhar, A. Raza, N. Begum, A. Tahamina, H. Syed, M. Khan and J. Wang, *LWT*, 2020, **134**, 110243.

43 H. V. Annegowda, R. Bhat, L. Min-Tze, A. A. Karim and S. M. Mansor, *J Food Sci Technol*, 2012, **49**, 510–514.

44 R. P. Putra, S. I. Aisyah, P. A. Kurniatin and W. Nurcholis, *Trop J Nat Prod Res*, 2024, **8**, 6552–6557.

45 Y.-Y. Tang, X.-M. He, J. Sun, C.-B. Li, L. Li, J.-F. Sheng, M. Xin, Z.-C. Li, F.-J. Zheng, G.-M. Liu, J.-M. Li and D.-N. Ling, *Molecules*, 2019, **24**, 1186–1186.

46 R. Fang, W. Sang, Q. Cheng and S. Liu, *Food Science*, 2024, **45**, 41–48.

47 T. Bao, Y. Wang, Y. Li, V. Gowd, X. Niu, H. Yang, L. Chen, W. Chen and C. Sun, *Journal of Zhejiang University-SCIENCE B*, 2016, **17**, 941–951.

48 H. N. E. Kessy, K. Wang, L. Zhao, M. Zhou and Z. Hu, *LWT*, 2018, **87**, 301–309.

49 X. Zhu, H. Wang, J. Sun, B. Yang, X. Duan and Y. Jiang, *J. Zhejiang Univ. Sci. B*, 2019, **20**, 503–512.

[View Article Online](#)
DOI: 10.1039/D5FB00879D



Data availability

Data is available from the corresponding author upon reasonable request. The cited references in the supplementary information are provided in the reference list section.