



Cite this: *Food Funct.*, 2025, **16**, 3227


### Correction: The comparative effects of $\omega$ -7 fatty acid-rich sea buckthorn oil and $\omega$ -3 fatty acid-rich DHA algal oil on improving high-fat diet-induced hyperlipidemia

Jing Li,<sup>a</sup> Jiahn Guo,<sup>a</sup> Michael Yuen,<sup>b</sup> Hywel Yuen<sup>b</sup> and Qiang Peng<sup>\*a</sup>

DOI: 10.1039/d5fo90023a  
rsc.li/food-function

Correction for 'The comparative effects of  $\omega$ -7 fatty acid-rich sea buckthorn oil and  $\omega$ -3 fatty acid-rich DHA algal oil on improving high-fat diet-induced hyperlipidemia' by Jing Li *et al.*, *Food Funct.*, 2025, **16**, 1241–1253, <https://doi.org/10.1039/D4FO04961F>.

In the original article, the authors regret that the Fig. 3 caption was incorrectly presented. The correct version of Fig. 3 is shown below.



**Fig. 3** Effects of SBT and DHA on the blood lipid levels of HFD-induced hyperlipidemia mice. (A) Serum TG, TC, LDL-C, HDL-C content of mice after modeling. (B) Serum TG content of mice after intervention. (C) Serum TC content in mice after intervention. (D) Serum LDL-C content in mice after intervention. (E) Serum HDL-C content in mice after intervention. (F) The atherosclerosis index (AI) of mice after intervention. (G) Coronary index (R-CHR) of mice after intervention. Data are expressed as mean  $\pm$  SEM,  $n = 6-8$ . Compared with CON group  $*p < 0.05$ ,  $**p < 0.01$ ; compared with HFD group  $\#p < 0.05$ ,  $###p < 0.01$ .

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

<sup>a</sup>College of Food Science and Engineering, Northwest A&F University, Yangling 712100, China. E-mail: pengqiang@nwsuaf.edu.cn

<sup>b</sup>Puredia Limited, Xining 810003, China

