

CORRECTION

[View Article Online](#)

[View Journal](#) | [View Issue](#)



Cite this: *Food Funct.*, 2025, **16**, 7959

Correction: Tropical postbiotics alleviate the disorders in the gut microbiota and kidney damage induced by ochratoxin A exposure

Shuaiming Jiang,^a Lingwei Du,^a Qian Zhao,^a Shunyong Su,^a Shi Huang^{*b} and Jiachao Zhang^{*a,c}

DOI: 10.1039/d5fo90079d

rsc.li/food-function

Correction for 'Tropical postbiotics alleviate the disorders in the gut microbiota and kidney damage induced by ochratoxin A exposure' by Shuaiming Jiang *et al.*, *Food Funct.*, 2024, **15**, 3980–3992, <https://doi.org/10.1039/D3FO05213C>.

The authors regret that in the original article panel F in Fig. 3 was incorrect. The correct version of Fig. 3 is shown below.

^aKey Laboratory of Food Nutrition and Functional Food of Hainan Province, School of Food Science and Engineering, Hainan University, Haikou 570228, China.

E-mail: zhjch321123@163.com

^bFaculty of Dentistry, The University of Hong Kong, Hong Kong SAR, China. E-mail: shihuang@hku.hk

^cOne Health Institute, Hainan University, Haikou, Hainan 570228, China






Fig. 3 Postbiotics alleviate the oxidative damage of the kidney induced by ochratoxin A. (A) The MDA level ($\text{nmol mg}_{\text{prot}}^{-1}$) of the kidney was measured. (B) The SOD level ($\text{U mg}_{\text{prot}}^{-1}$) of the kidney was measured. (C) The GSH level ($\mu\text{mol L}^{-1}$) of the kidney was measured. (D and E) The immunofluorescence and the expression levels of HO-1 are shown. (F and G) The immunofluorescence and the expression levels of Nrf2 are shown. The mean \pm standard deviation (SD) was used to represent the experimental values. The significance is shown as $*P < 0.05$, $**P < 0.01$, and $***P < 0.001$.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

