

Cite this: *Lab Chip*, 2025, **25**, 2795

Correction: The effect of cyclic fluid perfusion on the proinflammatory tissue environment in osteoarthritis using equine joint-on-a-chip models

Johannes Heidenberger,^a Eva I. Reihs,^{ab} Jonathan Strauss,^{ab} Martin Frauenlob,^b Sinan Gültekin,^c Iris Gerner,^c Stefan Toegel,^{ad} Peter Ertl,^b Reinhard Windhager,^a Florien Jenner^c and Mario Rothbauer^{*ab}

DOI: 10.1039/d5lc90048d

rsc.li/loc

Correction for ‘The effect of cyclic fluid perfusion on the proinflammatory tissue environment in osteoarthritis using equine joint-on-a-chip models’ by Johannes Heidenberger *et al.*, *Lab Chip*, 2025, **25**, 2256–2269, <https://doi.org/10.1039/d4lc01078g>.

The authors sincerely apologise for the incorrect duplication of Fig. 6A in the position of Fig. 8A. The corrected version of Fig. 8 is supplied here.



^a Department of Orthopedics and Trauma Surgery, Karl Chiari Lab for Orthopaedic Biology, Medizinische Universität Wien, Währinger Gürtel 18-20, 1090 Vienna, Austria.
E-mail: mario.rothbauer@mu.ac.at

^b Faculty of Technical Chemistry, Technische Universität Wien, Getreidemarkt 9/163, 1060 Vienna, Austria. E-mail: mario.rothbauer@tuwien.ac.at

^c Department for Small Animals and Horses, Centre for Equine Health and Research, Equine Surgery Unit, Veterinary Regenerative Medicine Laboratory, University of Veterinary Medicine Vienna, Veterinärplatz 1, 1210 Vienna, Austria

^d Ludwig Boltzmann Institute for Arthritis and Rehabilitation, Vienna, Austria



A**B**

Fig. 8 Effect fluid flow on inflamed mono- and cocultures of chondrocytes and synoviocytes. (A) Gene expression levels of perfused and IL-1 β -treated chondrocytes and synoviocyte biochip mono- and cocultures at day 10 post-seeding for MMP1 and IL6. The data is expressed as relative expression (fold change) to the housekeeping gene SDHA with $n = 3\text{--}5$ biological replicates (Mann–Whitney test, $*p < 0.05$). (B) Extended gene expression panel of fluid mechanically actuated IL-1 β -treated chondrocyte biochip mono- and cocultures for SOX9, RUNX2, COL10A1, SPARC, SPP1, COX2 and COMP genes. The data is expressed as relative expression (fold change) to the housekeeping gene SDHA with $n = 3\text{--}5$ biological replicates (Mann–Whitney test, $*p < 0.05$).

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

