ROYAL SOCIETY OF CHEMISTRY

CORRECTION

Check for updates

Correction: Negatively charged gold nanoclusters protect against diabetic cardiomyopathy by inhibiting mitophagy

Xiaolei Shen,[a] Dan Li,[b] Pengfei Zhuang,[b] Yang Yu,[a] Zuqiang Shi,[a] Xifan Mei*[a] and Chang Liu*[a]

Correction for 'Negatively charged gold nanoclusters protect against diabetic cardiomyopathy by inhibiting mitophagy' by Xiaolei Shen et al., New J. Chem., 2022, **46**, 10878–10886, https://doi.org/10.1039/D2NJ01536F.

The authors regret that an incorrect image was included for PEG-M in panel D of Fig. 3. The correct version of Fig. 3 is presented below.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

[a] The First Affiliated Hospital of Jinzhou Medical University, Jinzhou, China. E-mail: meixifan@jzmu.edu.cn, liuchang1971mei@jzmu.edu.cn
[b] Department of Basic Science, Jinzhou Medical University, Jinzhou, China

Fig. 3 MMP, ROS, and metabolic change caused by different AuNCs. (A) JC-1 assay was examined by an inverted fluorescence microscope (scale bar: 100 μm). Shown are the representative images from three independent H9C2 cell samples. (B and C) The fluorescence intensity was measured by ImageJ. Data were expressed as mean ± SD (n = 3). (D) Representative microscopy images of H9C2 cells incubated with AuNCs and palmitate acid for 24 h. H9C2 cells were stained by mitotracker red and green DCFH-DA (scale bar: 100 μm). (E) A representative picture of CPT1B expression was shown by WB and (F) quantified by ImageJ as the rate of CPT1B/GAPDH. GAPDH as the inner reference. Data were expressed as mean ± SD (n = 3). Significant differences (*p < 0.05, **p < 0.01, ***p < 0.005).