



Cite this: *New J. Chem.*, 2025, 49, 6495

DOI: 10.1039/d5nj90046h

rsc.li/njc

Correction: Synthesis of chiral hexynones for use as precursors to native photosynthetic hydroporphyrins

Khiem Chau Nguyen,^a Duy T. M. Chung,^a Phattananawee Nalaoh^b and Jonathan S. Lindsey^{*a}

Correction for 'Synthesis of chiral hexynones for use as precursors to native photosynthetic hydroporphyrins' by Khiem Chau Nguyen *et al.*, *New J. Chem.*, 2024, **48**, 2097–2117, <https://doi.org/10.1039/D3NJ03900E>.


The authors regret that the line drawing of known compound tributyl(1,1-dimethoxymethyl)stannane, termed Sn-(OMe)₂, was incorrectly displayed in Scheme 6 in the original manuscript. The corrected version of Scheme 6 is shown here.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

^a Department of Chemistry, North Carolina State University, Raleigh, NC 27695-8204, USA. E-mail: jlindsey@ncsu.edu

^b Department of Chemistry, University of Tennessee, Knoxville, TN 37996, USA





Scheme 6 Diversification to form chiral 4-hexyn-2-ones (lower panel) with use of tributyltin-one-carbon synthons (top panel).

