



Cite this: *Org. Biomol. Chem.*, 2025, **23**, 4131

## Total synthesis of the HDAC inhibitor (+)-(R)-trichostatin A via *O*-directed dialkylacetylene free radical hydrostannation with $\text{Ph}_3\text{SnH}/\text{Et}_3\text{B}$ . The unusual inhibitory effect of a proximal $\alpha$ -OPv group on the course of a vinyl iodide Stille cross-coupling†‡

Ke Pan, Soraya Manaviazar§ and Karl J. Hale ID \*§

In this paper, a new asymmetric total synthesis of optically pure (+)-trichostatin A (**1a**) is described via a route that utilises a Marshall chiral allenylzinc addition between **9** and 4-dimethylaminobenzaldehyde (**10**) and an *O*-depivaloylation at its early stages. *O*-Directed free radical hydrostannation of the resulting propargylic alcohol **15** with  $\text{Ph}_3\text{SnH}/\text{cat. Et}_3\text{B}/\text{O}_2$  in PhMe at rt thereafter provided the (*Z*)- $\alpha$ -triphenylstannylvinyltin **16** in 80–89% yield, with complete stereocontrol and very high  $\alpha:\beta$  regioselectivity (25:1). A stereoretentive  $\text{I-Sn}$  exchange reaction between **16** and  $\text{I}_2$  (1.4 equiv.) in  $\text{CH}_2\text{Cl}_2$  ( $-78^\circ\text{C}$  to rt, 1 h) subsequently secured the vinyl iodide **18** in 84–96% yield. The latter was transformed into the enal **4** by successive TPAP/NMO (Ley–Griffith) oxidation and a high yielding (80%) Stille reaction between the  $\alpha$ -iodo enal **20** and  $\text{Me}_4\text{Sn}$ , catalysed by  $\text{Pd}(\text{PPh}_3)_4$  in DMF at  $60^\circ\text{C}$ , under the Baldwin–Lee conditions, which use  $\text{CsF}$  and  $\text{CuI}$  as promoters. A Wittig reaction between **4** and  $\text{Ph}_3\text{P}=\text{CHCO}_2\text{Et}$  (**5**), saponification, and DDQ oxidation next afforded (+)-trichostatic acid (**22**). Helquist's ethyl chloroformate mixed-anhydride/TBSONH<sub>2</sub> coupling procedure (ref. 17e) thereafter secured (+)-trichostatin A (**1a**) in good yield. This new total synthesis of **1a** is the first-ever successful application of the *O*-directed dialkylacetylene free radical hydrostannation with  $\text{Ph}_3\text{SnH}/\text{cat. Et}_3\text{B}/\text{O}_2$  in a dialkylaniline *N*-containing disubstituted alkynol system, and it now provides a convenient means of accessing many novel trichostatin analogues for future biological screening.

Received 14th November 2024,  
Accepted 7th March 2025

DOI: 10.1039/d4ob01848f  
[rsc.li/obc](http://rsc.li/obc)



The School of Chemistry and Chemical Engineering, Queen's University Belfast, Stranmillis Road, Belfast BT9 5AG, UK. E-mail: [k.hale120@btinternet.com](mailto:k.hale120@btinternet.com)

† Dedicated with admiration and respect to the memory of the great Professor Amos B. Smith III (2014 William H. Nichols Gold Medallist, 2015 RSC Perkin Prize Winner, 2009 RSC Simonsen Medallist, and 2002 RSC Centenary Prize Medallist) in recognition of his numerous magnificent achievements in complex natural product total synthesis, new synthetic methodology development, materials science, and his rational design of a totally new class of HIV-1-neutra-lising drugs. Sadly, Professor Smith passed away on the morning of Monday February 3<sup>rd</sup>, 2025, aged 80 years. His landmark contributions to the fields of organic synthesis and medicinal chemistry will continue to serve as a source of much future inspiration to us all. He will forever be missed by his many friends, former students, postdoctoral fellows, Associate Editors, and admirers within the world of organic chemistry. Professor Smith was a recipient of the Order of the Rising Sun of Japan.


‡ Electronic supplementary information (ESI) available: experimental procedures for the synthesis of **15**, copies of the NMR, IR, and HR mass spectra of all compounds, TLC photos, and additional experimental discussions. See DOI: <https://doi.org/10.1039/d4ob01848f>

§ Current address: Halazair Pharma Ltd, Edgware, Middlesex, HA8 7RB, UK.

## Introduction

For some time now, we have been studying the scope,<sup>1,2a</sup> mechanism,<sup>3,4</sup> and utility<sup>1</sup> of the *O*-directed free radical hydrostannation reaction of various propargylically-oxygenated dialkylacetylenes with stannanes,<sup>1–4</sup> and the protocol that has consistently emerged best for most acyclic synthetic applications is the room temperature  $\text{Ph}_3\text{SnH}/\text{cat. Et}_3\text{B}$  variant of this reaction performed in PhMe (Scheme 1).<sup>2a</sup>

But that is not to say that the thermally-mediated  $\text{Bu}_3\text{SnH}/\text{AIBN}$  counterpart<sup>4</sup> of this reaction does not have an equally important role to play in synthesis, particularly where tandem radical cyclisation is a cardinal requirement, as in Alabugin's traceless polyaromatic ring construction reactions,<sup>5</sup> where his team's powerful contributions have been particularly elegant, innovative and synthetically impactful. Those same studies<sup>5</sup> have also provided profound computational and experimental insights into the detailed mechanistic workings of the free radical hydrostannation reaction of propargylically-oxygenated



**Scheme 1** The rt *O*-directed free radical hydrostannylative synthesis of (*Z*)-trisubstituted alkenes with  $\text{Ph}_3\text{SnH}$ /cat.  $\text{Et}_3\text{B}$ .

disubstituted acetylenes, and affirmed the entirely free radical *O*-directed mechanism proposed for this process.<sup>3,4a-c</sup>

The *O*-directed  $\text{Ph}_3\text{SnH}$ /cat.  $\text{Et}_3\text{B}$  variant<sup>1,2</sup> of this reaction has now shown its versatility in a wide range of complex settings that have included total syntheses of the natural products (+)-pumiliotoxin B<sup>6</sup> and (−)-(3*R*)-inthomycin C.<sup>7</sup> A noteworthy double *O*-directed free radical hydrostannation has also been introduced for the simultaneous installation of two structurally distinct trisubstituted alkenes within target structures. Its synthetic utility was powerfully demonstrated by the fully stereocontrolled route that was developed to the C(7)–C(22)-sector of (+)-acutiphycin.<sup>8</sup> The Furstner team has likewise elegantly employed an *O*-directed free radical hydrostannation with  $\text{Ph}_3\text{SnH}$ , cat. AIBN and PhMe in their total synthesis of (+)-isomigrastatin.<sup>9</sup> In each of these synthetic applications, I–SnPh<sub>3</sub> exchange has played a central role in final trisubstituted alkene elaboration<sup>10</sup> and, for the synthesis of all-carbon branched alkene structures, the CuI/CsF Baldwin–Lee<sup>11</sup> and Farina  $\text{Ph}_3\text{As}$  variants<sup>12</sup> of the Stille reaction have both proven themselves immensely valuable.<sup>6–8,10</sup> Other Pd(0)-mediated cross-coupling methods (e.g. Suzuki, Negishi, and Cuprate) and carbonylations have shown themselves to be equally applicable in this sphere.<sup>10</sup>

While there have now been many impressive displays of the utility of the rt  $\text{Ph}_3\text{SnH}$ /cat.  $\text{Et}_3\text{B}$  *O*-directed hydrostannation in synthesis,<sup>6–8,10</sup> there remain some classes of alkynes on which this reaction has yet to be successfully applied. Prominent amongst these substrates are propargyloxy dialkylacetylenes with an *unprotected* amine functionality. These generally form a complex with the  $\text{Et}_3\text{B}$  initiator, to prevent the  $\text{O}_2$ -mediated  $\text{S}_\text{H}2$  radical initiation event from ever taking place. Although a stoichiometric excess of  $\text{Et}_3\text{B}$  can sometimes allow radical initiation to occur, frequently the outcomes of these processes are disappointing.


Despite all of the past difficulties, we recently decided to investigate whether weakly basic amines, such as anilines, might prove compatible with the  $\text{Ph}_3\text{SnH}$ /cat.  $\text{Et}_3\text{B}$  *O*-directed hydrostannation method. In this paper, we now report that the 4-dimethylanilino functionality is indeed well tolerated in this process, as demonstrated by our new asymmetric total synthesis of (+)-trichostatin A, where this reaction protocol was very effectively deployed alongside a Marshall chiral allenylzinc addition process.<sup>13</sup>

We also document here, for the first time ever, the profound inhibitory effect that an allylic OPv group can have on

the course of a Pd-catalysed cross-coupling of a vinyl iodide. The latter observation was made quite by chance, whilst we were attempting to apply such a coupling to a trisubstituted vinyl iodide with such a functionality. Given this synthetic impasse, we thought it important to record this difficulty here, to prevent others from becoming similarly ensnared in the future.

(+)-Trichostatin A (**1a**) is a highly potent, naturally-occurring, histone deacetylase (HDAC) inhibitor that has elicited considerable medicinal interest as a possible antifungal,<sup>14a</sup> anticancer,<sup>14b,c</sup> immunosuppressive,<sup>14d</sup> and anti-Duchenne muscular dystrophy drug,<sup>15</sup> but its clinical use has so far been precluded by its remarkably short plasma half-life (<6.3 min at 80 mg kg<sup>−1</sup>)<sup>14c</sup> and its pronounced genotoxicity. It is widely thought that if these undesirable properties could somehow be ablated by careful structural modification, as was recently found for glycosylated (+)-trichostatin C (**1b**),<sup>16</sup> and HDAC specificity could also be improved, such trichostatin analogues might potentially become useful new treatments for a host of diseases. It was with such pharmaceutical and synthetic objectives in mind that we first embarked on the development of a new improved total synthesis<sup>17</sup> of (+)-trichostatin A (**1a**) ( $\text{R} = \text{OH}$ ), for the purpose of providing novel C(4)-analogues<sup>17e</sup> for drug screening and X-ray crystallographic/NMR studies. It was hoped that such studies might provide useful new insights into how individual HDAC isozymes work, as well as new therapeutic drugs.

Accordingly, we duly formulated the retrosynthetic plan of Scheme 2 for the synthetic acquisition<sup>17</sup> of (+)-(*R*)-trichostatin



**Scheme 2** Retrosynthetic planning for (+)-(6*R*)-trichostatin A (**1a**).



**A (1a).** Underpinning our approach would be the *O*-directed free radical hydrostannation of alkyne **8** with  $\text{Ph}_3\text{SnH}$ / $\text{Et}_3\text{B}$ , which would be allied with an I-Sn exchange and a Stille cross-coupling with  $\text{Me}_4\text{Sn}$ . An *O*-depivaloylation and an alcohol oxidation would thereafter procure **4**. The latter would then be chain extended with the ylide **5**, the C(1)-carboxyl subsequently unmasked, and the C(7)-ketone thereafter elaborated, prior to C(1)-oxyamidation with  $\text{TBSONH}_2$ . An *O*-desilylation would then complete the synthesis of **1a**.<sup>17</sup> Alkyne **8** would itself be created through a new Marshall chiral allenylzinc addition between the known chiral allenylzinc **9**<sup>13</sup> and 4-dimethylaminobenzaldehyde (**10**) under  $\text{Pd}(0)$ -catalysis.

## Results and discussion


Our first objective was the preparation of the chiral allenylzinc **9** from the known propargyl *O*-mesylate **11** (Scheme 3)<sup>13</sup> by the method of Marshall and Xie.<sup>13</sup> Specifically, **11** was added dropwise to a solution of  $\text{Pd}(\text{OAc})_2$  (0.05 equiv.) and  $\text{Ph}_3\text{P}$  (0.05 equiv.) that had been pre-aged for 20 min in dry THF at  $-20\text{ }^\circ\text{C}$ , under  $\text{N}_2$ . The reactants were then stirred for a further 5 min post-addition, before 4-dimethylaminobenzaldehyde (**10**) was added dropwise, followed by  $\text{Et}_2\text{Zn}$  (3 equiv.). Stirring was then continued at  $-20\text{ }^\circ\text{C}$  for 48 h to give an essentially inseparable 1 : 1 mixture of two alcohol diastereomers at C(5). This was inconsequential for further synthetic progression, since the C(5)-alcohols of **12** would ultimately be oxidized to the C(7) ketone in **1a**, at the penultimate step, and this same reaction had fully controlled the stereochemistry of the key C(4)-Me group in **12** (this would be C(6) in **1a**). After *O*-silylation with  $\text{TBSCl}$ , the alkynyl *O*-pivaloate **8** was subjected

to an *O*-directed free radical hydrostannation with  $\text{Ph}_3\text{SnH}$  (1.5 equiv.) and  $\text{Et}_3\text{B}$  (0.1 equiv.) in  $\text{PhMe}$  under our standard rt conditions. This reaction proceeded efficiently, affording **7** as two C(5)-diastereomeric products in 78% yield after 18 h. With **7** in hand, the key I-Sn exchange was investigated to obtain **6**. It was found that when this reaction was performed with just 1.5 equiv. of NIS at  $0\text{ }^\circ\text{C}$ , it did not proceed to completion, even after 2 h. However, when an additional 4 equiv. of NIS was added at rt,<sup>7</sup> and the reactants were stirred for a further 2 h, the reaction did deliver the slightly impure vinyl iodide **6**, as a much *slower-moving* product spot on TLC analysis. The necessity for such a large excess of the NIS was indicative of strong internal O-Sn coordination occurring within **7**, between the pivaloyloxy-carbonyl-O and the  $\text{SnPh}_3$  substituent, which preferentially enhanced the reactivity of the Ph groups.<sup>7</sup> As a result, a highly polar vinyltin triiodide initially needed to form, which was then eventually replaced by the desired vinyl iodide after prolonged stirring at rt. Iodide **6** was next subjected to a Stille cross-coupling with  $\text{Me}_4\text{Sn}$  under the normally successful Baldwin–Lee conditions.<sup>11</sup> Surprisingly, this reaction performed very poorly; an outcome that we have attributed to an analogous strong internal coordinative effect from the nearby OPv group, which we believe displaces the iodide from the initially formed vinyl-palladium(II) iodide to give **13**,<sup>18</sup> whose high stability then halts iodide return, to prevent the key transmetalation step from ever occurring with the  $\text{Me}_4\text{Sn}$  or hypervalent  $[\text{Me}_4\text{SnF}]$ .


In light of this setback, we decided to cleave the OPv group from **8** with  $\text{i-Bu}_2\text{AlH}$  (2 equiv.) in  $\text{CH}_2\text{Cl}_2$  at  $-78\text{ }^\circ\text{C}$ . The reaction took 2 h to reach completion, affording the desired alkynol **15** in 83% yield after extractive work up and  $\text{SiO}_2$  flash chromatography. It was later found more convenient and higher yielding (96–98% yield) to use  $\text{K}_2\text{CO}_3$  in  $\text{MeOH}$  to accomplish this transformation over 5 h at rt (Scheme 4). The resulting alkynol **15** was then subjected to *O*-directed hydrostannation with  $\text{Ph}_3\text{SnH}$  (1.7 equiv.) and  $\text{Et}_3\text{B}$  (0.1 equiv.) and  $\text{O}_2$  in  $\text{PhMe}$  (1 M in **15**) for 19 h at rt. This afforded a 25 : 1 mixture of  $\alpha$  :  $\beta$  vinyltriphenylstannanes **16** and **17**, from which the two  $\alpha$ -stannylated C(5)-diastereomers **16** could typically be purified with 80–89% yield after  $\text{SiO}_2$  flash chromatography.

Now that the Pv protecting group had been detached, the requisite I-Sn exchange reaction proceeded successfully with **16** to give **18** in >90% yield using only 1.4 equiv. of  $\text{I}_2$ . This result provided good confirmatory evidence that the OPv group had been adversely involved in strong internal O-coordination to the metal centre in both **7** and **13**, and our combined findings now pointed to a new strategic way forward.

The iodo-allylic alcohol **18** was next subjected to a Baldwin–Lee Stille cross-coupling with  $\text{Me}_4\text{Sn}$ .<sup>11</sup> It provided **19** efficiently in 80% yield after 22 h at rt. Importantly, this reaction had to be performed in complete darkness, inside a sealed vessel under  $\text{N}_2$ , to prevent light-induced vinyl iodide decomposition. Notwithstanding this success, difficulties soon arose when attempts were made to oxidise alcohol **19** under Swern, PCC, or  $\text{MnO}_2$  oxidation conditions, due to the presence of the di-



**Scheme 3** Initially investigated route to **7** and **14**.



**Scheme 4** The *O*-directed hydrostannylative route to (+)-(*R*)-trichostatin A (**1**).

methylamino group. Only cat. TPAP/NMO<sup>19</sup> afforded the desired enal **4**, but in a disappointing 20% yield.

Even so, this positive outcome did prompt us to evaluate the Ley–Griffith TPAP (0.05 equiv.)/NMO (2 equiv.) oxidising system<sup>19</sup> on the iodoallylic alcohol **18** in CH<sub>2</sub>Cl<sub>2</sub> in the presence of 4 Å sieves. This reaction worked reasonably well at rt, providing the desired  $\alpha$ -iodo-enal **20** in 54–64% yield after just 15 min. To our surprise, a much slower-moving by-product was also formed alongside **20**; this is suspected to be the *N*-oxide based upon <sup>1</sup>H and <sup>13</sup>C NMR analysis. Fortunately, it could be separated from **20** by SiO<sub>2</sub> flash chromatography. The unstable iodo-enal **20** was thereafter used quickly for a Baldwin–Lee Stille cross-coupling with Me<sub>4</sub>Sn (15 equiv.), CsF (2.5 equiv.) and CuI (0.25 equiv.) in DMF at 60 °C; a reaction that needed to be conducted in darkness. The success of this coupling (76–80% yield) was confirmed by the presence of two new allylic methyl doublets at  $\delta$  1.62 ppm ( $J$  = 1.2 Hz) and 1.61 ppm ( $J$  = 1.6 Hz) in CDCl<sub>3</sub>. Due to **20** and **4** both having

near identical TLC plate mobilities after a single elution, it was necessary to follow the progress of this reaction by multi-elution TLC with 3 : 1 petrol/CH<sub>2</sub>Cl<sub>2</sub> as the eluent. The product enal **4** moved as a slightly slower single spot, when compared with the iodo-enal **20**; **4** also stained dark blue when the TLC plate was heated with the anisaldehyde/H<sub>2</sub>SO<sub>4</sub> stain. The iodo-enal **20** stained brown. It is worth noting that performing this coupling at a temperature significantly below 60 °C led to a marked diminution in the yield of **4**; so 60 °C is essential and optimal.

Surprisingly the Wittig condensation of aldehyde **4** with stabilised P-ylide **5** took 6–10 d to reach completion at 38–40 °C in PhMe/CH<sub>2</sub>Cl<sub>2</sub> (1.25 : 1), and again, **4** and **3** had near identical TLC mobilities. When complete, the reaction furnished the dienoate **3** in 86–90% yield with total geometric control.

Unfortunately, the subsequent *O*-desilylation of **3** with *n*-Bu<sub>4</sub>NF in THF, or CsF in DMF, or HF/pyridine did not lead to the expected alcohol but, instead, a multitude of products. With *n*-Bu<sub>4</sub>NF, 4-dimethylaminobenzaldehyde (**10**) was observed amongst the products, confirming that a retro-vinylous aldol cleavage occurred at C(6)/C(7). This unexpected result led to an immediate change in our synthetic planning.

Without separation, the two C(7)-diastereoisomers of ester **3** were saponified with 90–98% yield with LiOH (4 equiv.) in 1 : 1 : 1 THF/MeOH/H<sub>2</sub>O. The known acid **21**<sup>17f</sup> was then oxidatively converted into (+)-trichostatic acid (**22**) by brief exposure to 1.5 equiv. of DDQ in 20 : 1 CH<sub>2</sub>Cl<sub>2</sub>/H<sub>2</sub>O at 0 °C for just 7 min. Such precise timing was essential to avoid extensive decomposition of the (+)-trichostatic acid **22** so formed. Although, in principle, the viability of this reaction had already been demonstrated by Hosokawa and Tatsuta in 2005,<sup>17f</sup> these workers never supplied an experimental procedure for this reaction, which meant that considerable experimentation was needed on our part to obtain a successful outcome. Nonetheless, with the new procedure reported here, the above reaction can now deliver **22** in 54–71% yield, with only a small amount of the starting material ever remaining at the reaction end.

With acid **22** in hand, it could be converted into the mixed anhydride **23** by treatment with 2 equiv. of ethyl chloroformate at 0 °C in THF.<sup>17b,e</sup> Exposure to 2 equiv. of TBSO-NH<sub>2</sub> according to Helquist's procedure<sup>17e</sup> thereafter afforded **24**, which was deprotected with CsF in MeOH to obtain (+)-(*R*)-trichostatin A (**1**) in 40–60% yield over 2 steps, without loss of chirality.<sup>17</sup> The latter was proven by our synthetic conversion of **22**<sup>20</sup> into its naturally-occurring  $\beta$ -glucoside, (+)-trichostatin C (**1b**) (NMR data in the ESI†), which also, simultaneously, unambiguously confirmed the assigned structure of that natural product.<sup>16,20</sup>

## Conclusions

With this new enantioselective total synthesis of (+)-trichostatin A that has been developed,<sup>21</sup> we have provided yet another




powerful demonstration of the utility of the room temperature *O*-directed free radical hydrostannation of propargylically-oxygenated dialkylacetylenes with  $\text{Ph}_3\text{SnH}$  and  $\text{Et}_3\text{B}/\text{O}_2$  in complex molecule total synthesis,<sup>1–3,6–8,22</sup> and we have exemplified how it can be successfully deployed in an alkyne system that bears the dialkylaniline *N*-functionalities. We have also shown how vinyl iodides with a proximal  $-\text{CH}_2\text{OPv}$  group can internally *O*-coordinate to a vinyl palladium(II) intermediate in a manner that prevents it from successfully engaging in trans-metallation<sup>18</sup> and reductive elimination.

## Experimental

### Procedures for the total synthesis of (+)-(R)-trichostatin A (1)

For the synthesis of propargyl alcohol **15** see the accompanying ESI.†

### O-Directed hydrostannation of **15**: synthesis of vinyl triphenyltin **16**




Inside a sealed glove bag filled with an  $\text{N}_2$  atmosphere,  $\text{Ph}_3\text{SnH}$  (5.11 g, 14.56 mmol, 1.7 equiv.) was quickly weighed into a pear-shaped flask fitted with a rubber septum. When the weighing process was complete, the septum-sealed flask was removed from the glove bag and fitted with an  $\text{N}_2$ -filled balloon connected to a Luer-locked needle. Dry  $\text{PhMe}$  (8.5 mL) was added to the neat  $\text{Ph}_3\text{SnH}$  and that solution was cannulated into alkynol **15** (2.98 g, 8.57 mmol) under  $\text{N}_2$  at rt. A solution of  $\text{Et}_3\text{B}$  (1.0 M in hex, 1.30 mL, 1.30 mmol, 0.15 equiv.) was then added dropwise to the two reactants. Air (20 mL) was thereafter injected into the reaction mixture twice at 5 min and then at 1 h. The reaction mixture was stirred at rt for 19 h with the  $\text{N}_2$  atmosphere being maintained throughout. The reaction mixture was then diluted with  $\text{EtOAc}$  (20 mL) and quenched with  $\text{H}_2\text{O}$  (20 mL). The aqueous layer was extracted with  $\text{EtOAc}$  (20 mL  $\times$  5). The combined organic extracts were then dried using  $\text{MgSO}_4$ , filtered and concentrated under reduced pressure. The crude residue, which consisted essentially of a 25:1  $\alpha:\beta$  mixture of **16**:**17**, was purified by gradient elution  $\text{SiO}_2$  flash chromatography initially using petrol:  $\text{CH}_2\text{Cl}_2$  (3:1  $\rightarrow$  2:1  $\rightarrow$  1:1) to remove the tin residues and other impurities. Gradient elution with petrol:  $\text{EtOAc}$  (18:1  $\rightarrow$  15:1) thereafter gave the entire product **16** (4.76 g, 80%) as a mixture of C5-epimers and as a colourless oil. A small amount of the  $\beta$ -vinyltin adduct **17** was subsequently eluted with petrol:  $\text{EtOAc}$  (12:1  $\rightarrow$  8:1) as the eluent. It was characterised and identified by  $^1\text{H}$  NMR spectroscopy (see the ESI† for a copy of the 600 MHz  $^1\text{H}$  NMR spectrum of **17**). Data for **16**: IR of **16** (neat film): 3423 (m), 3063 (m), 3050 (m), 2959 (s), 2926 (s), 2856 (s), 1730 (w), 1615 (s), 1524 (s), 1431 (s), 1385 (m), 1254 (m), 1073 (s), 864 (m), 837 (m), 779 (m), 731 (s), 698 (s)  $\text{cm}^{-1}$ .

$^1\text{H}$  NMR of pure **16** (pure  $\alpha$ -diastereoisomer 1) (399.9 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.60 (m, 6H,  $^3J$   $^{119/117}\text{Sn}-^1\text{H}$  = ca. 48.0 Hz, *o*-CH<sub>2</sub>,  $-\text{SnPh}_3$ ), 7.35 (m, 9H, *p*- and *m*-CH<sub>2</sub>,  $-\text{SnPh}_3$ ), 6.78 (d,  $J$  = 8.4 Hz, 2H, H7), 6.35 (d,  $J$  = 8.7 Hz, 2H, H8), 6.43 (d,  $J$  = 10.0 Hz,  $^3J$   $^{119}\text{Sn}-^1\text{H}$  = 173.6 Hz,  $^3J$   $^{117}\text{Sn}-^1\text{H}$  = 153.6 Hz, 1H, H3), 4.32 (s, 2H,  $\text{CH}_2$ ,  $^3J$   $^{119/117}\text{Sn}-^1\text{H}$  = 54.0 Hz, H1a,b), 4.22 (d, 1H,  $J$  = 6.8 Hz, H5), 2.92 (s, 6H,  $\text{N}(\text{CH}_3)_2$ ), 2.47 (m, 1H,  $J$  = 6.8 Hz, H4), 1.32 (br, 1H, OH), 0.83 (s, 9H, *t*-Bu of OTBS), 0.56 (d,  $J$  = 6.8 Hz, C4-Me),  $-0.09$  (s, 3H,  $\text{CH}_3\text{Si}$ ),  $-0.27$  (s, 3H,  $\text{CH}_3\text{Si}$ ) ppm.  $^{13}\text{C}$  NMR of **16** (pure  $\alpha$ -diastereoisomer 1) (100.57 MHz,  $\text{CDCl}_3$ ):  $\delta$  149.6 (C9), 149.2 (C3,  $^2J$   $^{119/117}\text{Sn}-^{13}\text{C}$  = 32.2 Hz), 139.7 (C2), 139.4 (quaternary C,  $-\text{SnPh}_3$ ), 137.2 (*o*-C,  $-\text{SnPh}_3$ ,  $^3J$   $^{119/117}\text{Sn}-^{13}\text{C}$  = 38.2 Hz), 131.4 (C6), 128.7 (*p*-C,  $-\text{SnPh}_3$ ,  $^4J$   $^{119/117}\text{Sn}-^{13}\text{C}$  = 12.1 Hz), 128.5 (*m*-C,  $-\text{SnPh}_3$ ,  $^3J$   $^{119}\text{Sn}-^{13}\text{C}$  = 52.3 Hz), 127.9 (C7), 111.8 (C8), 79.3 (C5), 70.4 (C1,  $^2J$   $^{119/117}\text{Sn}-^{13}\text{C}$  = 46.3 Hz), 47.4 (C4,  $^3J$   $^{119/117}\text{Sn}-^{13}\text{C}$  = 38.2 Hz), 40.6 ( $\text{N}(\text{CH}_3)_2$ ), 25.9 ( $(\text{CH}_3)_3\text{CSi}$ ), 18.3 ( $(\text{CH}_3)_3\text{CSi}$ ), 17.6 (C4-Me),  $-4.4$  ( $\text{CH}_3\text{Si}$ ),  $-5.1$  ( $\text{CH}_3\text{Si}$ ) ppm.

$^1\text{H}$  NMR of the other diastereoisomer of **16** ( $\alpha$ -diastereoisomer 2) (399.9 MHz,  $\text{CDCl}_3$ ). Resonances and multiplicities have been reported, where determinable, from the purified but diastereomerically enriched 3:1 mixture of C5 epimers:  $\delta$  7.61 (m, 6H, *o*-CH<sub>2</sub>,  $-\text{SnPh}_3$ ,  $^3J$   $^{119/117}\text{Sn}-^1\text{H}$  = 48.4 Hz), 7.38 (m, 9H,  $-\text{SnPh}_3$ ), 6.44 (d,  $J$  = 8.8 Hz, 2H, H7) superimposed upon 6.42 (m, 1H, H3), 6.40 (d,  $J$  = 8.8 Hz, 2H, H8), 4.43 (d,  $J$  = 3.6 Hz, 1H, H5), 4.32 (s, 2H, H1), 2.88 (s, 6H,  $\text{N}(\text{CH}_3)_2$ ), 2.27 (m, 1H, H4), 1.32 (br, 1H, OH), 0.91 (s, 9H, *t*-Bu of OTBS), 0.74 (d,  $J$  = 6.8 Hz, 3H, C4-Me),  $-0.00$  (s, 3H,  $\text{CH}_3\text{Si}$ ),  $-0.23$  (s, 3H,  $\text{CH}_3\text{Si}$ ) ppm.  $^{13}\text{C}$  NMR (100.57 MHz,  $\text{CDCl}_3$ ) of the other diastereoisomer of **16** ( $\alpha$ -diastereoisomer 2). Resonances and  $J$  values have been reported, where determinable, from the purified but diastereomerically enriched 3:1 mixture:  $\delta$  150.1 (C9), 149.2 (C3), 139.2 (*Sn*-C-CH<sub>2</sub>,  $-\text{SnPh}_3$ ), 138.8 (C2), 137.1 (*o*-C,  $-\text{SnPh}_3$ ,  $^3J$   $^{119/117}\text{Sn}-^{13}\text{C}$  = 38.2 Hz), 131.6 (C6), 128.9 (*p*-C,  $-\text{SnPh}_3$ ,  $^4J$   $^{119/117}\text{Sn}-^{13}\text{C}$  = 12.1 Hz), 128.6 (*m*-C,  $-\text{SnPh}_3$ ,  $^4J$   $^{119/117}\text{Sn}-^{13}\text{C}$  = 52.3 Hz), 126.8 (C7), 111.7 (C8), 77.1 (C5), 70.3 (C1,  $^2J$   $^{119/117}\text{Sn}-^{13}\text{C}$  = 46.3 Hz), 47.12 (CH-Me), 40.7 ( $\text{N}(\text{CH}_3)_2$ ), 25.9 ( $(\text{CH}_3)_3\text{CSi}$ ), 18.3 ( $(\text{CH}_3)_3\text{CSi}$ ), 13.8 (CH-CH<sub>3</sub>),  $-4.5$  ( $\text{CH}_3\text{Si}$ ),  $-5.0$  ( $\text{CH}_3\text{Si}$ ) ppm.

TOF ES<sup>+</sup> HRMS of **16**: calcd for  $\text{C}_{38}\text{H}_{50}\text{NO}_2\text{SiSn}$  [ $\text{M} + \text{H}$ ]<sup>+</sup>: 700.2640. Found: 700.2611.

### I-Sn exchange of **16**: preparation of vinyl iodide **18**



Before commencing this experiment, the reaction flask was covered with Al-foil to protect it from the adverse effects of light. Thereafter, to a stirred  $-78$  °C solution of the vinyl triphenyltin **16** (4.76 g, 6.81 mmol) in dry  $\text{CH}_2\text{Cl}_2$  (68 mL) under  $\text{N}_2$  was added  $\text{I}_2$  (2.42 g, 9.52 mmol, 1.4 equiv.) in one portion. Stirring was continued at  $-78$  °C for 10 min, after which the cooling bath was removed, and the reactants were allowed to stir at rt for a further 1 h. The reaction mixture was then


diluted with  $\text{CH}_2\text{Cl}_2$  (50 mL) and quenched with  $\text{H}_2\text{O}$  (50 mL). The aqueous layer was extracted with EtOAc (50 mL  $\times$  3) and the combined organic extracts were successively washed with saturated aq.  $\text{Na}_2\text{S}_2\text{O}_3$  (100 mL) and  $\text{H}_2\text{O}$  (50 mL  $\times$  2). The organic extract was then dried over  $\text{MgSO}_4$ , filtered and concentrated *in vacuo*. The crude residue was purified by gradient elution  $\text{SiO}_2$  flash chromatography using petrol :  $\text{CH}_2\text{Cl}_2$  (3 : 1  $\rightarrow$  2 : 1  $\rightarrow$  1 : 1) as the eluent to remove tin residues, and thereafter petrol : EtOAc (15 : 1  $\rightarrow$  10 : 1) to elute iodide **18** (3.0 g, 93%) as a C5-mixture of epimers; it was obtained as an amber oil. IR for the **18** mixture (neat film): 3383 (m), 3101 (w), 3075 (w), 2959 (s), 2927 (s), 2856 (s), 2800 (m), 1738 (w), 1617 (s), 1521 (s), 1471 (m), 1357 (m), 1254 (m), 1074 (s), 941 (m), 876 (s), 835 (s), 775 (s)  $\text{cm}^{-1}$ .

Data for pure **18**:  $^1\text{H}$  NMR of **18** (pure isomer 1) (399.9 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.16 (d,  $J$  = 8.6 Hz, 2H, H7), 6.68 (d,  $J$  = 8.6 Hz, 2H, H8), 5.81 (d,  $J$  = 8.9 Hz, 1H, H3), 4.64 (d, 1H,  $J$  = 4.2 Hz, H5), 4.19 (s, 2H, H1a,b), 2.94 (s, 6H,  $\text{N}(\text{CH}_3)_2$ ), 2.65 (m, 1H, H4), 1.79 (br, 1H, OH), 0.94 (d,  $J$  = 7.0 Hz, 3H, C4-Me), 0.90 (s, 9H, *t*-Bu of OTBS), 0.02 (s, 3H,  $\text{CH}_3\text{Si}$ ), -0.20 (s, 3H,  $\text{CH}_3\text{Si}$ ) ppm.  $^{13}\text{C}$  NMR of **18** (pure isomer 1) (100.57 MHz,  $\text{CDCl}_3$ ):  $\delta$  149.8 (C9), 139.4 (C3), 130.9 (C6), 127.6 (C7), 111.9 (C8), 107.8 (C2), 76.8 ( $\text{CH}$ -OTBS), 71.9 (C1), 49.3 (C4), 40.7 ( $\text{N}(\text{CH}_3)_2$ ), 25.8 (( $\text{CH}_3$ )<sub>3</sub>CSi), 18.2 (( $\text{CH}_3$ )<sub>3</sub>CSi), 15.3 (C4-Me), -4.5 ( $\text{CH}_3\text{Si}$ ), -5.0 ( $\text{CH}_3\text{Si}$ ) ppm.

$^1\text{H}$  NMR of **18** (isomer 2) (399.9 MHz,  $\text{CDCl}_3$ ). Resonances and  $J$  values have been reported, where determinable, from the purified but diastereomerically enriched 3 : 1 mixture:  $\delta$  7.10 (d,  $J$  = 8.8 Hz, 2H, H7), 6.66 (d,  $J$  = 8.0 Hz, 2H, H8), 5.72 (d,  $J$  = 8.8 Hz, 1H, H3), 4.49 (d, 1H,  $J$  = 6.0 Hz, H5), 4.19 (s, 2H, H1), 2.94 (s, 6H,  $\text{N}(\text{CH}_3)_2$ ), 2.74 (m, 1H, H4), 1.58 (br, 1H, OH), 0.89 (d,  $J$  = 6.8 Hz, 3H, C4-Me), 0.87 (s, 9H, *t*-Bu of OTBS), -0.02 (s, 3H,  $\text{CH}_3\text{Si}$ ), -0.20 (s, 3H,  $\text{CH}_3\text{Si}$ ) ppm.  $^{13}\text{C}$  NMR of **18** (isomer 2) (100.57 MHz,  $\text{CDCl}_3$ ):  $\delta$  149.8 (C9), 139.4 (C3), 130.9 (C6), 127.6 (C7), 111.9 (C8), 107.7 (C2), 77.5 (C5), 71.9 (C1), 49.3 (C4), 40.7 ( $\text{N}(\text{CH}_3)_2$ ), 25.9 (( $\text{CH}_3$ )<sub>3</sub>CSi), 18.2 (( $\text{CH}_3$ )<sub>3</sub>CSi), 15.3 (C4-Me), -4.5 ( $\text{CH}_3\text{Si}$ ), -5.0 ( $\text{CH}_3\text{Si}$ ) ppm.

TOF ES<sup>+</sup> HRMS for **18**: calcd for  $\text{C}_{20}\text{H}_{35}\text{NO}_2\text{Si} [\text{M} + \text{H}]^+$ : 476.1482. Found: 476.1500.

### Preparation of allylic alcohol **19**




To the iodovinylic alcohol **18** (0.3065 g, 0.645 mmol) inside a small pear-shaped flask capped with a rubber septum and an  $\text{N}_2$ -filled balloon was added dry DMF (1.28 mL) *via* a syringe. The outside of the flask was wrapped in Al-foil to protect it from daylight. While maintaining the  $\text{N}_2$  atmosphere inside the flask,  $\text{Me}_4\text{Sn}$  (1.34 mL, 9.67 mmol, 15 equiv.) was added in one portion *via* a syringe.  $\text{CsF}$  (0.2448 g, 1.6 mmol, 2.5 equiv.),  $\text{CuI}$  (30.7 mg, 0.16 mmol, 0.25 equiv.), and  $(\text{Ph}_3\text{P})_4\text{Pd}$  (37.2 mg, 0.032 mmol, 0.05 equiv.) were then sequentially and successively added to the reaction flask in that order, maintaining

the  $\text{N}_2$  atmosphere throughout the various additions. The reaction mixture was then left to stir at rt for 22 h, whereafter TLC analysis revealed that two C5-diastereomeric products **19** had formed that had exactly the same TLC mobility (in 5 : 1 hexane : EtOAc) as the starting iodovinylic alcohol **18**. Staining and heating of this TLC plate with the anisaldehyde/ $\text{H}_2\text{SO}_4$  stain did, however, show that the two newly formed products of structure **19** both stained with a purple/blue colour, which allowed them to be readily distinguished from their precursor **18**; this revealed that the reaction was complete, and that no starting **18** remained. The reaction mixture was thereupon diluted with EtOAc (10 mL), transferred to a separatory funnel and  $\text{H}_2\text{O}$  (15 mL) was added. After extraction and separation, the aqueous later was extracted with additional EtOAc (6  $\times$  20 mL), and the combined organic extracts were washed with  $\text{H}_2\text{O}$  (2  $\times$  50 mL). The organic layer was then dried over  $\text{MgSO}_4$ , filtered, and concentrated *in vacuo*. The resulting oil was then purified by gradient elution  $\text{SiO}_2$  flash chromatography with petrol : EtOAc 20 : 1  $\rightarrow$  15 : 1 initially, to remove reagent-related impurities, followed by petrol : EtOAc 12 : 1  $\rightarrow$  10 : 1 to secure the entire allylic alcohol **19** (0.189 g, 80%) product as an oil, and as a C5-mixture of epimers.

Data for pure **19**:  $^1\text{H}$  NMR of **19** (pure isomer 1) (600.13 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.09 (d,  $J$  = 8.4 Hz, 2H, H7), 6.66 (d,  $J$  = 7.8 Hz, 2H, H8), 5.26 (ddd,  $J$  = 9.6 Hz, 1H, H3), 4.35 (d,  $J$  = 6.0 Hz, 1H, H5), 3.98 (s, 2H, H1a,b), 2.93 (s, 6H,  $\text{N}(\text{CH}_3)_2$ ), 2.616 (m, 1H, H4), 1.557 (d,  $J$  = 1.2 Hz, C2-Me), 1.25 (br s, 1H, OH), 0.85 (s, 9H, *t*-Bu of OTBS), 0.82 (d,  $J$  = 6.6 Hz, 3H C4-Me), -0.02 (s, 3H,  $\text{CH}_3\text{Si}$ ), -0.22 (s, 3H,  $\text{CH}_3\text{Si}$ ) ppm.  $^{13}\text{C}$  NMR of **19** (pure isomer 1) (150.92 MHz,  $\text{CDCl}_3$ ):  $\delta$  C9 not detected, 134.5 (C3), 129.9 (C6), 127.6 (C7 and C2), 111.9 (C8), 78.9 (C5), 69.4 (C1), 41.1 ( $\text{N}(\text{CH}_3)_2$ ), 40.7 (C4), 25.8 (( $\text{CH}_3$ )<sub>3</sub>CSi), 18.2 (( $\text{CH}_3$ )<sub>3</sub>CSi), 17.0 (C2-Me), 13.9 (C4-Me), -4.5 ( $\text{CH}_3\text{Si}$ ), -5.0 ( $\text{CH}_3\text{Si}$ ) ppm.

$^1\text{H}$  NMR of **19** (pure isomer 2) (600.13 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.09 (m, 2H, H7), 6.64 (m, 2H, H8), 5.23 (ddd,  $J$  = 10.2, 3.0, 1.2 Hz, 1H, H3), 4.35 (d,  $J$  = 6.0 Hz, 1H, H5), 3.89 (s, 2H, H1a,b), 2.92 (s, 6H,  $\text{N}(\text{CH}_3)_2$ ), 2.59 (m,  $J$  = 9.6, 6.0 Hz, 1H, H4), 1.464 (d,  $J$  = 1.2 Hz, C2-Me), 1.25 (br s, 1H, OH), 0.96 (d,  $J$  = 7.2 Hz, C4-Me), 0.85 (s, 9H, *t*-Bu of OTBS), -0.013 (s, 3H,  $\text{CH}_3\text{Si}$ ), -0.21 (s, 3H,  $\text{CH}_3\text{Si}$ ) ppm.  $^{13}\text{C}$  NMR of **19** (pure isomer 2) (150.92 MHz,  $\text{CDCl}_3$ ):  $\delta$  149.4 (C9), 134.2 (C3), 130.0 (C6), 127.5 (C7 and C2), 111.7 (C8), 78.7 (C5), 69.3 (C1), 41.2 ( $\text{N}(\text{CH}_3)_2$ ), 40.7 (C4), 25.9 (( $\text{CH}_3$ )<sub>3</sub>CSi), 18.3 (( $\text{CH}_3$ )<sub>3</sub>CSi), 16.0 (C2-Me), 13.8 (C4-Me), -4.5 ( $\text{CH}_3\text{Si}$ ), -5.1 ( $\text{CH}_3\text{Si}$ ) ppm.

### TPAP/NMO oxidation<sup>19</sup> of iodo-allylic alcohol **18** to iodoenal **20**



Before commencing this experiment, the reaction flask was covered with Al-foil to protect it from the adverse effects of visible light. Thereafter, to a stirred rt solution of the iodovinylic alcohol **18** (3.55 g, 7.47 mmol) in dry  $\text{CH}_2\text{Cl}_2$  (74.7 mL) under  $\text{N}_2$  was successively added powdered 4  $\text{\AA}$  molecular sieves (flame dried,



3.60 g), *N*-methylmorpholine-*N*-oxide (NMO, 1.75 g, 14.93 mmol, 2 equiv.) and tetrapropylammonium perruthenate<sup>19</sup> (TPAP, *n*-Pr<sub>4</sub>NRuO<sub>4</sub>, 0.13 g, 0.37 mmol, 0.05 equiv.), each in one portion. Stirring was continued at rt for 15 min. The reaction mixture was then filtered to remove the sieves, the filtrate was concentrated *in vacuo*, and the resulting crude residue was purified by SiO<sub>2</sub> flash chromatography with 40 : 1 petrol : EtOAc as the eluent. Following purification, aldehyde **20** (2.10 g, 59%) was obtained as a C5-epimeric mixture (*ca.* 3 : 1) as an amber oil.


Data for **20** (*ca.* 3 : 1 mixture of isomers of tentatively assigned stereochemistry at C5): IR of the **20** mixture (neat film): 3393 (w), 2959 (s), 2929 (s), 2856 (s), 2806 (w), 1701 (s), 1615 (s), 1524 (s), 1461 (m), 1388 (s), 1355 (m), 1251 (m), 1186 (w), 1168 (m), 1080 (s), 1027 (m), 1007 (m), 944 (w), 858 (m), 838 (m), 780 (m) cm<sup>-1</sup>.

<sup>1</sup>H NMR of **20** (minor isomer 1 – *anti*) (399.9 MHz, CDCl<sub>3</sub>):  $\delta$  8.64 (s, 1H, H1), 7.10 (d, *J* = 8.4 Hz, 2H, H7), 7.09 (d, *J* = 9.2 Hz, 1H, H3), 6.66 (d, *J* = 8.4 Hz, 2H, H8), 4.61 (d, 1H, *J* = 6.0 Hz, H5), 3.18 (m, 1H, H4), 2.94 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 1.02 (d, *J* = 6.8 Hz, 3H, C4-Me), 0.86 (s, 9H, *t*-Bu of OTBS), 0.028 (s, 3H, CH<sub>3</sub>Si), –0.20 (s, 3H, CH<sub>3</sub>Si) ppm. <sup>13</sup>C NMR of **20** (minor isomer 1 – *anti*) (100.57 MHz, CDCl<sub>3</sub>):  $\delta$  188.1 (C1), 165.3 (C3), 150.0 (C9), 130.1 (C6), 127.33 (C7), 112.0 (C8), 111.3 (C2), 77.6 (C5), 50.2 (C4), 40.5 (N(CH<sub>3</sub>)<sub>2</sub>), 25.79 ((CH<sub>3</sub>)<sub>3</sub>CSi), 18.1 ((CH<sub>3</sub>)<sub>3</sub>CSi), 15.2 (C4-Me), –4.5 (CH<sub>3</sub>Si), –5.1 (CH<sub>3</sub>Si) ppm.

<sup>1</sup>H NMR of **20** (major isomer 2 – *syn*) (399.9 MHz, CDCl<sub>3</sub>):  $\delta$  8.59 (s, 1H, H1), 7.16 (d, *J* = 8.4 Hz, 2H, H7), 7.05 (d, *J* = 9.6 Hz, 1H, H1), 6.68 (d, *J* = 8.4 Hz, 2H, H8), 4.73 (d, 1H, *J* = 4.4 Hz, H5), 3.12 (m, 1H, H4), 2.95 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 1.07 (d, *J* = 6.8 Hz, 3H, C4-Me), 0.90 (s, 9H, *t*-Bu of OTBS), –0.012 (s, 3H, CH<sub>3</sub>Si), –0.17 (s, 3H, CH<sub>3</sub>Si) ppm. <sup>13</sup>C NMR of **20** (major isomer 2) (100.57 MHz, CDCl<sub>3</sub>):  $\delta$  188.09 (C1), 165.3 (C3), 149.9 (C9), 129.8 (C6), 127.29 (C7), 111.9 (C8), 110.4 (C2), 76.4 (C5), 49.7 (C4), 40.5 (N(CH<sub>3</sub>)<sub>2</sub>), 25.84 ((CH<sub>3</sub>)<sub>3</sub>CSi), 18.2 ((CH<sub>3</sub>)<sub>3</sub>CSi), 13.3 (C4-Me), –4.5 (CH<sub>3</sub>Si), –5.1 (CH<sub>3</sub>Si) ppm.

TOF ES<sup>+</sup> HRMS of **20**: calcd for C<sub>20</sub>H<sub>33</sub>NO<sub>2</sub>Si [M + H]<sup>+</sup>: 474.1325. Found: 474.1317.

### Baldwin–Lee Stille cross-coupling of iodo enal **20** to obtain trisubstituted enal **4**



A solution of the iodoenal **20** (0.98 g, 2.06 mmol) in dry DMF (4.12 mL) was transferred *via* a cannula into a Teflon-screw-capped sealed tube temporarily fitted with a rubber septum under N<sub>2</sub>. Me<sub>4</sub>Sn (4.29 mL, 30.90 mmol, 15 equiv.), CsF (0.78 g, 5.15 mmol, 2.5 equiv.), CuI (0.10 g, 0.52 mmol, 0.25 equiv.), and (Ph<sub>3</sub>P)<sub>4</sub>Pd (0.12 g, 0.10 mmol, 0.05 equiv.) were then successively added in that order, and the septum was replaced with the Teflon-screw cap, maintaining the N<sub>2</sub> atmosphere throughout the addition and sealing process. The sealed tube was then covered with Al-foil, placed inside an oil


bath, and heated at 60 °C overnight (18 h) with vigorous magnetic stirring. The reaction mixture was then cooled to rt, carefully opened, and diluted with EtOAc (5 mL) before being fully quenched with H<sub>2</sub>O (10 mL). The sealed tube was rinsed multiple times with EtOAc (20 mL × 5) and H<sub>2</sub>O (20 mL × 3) and the combined reaction washings were filtered through Celite, before being fractionated in a separatory funnel. The aqueous layer was further extracted with EtOAc (50 mL × 3), and the combined organic extracts were washed with H<sub>2</sub>O (100 mL × 2), dried over MgSO<sub>4</sub>, filtered and concentrated under reduced pressure. The crude residue was purified by SiO<sub>2</sub> flash chromatography using 45 : 1 petrol : EtOAc as an eluent to afford **4** (0.59 g, 79%) as a mixture of C5-epimers and as a colourless oil. IR of the **4** mixture (neat film): 3353 (w), 2959 (s), 2932 (s), 2858 (s), 2808 (m), 2707 (w), 1690 (s), 1617 (s), 1524 (s), 1466 (m), 1385 (m), 1355 (m), 1257 (m), 1186 (w), 1072 (m), 1022 (m), 949 (w), 863 (m), 835 (m), 777 (m), 671 (w), 565 (w) cm<sup>-1</sup>.

Data for **4**: <sup>1</sup>H NMR of **4** (*ca.* 3 : 1 mixture of isomers at C5) (isomer 1 – major) (399.9 MHz, CDCl<sub>3</sub>):  $\delta$  9.40 (s, 1H, H1), 7.09 (d, *J* = 8.8 Hz, 2H, H7), 6.65 (d, *J* = 8.8 Hz, 2H, H8), 6.43 (dd, *J* = 10.0 Hz, 1.2 Hz, 1H, H3), 4.47 (d, 1H, *J* = 6.0 Hz, H5), 2.93 (br m, 1H, H4) superimposed upon 2.93 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 1.62 (d, *J* = 1.2 Hz, 3H, C2-Me), 0.95 (d, *J* = 6.8 Hz, 3H, C4-Me), 0.84 (s, 9H, *t*-Bu of OTBS), –0.011 (s, 3H, CH<sub>3</sub>Si), –0.23 (s, 3H, CH<sub>3</sub>Si) ppm. <sup>13</sup>C NMR of **4** (isomer 1 – major) (100.57 MHz, CDCl<sub>3</sub>):  $\delta$  195.6 (C1), 158.0 (C3), 149.9 (C9), 139.1 (C2), 130.9 (C6), 127.3 (C7), 112.0 (C8), 78.6 (C5), 42.9 (C4), 40.55 (N(CH<sub>3</sub>)<sub>2</sub>), 25.75 ((CH<sub>3</sub>)<sub>3</sub>CSi), 18.1 ((CH<sub>3</sub>)<sub>3</sub>CSi), 16.5 (C4-Me), 9.31 (C2-Me), –4.5 (CH<sub>3</sub>Si), –5.1 (CH<sub>3</sub>Si) ppm.

<sup>1</sup>H NMR of **4** (isomer 2 – minor) (399.9 MHz, CDCl<sub>3</sub>):  $\delta$  9.31 (s, 1H, H1), 7.08 (d, *J* = 8.8 Hz, 2H, H7), 6.63 (d, *J* = 8.7 Hz, 2H, H8), 6.30 (dd, *J* = 10.0 Hz, 1.2 Hz, 1H, H3), 4.50 (d, 1H, *J* = 5.6 Hz, H5), 2.93 (br m, 1H, H4) superimposed upon 2.93 (s, 6H, N(CH<sub>3</sub>)<sub>2</sub>), 1.61 (d, *J* = 1.6 Hz, 3H, C2-Me), 1.07 (d, *J* = 6.4 Hz, 3H, C4-Me), 0.89 (s, 9H, *t*-Bu of OTBS), –0.04 (s, 3H, CH<sub>3</sub>Si), –0.20 (s, 3H, CH<sub>3</sub>Si) ppm. <sup>13</sup>C NMR of **4** (isomer 2 – minor) (100.57 MHz, CDCl<sub>3</sub>):  $\delta$  195.6 (C1), 157.4 (C3), 149.8 (C9), 138.5 (C2), 130.6 (C6), 127.3 (C7), 111.8 (C8), 77.8 (C5), 42.6 (C4), 40.53 (N(CH<sub>3</sub>)<sub>2</sub>), 25.83 ((CH<sub>3</sub>)<sub>3</sub>CSi), 18.2 ((CH<sub>3</sub>)<sub>3</sub>CSi), 15.3 (C4-Me), 9.3 (C2-Me), –4.5 (CH<sub>3</sub>Si), –5.1 (CH<sub>3</sub>Si) ppm.

TOF ES<sup>+</sup> HRMS of **4**: calcd for C<sub>21</sub>H<sub>36</sub>NO<sub>2</sub>Si [M + H]<sup>+</sup>: 362.2515. Found: 362.2517.

### Wittig olefination of enal **4** to obtain dienoate **3**



To a stirred rt solution of the enal **4** (0.5777 g, 1.60 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (1.6 mL) and PhMe (2 mL) under N<sub>2</sub> was added carbethoxymethylenetriphenylphosphorane (0.835 g, 2.40 mmol, 1.5 equiv.) and the reactants were stirred at 38 °C for 10 d. The reaction mixture was thereafter evaporated to dryness before petrol (5 mL) was added and the resulting mixture was stirred for 2 h. The reaction mixture was filtered


to remove  $\text{Ph}_3\text{P}=\text{O}$  and concentrated under reduced pressure. The crude residue was purified by  $\text{SiO}_2$  flash chromatography using petrol :  $\text{EtOAc}$  (50 : 1) as the eluent to give **3** (0.62 g, 90%) as a colourless oil and mixture of C7-epimers (N.B.: (+)-(R)-trichostatin A numbering is now being used here onwards). IR of the **3** mixture (neat film): 2959 (m), 2934 (m), 2856 (m), 1721 (m), 1620 (s), 1521 (m), 1458 (m), 1388 (s), 1310 (w), 1259 (w), 1168 (m), 1097 (w), 1070 (w), 1029 (w), 871 (w), 780 (w)  $\text{cm}^{-1}$ .

$^1\text{H}$  NMR of **3** (isomer 1 – minor) (399.9 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.32 (dd,  $J$  = 15.6, 0.4 Hz, 1H, H2), 7.07 (d,  $J$  = 8.8 Hz, 2H, H9), 6.64 (d,  $J$  = 8.8 Hz, 2H, H10), 5.79 (d, 1H,  $J$  = 10.0 Hz, H5), 5.75 (d,  $J$  = 15.2 Hz, 1H, H3), 4.39 (d,  $J$  = 6.0 Hz, 1H, H7), 4.20 (m,  $J$  = 7.2 Hz, 2H,  $\text{CH}_2$  of OEt), 2.93 (s, 6H,  $\text{N}(\text{CH}_3)_2$ ), 2.76 (m, 1H, H6), 1.64 (d,  $J$  = 1.2 Hz, 3H, C4-Me), 1.30 (t,  $J$  = 7.2 Hz, 3H, Me of OEt), 0.87 (d, C6-Me) 0.83 (s, 9H,  $t\text{-Bu}$  of OTBS), –0.03 (s, 3H,  $\text{CH}_3\text{Si}$ ), –0.23 (s, 3H,  $\text{CH}_3\text{Si}$ ) ppm.  $^{13}\text{C}$  NMR of **3** (isomer 1 – minor) (100.57 MHz,  $\text{CDCl}_3$ ):  $\delta$  167.8 (C1), 150.0 (C2), 149.8 (C11), 145.4 (C5), 132.6 (C4), 131.6 (C8), 127.5 (C9), 115.4 (C3), 111.94 (C10), 78.8 (C7), 60.1 ( $\text{CH}_2$  of OEt), 42.4 (C6), 40.6 ( $\text{N}(\text{CH}_3)_2$ ), 25.8 ( $(\text{CH}_3)_3\text{CSi}$ ), 18.1 ( $(\text{CH}_3)_3\text{CSi}$ ), 16.9 (C6-Me), 14.3 (Me of OEt), 12.4 (C4-Me), –4.6 ( $\text{CH}_3\text{Si}$ ), –5.1 ( $\text{CH}_3\text{Si}$ ) ppm.

$^1\text{H}$  NMR of **3** (isomer 2 – major) (399.9 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.23 (d,  $J$  = 15.6, 0.4 Hz, 1H, H2), 7.06 (d,  $J$  = 8.8 Hz, 2H, H9), 6.63 (d,  $J$  = 8.4 Hz, 2H, H10), 5.71 (d,  $J$  = 15.6 Hz, 1H, H3), 5.70 (d, 1H,  $J$  = 10.0 Hz, H5), 4.41 (d,  $J$  = 6.0 Hz, 1H, H7), 4.20 (m,  $J$  = 7.2 Hz, 2H,  $\text{CH}_2$  of OEt), 2.92 (s, 6H,  $\text{N}(\text{CH}_3)_2$ ), 2.76 (m, 1H, H6), 1.60 (d,  $J$  = 1.2 Hz, 3H, C4-Me), 1.29 (t,  $J$  = 7.2 Hz, 3H, Me of OEt), 1.00 (d,  $J$  = 6.8 Hz, 3H, C6-Me), 0.88 (s, 9H,  $t\text{-Bu}$  of OTBS), –0.015 (s, 3H,  $\text{CH}_3\text{Si}$ ), –0.22 (s, 3H,  $\text{CH}_3\text{Si}$ ) ppm.  $^{13}\text{C}$  NMR of **3** (isomer 2 – major) (100.57 MHz,  $\text{CDCl}_3$ ):  $\delta$  167.7 (C1), 149.9 (C2), 149.7 (C11), 145.0 (C5), 132.1 (C4), 131.4 (C8), 127.4 (C9), 115.5 (C3), 111.9 (C10), 78.3 (C7), 60.1 ( $\text{CH}_2$  of OEt), 42.4 (C6), 40.6 ( $\text{N}(\text{CH}_3)_2$ ), 25.9 ( $(\text{CH}_3)_3\text{CSi}$ ), 18.2 ( $(\text{CH}_3)_3\text{CSi}$ ), 15.9 (C6-Me), 14.3 (Me of OEt), 12.3 (C4-Me), –4.5 ( $\text{CH}_3\text{Si}$ ), –5.1 ( $\text{CH}_3\text{Si}$ ) ppm.

TOF ES<sup>+</sup> HRMS of **3**: calcd for  $\text{C}_{25}\text{H}_{42}\text{NO}_3\text{Si}$  [M + H]<sup>+</sup>: 432.2934. Found: 432.2942.

### Preparation of 7-OTBS trichostatic acid **21**



To a stirred rt solution of ester **3** (1.06 g, 2.46 mmol) in  $\text{THF}/\text{H}_2\text{O}/\text{MeOH}$  (2 mL : 2 mL : 2 mL) was added  $\text{LiOH}$  monohydrate (0.41 g, 9.82 mmol, 4 equiv.) and the mixture was thereafter allowed to stir vigorously for 48 h. The reaction mixture was then diluted with  $\text{EtOAc}$  (10 mL) and acidified with 10% aq.  $\text{HCl}$  until pH 5 was attained. The aqueous layer was extracted with  $\text{EtOAc}$  (20 mL  $\times$  3). The combined organic layer was washed with  $\text{H}_2\text{O}$  (50 mL  $\times$  2), dried over  $\text{MgSO}_4$ , filtered and concentrated under reduced pressure. The crude mixture was purified by gradient elution  $\text{SiO}_2$  flash chromatography with 6 : 1  $\rightarrow$  4 : 1 petrol :  $\text{EtOAc}$  as an eluent to obtain **21** (0.97 g, 98%) as a colourless oil.


Data for **21** (for a 3 : 1 mixture of C7-epimers): IR of the **21** mixture (neat film): 3376 (extremely br), 2959 (s), 2932 (s), 2858 (m), 2806 (w), 1688 (s), 1617 (s), 1524 (s), 1468 (m), 1383 (m), 1282 (m), 1254 (m), 1209 (w), 1072 (s), 1029 (m), 987 (w), 871 (s), 777 (m)  $\text{cm}^{-1}$ .

$^1\text{H}$  NMR of **21** (isomer 1 – minor) (399.9 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.58 (very br, 1H,  $-\text{CO}_2\text{H}$ ), 7.41 (d,  $J$  = 15.6 Hz, 1H, H2), 7.08 (d, 2H, H9), 6.65 (d, 2H, H10), 5.86 (d, 1H,  $J$  = 9.6 Hz, H5), 5.72 (d,  $J$  = 15.6 Hz, 1H, H3), 4.40 (d,  $J$  = 6.0 Hz, 1H, H7), 2.93 (s, 6H,  $\text{N}(\text{CH}_3)_2$ ), 2.78 (m, 1H, H6), 1.66 (s, 3H, C4-Me), 0.89 (d, 3H, C6-Me obscured by a large singlet for *t*-Bu for the major isomer of **21**), 0.84 (s, 9H, *t*-Bu of OTBS), –0.02 (s, 3H,  $\text{CH}_3\text{Si}$ ), –0.23 (s, 3H,  $\text{CH}_3\text{Si}$ ) ppm.  $^{13}\text{C}$  NMR of **21** (isomer 1 – minor) (100.57 MHz,  $\text{CDCl}_3$ ):  $\delta$  172.9 (C1), 152.3 (C2), 149.8 (C11), 146.8 (C5), 132.7 (C4), 131.6 (C8), 127.5 (C9), 114.5 (C3), 112.0 (C10), 78.8 (C7), 42.6 (C6), 40.7 ( $\text{N}(\text{CH}_3)_2$ ), 25.8 ( $(\text{CH}_3)_3\text{CSi}$ ), 18.1 ( $(\text{CH}_3)_3\text{CSi}$ ), 16.9 (C6-Me), 12.4 (C4-Me), –4.6 (C4-Me), –5.1 (C4-Me) ppm.

$^1\text{H}$  NMR of **21** (isomer 2 – major) (399.9 MHz,  $\text{CDCl}_3$ ):  $\delta$  10.58 (very br, 1H,  $-\text{CO}_2\text{H}$ ), 7.32 (d,  $J$  = 15.6 Hz, 1H, H2), 7.07 (d,  $J$  = 8.4 Hz, 2H, H9), 6.65 (d,  $J$  = 8.4 Hz, 2H, H10), 5.76 (d, 1H,  $J$  = 10.4 Hz, H5), 5.71 (d,  $J$  = 15.6 Hz, 1H, H3), 4.43 (d,  $J$  = 6.0 Hz, 1H, H7), 2.93 (s, 6H,  $\text{N}(\text{CH}_3)_2$ ), 2.78 (m, 1H, H6), 1.62 (s, 3H, C4-Me), 1.01 (d,  $J$  = 6.8 Hz, 3H, C6-Me), 0.89 (s, 9H,  $\text{SiC}(\text{CH}_3)_3$ ), –0.01 (s, 3H,  $\text{CH}_3\text{Si}$ ), –0.21 (s, 3H,  $\text{CH}_3\text{Si}$ ) ppm.  $^{13}\text{C}$  NMR of **21** (isomer 2 – major) (100.57 MHz,  $\text{CDCl}_3$ ):  $\delta$  172.8 (C=O), 152.2 (C2), 149.7 (C11), 146.4 (C5), 132.2 (C4), 131.4 (C8), 127.4 (C9), 114.7 (C3), 112.0 (C10), 78.2 (C7), 42.5 (C6), 40.7 ( $\text{N}(\text{CH}_3)_2$ ), 25.9 ( $(\text{CH}_3)_3\text{CSi}$ ), 18.2 ( $(\text{CH}_3)_3\text{CSi}$ ), 15.8 (C6-Me), 12.3 (C4-Me), –4.5 (C4-Me), –5.1 (C4-Me) ppm.

TOF ES<sup>+</sup> HRMS of **21**: calcd for  $\text{C}_{23}\text{H}_{38}\text{NO}_3\text{Si}$  [M + H]<sup>+</sup>: 404.2621. Found: 404.2613.

### Conversion of 7-OTBS trichostatic acid **21** into (+)-(R)-trichostatic acid (**22**)



To a stirred 0 °C solution of 7-OTBS trichostatic acid **21** (0.61 g, 1.51 mmol) in  $\text{CH}_2\text{Cl}_2/\text{H}_2\text{O}$  (20 : 1) (9.88 mL; made up from 9.41 mL  $\text{CH}_2\text{Cl}_2$ : 0.47 mL  $\text{H}_2\text{O}$ ) was added 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ, 0.51 g, 2.26 mmol, 1.5 equiv.) in one portion and the cold reaction mixture was vigorously stirred for just 7 min (*important note: stirring for longer than this time caused significant decomposition*). The reaction mixture was then diluted with  $\text{CH}_2\text{Cl}_2$  (10 mL) and quenched with  $\text{H}_2\text{O}$  (10 mL). The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  (50 mL  $\times$  3). The combined organic extracts were washed with  $\text{H}_2\text{O}$  multiple times (50 mL  $\times$  8), dried over  $\text{MgSO}_4$ , filtered and concentrated under reduced pressure. The crude mixture was purified by gradient elution  $\text{SiO}_2$  flash chromatography with petrol :  $\text{EtOAc}$  (1 : 1  $\rightarrow$  1 : 2) as an eluent with **21** (0.31 g, 71%) being obtained as an amber oil.

Data for **22**:  $[\alpha]_D$  = +150.7° (c 0.416,  $\text{CH}_2\text{Cl}_2$ ), mp 89–90 °C [literature mp<sup>17</sup> 88–89 °C].




IR of **22** (neat film): 3600–2300 (extremely br), 2980 (m), 2932 (m), 2874 (m), 2669 (w), 2596 (w), 1685 (s), 1597 (s), 1552 (m), 1438 (m), 1380 (s), 1272 (m), 1244 (m), 1191 (m), 1171 (m), 1004 (w), 979 (w), 858 (w), 828 (w), 737 (m)  $\text{cm}^{-1}$ .

$^1\text{H}$  NMR of pure **22** (400.11 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.84 (d,  $J$  = 9.2 Hz, 2H, H9), 7.36 (d,  $J$  = 15.6 Hz, 1H, H2), 6.64 (d,  $J$  = 9.2 Hz, 2H, H10), 6.06 (d,  $J$  = 9.6 Hz, 1H, H5), 5.85 (d, 1H,  $J$  = 15.6 Hz, H3), 4.38 (m, 1H, H6), 3.05 (s, 6H,  $\text{N}(\text{CH}_3)_2$ ), 1.92 (d,  $J$  = 1.2 Hz, 3H, C4-Me), 1.31 (d,  $J$  = 6.8 Hz, 3H, C6-Me) ppm.  $^{13}\text{C}$  NMR of **22** (100.57 MHz,  $\text{CDCl}_3$ ):  $\delta$  198.3 (C7), 171.7 (C1), 153.5 (C11), 151.4 (C2), 143.0 (C5), 132.6 (C4), 130.6 (C9), 123.9 (C8), 115.7 (C3), 110.8 (C10), 40.9 (C6), 40.0 ( $\text{N}(\text{CH}_3)_2$ ), 17.7 (C6-Me), 12.5 (C4-Me) ppm.

TOF ES<sup>+</sup> HRMS of **22**: calcd for  $\text{C}_{17}\text{H}_{22}\text{NO}_3$  [ $\text{M} + \text{H}$ ]<sup>+</sup>: 288.1600. Found: 288.1584.

### Conversion of (+)-(6*R*)-trichostatic acid (22) into (+)-(6*R*)-trichostatin A (1a)



Following Helquist's procedure,<sup>17e</sup> a stirred  $0^\circ\text{C}$  solution of (+)-(6*R*)-trichostatic acid **22** (56.2 mg, 0.195 mmol) in dry  $\text{THF}$  (1 mL) under  $\text{N}_2$  was treated with freshly distilled dry  $\text{Et}_3\text{N}$  (0.03 mL, 0.23 mmol, 1.2 equiv.) followed by dropwise addition of ethyl chloroformate (0.04 mL, 0.40 mmol, 2 equiv.). The reactants were allowed to stir at  $0^\circ\text{C}$  for 2 h, whereafter a solution of  $\text{TBSONH}_2$  (57.6 mg, 0.39 mmol, 1.95 equiv.) in dry  $\text{THF}$  (0.5 mL) was added *via* a microcannula. The reaction mixture was stirred at  $0^\circ\text{C}$  for 0.5 h and then warmed to rt for 1.5 h. It was then diluted with  $\text{EtOAc}$  (1 mL) and quenched with  $\text{H}_2\text{O}$  (1 mL). The aqueous layer was extracted with  $\text{EtOAc}$  (10 mL  $\times$  3). The organic layer was washed with  $\text{H}_2\text{O}$  (10 mL  $\times$  2), dried over  $\text{MgSO}_4$ , filtered and concentrated under reduced pressure and the crude residue of **24** was used directly without further purification.

To a stirred rt solution of the aforementioned TBS-protected crude trichostatin A **24** in dry  $\text{MeOH}$  (2 mL) under  $\text{N}_2$  was added solid  $\text{CsF}$  (35.6 mg, 0.23 mmol, 1.2 equiv.) (pre-dried at  $120^\circ\text{C}$  under high vacuum for 2 h), and the reactants were allowed to stir for 1 h. The reaction mixture was then diluted with  $\text{CH}_2\text{Cl}_2$  (5 mL) and washed with  $\text{H}_2\text{O}$  (2 mL). The aqueous layer was then further extracted with  $\text{CH}_2\text{Cl}_2$  (10 mL  $\times$  5). The combined organic extracts were washed with  $\text{H}_2\text{O}$  (20 mL  $\times$  1), dried over  $\text{MgSO}_4$ , filtered and concentrated under reduced pressure. The crude mixture was purified by gradient elution  $\text{SiO}_2$  flash chromatography with  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  (25 : 1  $\rightarrow$  20 : 1) to obtain (+)-(R)-trichostatin A (**1a**) (35 mg, 59% over 2 steps) as a white solid.

Data for (*R*)-(+)-trichostatin A (**1a**):  $[\alpha]_D$  = +88.8° (*c* 0.26,  $\text{MeOH}$ ), mp 140–143 °C [lit.<sup>17</sup> 140–143 °C]. IR of (+)-TSA (**1a**) (neat film): 3234 (br, s), 2927 (s), 2853 (s), 1650 (m), 1595 (s), 1547 (m), 1383 (s), 1249 (m), 1189 (s), 1059 (m), 976 (m)  $\text{cm}^{-1}$ .

$^1\text{H}$  NMR of (+)-TSA (**1a**) (600.13 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  7.87 (d,  $J$  = 9.0 Hz, 2H, H9), 7.18 (d,  $J$  = 15.0 Hz, 1H, H2), 6.72 (d,  $J$  = 9.6 Hz, 2H, H10), 5.91 (d,  $J$  = 9.6 Hz, 1H, H5), 5.87 (d, 1H,  $J$  = 15.6 Hz, H3), 4.53 (dq,  $J$  = 9.0, 6.6 Hz, 1H, H6), 3.06 (s, 6H,  $\text{N}(\text{CH}_3)_2$ ), 1.92 (s, 3H, C4-Me), 1.27 (d,  $J$  = 6.6 Hz, 3H, C6-Me) ppm.

$^{13}\text{C}$  NMR of (+)-TSA (**1a**) (150.92 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  201.4 (C7), 166.8 (C1), 155.5 (C11), 145.9 (C2), 141.3 (C5), 134.3 (C8), 131.9 (C9), 124.7 (C4), 117.1 (C3), 111.9 (C10), 41.7 (C6), 40.1 ( $\text{N}(\text{CH}_3)_2$ ), 18.3 (C6-Me), 12.7 (C4-Me) ppm.

TOF ES<sup>+</sup> HRMS of (+)-TSA (**1a**): [ $\text{M} + \text{H}$ ]<sup>+</sup> calcd for  $\text{C}_{17}\text{H}_{23}\text{N}_2\text{O}_3$ : 303.171. Found: 303.177.

### Data availability

All data supporting this article are included in the ESI.<sup>†</sup>

### Conflicts of interest

There are no conflicts to declare.

### Acknowledgements

We thank the ACS and QUB for financial support of this work.

### References

- Review: K. J. Hale, S. Manaviazar and H. A. Watson, *Chem. Rec.*, 2019, **19**, 238.
- (a) P. Dimopoulos, A. Athlan, S. Manaviazar, J. George, M. Walters, L. Lazarides, A. Aliev and K. J. Hale, *Org. Lett.*, 2005, **7**, 5369; (b) R. Willem, A. Delmotte, I. De Borger, I. Biesemans, M. Gielen and F. Kayser, *J. Organomet. Chem.*, 1994, **480**, 255; (c) P. Dussault, C. T. Eary, R. J. Lee and U. R. Zope, *J. Chem. Soc., Perkin Trans. 1*, 1999, 2189.
- Mechanistic studies: (a) P. Dimopoulos, J. George, D. A. Tocher, S. Manaviazar and K. J. Hale, *Org. Lett.*, 2005, **7**, 5377; (b) H. A. Watson, S. Manaviazar, H. G. Steeds and K. J. Hale, *Chem. Commun.*, 2019, **55**, 14454; (c) H. A. Watson, S. Manaviazar, H. G. Steeds and K. J. Hale, *Tetrahedron*, 2020, **76**, 131061; (d) H. A. Watson, A. J. Fielding and K. J. Hale, *Chem. Commun.*, 2021, **57**, 7449.
- (a) E. J. Corey and R. H. Wollenberg, *J. Org. Chem.*, 1975, **40**, 2265; (b) H. E. Ensley, R. R. Buescher and K. Lee, *J. Org. Chem.*, 1982, **47**, 404; (c) C. Nativi and M. Taddei, *J. Org. Chem.*, 1988, **58**, 820; (d) M. Lautens and A. H. Hudson, *Tetrahedron Lett.*, 1990, **31**, 3105.



5 (a) K. Pati, G. dos Passos Gomes, T. Harris, A. Hughes, H. Phan, T. Banerjee, K. Hanson and I. V. Alabugin, *J. Am. Chem. Soc.*, 2015, **137**, 1165; (b) N. P. Tsvetkov, E. Gonzales-Rodriguez, A. Hughes, G. dos Passos Gomes, F. D. White, F. Kuriakose and I. V. Alabugin, *Angew. Chem., Int. Ed.*, 2018, **57**, 3651; (c) E. Gonzalez-Rodriguez, M. A. Abdo, G. dos Passos Gomes, S. Ayad, F. D. White, N. P. Tsvetkov, K. Hanson and I. V. Alabugin, *J. Am. Chem. Soc.*, 2020, **142**, 8352; (d) C. Hu, L. Kuhn, F. D. Makurvet, E. S. Knorr, X. Lin, R. K. Kawade, F. Mentink-Vigier, K. Hanson and I. V. Alabugin, *J. Am. Chem. Soc.*, 2024, **146**, 4187; (e) I. V. Alabugin, G. dos Passos Gomes and M. A. Abdo, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2019, **9**, e1389; (f) I. V. Alabugin, K. M. Gilmore and P. W. Peterson, *Wiley Interdiscip. Rev.: Comput. Mol. Sci.*, 2011, **1**, 109; (g) I. V. Alabugin, *Stereoelectronic Effects: A Bridge Between Structure and Reactivity*, John Wiley & Sons Ltd, Chichester, UK, 2016, ch. 3, p. 42; (h) C. Hu, J. Mena and I. V. Alabugin, *Nat. Rev. Chem.*, 2023, **7**, 405.

6 S. Manaviazar, K. J. Hale and A. LeFranc, *Tetrahedron Lett.*, 2011, **52**, 2080.

7 K. J. Hale, M. Grabski, S. Manaviazar and M. Maczka, *Org. Lett.*, 2014, **16**, 1164.

8 K. J. Hale, M. Maczka, A. Kaur, S. Manaviazar, M. Ostovar and M. Grabski, *Org. Lett.*, 2014, **16**, 1168.

9 K. Micoine, P. Persich, J. Llaveria, M.-H. Lam, A. Marderna, F. Loganzo and A. Furstner, *Chem. – Eur. J.*, 2013, **19**, 7370.

10 P. Dimopoulos, A. Athlan, S. Manaviazar and K. J. Hale, *Org. Lett.*, 2005, **7**, 5373.

11 (a) S. P. H. Mee, V. Lee and J. E. Baldwin, *Angew. Chem., Int. Ed.*, 2004, **43**, 1132; (b) V. Lee, *Org. Biomol. Chem.*, 2019, **17**, 9095.

12 V. Farina, S. Kapadia, B. Krishnan, C. Wang and L. S. Liebeskind, *J. Org. Chem.*, 1994, **59**, 5909.

13 J. A. Marshall and S. Xie, *J. Org. Chem.*, 1995, **60**, 723.

14 (a) N. Tsuji, M. Kobayashi, K. Nagashima, Y. Wakisawa and K. Koizumi, *J. Antibiot.*, 1976, **29**, 1; (b) D. M. Vigushin, S. Ali, N. Mirsaldi, K. Ito, I. Adcock and R. C. Coombes, *Clin. Cancer Res.*, 2001, **7**, 971; (c) L. Sanderson, G. W. Taylor, E. O. Aboagye, J. P. Alao, J. R. Latigo, R. C. Coombes and D. M. Vigushin, *Drug Metab. Dispos.*, 2004, **32**, 1132; (d) A. M. Grabiec, S. Krausz, W. de Jager, T. Burakowski, D. Groot, M. E. Sanders, B. J. Prakken, W. Maslinski, E. Eldering, P. P. Tak and K. A. Reedquist, *J. Immunol.*, 2010, **184**, 2718.

15 C. Mozzetta, S. Consalvi, V. Saccone, M. Tierney, A. Diamantini, K. J. Mitchell, G. Marazzi, G. Borsellino, L. Battistini, D. Sasoon, A. Sacco and P. L. Puri, *EMBO Mol. Med.*, 2013, **5**, 626.

16 C. Wang, L. Lei, Y. Xu, Y. Li, J. Zhang, Y. Xu and S. Si, *Pharmaceuticals*, 2024, **17**, 425.

17 For previous trichostatin A total syntheses, see: (a) I. Fleming, J. Iqbal and E.-P. Krebs, *Tetrahedron*, 1983, **39**, 841; (b) K. Mori and K. Koseki, *Tetrahedron*, 1988, **44**, 6013; (c) S. Zhang, W. Duan and W. Wang, *Adv. Synth. Catal.*, 2006, **348**, 1228; (d) C. C. Cosner and P. Helquist, *Org. Lett.*, 2011, **13**, 3564; (e) C. C. Cosner, V. B. R. Iska, A. Chatterjee, J. T. Markiewicz, S. J. Corden, J. Lofstedt, T. Anker, J. Richer, T. Hulett, D. J. Schauer, O. Wiest and P. Helquist, *Eur. J. Org. Chem.*, 2013, 162; (f) Trichostatin D total synthesis: S. Hosokawa, T. Ogura, H. Togashi and K. Tatsuta, *Tetrahedron Lett.*, 2005, **46**, 333; (g) Synthesis of trichostatic acid (22): J. T. Markiewicz, D. J. Schauer, J. Lofstedt, S. J. Corden, O. Wiest and P. Helquist, *J. Org. Chem.*, 2010, **75**, 2061.

18 For a related allenyl-Pd(II) intermediate also being prevented from undergoing transmetallation by an  $\alpha$ -allylic OPv, see: J. A. Marshall and J. J. Mulhearn, *Org. Lett.*, 2005, **7**, 1593.

19 (a) W. P. Griffith, S. V. Ley, G. P. Whitcombe and A. D. White, *J. Chem. Soc., Chem. Commun.*, 1987, **21**, 1625; (b) S. V. Ley, J. Norman, W. P. Griffith and S. P. Marsden, *Synthesis*, 1994, 639; (c) For a previous application of the Ley and Griffith cat. TPAP/NMO oxidation system in the total synthesis of the macrolide immunosuppressants (+)-prunostatin A and (+)-SW-163A, see: S. Manaviazar, P. Nockemann and K. J. Hale, *Org. Lett.*, 2016, **18**, 2902.

20 (a) N. Tsuji and M. Kobayashi, *J. Antibiot.*, 1978, **31**, 939; (b) R. R. Hughes, K. A. Shaaban, J. Zhang, H. Cao, G. N. Phillips and J. S. Thorsen, *ChemBioChem*, 2017, **18**, 363; (c) The details of our synthesis of (+)-trichostatin C from (+)-trichostatic acid (22) will be reported in due course.

21 A reviewer of our paper has requested that we briefly compare our new synthetic route with the other total syntheses of trichostatin A that are listed in ref. 17. We will duly do this here. (a) The first total synthesis of  $(\pm)$ -trichostatin A to be achieved was that of Fleming and coworkers in 1983. It provided the natural product in racemic form in 5 steps (ref. 17a), but according to Mori and Koseki (ref. 17b), its final KOH/NH<sub>2</sub>OH-mediated ester N-hydroxyammonolysis step cannot be used to secure biologically-active (+)-trichostatin A; (b) Although Mori and Koseki's later synthesis of (+)-trichostatin A (**1a**) (ref. 17b) did deliver the natural product in  $\geq 98\%$  ee from *(R)*-*(–)*-3-hydroxy-2-methyl-propionate, it did require 18 steps to be implemented overall, and it had a longest linear sequence of 16 steps. While Mori's synthesis was perfectly stereocontrolled, with respect to installation of the *(6R)*-stereocentre and the C(2)-C(5)-dienoate array, it did encounter low yields during its final stages; its penultimate steps 14 and 15 proceeded with a combined yield of 11%; (c) Helquist later published three synthetic routes to  $(\pm)$ -trichostatin A (ref. 17d, e and g), one of which (ref. 17e) was subsequently rendered enantioselective. The latter route required 17 steps to be implemented overall, when the more reliable and higher yielding 7-step *(S)*-ethyl lactate pathway was used to access its key *(S)*-3-butyn-2-yl O-mesylate starting material. Importantly, Helquist's asymmetric route to **1a** had a longest linear sequence of 10 steps, and its yields were largely good throughout. It was also fully stereocontrolled with regard to installation of the C(6)-Me group and



the C(2)–C(5)-dienyl array. It did, however, produce (+)-(R)-trichostatin A (**1a**) in only 81% ee; (*d*) Contrastingly, our new *O*-directed hydrostannylation route to (+)-trichostatin A proceeds in 18 steps overall (which is one step more than Helquist's), and it has a longest linear sequence of 12 steps. It also proceeds with a maximal overall yield of 8%. However, most critically, it does provide (+)-(R)-trichostatin A in  $\geq 98\%$  ee, as evidenced by our conversion of **1a** to (+)-trichostatin C, without the accompanying formation of the C(6)-(S)- $\beta$ -glycoside diastereoisomer. Importantly, our prior derivatisation of **15** to obtain **25** (see ESI) also confirmed that **15** was of  $\geq 98\%$  ee; (*e*) Although Wang's 2006 report (ref. 17c) did describe a 10 step L-proline-catalysed aldol route to (+)-(R)-trichostatin A, which delivered a material of  $\geq 99\%$  ee in an apparently good overall yield (17.4%), its final benzylic alcohol oxidation step had to be conducted with just 0.59 equiv. of DDQ in dioxane. Presumably this was done to minimise or prevent competing *N*-oxidation of the hydroxamic acid unit to give a highly reactive *N*-acyl-nitroso intermediate, which would almost certainly self-condense, if generated. We note here that the ref. 17c team were unable to recover any of their unreacted starting hydroxamic acid amide precursor from this DDQ

oxidation, which proceeded with 49% yield for these last two steps. Contrastingly, each of the other enantioselective routes to (+)-trichostatin A have utilised an *O*-alkoxy/siloxamine coupling strategy with a mixed anhydride derived from (+)-trichostatic acid to install the hydroxamic acid motif of the natural product.

22 (*a*) For a recent outstanding review on alkyne hydrometallation with Group IV metal hydrides, see the following book chapter by: T. Wiesner and M. Haas, *Reference Module in Chemistry, Molecular Sciences and Chemical Engineering*, Elsevier, 2024, DOI: [10.1016/B978-0-323-96025-0.00125-3](https://doi.org/10.1016/B978-0-323-96025-0.00125-3) (*b*) For McLaughlin and Roberts' 2023 report on the highly regiocontrolled PtCl<sub>2</sub>/XPhos-catalysed hydrostannation of terminal aryl acetylenes and propargylic alcohols, see: D. D. Roberts and M. G. McLaughlin, *Adv. Synth. Catal.*, 2023, **365**, 1602. This paper lists much valuable new metal-catalysed hydrostannation literature that has recently appeared; (*c*) For McLaughlin's landmark application of the PtCl<sub>2</sub>/XPhos/Et<sub>3</sub>SiH-catalyst system to mediate an analogous highly regiocontrolled hydroboration of terminal alkyl, aryl and heteroaryl acetylenes with HBPin, see: K. L. E. Hale, D. D. Roberts and M. G. McLaughlin, *Eur. J. Org. Chem.*, 2025, e202401355.

