

COMMUNICATION

View Article Online
View Journal | View Issue



Cite this: *Org. Biomol. Chem.*, 2025, **23**, 2802

Received 15th January 2025,
Accepted 10th February 2025

DOI: 10.1039/d5ob00070j

rsc.li/obc

Stereospecific access to α -haloalkyl esters *via* enol ester epoxides and synthesis of a C3–C21 fragment of bastimolide A†

Lucas W. Howell, Jacob N. Hackbarth, Jeffrey S. Abraham and
Gregory K. Friestad *

We report a 14-step synthesis of a C3–C21 fragment of bastimolides A and B, antimalarial macrocyclic polyketides. A crucial ring-opening reaction of an enol ester epoxide showed previously unexplored reactivity, leading to an asymmetric synthesis of α -haloalkyl esters. The α -haloalkyl ester synthesis was shown to be stereospecific, and provided access to a key α -silyloxyaldehyde to initiate application of configuration-encoded 1,5-polyol synthesis. This strategy established the C11/C15 and C15/C19 remote stereochemical relationships of the bastimolides. The potential of this C3–C21 fragment for coupling to C22–C41 was established using a Mukaiyama aldol reaction with a simple enolsilane.

Introduction

Malaria continues to have a significant global impact causing an estimated 249 million malaria cases in 2022,¹ and the threat of resistance emphasizes the continuing need for new treatment options. Marine cyanobacteria offer structurally interesting natural products that offer insights for drug discovery.² Among these are bastimolide A (**1**),³ a 40-membered macrolide that has proven to exhibit potent antimalarial properties against four resistant strains of *Plasmodium falciparum* (IC_{50} = 80–270 nM), and its 24-membered macrolactone isomer, bastimolide B (**2**).⁴ Both 1,3- and 1,5-polyol motifs are combined in its structure (Fig. 1), which bears little apparent resemblance to any clinical antimalarial drugs or preclinical candidates.⁵ This suggests the possibility that biological evaluation of **1**, **2**, or analogs could uncover a new mode of action *via* a novel *Plasmodium* drug target.

Synthesis of the bastimolides is therefore a high priority, and various strategies to access structural subunits⁶ and halogenated analogues⁷ have appeared. Smith⁸ and Aggarwal⁹ have

reported the first total syntheses of **1** and **2**, respectively, and Kirsch *et al.* reported a formal synthesis.¹⁰ We recently reported asymmetric synthesis of a 1,5-polyol comprising the C22–C41 fragment of the bastimolides.¹¹

Polyols bearing 1,5-relationships between hydroxyl groups often cause complications for configurational assignments,¹² stereocontrolled synthesis, and diastereomer separations.¹³ This prompted our development of a *configuration-encoded* synthetic strategy¹⁴ (Fig. 2) utilizing building blocks of defined hydroxyl configuration that are linked iteratively *via* Julia–Kocienski olefination.¹⁵ Subjecting α -silyloxyaldehydes to Julia–Kocienski olefination with γ -sulfononitrile building blocks (*R*)-**3** or (*S*)-**3** establishes *syn*- or *anti*-1,5-diol relationships, and subsequent reduction of the nitrile regenerates α -silyloxyaldehyde functionality at the chain terminus for another iteration. The programmed assembly allows synthesis of all possible diastereomers of 1,5-polyols with equal facility, and obviates analytical or preparative separations of diastereomers. Here we disclose the configuration-encoded synthesis of the C3–C21 subunit of the bastimolides (**4**), aided by the discovery of mild conditions for stereospecific transformation of enol ester epoxides into 1-haloalkyl esters.

Results and discussion

Our retrosynthetic analysis (Fig. 2) involves two iterations of the configuration-encoded 1,5-polyol synthesis strategy. This




Fig. 1 Structures of bastimolides A (**1**) and B (**2**).

Department of Chemistry, University of Iowa, Iowa City, Iowa 52242, USA.

E-mail: gregory-friestad@uiowa.edu

† Electronic supplementary information (ESI) available. CCDC 2403648. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d5ob00070j>







Fig. 2 Two iterations of configuration-encoded 1,5-polyol synthesis to access an *anti,syn*-1,5,9 triol stereotriad.

suggesting α -silyloxyaldehyde **5** as a key precursor, which would be coupled successively with (*S*)-**3** and (*R*)-**3** to unambiguously establish C15 and C19 configurations. Our earlier successes in three-step synthesis of α -silyloxyaldehydes from alkynes *via* enol ester epoxides¹⁶ prompted a similar approach to **5** from alkyne **6**.

The synthetic sequence began with preparation of epoxide **8**¹⁷ (Scheme 1), obtained in 92% ee by enantioselective Katsuki epoxidation of alkene **7** with the Berkessel Ti catalyst **A**.¹⁸ Reaction of **8** with lithiated trimethylsilylacetylene, followed by alkyne desilylation and TBS protection of the 2° alcohol, furnished alkyne **6** in 72% yield over five steps, with only two column chromatography purifications.



Scheme 1 Synthesis of enol ester epoxide **10** and its anomalous reactivity.

The next task was to implement our three-step route from alkynes to α -silyloxyaldehydes. Ruthenium-catalyzed addition¹⁹ of anisic acid to alkyne **6** (Scheme 1) gave (*Z*)-enol ester **9** in 90% yield, and another Berkessel–Katsuki epoxidation furnished enol ester epoxide **10** with excellent yield and selectivity (94%, 96:4 dr). The epoxide ring-opening, which had normally provided smooth access to α -silyloxyaldehydes upon treatment with silyl triflates and lutidine,^{11,14c,16} failed in this case; only traces of the aldehyde **5** were observed. In an effort to understand this anomaly, simplified enol ester substrate **11a** (Fig. 3a) was subjected to the ring opening. Instead of α -silyloxyaldehyde **12a**, a dimeric hydroxyfuran structure **13** was obtained. This could be rationalized by an oxocarbenium ion (or its equivalent) undergoing nucleophilic attack by the nearby silyloxy substituent as implied by structure **C**, followed by some combination of silyl transfer events and dimerization. The dimerization finds precedent in a similar structure formed from furanoses.²⁰ Replacement of OTBS with less nucleophilic OBz would be expected to suppress the formation of dimer **13**, and indeed with substrate **11b** the expected ring-opening pathway to the α -silyloxyaldehyde **12b** was restored (64% yield).¹¹

Further comment about unexpected structure **13** is warranted. The *cis* configuration at both ring junctions of **13** was assigned by its apparent C2 symmetry (4 signals in its ¹³C NMR spectrum) and small coupling constants at the ring junction (*J* = 3.7 Hz observed at the anomeric C–H). A boatlike central ring would be accompanied by high torsional strain and lack of anomeric stabilization; we propose a chairlike conformation for the central ring of **13**, with pseudo-C2 symmetry attributed to a rapid chair–chair conformational equilibrium.




Fig. 3 (a) Simplified analogs **11a** and **11b** reveal the reason for anomalous reactivity of **10**. (b) Stereospecific enol ester epoxide ring-opening to 1-haloalkyl esters **15**.




Although changing to ester protection at the C9-OH could circumvent the anomalous reactivity of **10** and **11a**, we sought to address the problem head-on, with the hypothesis that a large concentration of an external nucleophile could suppress the offending cyclization. Indeed, upon treatment of enol ester epoxide **14a** with LiCl and TsOH, complete conversion to 1-haloalkyl ester **15a** was observed within 5 min. Similar results were observed with enol ester epoxides **14b**, **14c**, **11a**, and **11b**, producing 1-haloalkyl esters **15b–15e** respectively. All of these 1-haloalkyl esters were formed with complete stereospecificity despite the potential intermediacy of a planar oxocarbenium ion. Enol ester epoxide *trans*-**14a**, prepared *via* the corresponding *E*-enol ester, gave diastereomer *epi*-**15a**, providing further evidence of the stereospecific nature of the ring-opening. Crystallographic analysis of the 3,5-dinitrobenzoate derivative of **15b** (see ESI†) confirmed the structural assignment suggesting inversion of configuration by the chloride nucleophile. To our knowledge there is only one related example leading from an enol ester epoxide to a 1-haloalkyl ester (SnCl₄, 50% yield).²¹ Here we provide for the first time (a) evidence of stereospecificity, (b) preliminary evaluation of scope, and (c) subsequent reactivity studies in application to target-oriented synthesis. Racemic 1-haloalkyl esters²² have been used in various bond constructions,²³ and this access to enantiopure samples presents new opportunities for reaction development and mechanistic study.


Turning our attention back to the synthetic route, we put the new method to the test in conversion of enol ester epoxide **10** to α -silyloxyaldehyde **5**. Treatment of **10** with a chloride source (NBu₄Cl was used for improved solubility) and camphorsulfonic acid, followed by *O*-silylation, cleanly furnished the α -haloalkyl ester **16** in 84.5% yield (Scheme 2). Transesterification with concomitant elimination of chloride then completed the alternative method for conversion of **10** to aldehyde **5**. While the typical K₂CO₃/MeOH was effective for this transesterification on microscale, scale-up was proble-

matic. A more reliable modification employed phase-transfer catalysis: in an immiscible mixture of benzene (or toluene) and ethylene glycol, exposure to K₂CO₃ along with phase-transfer catalyst NBu₄HSO₄ furnished aldehyde **5** in 74% yield, along with recovered **10** (88% yield based on conversion).²⁴

With aldehyde **5** in hand, we applied our configuration-encoded 1,5-polyol synthesis. Julia–Kocienski olefination with (*S*)-**3** smoothly established the *syn*-1,5-diol relationship between C11 and C15 with a quantitative yield of nitrile **17**. Reduction of the nitrile with DIBAL-H and a second iteration of the Julia–Kocienski reaction with alternative building block **18** then established the desired *anti*,*syn*-1,5,9-triol stereotriad in **19**, while placing masked aldehyde functionality at the terminus.²⁵ Sulfone **18** was prepared in 6 steps from acrolein (see ESI†), using Keck allylation (95% ee) to encode the desired configuration at the carbon destined to become C19 of the bastimolides. Hydrogenation of the alkenes of **19** afforded acetal **20**, which was converted to β -silyloxyaldehyde **4** through Fujioka–Kita acetal hydrolysis.²⁶ This aldehyde is the C3–C21 fragment of bastimolides, ready for Mukaiyama aldol coupling to the C22–C41 subunit.



An initial assessment of Mukaiyama aldol conditions for coupling to the C3–C21 subunit (**4**) employed the trimethylsilyl enol ether derived from pinacolone (eqn (1)). In the presence of BF₃·OEt₂, this enolsilane added smoothly to aldehyde **4** to afford aldol **21** (dr 88:12) with close correspondence to the 1,3-diastereocontrol we previously observed in similar polyacetate-type aldol adducts.²⁷ Diagnostic ¹H NMR data for several closely related aldol products in that prior report enabled



Scheme 2 Access to α -silyloxyaldehyde **5** and configuration-encoded assembly of the C3–C21 subunit of bastimolides.



assignment of *anti* configuration to the major diastereomer **21** and demonstrated the potential of **4** as a viable intermediate *en route* to bastimolides and analogs.

Conclusions

We have developed a synthesis of the C3–C21 fragment of bastimolides using the configuration-encoded approach to 1,5-polyol assembly. The synthetic sequence encountered an unexpected structural incompatibility of our previously established three-step conversion of alkynes to α -silyloxyaldehydes. Solving this problem led to the discovery of a stereospecific synthesis of enantiopure 1-haloalkyl esters; these are richly functionalized synthetic building blocks with further synthetic potential.

Data availability

Crystallographic data are available in CCDC 2403648. Preparative procedures and characterization data are provided in ESI.†

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

With great appreciation, this article is dedicated to the memory of Professor Amos B. Smith, III. We thank the National Institutes of Health (AI-171379) for generously supporting this research. We gratefully acknowledge seed funding and fellowship support (LWH) from the University of Iowa Department of Chemistry and College of Liberal Arts and Sciences. We thank Prof. James Gloer and Dr Chris Knutson of the University of Iowa for assistance with HPLC instrumentation, and the University of Iowa NMR (G. Crull) and MATFab (D. Unruh) facilities for analytical support.

References

- World Health Organization (WHO), World Malaria Report 2023. <https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2023>, accessed 10/15/24.
- L. T. Tan, Bioactive natural products from marine cyanobacteria for drug discovery, *Phytochemistry*, 2007, **68**, 954–979.
- C.-L. Shao, R. G. Linington, M. J. Balunas, A. Centeno, P. Boudreau, C. Zhang, N. Engene, C. Spadafora, T. S. Mutka, D. E. Kyle, L. Gerwick, C.-Y. Wang and W. H. Gerwick, Bastimolide A, a Potent Antimalarial Polyhydroxy Macrolide from the Marine Cyanobacterium *Okeania hirsuta*, *J. Org. Chem.*, 2015, **80**, 7849–7855.
- C.-L. Shao, X.-F. Mou, F. Cao, C. Spadafora, E. Glukhov, L. Gerwick, C.-Y. Wang and W. H. Gerwick, Bastimolide B, an Antimalarial 24-Membered Marine Macrolide Possessing a tert-Butyl Group, *J. Nat. Prod.*, 2018, **81**, 211–215.
- (a) T. Umumarungu, J. B. Nkuranga, G. Habarurema, J. B. Nyandwi, M. J. Mukazayire, J. Mukiza, R. Muganga, I. Hahirwa, M. Mpenda, A. N. Katembezi, E. O. Olawode, E. Kayitare and P. C. Kayumba, Recent developments in antimalarial drug discovery, *Bioorg. Med. Chem.*, 2023, **88–89**, 117339; (b) M. A. Phillips, J. N. Burrows, C. Manyando, R. H. van Huijsduijnen, W. C. Van Voorhis and T. N. C. Wells, Malaria, *Nat. Rev. Dis. Primers*, 2017, **3**, 17050.
- N. S. Kumar, B. J. Ramulu and S. Ghosh, Stereoselective Synthesis of the C19–C39 Fragment of Bastimolide A, *SynOpen*, 2021, **5**, 285–290.
- A. Quintard, C. Sperandio and J. Rodriguez, Modular Enantioselective Synthesis of an Advanced Pentahydroxy Intermediate of Antimalarial Bastimolide A and of Fluorinated and Chlorinated Analogues, *Org. Lett.*, 2018, **20**, 5274–5277.
- J. B. Cox, A. A. Kellum, Y. Zhang, B. Li and A. B. Smith, Total Synthesis of (–)-Bastimolide A: A Showcase for Type I Anion Relay Chemistry, *Angew. Chem., Int. Ed.*, 2022, e202204884.
- D. Fiorito, S. Keskin, J. M. Bateman, M. George, A. Noble and V. K. Aggarwal, Stereocontrolled Total Synthesis of Bastimolide B Using Iterative Homologation of Boronic Esters, *J. Am. Chem. Soc.*, 2022, **144**, 7995–8001.
- I.-E. Celik, F. Mittendorf, A. Gomez-Suarez and S. F. Kirsch, Formal synthesis of bastimolide A using a chiral Horner-Wittig reagent and a bifunctional aldehyde as key building blocks, *Tetrahedron Chem*, 2024, **9**, 100065.
- J. N. Hackbarth and G. K. Friestad, A Three-Step Catalytic Asymmetric Sequence from Alkynes to α -Silyloxyaldehydes and Its Application to a C22–C41 Fragment of Bastimolide A, *Org. Lett.*, 2024, **25**, 4492–4496.
- Assignment of the stereostructure of bastimolide A required X-ray diffraction analysis of the corresponding nona-*p*-nitrobenzoate derivative (see ref. 3).
- Review: R. M. Friedrich and G. K. Friestad, Inspirations from Tetrafibricin and Related Polyketides: New Methods and Strategies for 1,5-Polyol Synthesis, *Nat. Prod. Rep.*, 2020, **37**, 1229–1261.
- (a) G. K. Friestad and G. Sreenilayam, Versatile configuration-encoded strategy for rapid synthesis of 1,5-polyol stereoisomers, *Org. Lett.*, 2010, **12**, 5016–5019; (b) R. M. Friedrich and G. K. Friestad, Access to an anti, syn-1,5,7-Triol by Configuration-Encoded 1,5-Polyol Synthesis: The C15–C25 Fragment of Tetrafibricin, *Eur. J. Org. Chem.*, 2017, 1961–1964; (c) R. M. Friedrich, G. Sreenilayam, J. Hackbarth and G. K. Friestad, Unified Strategy for 1,5,9- and 1,5,7-Triols via Configuration-Encoded 1,5-Polyol Synthesis: Enantioselective Preparation of γ -Sulfonyl- α -silyloxyaldehydes and Iterative Julia-



- Kocienski Coupling, *J. Org. Chem.*, 2018, **83**, 13636–13649; (d) R. M. Friedrich, J. Q. Bell, A. Garcia, Z. Shen and G. K. Friestad, Unified Strategy for 1,5,9- and 1,5,7-Triols via Configuration-Encoded 1,5-Polyol Synthesis: Preparation and Coupling of C15–C25 and C26–C40 Fragments of Tetrafibricin, *J. Org. Chem.*, 2018, **83**, 13650–13669.
- 15 (a) Use of Julia–Kocienski reactions to establish 1,5-diol relationships has also been reported for late-stage fragment couplings (see ref. 13). Meanwhile, our iterative strategy is uniquely positioned for flexible access to various analogs including stereoisomers (b) See also. L. C. Dias and E. C. de Lucca, Total Synthesis of (–)-Marinisorolide C, *J. Org. Chem.*, 2017, **82**, 3019–3045.
 - 16 G. K. Friestad, G. Sreenilayam, J. C. Cannistra and L. M. Slominski, Preparation of enol ester epoxides and their ring-opening to α -silyloxyaldehydes, *Tetrahedron Lett.*, 2012, **53**, 5064–5067. Also see ref. 11 and 14c.
 - 17 J. E. Robinson and M. A. Brimble, The first enantioselective total synthesis of the anti-*Helicobacter pylori* agent (+)-spiroxaline methyl ether, *Chem. Commun.*, 2005, 1560–1562.
 - 18 (a) K. Matsumoto, C. Feng, S. Handa, T. Oguma and T. Katsuki, Asymmetric epoxidation of (Z)-enol esters catalyzed by titanium(salalen) complex with aqueous hydrogen peroxide, *Tetrahedron*, 2011, **67**, 6474–6478; (b) A. Berkessel, T. Gunther, Q. Wang and J.-M. Neudörfl, Titanium Salalen Catalysts Based on cis-1,2-Diaminocyclohexane: Enantioselective Epoxidation of Terminal Non-Conjugated Olefins with H₂O₂, *Angew. Chem., Int. Ed.*, 2013, **52**, 8467–8471; (c) H. Engler, M. Lansing, C. P. Gordon, J.-M. Neudörfl, M. Schäfer, N. E. Schlörner, C. Copéret and A. Berkessel, Olefin Epoxidation Catalyzed by Titanium–Salalen Complexes: Synergistic H₂O₂ Activation by Dinuclear Ti Sites, Ligand H-Bonding, and π -Acidity, *ACS Catal.*, 2021, **11**, 3206–3217.
 - 19 H. Doucet, B. Martin-Vaca, C. Bruneau and P. H. Dixneuf, General Synthesis of (Z)-Alk-1-en-1-yl Esters via Ruthenium-Catalyzed anti-Markovnikov trans-Addition of Carboxylic Acids to Terminal Alkynes, *J. Org. Chem.*, 1995, **60**, 7247–7255.
 - 20 (a) S. Mukherjee, B. G. Roy, S. N. Das and S. B. Mandal, Deoxygenation/dimerization of sugar derivatives with BF₃·Et₂O–Et₃SiH: synthesis of a β -isonucleoside, *Tetrahedron Lett.*, 2012, **53**, 4929–4932; (b) L. Di, K.-J. Wang, C.-C. Zhu and N. Li, Chemical Constituents from Rhizomes of *Curculigo capitulata*, *Bull. Korean Chem. Soc.*, 2010, **31**, 2999–3002.
 - 21 B. Peschke, J. Lussmann, M. Dyrbusch and D. Hoppe, Diastereo- and Enantioselective Synthesis of 1,2:5,6-Diepoxy- δ -hydroxyalkyl Carbamates. Regioselective Ring Opening and Their Transformation into Doubly C-Branched Deoxy Sugar Analogues, *Chem. Ber.*, 1992, **125**, 1421–1430.
 - 22 For preparations of racemic 1-haloalkyl esters, see: F. Fache, I. de Azpiazu, B. Pelotier, O. Piva and C. Gozzi, Green Access to α -Haloalkyl and α -Halobenzyl Esters, Versatile Intermediates for the One-Pot Two-Step Synthesis of O,O'-Diacyl Acetals Using Zinc-Based Ionic Liquid Catalyst, *Synthesis*, 2019, 2430–2434, references therein.
 - 23 Selected examples: (a) K. Sakata, D. Urabe and M. Inoue, Preparation and palladium-mediated cross-coupling of α -benzyloxyalkylzinc bromides, *Tetrahedron Lett.*, 2013, **54**, 4189–4192; (b) L. Wang, J. M. Lear, S. M. Rafferty, S. C. Fosu and D. A. Nagib, Ketyl radical reactivity via atom transfer catalysis, *Science*, 2018, **362**, 225–229; (c) Z.-P. Yang and G. C. Fu, Convergent Catalytic Asymmetric Synthesis of Esters of Chiral Dialkyl Carbinols, *J. Am. Chem. Soc.*, 2020, **142**, 5870–5875; (d) S. M. Rafferty, J. E. Rutherford, L. Zhang, L. Wang and D. A. Nagib, Cross-Selective Aza-Pinacol Coupling via Atom Transfer Catalysis, *J. Am. Chem. Soc.*, 2021, **143**, 5622–5628; (e) H. Ji, D. Lin, L. Tai, X. Li, Y. Shi, Q. Han and L.-A. Chen, Nickel-Catalyzed Enantioselective Coupling of Acid Chlorides with α -Bromobenzoates: An Asymmetric Acyloin Synthesis, *J. Am. Chem. Soc.*, 2022, **144**, 23019–23029; (f) C. Li, J. Cheng, X. Wan, J. Li, W. Zu, Y. Xu, Y. Huang and H. Huo, Ni/Photoredox-Catalyzed Enantioselective Acylation of α -Bromobenzoates with Aldehydes: A Formal Approach to Aldehyde–Aldehyde Cross-Coupling, *J. Am. Chem. Soc.*, 2024, **146**, 19909–19918.
 - 24 Best results were obtained at incomplete conversion; longer reaction times led to partial decomposition of **5**.
 - 25 Coupling with (R)-**3** was also promising, but the modified building block **18** enabled more direct access to the C21 aldehyde terminus needed for fragment coupling.
 - 26 (a) H. Fujioka, T. Okitsu, Y. Sawama, N. Murata, R. Li and Y. Kita, Reaction of the Acetals with TESOTf–Base Combination; Speculation of the Intermediates and Efficient Mixed Acetal Formation, *J. Am. Chem. Soc.*, 2006, **128**, 5930–5938.
 - 27 L. W. Howell and G. K. Friestad, On the Role of β -Silyloxy- and β -Alkoxyaldehyde Protecting Groups in Mukaiyama Aldol 1,3-Diastereocontrol, *Tetrahedron Lett.*, 2022, **111**, 154203.

