



Cite this: *Org. Biomol. Chem.*, 2025, **23**, 3443

## Efficient chemical synthesis of mirror-image DNA polymerase Dpo4 assisted by one-pot multi-segment condensation†

Miao Wang,<sup>‡a</sup> Tingting Cui,<sup>‡c</sup> Shuqing Huang,<sup>a</sup> Dongyang Han,<sup>b</sup> Xiangyu Deng,<sup>b</sup> Yanbo Liu<sup>id</sup><sup>b</sup> and Chaowei Shi<sup>id</sup><sup>\*a</sup>

The mirror-image DNA polymerase Dpo4 is a valuable enzymatic tool in biomedical research; however, its chemical synthesis has been hindered by low efficiency. Here, we describe an efficient chemical synthesis of D-Dpo4 using a one-pot multi-segment condensation strategy, which was achieved using Fmoc-masked peptide thioester. By minimizing the need for isolating and handling intermediates through one-pot three- and four-segment ligations, we were able to obtain D-Dpo4 with higher efficiency, achieving an improved overall yield of 9.8%, representing a twofold improvement from previous work. Our work provides a practical and streamlined approach for the synthesis of mirror-image proteins with larger sizes and underscores the utility of one-pot multi-segment condensation in chemical protein synthesis.

Received 11th February 2025,  
Accepted 6th March 2025

DOI: 10.1039/d5ob00243e

rsc.li/obc

### Introduction

Mirror-image proteins (D-proteins), composed entirely of D-amino acids and achiral glycine, have emerged as valuable tools in biomedical research due to their unique biochemical properties.<sup>1–3</sup> These unique biochemical properties make mirror-image proteins useful biological tools with applications in drug discovery, protein crystallography, and the development of mirror-image biochemical systems.<sup>4–7</sup> The original studies on mirror-image proteins began with Kent *et al.*<sup>8</sup> who pioneered the chemical synthesis of mirror image enzymes to develop and explore biochemical systems based on biomolecules of opposite chirality. Unlike natural L-proteins, these chirally inverted D-enantiomers can only be obtained by chemical synthesis.<sup>3,8–21</sup> While recent advances in chemical protein synthesis have enabled the routine production of moderately sized D-proteins (typically 100–200 residues),<sup>4,10,21–39</sup> the preparation of larger D-proteins (*e.g.*, those containing more than 300 amino acids) still often suffers from low efficiency.<sup>40,41</sup> A representative example is the 352-residue D-enantiomer of *Sulfolobus solfataricus* P2 DNA polymerase IV

(D-Dpo4),<sup>42</sup> which can be used as a tool to develop therapeutically promising nuclease-resistant L-nucleic acid aptamers through mirror-image systematic evolution of ligand by exponential enrichment (miSELEX) technology.<sup>43</sup> In a recent study, D-Dpo4 was assembled from nine peptide segments, requiring more than 15 steps of intermediate purification during its preparation.<sup>44</sup> These tedious and time-consuming purification procedures heavily complicated the synthetic workflows and compromised the overall efficiency (with a low yield of less than 5%), limiting the practical accessibility of this molecular tool.

In this context, here we report the study on the use of a one-pot multi-segment condensation strategy to facilitate the chemical synthesis of D-Dpo4. Through the strategic implementation of one-pot three-segment and one-pot four-segment ligations to minimize the need for isolation and handling of intermediates, D-Dpo4 was obtained with an improved overall yield of 9.8%, which is 2-fold higher than that of the previous study.<sup>44</sup> Our work provides a practical synthetic route for obtaining a useful mirror-image protein tool for biomedical research and underlines the utility of one-pot multi-segment condensation in streamlining the chemical synthesis of D proteins with larger sizes.

### Results and discussion

The D-Dpo4 protein (352 amino acids) with an N-terminal His<sub>6</sub>-tag was strategically divided into nine peptide segments (segments 1–9), each spanning 20 to 50 residues (Fig. 1A). To

<sup>a</sup>Hefei National Research Center for Interdisciplinary Sciences at the Microscale, University of Science and Technology of China, 230027 Hefei, China.

E-mail: scwei@ustc.edu.cn

<sup>b</sup>Department of Chemistry, Tsinghua University, Beijing 100084, China

<sup>c</sup>School of Food and Biological Engineering, Engineering Research Center of Bioprocess, Ministry of Education, Hefei University of Technology, 230009 Hefei, China

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d5ob00243e>

‡ These authors contributed equally to this work.







Fig. 1 (A) The amino acid sequence of His<sub>6</sub>-D-Dpo4. The full-length protein is divided into nine segments and ligation sites are highlighted in bold. (B) Synthetic route of His<sub>6</sub>-D-Dpo4.

enable native chemical ligation (NCL), seven cysteine residues (42, 123, 155, 207, 229, 268, and 313) were introduced as temporary ligation sites (which were later reverted to alanine after ligation), with a subsequent plan to revert these positions to alanine post ligation.<sup>45–48</sup> A His<sub>6</sub> tag was incorporated at the N-terminus of D-Dpo4 to facilitate final protein purification (Fig. 1A).

We synthesized all segments (1–9) using standard Fmoc (9-fluorenylmethoxycarbonyl)-based solid-phase peptide synthesis (SPPS), purified by reversed-phase high-performance

liquid chromatography (RP-HPLC) and verified by electrospray ionization mass spectrometry (ESI-MS) analysis (Fig. 2). To facilitate the one-pot C-to-N sequential ligation strategy, segments 4, 5, 7, and 8 were designed with C-terminal aryl thioester precursors to facilitate one pot ligation.<sup>49,50</sup> The N-terminal cysteine of segment 5 was protected with the Fmoc group for preventing cyclization,<sup>51</sup> while segments 7 and 8 similarly carried Fmoc-protected N-terminal cysteines to enable sequential ligation with segment 9. Additionally, orthogonal protection was implemented for Cys86 and Cys123





**Fig. 2** Analytical RP-HPLC chromatogram ( $\lambda = 214$  nm) and ESI-MS spectra of segments 1–9. (A) Segment 1. (B) Segment 2. (C) Segment 3. (D) Segment 4. (E) Segment 5. (F) Segment 6. (G) Segment 7, "\*" corresponds to the trifluoroacetate of 7. (H) Segment 8. (I) Segment 9.

using acetamidomethyl (Acm) groups,<sup>52</sup> ensuring compatibility with the Fmoc-based ligation strategy (Fig. 1B).


We have successfully achieved the efficient synthesis of the 105-amino-acid fragment **14** by adopting a one-pot ligation–Fmoc deprotection–thioester transfer strategy (Fig. 3A). Initially, three functional modules were meticulously prepared through Fmoc-SPPS: a 32-amino-acid (AA) fragment **4** (Cys(Acm)123-Ala154-COSC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>COOH) with a C-terminal thioester precursor, a 51-AA thioesterified fragment **5** (Fmoc-Cys155-Leu206-COSC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>COOH) with the N-terminal Fmoc protection intact, and a 22-AA fragment **6**. Considering the relatively hydrophobic nature of its sequence, Acm was selected as the cysteine protecting group in fragment **4** to minimize its impact on solubility. Notably, the N-terminal Fmoc group on Cys155 of fragment **5** was retained as a removable protecting group for the terminal cysteine, which effectively suppressed the thioester-induced intramolecular cyclization of fragment **5**. Subsequently, an acetylaceton-mediated activation method was employed to directly transform the crude peptides of fragments **4** and **5** into MPAA thioesters (COSC<sub>6</sub>H<sub>4</sub>CH<sub>2</sub>COOH).<sup>49</sup> This approach remarkably reduced the separation and purification steps in precursor preparation, thus facilitating a more streamlined one-pot ligation process.

With all three peptides in hand, we commenced the process by conducting native chemical ligation (NCL) between fragment **5** (1.0 equiv.) and fragment **6** (1.0 equiv.) (Fig. 3B). The ligation reaction was carried out in a ligation buffer solution with a pH of 6.9, containing 100 mM MPAA. The reaction progress was monitored by RP-HPLC. After 10 hours, the lig-

ation reaction reached completion. Immediately following the NCL reaction, without any intermediate separation steps, 20% (by volume) of piperidine was directly added to the ligation system. This addition enabled the efficient removal of the Fmoc group at the N-terminus of the ligation product by adjusting the pH of the system to approximately 11. The reaction progress was continuously monitored by RP-HPLC, and within 10 minutes, all Fmoc groups were completely removed, yielding fragment **12** (Fig. 3B and C). Subsequently, the pH of the system was precisely adjusted back to 6.9. This step protonated the piperidine in the system, ensuring that it would not interfere with the subsequent second NCL reaction. Then, fragment **4** was added to the reaction system, and the reaction progress was continuously monitored using RP-HPLC. After another 10 hours, the reaction concluded, resulting in the formation of the ligation product **13** (Fig. 3D). After this reaction, without any purification procedures, the pH of the system was adjusted to the range of 2–3, and 5 equivalents of acetylaceton(acac) were added. The reaction mixture was then incubated at 37 °C for 1 hour. Subsequently, 100 mM MesNa was introduced, and the pH was adjusted to 5.0. The reaction was maintained at 37 °C for an additional hour. Under these conditions, the majority of the peptide hydrazide **13** was successfully converted into product **14** (Fig. 3E).

The overall isolated yield of this four-step one-pot process, involving three-fragment ligation, Fmoc-deprotection, a second ligation, and thioesterification, was 18.1%. This innovative scheme effectively avoided the use of custom-made, non-commercial TFA–Thz protecting groups, which were uti-





**Fig. 3** One-pot three segment ligation of segment 14. (A) Synthetic route of segment 14. (B) Analytical RP-HPLC trace ( $\lambda = 214$  nm) of segment 12. (C) ESI-MS spectrum of segment 12. (D) ESI-MS spectrum of segment 13. (E) Analytical RP-HPLC trace ( $\lambda = 214$  nm) of segment 14 and the ESI-MS deconvoluted spectrum of segment 14. “##” corresponds to the dibenzofulvene–piperidine adduct.

lized in previous methodologies. Moreover, it streamlined the previously cumbersome synthetic route. The original route, which involved 8 reaction steps, 3 HPLC purification steps, and 3 lyophilization treatments, was simplified to just four reaction steps, along with only 1 HPLC purification and 1 lyophilization.<sup>44</sup> This substantial simplification of the synthetic process led to a remarkable reduction in the total time required for fragment acquisition. It took merely two days to obtain fragment 14, which was three to four times faster than the original synthetic route, vividly demonstrating the distinct superiority of the one-pot method.


Inspired by the outcome achieved with segment 14, we further explored the one-pot strategy for the synthesis of compound 15 (Fig. 4A). The sequence of 15 was strategically partitioned into 4 segments, as depicted in Fig. 1: 7 (Cys229-Arg267, 38AA), 8 (Cys268-Tyr312, 44AA), 9 (Cys313-Thr352, 39AA), and 14 (Cys123-Leu228, 105AA). These segments were successively ligated to generate the target polypeptide. Both segments 7 and 8, prepared as MPAA thioesters, were synthesized through the acetylaceton-mediated activation of peptidyl hydrazides,<sup>49</sup> which were derived from the crude peptides following Fmoc SPPS and cleavage. Segment 14, a MesNa thioester, was obtained *via* a previously described one-pot ligation process (Fig. 3). The synthesis was initiated by the ligation of peptides 8 (1.0 equiv.) and 9 (1.0 equiv.) to yield peptide 10 (Fig. 4B). The reaction was conducted in pH 6.9 buffer containing 100 mM MPAA, and its progress was monitored by

RP-HPLC. After 10 hours, the ligation reaction reached completion, and then the *in situ* removal of the Fmoc group at the N-terminus was carried out in a one-pot manner. The Fmoc group was smoothly removed within 10 minutes by adding 20% (by volume) piperidine by adjusting the final pH to approximately 11, and the reaction progress was monitored by RP-HPLC. Subsequently, the pH of the reaction mixture was adjusted to 6.9 for the next ligation step. Without any purification, peptide 7 (1.0 equiv.) was directly added for ligation with 10 to obtain 11 in a one-pot reaction (Fig. 4C). The subsequent Fmoc deprotection and pH adjustment were carried out following the same procedure as that for fragment 10. Finally, 14 (1.3 equiv.) was added for the final ligation under a pH of 6.9, affording polypeptide 15 with an isolated yield of 41.7% (Fig. 4D).

During this one-pot process, the formation of the main by-product was ascribed to the reaction of 9 with an excess of 14. Notably, the overall yield (41.7% in one-pot) was significantly enhanced compared to the reported multi-step synthetic route (18.9%). Furthermore, the purification and lyophilization steps has been substantially reduced from six times to two times, thereby greatly improving the overall synthetic efficiency of *D*-Dpo4.

Hydrazide-based native chemical ligation (NCL) between 1 (1.1 equiv.) and 2 (1 equiv.) was carried out at pH 6.6 for 12 hours. Subsequently, desulfurization was performed to yield 20 (Fig. S2 and S3†). After activation with NaNO<sub>2</sub> at pH 3,






**Fig. 4** One-pot three segment ligation of segment 15. (A) Synthetic route of segment 15. (B) Analytical RP-HPLC trace ( $\lambda = 214$  nm) of segment 10. (C) ESI-MS spectrum of segment 11. (D) Analytical RP-HPLC trace ( $\lambda = 214$  nm) of segment 15 and the ESI-MS deconvoluted spectrum of segment 15. “#” corresponds to the excess of 14 ligated with 9. “##” corresponds to the dibenzofulvene–piperidine adduct.

a clean conversion of the hydrazide **20** to the MesNa thioester **21** was detected (Fig. S4†). Similarly, **3** was transformed into **17** through the same activation and thiolysis procedure (Fig. S1†). Then, **17** was ligated with **16** at pH 6.5 for 16 hours, smoothly and cleanly forming **18** with an isolated yield of 78% (Fig. S6†). Subsequently, free-radical-based desulfurization of **18** was conducted using the VA-044-based method for 14 hours. This process successfully converted all six ligation-site Cys residues back to Ala with an isolated yield of 79% (Fig. S7†). Thereafter, the Acm protecting group on Cys86 was removed by PdCl<sub>2</sub> with an isolated yield of 75% (Fig. S8†).

Finally, the purified **19** (1.0 equiv.) was ligated with **21** (2.0 equiv.) at pH 6.5 for 18 hours, affording the full-length product *D*-Dpo4 with an isolated yield of 70% (Fig. S9†). The final product was characterized by analytical RP-HPLC and ESI-MS (Fig. 5A). Then the *D*-protein was folded by successive dialysis against a series of renaturation buffer solutions containing 4 M, 2 M, 1 M, 0.5 M, 0.25 M, and 0 M Gn-HCl, respectively. After dialysis, the folded *D*-Dpo4 was heated and purified using Ni-NTA. Additionally, the synthetic *D*-Dpo4 was characterized by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), which showed a single band at the expected molecular weight of approximately 40 kDa, identical to that of recombinant *L*-Dpo4 (Fig. 5B).

With the synthetic mirror-image Dpo4 obtained, we utilized the miPCR system to test its enzyme activity. The miPCR reactions were carried out in 25  $\mu$ L reaction systems containing 25 mM HEPES (pH 7.5), 5 mM MgCl<sub>2</sub>, 50 mM NaCl, 0.1 mM EDTA, 5 mM DTT, 10% glycerol, 3% DMSO, 0.1 mg/mL BSA, 160  $\mu$ M (each) *L*-dNTPs, 0.2  $\mu$ M (each) *L*-primers, 60 nM



**Fig. 5** Chemical synthesis and characterization of *D*-Dpo4. (A) Analytical RP-HPLC chromatogram ( $\lambda = 214$  nm) and ESI-MS deconvoluted spectra of *D*-Dpo4. (B) SDS-PAGE analysis of synthetic *D*-Dpo4 and recombinant *L*-Dpo4, stained with Coomassie Brilliant Blue. M, protein maker. (C) PCR and miPCR amplification of a 128 bp *D/L*-DNA sequence using *L/D*-Dpo4, analyzed by 3% agarose gel electrophoresis and stained with 4S Gelred. M, DNA maker. (D) The products of miPCR and PCR were digested with DNase I and analysed by 3% agarose gel electrophoresis. M, DNA maker.

*L*-ssDNA template, and  $\sim 1$   $\mu$ M *D*-Dpo4 polymerase. Considering that the mirror-image DNA polymerization was less efficient than the corresponding natural system, the PCR reaction was



carried out in two phases of 15 cycles each, with the addition of fresh enzyme between the phases. The PCR program settings were as follows: 86 °C for 30 s (initial denaturation); 86 °C for 15 s (denaturation), 54 °C for 15 s (annealing), and 65 °C for 2 min (extension) for 30 cycles; 65 °C for 5 min (final extension). The miPCR products exhibited a clear band in the agarose gel with the expected length of 128 bp, and the intensity of the band increased with the number of cycles (Fig. 5C). To test the chirality of the miPCR product, we used an endonuclease (DNase I) to digest the products of the miPCR and PCR. Both the products of miPCR and PCR were digested with 5 U DNase I at 37 °C for 30 min and then analyzed by 3% agarose gel electrophoresis. In this case, the products of miPCR were completely resistant to digestion by the endonuclease of natural chirality (Fig. 5D). It shows promising potential for generating L-nucleic acid aptamers for research and therapeutic purposes.

## Conclusions

In summary, we reported the efficient total chemical synthesis of the 352-residue D-Dpo4 with 6 His tag using a one-pot multi-segment condensation strategy. To simplify the intermediate purification, peptides **4**, **5** and **6** were assembled into **14** by one-pot three-segment ligation, and peptides **7**, **8**, **9** and **14** were then assembled into **15** by one-pot four-segment ligation. In this way, D-Dpo4 can be obtained more efficiently with an overall yield of 9.8%. Our work provides a practical means to chemically synthesize mirror-image proteins with larger sizes.

## Author contributions

Miao Wang: most part of synthesis, original draft, data analysis, investigation. Tingting Cui: data analysis, manuscript review, investigation. Shuqing Huang: writing – review & editing. Dongyang Han: writing – review & editing. Xiangyu Deng: writing – review & editing. Yanbo Liu: writing – review & editing. Chaowei Shi: resources, supervision.

## Data availability

The data supporting this article have been included as part of the ESI.†

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This work was supported by the National Key R&D Program of China (2021YFA0909400) and the National Natural Science Foundation of China (22422411).

## References

- 1 L. Zhao and W. Lu, *Curr. Opin. Chem. Biol.*, 2014, **22**, 56–61.
- 2 M. Muttenthaler, G. F. King, D. J. Adams and P. F. Alewood, *Nat. Rev. Drug Discovery*, 2021, **20**, 309–325.
- 3 K. Harrison, A. S. Mackay, L. Kambanis, J. W. C. Maxwell and R. J. Payne, *Nat. Rev. Chem.*, 2023, **7**, 383–404.
- 4 X. Zhou, C. Zuo, W. Li, W. Shi, X. Zhou, H. Wang, S. Chen, J. Du, G. Chen, W. Zhai, W. Zhao, Y. Wu, Y. Qi, L. Liu and Y. Gao, *Angew. Chem., Int. Ed.*, 2020, **59**, 15114–15118.
- 5 A. J. Callahan, S. Gandhesiri, T. L. Travaline, R. M. Reja, L. L. Salazar, S. Hanna, Y. C. Lee, K. Li, O. S. Tokareva, J. M. Swiecicki, A. Loas, G. L. Verdine, J. H. McGee and B. L. Pentelute, *Nat. Commun.*, 2024, **15**, 1813.
- 6 C. Zuo, W. W. Shi, X. X. Chen, M. Glatz, B. Riedl, I. Flamme, E. Pook, J. W. Wang, G. M. Fang, D. Bierer and L. Liu, *Sci. China: Chem.*, 2019, **62**, 1371–1378.
- 7 Y.-K. Qi, J.-S. Zheng and L. Liu, *Chem*, 2024, **10**, 2390–2407.
- 8 R. C. Milton, S. C. Milton and S. B. Kent, *Science*, 1992, **256**, 1445–1448.
- 9 M. T. Weinstock, M. T. Jacobsen and M. S. Kay, *Proc. Natl. Acad. Sci. U. S. A.*, 2014, **111**, 11679–11684.
- 10 A. M. Levinson, J. H. McGee, A. G. Roberts, G. S. Creech, T. Wang, M. T. Peterson, R. C. Hendrickson, G. L. Verdine and S. J. Danishefsky, *J. Am. Chem. Soc.*, 2017, **139**, 7632–7639.
- 11 A. C. Conibear, E. E. Watson, R. J. Payne and C. F. W. Becker, *Chem. Soc. Rev.*, 2018, **47**, 9046–9068.
- 12 J. S. Zheng, S. Tang, Y. K. Qi, Z. P. Wang and L. Liu, *Nat. Protoc.*, 2013, **8**, 2483–2495.
- 13 M. Pan, S. Gao, Y. Zheng, X. Tan, H. Lan, X. Tan, D. Sun, L. Lu, T. Wang, Q. Zheng, Y. Huang, J. Wang and L. Liu, *J. Am. Chem. Soc.*, 2016, **138**, 7429–7435.
- 14 Z. Wang, W. Xu, L. Liu and T. F. Zhu, *Nat. Chem.*, 2016, **8**, 698–704.
- 15 A. Pech, J. Achenbach, M. Jahnz, S. Schulzchen, F. Jarosch, F. Bordusa and S. Klussmann, *Nucleic Acids Res.*, 2017, **45**, 3997–4005.
- 16 J. B. Blanco-Canosa and P. E. Dawson, *Angew. Chem., Int. Ed.*, 2008, **47**, 6851–6855.
- 17 G. M. Fang, Y. M. Li, F. Shen, Y. C. Huang, J. B. Li, Y. Lin, H. K. Cui and L. Liu, *Angew. Chem., Int. Ed.*, 2011, **50**, 7645–7649.
- 18 G. M. Fang, J. X. Wang and L. Liu, *Angew. Chem., Int. Ed.*, 2012, **51**, 10347–10350.
- 19 Y. C. Huang, G. M. Fang and L. Liu, *Natl. Sci. Rev.*, 2016, **3**, 107–116.
- 20 H. Ai, M. Sun, A. Liu, Z. Sun, T. Liu, L. Cao, L. Liang, Q. Qu, Z. Li, Z. Deng, Z. Tong, G. Chu, X. Tian, H. Deng, S. Zhao, J. B. Li, Z. Lou and L. Liu, *Nat. Chem. Biol.*, 2022, **18**, 972–980.



- 21 S. Tang, Y. Y. Si, Z. P. Wang, K. R. Mei, X. Chen, J. Y. Cheng, J. S. Zheng and L. Liu, *Angew. Chem., Int. Ed.*, 2015, **54**, 5713–5717.
- 22 R. Ding, W. W. Shi and J. S. Zheng, *Org. Lett.*, 2023, **25**, 4857–4861.
- 23 H. N. Chang, B. Y. Liu, Y. K. Qi, Y. Zhou, Y. P. Chen, K. M. Pan, W. W. Li, X. M. Zhou, W. W. Ma, C. Y. Fu, Y. M. Qi, L. Liu and Y. F. Gao, *Angew. Chem., Int. Ed.*, 2015, **54**, 11760–11764.
- 24 P. S. Marinec, K. E. Landgraf, M. Uppalapati, G. Chen, D. Xie, Q. Jiang, Y. Zhao, A. Petriello, K. Deshayes, S. B. H. Kent, D. Ault-Riche and S. S. Sidhu, *ACS Chem. Biol.*, 2021, **16**, 548–556.
- 25 M. Uppalapati, D. J. Lee, K. Mandal, H. Li, L. P. Miranda, J. Lowitz, J. Kenney, J. J. Adams, D. Ault-Riche, S. B. Kent and S. S. Sidhu, *ACS Chem. Biol.*, 2016, **11**, 1058–1065.
- 26 S. W. Dong, J. S. Zheng, Y. M. Li, H. Wang, G. Chen, Y. X. Chen, G. M. Fang, J. Guo, C. M. He, H. G. Hu, X. C. Li, Y. M. Li, Z. G. Li, M. Pan, S. Tang, C. L. Tian, P. Wang, B. Wu, C. L. Wu, J. F. Zhao and L. Liu, *Sci. China: Chem.*, 2024, **67**, 1060–1096.
- 27 Y. Zheng, B. Zhang, W. W. Shi, X. Deng, T. Y. Wang, D. Han, Y. Ren, Z. Yang, Y. K. Zhou, J. Kuang, Z. W. Wang, S. Tang and J. S. Zheng, *Angew. Chem., Int. Ed.*, 2024, **63**, e202318897.
- 28 P. E. Dawson, T. W. Muir, I. Clark-Lewis and S. B. Kent, *Science*, 1994, **266**, 776–779.
- 29 Y. Zhang, C. Xu, H. Y. Lam, C. L. Lee and X. Li, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**, 6657–6662.
- 30 J. S. Zheng, M. Yu, Y. K. Qi, S. Tang, F. Shen, Z. P. Wang, L. Xiao, L. Zhang, C. L. Tian and L. Liu, *J. Am. Chem. Soc.*, 2014, **136**, 3695–3704.
- 31 J. X. Wang, G. M. Fang, Y. He, D. L. Qu, M. Yu, Z. Y. Hong and L. Liu, *Angew. Chem., Int. Ed.*, 2015, **54**, 2194–2198.
- 32 C. Zuo, R. Ding, X. Wu, Y. Wang, G. C. Chu, L. J. Liang, H. Ai, Z. B. Tong, J. Mao, Q. Zheng, T. Wang, Z. Li, L. Liu and D. Sun, *Angew. Chem., Int. Ed.*, 2022, **61**, e202201887.
- 33 W. W. Shi, C. Shi, T. Y. Wang, Y. L. Li, Y. K. Zhou, X. H. Zhang, D. Bierer, J. S. Zheng and L. Liu, *J. Am. Chem. Soc.*, 2022, **144**, 349–357.
- 34 Y. M. Li, Y. T. Li, M. Pan, X. Q. Kong, Y. C. Huang, Z. Y. Hong and L. Liu, *Angew. Chem., Int. Ed.*, 2014, **53**, 2198–2202.
- 35 W. Shi, T. Wang, Z. Yang, Y. Ren, D. Han, Y. Zheng, X. Deng, S. Tang and J. S. Zheng, *Angew. Chem., Int. Ed.*, 2024, **63**, e202313640.
- 36 J. Zhao, X. Liu, J. Liu, F. Ye, B. Wei, M. Deng, T. Li, P. Huang and P. Wang, *J. Am. Chem. Soc.*, 2024, **146**, 2615–2623.
- 37 F. Ye, J. Zhao, P. Xu, X. Liu, J. Yu, W. Shangguan, J. Liu, X. Luo, C. Li, T. Ying, J. Wang, B. Yu and P. Wang, *Angew. Chem., Int. Ed.*, 2021, **60**, 12904–12910.
- 38 T. Li, H. Liu and X. Li, *Org. Lett.*, 2016, **18**, 5944–5947.
- 39 Z. Sun and X. Li, *Chin. J. Chem.*, 2021, **39**, 2795–2800.
- 40 J. S. Zheng, J. Liang, W. W. Shi, Y. Li, H. G. Hu, C. L. Tian and L. Liu, *Sci. Bull.*, 2021, **66**, 1542–1549.
- 41 Y. Xu and T. F. Zhu, *Science*, 2022, **378**, 405–412.
- 42 J. Cramer and T. Restle, *J. Biol. Chem.*, 2005, **280**, 40552–40558.
- 43 J. Chen, M. Chen and T. F. Zhu, *Nat. Biotechnol.*, 2022, **40**, 1601–1609.
- 44 W. Jiang, B. Zhang, C. Fan, M. Wang, J. Wang, Q. Deng, X. Liu, J. Chen, J. Zheng, L. Liu and T. F. Zhu, *Cell Discovery*, 2017, **3**, 17037.
- 45 L. Z. Yan and P. E. Dawson, *J. Am. Chem. Soc.*, 2001, **123**, 526–533.
- 46 Z. Q. Sun, W. J. Ma, Y. H. Cao, T. Y. Wei, X. Y. Mo, H. Y. Chow, Y. Tan, C. H. P. Cheung, J. M. Liu, H. K. Lee, E. C. M. Tse, H. Liu and X. C. Li, *Chem*, 2022, **8**, 2542–2557.
- 47 D. Han, Y. Cui, X. Deng, C. Li, X. Zhu, B. Wang, G. C. Chu, Z. A. Wang, S. Tang, J. S. Zheng, L. J. Liang and L. Liu, *J. Am. Chem. Soc.*, 2025, **147**, 4135–4146.
- 48 K. Jin, T. Li, H. Y. Chow, H. Liu and X. Li, *Angew. Chem., Int. Ed.*, 2017, **56**, 14607–14611.
- 49 D. T. Flood, J. C. J. Hintzen, M. J. Bird, P. A. Cistrone, J. S. Chen and P. E. Dawson, *Angew. Chem., Int. Ed.*, 2018, **57**, 11634–11639.
- 50 B. Zhang, Y. Zheng, G. Chu, X. Deng, T. Wang, W. Shi, Y. Zhou, S. Tang, J. S. Zheng and L. Liu, *Angew. Chem., Int. Ed.*, 2023, **62**, e202306270.
- 51 A. Kar, J. Mannuthodikayil, S. Singh, A. Biswas, P. Dubey, A. Das and K. Mandal, *Angew. Chem., Int. Ed.*, 2020, **59**, 14796–14801.
- 52 S. K. Maity, M. Jbara, S. Laps and A. Brik, *Angew. Chem., Int. Ed.*, 2016, **55**, 8108–8112.

