# Organic & Biomolecular Chemistry



View Article Online

## PAPER

Check for updates

Cite this: Org. Biomol. Chem., 2025, 23, 4730

## Nonsilyl bicyclic secondary amine catalyzed Michael addition of nitromethane to $\beta$ , $\beta$ -disubstituted $\alpha$ , $\beta$ -unsaturated aldehydes†

Rohtash Kumar,‡ Avinash Avinash,‡ Ronak Mehta and Chandrakumar Appayee 吵 \*

The asymmetric Michael addition of nitroalkanes to  $\beta_i\beta$ -disubstituted  $\alpha_i\beta$ -unsaturated aldehydes is a useful method for the construction of all-carbon quaternary stereocenters. Nonsilyl bicyclic secondary amine organocatalysts were employed in reactions involving a wide range of  $\beta_i\beta_i$ -disubstituted  $\alpha_i\beta_i$ -unsaturated aldehydes with nitroalkanes to achieve products with all-carbon quaternary stereocenters in up to 69% yield and 95% ee. The scalability of this methodology was demonstrated at the 5.1 mmol scale. The synthetic utility of this methodology is showcased through the concise asymmetric synthesis of methsuximide, an anticonvulsant drug.

Received 15th February 2025, Accepted 14th April 2025 DOI: 10.1039/d5ob00277j

rsc.li/obc

## Introduction

The asymmetric synthesis of molecules bearing all-carbon quaternary stereocenters remains one of the challenging tasks in synthetic organic chemistry.<sup>1</sup> Interestingly, this methodology finds potential applications in the total synthesis of natural products and pharmaceutical drugs.<sup>2</sup> The Michael addition of carbon nucleophiles to  $\beta$ , $\beta$ -disubstituted  $\alpha$ , $\beta$ -unsaturated carbonyl compounds is one of the elegant methods to access molecules with all-carbon guaternary stereocenters.<sup>3</sup> Nevertheless, the Michael addition of nitromethane to  $\beta$ , $\beta$ -disubstituted  $\alpha$ , $\beta$ -unsaturated aldehydes has been comparatively underexplored (Scheme 1a).<sup>4</sup> Kudo and Akagawa reported an undecapeptide-catalyzed asymmetric Michael addition of nitromethane to  $\beta$ , $\beta$ -disubstituted  $\alpha$ , $\beta$ -unsaturated aldehydes, affording aldehydes bearing β-quaternary carbon stereocenters.<sup>4a</sup> Tu and co-workers developed a spiro-pyrrolidine catalyst for the same transformation, followed by reduction using NaBH<sub>4</sub>, yielding alcohols with all-carbon quaternary stereocenters.<sup>4b</sup> Although the above two methodologies offer products in moderate to good yields and excellent enantioselectivities, the synthesis of these catalysts involves multistep tedious processes. Hayashi and co-workers demonstrated an asymmetric Michael addition of nitromethane to  $\beta$ , $\beta$ -disubstituted  $\alpha$ , $\beta$ -unsaturated aldehydes using the Hayashi-Jørgensen catalyst, providing aldehydes with β-quaternary carbon stereocenters.<sup>4c,d</sup> Although the trimethyl-

†Electronic supplementary information (ESI) available. See DOI: https://doi.org/ 10.1039/d5ob00277j silyl ether group in Hayashi–Jørgensen catalysts plays a crucial role in achieving high enantioselectivity of products, its stability in the presence of acid additives<sup>5*a,b*</sup> and reagents<sup>5*c*</sup> is a concern. Recently, we reported the synthesis of a nonsilyl bicyclic secondary amine catalyst, derived from 4-hydroxy-Lproline in four steps, which was utilized for the asymmetric transfer hydrogenation of  $\beta$ , $\beta$ -disubstituted  $\alpha$ , $\beta$ -unsaturated aldehydes.<sup>6</sup> Herein, we disclose the nonsilyl bicyclic secondary amine-catalyzed asymmetric Michael addition of nitromethane



**Scheme 1** Asymmetric Michael addition o  $\beta$ , $\beta$ -disubstituted  $\alpha$ , $\beta$ -unsaturated aldehydes.

of nitromethane to

(an anticonvulsant drug)

40-69% yield

58-95% ee

Department of Chemistry, Indian Institute of Technology Gandhinagar, Palaj,

Gandhinagar, Gujarat 382055, India. E-mail: a.chandra@iitgn.ac.in

<sup>‡</sup>These authors contributed equally.

to  $\beta$ , $\beta$ -disubstituted  $\alpha$ , $\beta$ -unsaturated aldehydes. Furthermore, we demonstrate the synthetic utility of this methodology for the concise asymmetric synthesis of methsuximide,<sup>7</sup> an anticonvulsant drug (Scheme 1b).

#### **Results and discussion**

Initially,  $\beta$ -methyl cinnamaldehyde **1a** was treated with nitromethane **2a** (10 equiv.) in the presence of the nonsilyl bicyclic secondary amine catalyst **3a** (20 mol%) under neat reaction conditions at rt for 72 h to afford the Michael adduct. For characterization purposes, the crude product was reduced *in situ* using NaBH<sub>4</sub> in EtOH at -5 to 0 °C, furnishing alcohol **4a** in 20% isolated yield with 88% ee (Table 1, entry 1).

Among the solvents screened (Table 1, entries 2-12), protic solvents were found to be effective, offering shorter reaction times and improved product yields (Table 1, entries 5-12). Notably, the reaction carried out in IPA (isopropanol) was completed in 36 h, affording product 4a in 69% isolated yield with 95% ee (Table 1, entry 7). Attempts to either decrease (Table 1, entry 13) or increase (Table 1, entry 14) the equivalents of nitromethane 2a were not fruitful. The addition of benzoic acid (20 mol%) to the reaction led to a longer reaction time, which in turn resulted in decreased product yield and enantioselectivity (Table 1, entry 15). This decline is attributed to product decomposition occurring over the prolonged reaction time. Traces of product 4a were observed upon lowering the reaction temperature to 0 °C (Table 1, entry 16). The reaction with 10 mol% of catalyst 3a took a longer time (96 h), affording product 4a in 53% yield and 93% ee (Table 1, entry 17). When catalyst 3a was replaced with 3b or 3c under the same reaction conditions, lower product yield and enantioselectivity were observed (Table 1, entries 18 and 19).

After the successful reaction optimization for the asymmetric Michael addition of nitromethane 2a to β-methyl cinnamaldehyde 1a (Table 1, entry 7), the substrate scope was investigated, as shown in Scheme 2. The  $\beta$ -aryl  $\alpha$ ,  $\beta$ -unsaturated aldehydes 1b-1e containing electron-donating groups (p-OMe, p-Me, p-Cl and p-F) at the para-position of the aryl ring smoothly underwent Michael addition with nitromethane 2a to give the corresponding alcohols 4b-4e in 48-61% yields and 91-95% ees. Similarly, substrates 1f and 1g containing electron-withdrawing groups (p-NO<sub>2</sub> and p-CF<sub>3</sub>, respectively) at the para-position also participated effectively in the reaction, providing alcohols 4f and 4g in 61% yield and 91% ee and in 45% yield and 92% ee, respectively. Substrates 1h and 1k with metasubstituted aryl groups, including electron donating groups (*m*-OMe or *m*-Me) and electron-withdrawing groups (*m*-NO<sub>2</sub> or m-CF<sub>3</sub>), also participated in the transformation efficiently under the optimized reaction conditions to afford the corresponding alcohols 4h and 4k in moderate to good yields and excellent enantioselectivities.  $\beta$ -Aryl  $\alpha$ ,  $\beta$ -unsaturated aldehydes bearing biphenyl (11) and 2-naphthyl (1m) substituents at the  $\beta$ -position provided alcohols **4l** and **4m** in 50% yield and 93% ee and in 52% yield and 94% ee, respectively. Notably, a het-

Table 1 Optimization of the reaction conditions<sup>a</sup>



Entry	Cat.	Solvent	<i>t</i> (h)	$\operatorname{Yield}^{b}(\%)$	ee <sup>c</sup> (%)
1	3a	Neat	72	20	88
2	3a	$CH_2Cl_2$	120	21	90
3	3a	Toluene	72	25	93
4	3a	$CH_3CN$	120	16	45
5	3a	MeOH	24	30	91
6	3a	EtOH	54	59	93
7	3a	IPA	36	69	95
8	3a	t-BuOH	24	60	95
9	3a	$(CH_2OH)_2$	30	64	94
10	3a	Glycerol	48	65	94
11	3a	$HO(CH_2)_4OH$	48	62	91
12	3a	TFE	24	60	88
$13^d$	3a	IPA	30	40	91
$14^e$	3a	IPA	48	59	93
$15^{f}$	3a	IPA	92	40	91
$16^g$	3a	IPA	48	Trace	nd
$17^{h}$	3a	IPA	96	53	93
18	3b	IPA	48	35	91
19	3c	IPA	50	48	91

<sup>*a*</sup> Reactions were performed with **1a** (29 mg, 0.2 mmol, 1 equiv.), CH<sub>3</sub>NO<sub>2</sub> **2a** (107  $\mu$ L, 2.0 mmol, 10 equiv.) and catalyst 3 (0.04 mmol, 20 mol%) in solvent (0.2 mL, 1.0 M) at rt followed by reduction to the corresponding alcohol **4a** using NaBH<sub>4</sub> (76 mg, 2.0 mmol, 10 equiv.) in EtOH (2.5 mL), unless otherwise noted. <sup>*b*</sup> Isolated yield after column chromatography. <sup>*c*</sup> Determined by HPLC analysis on a chiral stationary phase. <sup>*d*</sup> CH<sub>3</sub>NO<sub>2</sub> (53  $\mu$ L, 1.0 mmol, 5 equiv.) was used. <sup>*e*</sup> CH<sub>3</sub>NO<sub>2</sub> (214  $\mu$ L, 4.0 mmol, 20 equiv.) was used. <sup>*f*</sup> Benzoic acid (5 mg, 0.04 mmol, 20 mol%) was added. <sup>*g*</sup> Reaction was performed at 0 °C. <sup>*h*</sup> Catalyst **3a** (6 mg, 0.02 mmol, 10 mol%) was used. IPA = isopropanol; nd = not determined; TFE = trifluoroethanol.

eroaromatic substrate bearing a 2-thiophenyl substituent (1n) was well tolerated, affording alcohol 4n in 47% yield and 95% ee. When the  $\beta$ -ethyl  $\beta$ -phenyl  $\alpha$ , $\beta$ -unsaturated aldehyde 1o was studied for the Michael addition of nitromethane 2a, alcohol 4o was obtained in 40% yield and 88% ee. The Michael addition of nitromethane 2a to the 2-oxoindoline-3-ylidene acetaldehyde 1p was not effective under the optimized reaction conditions.<sup>8</sup> Interestingly, the use of catalyst 3c (20 mol%) facilitated the reaction, affording alcohol 4p in 51% yield and 87% ee.

This methodology was also applied to the  $\beta$ , $\beta$ -dialkyl  $\alpha$ , $\beta$ -unsaturated aldehydes **1q–1s** to obtain alcohols **4q–4s** in 49–61% yields and 58–83% ees. The enantiomeric excesses of alcohols **4r** and **4s** were determined after converting them to



Scheme 2 Substrate scope. <sup>a</sup> Reactions were performed with 1 (0.2 mmol, 1 equiv.), nitroalkane 2 (2.0 mmol, 10 equiv.) and catalyst 3a (11 mg, 0.04 mmol, 20 mol%) in IPA (0.2 mL, 1.0 M) at rt followed by reduction to the corresponding alcohol 4 using NaBH<sub>4</sub> (76 mg, 2.0 mmol, 10 equiv.) in EtOH (2.5 mL), unless otherwise noted. <sup>b</sup> Catalyst 3c (22 mg, 0.04 mmol, 20 mol%) was used. <sup>c</sup> Enantiomeric excess was determined after converting alcohols 4r and 4s to the carbamate derivatives 6r and 6s, respectively, using 4-chlorophenyl isocyanate. <sup>d</sup> Reaction was performed in EtOH (1.0 M). <sup>e</sup> Enantiomeric excess was determined after converting alcohols 7t using ethyl(triphenylphosphoranylidene)acetate. <sup>f</sup> Reaction was performed under neat conditions.



Scheme 3 Scale-up synthesis of alcohol 4a.

the corresponding carbamate derivatives 6r (85% yield and 68% ee) and 6s (90% yield and 83% ee), respectively, using 4-chlorophenyl isocyanate and  $Et_3N$  in  $CH_2Cl_2$  (see the ESI<sup>†</sup>). The reaction of nitromethane 2a with substrate 1t containing an ester group at the  $\beta$ -position proceeded very slowly in IPA. However, when the reaction was carried out in EtOH, completion was achieved in 18 h, affording aldehyde 5t in 62% yield and 93% ee. The enantiomeric excess of aldehyde 5t was determined after converting it to diester 7t (85% vield and 93% ee) using ethyl(triphenylphosphoranylidene)acetate (see the ESI<sup> $\dagger$ </sup>). The reaction of nitropropane **2b** with  $\beta$ -methyl cinnamaldehyde 1a in the presence of catalyst 3a under neat conditions (as the reaction in IPA was very slow) yielded product 4u as an inseparable mixture of diastereomers in 29% yield with 1.4:1 dr and 83% ee. The absolute configuration of products 4a-4u was assigned based on the comparison of the optical rotation values of products 4a, 4b, 4f, 4p and 4q with those reported in the literature.<sup>4b</sup>

To showcase the scalability of the methodology, 5.1 mmol of  $\beta$ -methyl cinnamaldehyde **1a** was subjected to the optimized reaction conditions, affording alcohol **4a** in 66% yield and 95% ee (Scheme 3).

To demonstrate the application of this methodology, a concise asymmetric synthesis of methsuximide, an anticonvulsant drug, was envisaged starting from  $\beta$ -methyl cinnamaldehyde **1a**, as shown in Scheme 4. The Michael addition of nitromethane **2a** to  $\beta$ -methyl cinnamaldehyde **1a** catalyzed by **3a** in IPA furnished the corresponding Michael adduct, which was subsequently oxidized to carboxylic acid **8a** *via* Pinnick oxidation (56% yield over two steps). Furthermore, the nitro group in compound **8a** was converted to a carboxylic acid using NaNO<sub>2</sub> and acetic acid in DMSO, yielding the corresponding dicarboxylic acid. Condensation of this intermediate with methylamine in toluene under reflux conditions afforded methsuximide **9a** in 38% yield over two steps and 93% *ee*.

Based on the reaction outcomes and the observed stereochemistry of the Michael adducts, a possible reaction mechanism is proposed for the Michael addition of nitromethane **2a** to  $\beta$ , $\beta$ -disubstituted  $\alpha$ , $\beta$ -unsaturated aldehydes **1** catalyzed by the nonsilyl bicyclic secondary amine organocatalyst **3a**, as shown in Scheme 5. Accordingly, catalyst **3a** condenses with the  $\beta$ , $\beta$ -disubstituted  $\alpha$ , $\beta$ -unsaturated aldehyde **1** to form the corresponding *E*-iminium ion intermediate **A**. The nucleophilic nitromethide ion, generated from nitromethane **2a**, then



Scheme 4 Concise asymmetric synthesis of methsuximide 9a.



Scheme 5 Proposed reaction mechanism for the Michael addition of nitromethane 2a to  $\beta$ , $\beta$ -disubstituted  $\alpha$ , $\beta$ -unsaturated aldehydes 1 catalyzed by 3a.

attacks the  $\beta$ -position of the iminium ion **B** from the face opposite to the chiral bulky substituent of the catalyst, leading to the formation of the enamine intermediate **C**. Upon protonation, the enamine intermediate **C** forms the iminium intermediate **D**, which undergoes hydrolysis to yield the Michael adduct 5 along with the regeneration of catalyst **3a**.

## Conclusions

In summary, the nonsilyl bicyclic secondary amine organocatalysts 3a-3c were investigated for the enantioselective Michael addition of nitromethane 2a to  $\beta$ -methyl cinnamaldehyde 1a. A broad range of  $\beta$ , $\beta$ -disubstituted  $\alpha$ , $\beta$ -unsaturated aldehydes (1a-1s) were evaluated under the optimized reaction conditions, affording the corresponding products 4a-4s bearing all-carbon quaternary stereocenters in 40-69% yields and 58-95% ees. Substrate 1t, bearing an ester group at the  $\beta$ -position, was successfully converted to the corresponding Michael adduct 5t in 62% yield and 93% ee. Additionally, the use of nitropropane 2b as the nucleophile afforded product 4u in 29% yield with 1.4:1 dr and 83% ee. A scale-up reaction was successfully performed at the 5.1 mmol scale using  $\beta$ -methyl cinnamaldehyde 1a, delivering alcohol 4a in 66% vield and 95% ee. Furthermore, the synthetic utility of this methodology was exemplified through a concise asymmetric synthesis of methsuximide, an anticonvulsant drug, starting from  $\beta$ -methyl cinnamaldehyde **1a**. Based on the stereochemical outcomes of the Michael adducts, a possible reaction mechanism has been proposed. This study is expected to stimulate further exploration on the use of nonsilyl bicyclic secondary amine organocatalysts for enantioselective Michael addition of various nucleophiles to  $\alpha$ ,  $\beta$ -unsaturated aldehydes.

#### Data availability

Data supporting this article have been included as part of the ESI.†

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

Financial support from the Anusandhan National Research Foundation (ANRF), India, (SCP/2022/000823) is gratefully acknowledged. Financial support from the University Grants Commission (UGC), India (fellowship to R. K.) and the Indian Institute of Technology Gandhinagar (fellowship to A. A. and R. M.) is gratefully acknowledged.

#### References

 (a) K. W. Quasdorf and L. E. Overman, Nature, 2014, 516, 181–191; (b) E. A. Peterson and L. E. Overman, Proc. Natl. Acad. Sci. U. S. A., 2004, 101, 11943–11948; (c) M. Büschleb, S. Dorich, S. Hanessian, D. Tao, K. B. Schenthal and L. E. Overman, Angew. Chem., Int. Ed., 2016, 55, 4156–4186; (d) Y. Jia, K. Zhang, L.-Q. Lu, Y. Cheng and W.-J. Xiao, ACS Catal., 2024, 14, 13550–13556; (e) A. Y. Hong and B. M. Stoltz, Eur. J. Org. Chem., 2013, 2745–2759; (f) S. Goswami, K. Harada, M. F. El-Mansy, R. Lingampally and R. G. Carter, Angew. Chem., Int. Ed., 2018, 57, 9117– 9121; (g) T. S. Silva and F. Coelho, Beilstein J. Org. Chem., 2021, 17, 1565–1590; (h) X.-P. Zeng, Z.-Y. Cao, Y.-H. Wang, F. Zhou and J. Zhou, *Chem. Rev.*, 2016, **116**, 7330–7396;
(*i*) T. Ema, Y. Oue, K. Akihara, Y. Miyazaki and T. Sakai, *Org. Lett.*, 2009, **11**, 4866–4869;
(*j*) M. Tanaka, M. Imai, M. Fujio, E. Sakamoto, M. Takahashi, Y. Eto-Kato, X. M. Wu, K. Funakoshi, K. Sakai and H. Suemune, *J. Org. Chem.*, 2000, **65**, 5806–5816.

- 2 (a) C. Li, S. S. Ragab, G. Liu and W. Tang, Nat. Prod. Rep., 2020, 37, 276–292; (b) Y. Liu, S.-J. Han, W.-B. Liu and B. M. Stoltz, Acc. Chem. Res., 2015, 48, 740–751; (c) Z. Wang, Org. Chem. Front., 2020, 7, 3815–3841; (d) T. Ling and F. Rivas, Tetrahedron, 2016, 72, 6729–6777; (e) R. Long, J. Huang, J. Gong and Z. Yang, Nat. Prod. Rep., 2015, 32, 1584–1601; (f) D. C. Behenna, Y. Liu, T. Yurino, J. Kim, D. E. White, S. C. Virgil and B. M. Stoltz, Nat. Chem., 2012, 4, 130–133; (g) S.-J. Han, F. Vogt, S. Krishnan, J. A. May, M. Gatti, S. C. Virgil and B. M. Stoltz, Org. Lett., 2014, 16, 3316–3319; (h) R. M. Lemieux and A. I. Meyers, J. Am. Chem. Soc., 1998, 120, 5453–5457; (i) J. Deng, S. Zhou, W. Zhang, J. Li, R. Li and A. Li, J. Am. Chem. Soc., 2014, 136, 8185–8188; (j) R. Long, J. Huang, W. Shao, S. Liu, Y. Lan, J. Gong and Z. Yang, Nat. Commun., 2014, 5, 5707.
- 3 (a) A. G. Csakÿ, G. de la Herrán and M. C. Murcia, Chem. Soc. Rev., 2010, 39, 4080-4102; (b) C. Hawner and A. Alexakis, Chem. Commun., 2010, 46, 7295-7306; (c) E. Fillion and A. Wilsily, J. Am. Chem. Soc., 2006, 128, 2774-2775; (d) A. Wilsily and E. Fillion, Org. Lett., 2008, 10, 2801-2804; (e) Y. Hamashima, D. Hotta and M. Sodeoka, J. Am. Chem. Soc., 2002, 124, 11240-11241; (f) C. Hawner, D. Muller, L. Gremaud, A. Felouat, S. Woodward and A. Alexakis, Angew. Chem., Int. Ed., 2010, 49, 7769-7772; (g) K. Endo, D. Hamada, S. Yakeishi and T. Shibata, Angew.

*Chem., Int. Ed.*, 2013, **52**, 606–610; (*h*) R. Shintani, Y. Tstsumi, M. Nagaosa, T. Nishimura and T. Hayashi, *J. Am. Chem. Soc.*, 2009, **131**, 13588–13589.

- 4 (a) K. Akagawa and K. Kudo, Angew. Chem., Int. Ed., 2012,
  51, 12786-12789; (b) J.-M. Tian, Y.-H. Yuan, Y.-Q. Tu,
  F.-M. Zhang, X.-B. Zhang, S.-H. Zhang, S.-H. Wang and
  X.-M. Zhang, Chem. Commun., 2015, 51, 9979-9982;
  (c) Y. Hayashi, Y. Kawamoto, M. Honda, D. Okamura,
  S. Umemiya, Y. Noguchi, T. Mukaiyama and I. Sato, Chem. –
  Eur. J., 2014, 20, 12072-12082; (d) T. Mukaiyama, K. Ogata,
  I. Sato and Y. Hayashi, Chem. Eur. J., 2014, 20, 1358313588.
- 5 (a) M. H. Haindl, M. B. Schmid, K. Zeitler and R. M. Gschwind, *RSC Adv.*, 2012, 2, 5941–5943;
  (b) X. Companyó and J. Burés, *J. Am. Chem. Soc.*, 2017, 139, 8432–8435;
  (c) G. Hutchinson, C. Alamillo-Ferrer and J. Burés, *J. Am. Chem. Soc.*, 2021, 143, 6805–6809.
- 6 R. Kumar, V. Maurya, A. Avinash and C. Appayee, J. Org. Chem., 2024, 89, 8586–8600.
- 7 For the synthesis of methsuximide, see: (a) P.-W. Xu and Z.-X. Huang, *Nat. Chem.*, 2021, 13, 634–642; (b) Y. Kuroiwa and M. Tamura, *Adv. Synth. Catal.*, 2024, 366, 1996–2002; (c) J. Li, J. Wei, B. Zhu, T. Wang and N. Jiao, *Chem. Sci.*, 2019, 10, 9099–9103 For the biological activity of methsuximide, see: (d) J. Mifsud, J. S. Millership, P. S. Collier, R. Morrow, J. T. G. Hamilton and W. C. McRoberts, *Biopharm. Drug Dispos.*, 2001, 22, 129–136; (e) R. J. Porter, J. K. Penry, J. R. Lacy, M. E. Newmark and H. J. Kupferberg, *Neurology*, 1979, 29, 1509–1513.
- 8 Under the optimized reaction conditions, product **4p** was obtained in 80% yield and 27% ee after 16 h.