

Organic & Biomolecular Chemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: A. Chandra, S. C. Yadav, S. R. Cheekatla and A. Kumar, *Org. Biomol. Chem.*, 2025, DOI: 10.1039/D5OB00338E.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

REVIEW

A Review on Indoles Synthesis from Nitroarenes: Classical to Modern Approaches

Ajeet Chandra,^{*,[a]} Suresh C. Yadav,^{†[b]} Subba Rao Cheekatla^{‡[c]} and Abhijeet Kumar^[d]Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

Indole is a highly privileged and versatile heterocyclic pharmacophore that plays a crucial role in natural product synthesis, drug discovery, pharmaceuticals, and medicinal chemistry. This review provides a comprehensive analysis of various synthetic approaches to indole with a particular emphasis on nitroarenes as key precursors. Although indoles and their derivatives have been extensively explored for their bioactivity in natural and pharmacological contexts, several classical synthetic methodologies remain underutilized. Traditionally, indole synthesis from *ortho*-substituted nitroarenes has been achieved through methods such as Bartoli, Reissert, Cadogan, and Leimgruber–Batcho, etc. approaches. However, recent advancements have introduced novel one-pot and tandem strategies that effectively integrate redox and hydrogenation reactions to streamline indole formation. Emerging photochemical and electrochemical techniques have also enabled the selective conversion of nitroarenes into indoles bearing well-defined functional groups. Beyond their intrinsic biological activity, indoles serve as valuable intermediates for further derivatization into compounds such as isatins and oxindoles, expanding their synthetic potential. These advancements continue to enhance the synthetic toolkit for constructing biologically active indoles, with far-reaching applications in pharmaceuticals, agrochemicals, and advanced materials.

Introduction

The exploration, development, and synthesis of heterocyclic frameworks have long been captivating due to their remarkable biological and pharmacological properties.^{1–6} Heterocycles play a vital role in everyday life, offering various applications across various fields. Nitrogen-bearing heterocycles have long been a key focus in diverse fields due to their remarkable biological activities.^{7, 8} Over the past several decades, *N*-based heterocycles have garnered significant attention from synthetic chemists and chemical biologists for their unique ability to bind various receptors and their presence in numerous natural products and medicinally important compounds.^{9–12} Five-membered nitrogen-containing heterocyclic systems are widely distributed in nature and play a significant role in medicinal chemistry, pharmaceuticals, agrochemicals, dyes, chemosensors, and materials. Additionally, they serve as essential intermediates in organic synthesis.^{7, 13–21} Among these scaffolds, indole is a privileged pharmacophore, serving as a

core structure in many biologically active molecules.^{22–35}

Indole units are key structural motifs in natural products and synthetic macrocycles with distinct biological activities. Moreover, indoles interact with enzymes and receptors through non-covalent interactions and contribute to treatments for cancer, neurological disorders, and cardiovascular diseases.^{26, 36–43} They serve as versatile building blocks in heterocyclic synthesis and are present in alkaloids, hormones, and therapeutic agents. Indoles exhibit potent anticancer, antimicrobial, and anti-inflammatory properties.^{44–47} Structurally, they form the core of serotonin, a critical neurotransmitter.^{48–49} Their broad pharmacological relevance includes antiviral applications, such as SARS-CoV-2 3CL protease inhibition. Representative biologically active indole scaffolds are shown in Figure 1.^{50–53} In recent years, there has been growing interest in developing macrocyclic frameworks with heteroaryl systems.^{54–57} Notably, indole-based macrocyclic scaffolds are widely prevalent in biologically significant molecules and highlighting their central role in modern drug discovery and development.^{58–69} Indole-based drugs approved by the FDA play various roles in medical treatments (Figure 1).^{22, 28, 70–79} Numerous strategies have been developed and widely employed for constructing indoles and their macrocycles.⁸⁰

^a Department of Information Display, Kyung Hee University, Seoul 02447, Republic of Korea; Email ID: ajeet12300@gmail.com and ORCID: <https://orcid.org/0000-0003-3437-9605>

^b Department of Chemistry, Satish Chandra College, Ballia, Uttar Pradesh 277001, India

^c Department of Pharmacology, Korea University (KU), Seoul 02841, South Korea

^d Department of Chemistry, Mahatma Gandhi Central University, Motihari, Bihar, 845401, India

[†] Both the authors contributed equally.

DOI: 10.1039/x0xx00000x



REVIEW




Figure 1. Biologically relevant indole-based heterocycles.

To date, limited reactions have been reported to have easily accessible greener protocols with the formation of efficient yields.⁸¹ Herein, some of the important well-established classical name reactions for its synthesis, such as the Kanematsu indole synthesis,⁸² Mori indole synthesis,⁸³ Buchwald indole synthesis, Bartoli indole synthesis, Bischler indole synthesis, Fischer indole synthesis, Hemetsberger indole synthesis, Julia indole synthesis, Larock indole synthesis, Leimgruber indole synthesis, Madelung indole synthesis, Nenitzescu indole synthesis, Reissert indole synthesis, Fukuyama indole synthesis, Gassman indole synthesis, and Sundberg indole synthesis, etc.⁸⁴ Among these, various protocols lead to the formation of indoles in single-pot and multistep operations (Figure 2).⁸⁵ However, the various classical approaches are well known for the synthesis of indoles via single-step as well as multistep reactions. Among these, some of the reaction conditions afford high yields of the challenging indole derivatives. However, in addition to classical reactions, transition metal-catalyzed protocols have also been developed for indole synthesis, including those performed under microwave-assisted conditions.^{85a,e} In this review, we have attempted to cover the

most significant approaches for the synthesis of indoles from nitroarenes and their analogues. The report encompasses single-pot, multi-step, and multicomponent reactions. Additionally, we have included the synthesis of *N*-hydroxyindoles, owing to their facile transformation into indoles. All these methodologies proceed with the intramolecular annulation, where the five-membered ring is newly generated via the intramolecular as well as the coupling reaction, followed by intermolecular annulation.

2. Diverse Synthetic Approach of the Indoles from the Nitroarenes

2.1. Vicarious Nucleophilic Substitution Approach for Indoles Synthesis via *ortho*-Functionalization of Nitroarenes. In this synthetic process, the nitrobenzene was converted into 2-nitrophenylacetonitrile and 4-nitrophenylacetonitrile (10:1 ratio) with the aid of 2-chloroacetonitrile as a base. This is also called vicarious nucleophilic substitution (VNS) and is involved in the synthesis of indoles *via* the VNS followed by an annulation reaction commonly assigned as Mąkosza Indole synthesis, cf..



REVIEW




Figure 2. Classical Approaches for the Synthesis of Indoles.

Scheme 1.⁸⁶ However, the reductive annulation of the indole's synthesis with the corresponding intermediate 2-nitrobenzonitriles is called as Pschorr–Hoppe indole synthesis.⁸⁷ Furthermore, the isolated crude mixture was employed for the hydrogenation reaction using 10% Pd/C in ethyl acetate. The *ortho*-reduced nitro group undergoes annulation with the nitrile group and then abolishes ammonia *via* the reductive elimination process and produces an indole derivative with an overall two-step reaction yield of 24%.

The following heterocycles are highly important synthons for the synthesis of medicines,⁸⁸ natural products, perfumes, agrochemicals, etc.^{89–94} The synthesis of the 6-(pentafluorosulfanyl)-1*H*-indole from the pentafluoro(3-

nitrophenyl)-λ⁶-sulfane was a chaotic protocol due to the four-step synthesis and cost-ineffective method (Scheme 2a). Herein, a modified methodology has been accomplished for the synthesis of indoles from the pentafluoro(4/3-nitrophenyl)-λ⁶-sulfane within two steps with the help of 2-phenoxy acetonitrile in a basic medium and hydrogenation with the hydrogen gas and Pd/C (Scheme 2b).⁹⁵ Although the overall yield was lower compared to the previous method but this approach remains attractive due to its simplicity, reduced reaction time, and cost-effectiveness.⁹⁶ The reaction of pentafluoro(3-nitrophenyl)-λ⁶-sulfane with 2-phenoxy acetonitrile affords VNS products as the mixture of both *ortho*- and *para*-derivatives in the ratio of (85:15; 73%), whereas the pentafluoro(4-nitrophenyl)-λ⁶-sulfane ends up with only the *ortho*-VNS product with the high




REVIEW

Organic & Biomolecular Chemistry


yield of 74%. Among these, pentafluoro(4-nitrophenyl)- λ^6 -sulfane to 5-(pentafluorosulfanyl)-1*H*-indole leads to a high reaction yield due to the good to excellent yield of each step, cf. Scheme 2.

An alternative strategy for synthesizing 2-aryl 5/6-(pentafluorosulfanyl)indoles^{96, 97} has been explored by replacing the reagent chloromethyl phenyl sulfone instead of 2-phenoxy acetonitrile (Scheme 2b) under a basic medium (Scheme 3).⁹⁵ This approach involves a reduction with Raney Ni


or Pd/C catalysed hydrogenation, followed by imine formation with an aldehyde and annulation reaction leading to the formation of the indole analogues. The choice of base plays a crucial role in determining product yields. Herein, the high stoichiometric amounts of KOH or Cs₂CO₃ mediated annulation in the last step, resulting in the indole yields of 68-80%, and its formation depends on the involvement of the aryl group during the annulation and aromatization reaction (Scheme 3a).



Scheme 1. VNS Reaction Using Chloroacetonitrile Followed by Hydrogenation Approach.




Scheme 2. VNS Reaction Using 2-Phenoxyacetonitrile Followed by Hydrogenation Approach.



Scheme 3. VNS Reaction Using Chloromethyl Phenylsulfone as the Key Step for Indoles Synthesis.





Scheme 4. Higher Indoles Synthesis via the Cascade of Benzyl Fictionalization, Intramolecular Annulation and Deoxygenation Reactions.

Notably, the synthesis involving 4-nitro(pentafluorosulfonyl)benzene, annulation in the fourth step under inert conditions with KOH, resulted in a significantly higher yield compared to reactions conducted under aerobic conditions. Similarly, when benzaldehyde was subjected to annulation using NaOH, the product yields of 77% and 91% were observed under aerobic and inert conditions, respectively (Scheme 3b). These findings underscore the impact of reaction conditions on efficiency and selectivity in the synthesis of pentafluorosulfonyl-substituted indoles.

In the next reaction, the VNS has been fixed by the previous reaction protocol.⁸⁶ Further, it was subjected to the alkylation reaction in step i⁹⁸ to achieve the highly 2-substituted *N*-hydroxyindoles as the end product of step ii, and this process was utilized by Wróbel and Mąkosza in 1997.⁹⁹ During the explorations of step ii, the various basic conditions are employed for the annulation reaction. The NaOH protocol was found to be better than the ester functionality. Finally, the hydroxyindole was subjected to the synthesis of the deoxygenation reaction using the Zn/AcOH protocol under reflux conditions, offering moderate to excellent yields (Scheme 4).¹⁰⁰


The following examples also proceeded with a key step of VNS reactions, which was explored for the synthesis of *N*-hydroxyindoles as the end products. Wróbel and Mąkosza reported that the 4-halo-nitrobenzene on VNS reaction with chloroacetonitrile followed by condensation with ethanal afforded 2-(5-halo-2-nitrophenyl)but-2-enenitrile.¹⁰¹ The annulations in the strong basic media (5M NaOH) in the methanol lead to the formation of the product quinoline *N*-oxide derivatives, and to lowering the concentration of the base (0.1M NaOH) by using the mixture of solvent MeOH:DMSO, the major product was the 5-halo-*N*-hydroxy-2-(hydroxymethyl)-indole-3-carbonitrile derivative. Furthermore, they have made a successful attempt to get the

desired product, 5-halo-*N*-hydroxy-2-methyl-indole-3-carbonitrile, with the help of potassium carbonate in methanol at room temperature (Scheme 5a). Next, the approach of indoles of the tuning of similar indoles with the electron-deficient substituents as constructed at the position-2 of indoles product has been explored in the reaction of dimethyl 2-cyanocyclopropane-1,1-dicarboxylate in the mild basic condition at very low temperature, followed by 0 °C.¹⁰² The limitation of the reactions was found in the transesterification products (methoxyl exchanged with the *t*-butoxyl group) along with the desired products (Scheme 5b). Mechanistically, after the deprotonation of an acidic proton from the cyclopropane ring undergoes *ortho*-addition with nitrobenzene and it offers the sequence of intermediates 1 to 5, which further leads to the formation of *N*-hydroxyindoles derivatives as shown in step ii (Scheme 4). Due to the excess base and butanol, the transesterification products were also formed (Scheme 5c). Furthermore, the hydroxyindoles can easily be converted into the indoles as shown in step iii (Scheme 4).


Wróbel and Mąkosza explored a challenging approach for the synthesis of 7*H*-acenaphtho[1,2-*b*]indole from the 1-(2-nitrophenyl)-1,2-dihydroacenaphthylene using NaOH at ambient temperature.¹⁰⁰ During the reaction, they observed the formation of major products in a 4:1 ratio of 7*H*-acenaphtho[1,2-*b*]indol-7-ol to 7*H*-acenaphtho[1,2-*b*]indole, along with minor side products. To enhance the yield, they treated the reaction mixture with zinc dust and successfully increased the yield of the desired product to 75%. To simplify this process, an alternative route was developed starting from 2-(2-nitrophenyl)acenaphthylen-1(2*H*)-one, which underwent carbonyl reduction and dehydration to yield the key intermediate 1-(2-nitrophenyl)acenaphthylene.⁹⁹ Subsequent treatment with stannous chloride dihydrate resulted in the formation of 7*H*-acenaphtho[1,2-*b*]indole with a 57% yield (Scheme 6).



REVIEW




Scheme 5. Various Approaches for the Synthesis of 2,3-Disubstituted Indoles.




Scheme 6. Synthesis of Fused Indoles.



REVIEW



Scheme 7. VNS Reaction Involved in the Mimicking of Alkaloids.

Scheme 8. VNS Reaction using α -Chloroketone and Reductive-Annulation Followed by Deoxygenation for the *ortho*-Substituted Indoles.

Mąkosza and coworkers (1957) reported a novel approach for the synthesis of indole derivatives to mimic the naturally occurring nitrogen-containing alkaloids batzellines, isobatzellines, damirones, discorhabdins, etc.^{103, 104} They accomplished the indole ring starting from the synthesized 4,6-dinitroguaiacol with the help of 2-phenoxy acetonitrile *via* the VNS approach and a methylation reaction followed by the methylene functionalization on sequential treatment with

dimethyl sulfate and ethyl bromoacetate in basic conditions. Furthermore, the hydrogenation of the achieved product with the palladium chloride/iron reagent system and hydrogen gas in the presence of polar mixed-solvent systems afforded an indole derivative with a yield of 48% (Scheme 7). However, the same reaction condition based on the temperature deviations of two different intermediate products isolated, i.e.,



REVIEW

Organic & Biomolecular Chemistry

intermediate I (0 °C, 7 h) and intermediate II (30 °C, 7 h), was reported with the yield (88%).

Wojciechowski et al. have developed a novel protocol for the synthesis of indoles from the commercially available compound dinitrobenzene.¹⁰⁵ Interestingly, they prepared the hydroxyindoles by the VNS reaction with α -chloroketones in basic media followed by a reduction process using stannous chloride. For the deoxygenation of *N*-hydroxyindoles, lots of reducing reagents are well established. The smartness of the following deoxygenation is that it was reported with α -bromoacetophenone in the presence of the triethylamine (TEA) base as the mild protocol with excellent yields of 83–99% (Scheme 8a). Mechanistically, the nitro-hydroxyindoles undergo protection with α -bromoacetophenone and then further fragment into the desired product indoles and side products as the phenylglyoxal (Scheme 8b).


In 1997, Sutherland reported the unusual behavior of the DBU with methyl 3,5-dinitrobenzoate and trinitrobenzene in chloroform as well as in ethyl acetate at room temperature after two days.¹⁰⁶ It has been reported that, firstly, both the starting materials form Meisenheimer complexes as the intermediates I and II, which can be observed by the change of the deep red color of reaction mixtures. In the next step, these mixtures oxidized to form intermediates III and IV, which later undergo annulation reactions *via* the *ipso*-displacement offering indole analogues. Simultaneously, the intermediate also undergoes a six-membered isoquinolone product on cyclization with the ester group (Scheme 9).

In 2001, Chen and coworkers reported the synthesis of 2-methyl-7-methoxyindole starting from 3-methoxy-2-nitrobenzaldehyde.¹⁰⁷ The reaction involved a cascade sequence of reactions (Henry reaction, condensation reaction,


reductive annulation, and deaminative aromatization). This reaction has also been employed for the synthesis of 6,7-dimethoxy-2-methyl-1*H*-indole and 7-methoxy-2-methyl-1*H*-pyrrolo[2,3-*c*]pyridine from the 3,4-dimethoxy-2-nitrobenzaldehyde and 2-methoxy-3-nitroisonicotinaldehyde, respectively (Scheme 10).

Hossain et al. developed a straightforward two-step strategy for the synthesis of indoles. Each step of the reaction was convenient and associated with high yields.¹⁰⁸ In the first step of the reaction, 2-(2-nitroaryl)-3-hydroxypropenoic acid esters were synthesized on treatment with 2-nitrobenzaldehydes and ethyl diazoacetate in the presence of Lewis acid or Brønsted-Lowry acid, and the product formed by molecular rearrangement. In the final step, indole was achieved as the product after the hydrogenation of the first step associated with the reductive-annulation process (Scheme 11).

Soderberg et al. reported the synthesis of alkyl 2-(2-nitrophenyl)but-2-enoate by succeeding the Kosugi-Migita-Stille coupling and Barluenga coupling reactions, and the intermediate product was subjected to indole synthesis in the basic medium at 0 °C in butanol.¹⁰⁹ The reaction condition was well tolerated, and the library of the compounds was explored by this method and some of the approaches shown in Scheme 12a. Alternatively, the intermediate product ethyl 2-(2-nitrophenyl)but-2-enoate was also synthesized by the reaction between methyl 2-(2-nitrophenyl)acetate and acetaldehyde in a basic medium. During the mechanistic investigation, they found that the mono-substituted double bond must have a methyl group, which undergoes a fragmentation reaction that produces the potassium salt of 3-(methoxycarbonyl)-indol-1-olate, and then the alkylation produces methyl 1-methoxy-1*H*-




Scheme 9. DBU Participating in the Indoles Synthesis *via* VNS Reaction as the Key Step.




Scheme 10. Synthesis of Indoles *via* Henry Condensation Followed by Hydrogenation Reaction.





Scheme 11. Tandem Approach for the Synthesis of Indoles from the 2-nitrobenzaldehydes.



Scheme 12. Synthesis of *N*-Hydroxy/Alkoxyindoles.



indole-3-carboxylate on treatment with methyl iodide, cf. Scheme 12b. However, when the methyl of the vinyl group was substituted with the propyl group, the reaction condition was unable to produce the desired product, indole.

In 2021, Tsukamoto et al. reported the stepwise synthesis of indoles from the 2-nitrobenzaldehyde derivative.^{107, 110} They carried out the transformation through a three-step sequence involving a Henry reaction, dehydration, and amino-reductive annulation (Scheme 13a). The final step of the reaction can have the resemblance of the diamine derivative as the Fischer-Indole

intermediates, which leads to the formation of the five-membered heterocyclic compounds.¹¹¹ Later on, the Henry reaction was also employed for the synthesis of dihydroxy nitrostyrene from the corresponding aldehyde, which was employed for the nitration reaction with the help of tetranitromethane (TNM) with the Zn(II)-assisted condition.¹¹² In the next step, an effective approach was employed for the synthesis of the indole derivative with the help of the Zn-controlled reduction process, which led to the formation of the product with a good yield (Scheme 13b).¹¹³⁻¹¹⁶



REVIEW

Scheme 13. Alternative Approach of Indoles *via* Henry Condensation Followed by Hydrogenation Reaction.

Scheme 14. The Overall Comparative Yields of Gassman and Bartoli Indoles Synthesis from Nitroarenes.

Bartoli and Palmieri (1989) developed a facile and alternative substitute for Gassman indole synthesis.¹¹⁷ In this process, the nitroarenes can be directly converted into the indole derivative in a single step with the aid of vinyl magnesium bromide derivatives.^{118, 119} This method not only reduces reaction time and solvent usage while simplifying the protocol but also delivers excellent yields.^{120, 121} In the following cases, an example of Gassman indole synthesis was shown with a multistep path (Scheme 14a).^{122, 123} Herein, each step afforded a good to excellent yield of the reaction, and the total yield of the reaction was found to be 19%. However, the indole syntheses using 2/3/4-nitrobenzene with 3 moles of vinyl magnesium bromide derivatives (Grignard reagents) offered up to be good (Scheme 14b(i)). In their exploration, they found that the *ortho*-substituted offers a higher yield than the *meta*- and *para*-substituted nitroarenes. Although they achieved a poor yield of the reactions in most of the cases, the reaction

procedures are preferred due to the poor convenience and cheap protocol rather than the Gassman indole synthesis. Furthermore, Bartoli and coworkers also explored the synthesis of indole derivatives by employing the same reagent system with nitrosoarenes and reported moderate yields of the reactions. The substrates also worked well with the broad range of cyclic and acyclic salts of the olefinic magnesium bromides (Scheme 14b(ii)).


To illustrate the mechanism of indole synthesis, 2-nitrotoluene and vinylmagnesium bromide were selected as representative precursors. In which the first mol of vinyl magnesium bromide reacts with the nitro group to transform it to the reduced functional group nitrosobenzene, and then the second mol of vinyl magnesium bromide reacts with the nitroso group to form magnesium *o*-tolyl(vinyl)oxy)amide bromide followed by a [3,3]-sigmatropic shift and intramolecular annulation reaction to




offer magnesium 7-methylindolin-2-olate bromide. Finally, this intermediate reacts with the third mol of vinyl magnesium bromide to furnish magnesium 7-methyl-2-oxidoindolin-1-ide dibromide by the loss of an ethylene group, and then acidic hydrolysis of the di-magnesium bromide salt produces the desired product, such as 7-methyl indoles (Scheme 15).

Independently, the Bartoli indole and Barton-Zard reactions of 9-nitrophenanthrene offer two different types of isomeric products, such as 1*H*-dibenzo[*e,g*]indole and 2*H*-dibenzo[*e,g*]isoindole, cf. Scheme 16a. The reaction and mechanism of the Barton-Zard reaction are depicted in the Schemes 16b.¹²⁴⁻¹²⁷

In 2003, Knochel and his co-workers introduced a novel approach to the synthesis of indole analogues from the *o*- and *o*-nitrostyrenes *via* reaction with the Grignard reagent.¹²⁸ This transformation is recognized as the Knochel-Indole synthesis, an extension of the Bartoli indole reaction, and offers key advantages. Unlike the Bartoli reaction, this approach requires a lower stoichiometric amount of Grignard reagent due to intramolecular annulation, making it more efficient. Beyond facilitating indole synthesis, this methodology also enables the construction of benzimidazoles by selecting appropriate starting materials as both reactions proceed *via* a similar mechanism. In this process, two equivalents of phenyl magnesium chloride are



Scheme 15. Mechanism of the Bartoli Indole Synthesis.





Scheme 16. Synthesis of Dibenzo[*e,g*]indole/isoindole from the 9-Nitrophenanthrene and the Mechanistic Details of Barton-Zard Reaction.



REVIEW

Organic & Biomolecular Chemistry


View Article Online
DOI: 10.1039/D5OB00338E**Scheme 17.** Knochel-Indole Reactions in the Synthesis of Indoles and Benzimidazoles.

sufficient: the first generates the β -aryl *o*-nitrostyrene, while the second reduces it to form a nitrene intermediate. The resulting nitrene undergoes annulation, and subsequent aromatization generates either indoles or benzimidazoles as the final cyclized heterocyclic products (Scheme 17). This innovative approach broadens synthetic access to structurally numerous nitrogen-containing heterocycles.

2.2. Involvement of an *Ortho*-Group in the Synthesis of Nitroarenes to Indoles.

Batcho and Leimgruber have discovered a novel approach for the synthesis of indole¹²⁹ and their analogues from the corresponding 2-nitrotoluenes *via* the condensation of *N,N*-dimethylformamide dimethyl acetal in DMF at high temperature followed by hydrogenation with the aid of Pd/C in benzene *via* reductive-annulation reaction (Scheme 18a).¹³⁰


However, they also varied the secondary amines of formamide dimethyl acetal and found good to excellent yields of indoles. The other hydrogenation protocol was also explored with the Fe/AcOH and sodium dithionite, which afforded a poorer yield than the Pd/C. After the successful establishment of these reactions, various substrate scopes were explored in using this method. Hence, the reaction becomes more popular as the Batcho-Leimgruber indole reaction for the indole synthesis. Similarly, Ponticello and Baldwin have reported a common synthon, such as methyl indole-4-carboxylate, for the synthesis of alkaloids starting from the 2-methyl-3-nitrobenzoic acid in excellent yields of each step (Scheme 18b). Recently, Ley et al. modified the Leimgruber-Batcho enamine intermediate product synthesis by employing the microwave condition, and then they were subjected to the indole synthesis.¹³¹ This reaction protocol is a superior protocol to the previous in regards to the time and yields of the indoles (Scheme 18c).¹³²

**Scheme 18.** Batcho-Leimgruber Indoles Synthesis from 2-Nitrotoluenes.


REVIEW

Cadogan-Sundberg indole synthesis. Cadogan and Sundberg developed the progress of the direct indole synthesis with the reaction of nitrostyrene with triethyl phosphite or triphenylphosphine (Scheme 19a). Herein, the triethyl phosphite behaves as a deoxygenating reagent and transforms it into the nitroso derivative that undergoes a 6-electron thermal ring closure reaction and provides the aromatized indoles (Scheme 19b).¹³³⁻¹³⁷

Simultaneously, Sundberg, et al. also reported the deoxygenation-based annulation of *ortho*-nitrostyrenes to corresponding indoles. Furthermore, they also studied the terminal disubstituted 2-nitrostyrenes to corresponding indoles and proposed the various types of side products along with the major product (Scheme 20).¹³⁸⁻¹⁴⁰



Scheme 19. Reactions and Mechanism by the Cadogan and Coworkers.




Scheme 20. Sundberg Analysis in the Formation of Indoles Products and Side-Products.



REVIEW

Organic & Biomolecular Chemistry




Scheme 21. The Analysis of the Cadogan-Sundberg Indoles Reaction for the Information of *N*-Alkoxyindoles.

The combined reaction above is known as the Cadogan-Sundberg indole synthesis. Additionally, Peet et al. have studied that the reaction of 2-nitrostilbenes with triethyl phosphite leads to the formation of two types of major products, such as 2-arylindoles and the corresponding 2-aryl-*N*-ethoxyindoles.¹⁴¹ They also explored the mechanistic deoxidative investigation to deeply understand the formation of *N*-alkoxyindoles by using isotopically labeled ¹⁸O of the nitrostyrenes and found that the reactions are proceeding with the adopting of both the mechanistic paths as shown in Scheme 21.

Interestingly, Söderberg and coworkers have utilized a variety of palladium catalysts in combination with organic phosphines and nitrogenous organic ligands and developed modified approaches¹⁴² for the synthesis of indoles in the presence of CO (6 atm.). Herein, two important examples of the indole alkaloids noted as Hyrtinadine A and Fargesine were reported by the Söderbergs' group (Scheme 22a-b).^{143, 144} In 2003, Kueth and coworkers reported the synthesis of indoles from the nitrostyrenes in the presence of triethyl phosphite (TEP) at very high temperatures during the exploration of the total synthesis


of Tjipanazoles B, D, E, and I.¹⁴⁵ Instantaneously, they also explored the same target by modifying the protocol using Pd(OAc)₂/PPh₃ reagent conditions in a carbon monoxide environment at 6 atm in a refluxed condition. The latter approach of indoles afforded a superior yield with higher purity than the TEP condition (Scheme 22c).¹⁴⁶ The limitations of the CO atmosphere supported protocol offering the CO insertion product that leads to the formation of amides or lactams derivatives instead of the desired product as indoles. As a result, the yield of the product is dropping below than the expected yield of the indoles.

In the following successful attempt, Hamze et al. developed an excellent route for 2-arylindoles by treating equal stoichiometric amounts of *N*-tosylhydrazones with 2-nitrobenzyl bromide using Pd₂(dba)₃·CHCl₃ in a basic medium with P(2-furyl) generating the *ortho*-nitrostilbene intermediate followed by Cadogan annulation in the presence of an excessive phosphine (Scheme 23).¹⁴⁷




Scheme 22. Modified Approach of Cadogan-Sundberg Cyclization for the Syntheses of Indoles.





Scheme 23. The Synthesis of 2-Arylindoles from *N*-tosylhydrazones and 2-Nitrobenzyl.




Scheme 24. The Synthetic Approach of Indoles Using Pd(II) Reagents and Additives from 2-Nitroarenes.

In the following serendipitous examples, the indoles were explored using the Pd(II) catalysts with good to excellent yields. In Scheme 24a,¹⁴⁸ the reaction was attempted for the methoxycarbonylation reaction, but it offered the cyclized product indoles. Similarly, Scheme 24b was fully employed for Heck reactions to synthesize nitrostyrene analogues. However, this protocol was not efficient for the synthesis of indoles, but it worked within some of the limited scopes. Next, the Stille reaction was also attempted for the synthesis of carbonylated products, but the reaction led to the formation of indole derivatives.¹⁴⁹ A general reaction and mechanism of the Pd(II)-catalyzed reaction is shown with the cross experiments in Scheme 25. Similarly, Scheme 24c and Scheme 24d were

directly employed for the indoles with good to excellent yields of the reactions without any base.¹⁵⁰⁻¹⁵²

During the mechanistic investigations, it was reported that the terminally disubstituted and unsubstituted *o*-nitrostyrene on reaction with the above Pd(II) protocol, forms nitrene intermediates.¹⁵¹ Noticeably, the terminal disubstituted olefins undergo an annulation reaction that furnishes 2,3-disubstituted indoles *via* a cascade of steps such as [1,5]-sigmatropic shift and aromatization reaction. Whereas the *o*-nitrostyrene undergoes an insertion reaction *via* rearrangement leading to the formation of indoles. Furthermore, the experiment was confirmed by the deuterium labelling experiments (Scheme 25).




Scheme 25. Cross Experiments and Mechanism of the Formation of the Indoles.




REVIEW

In the following examples, the Cadogan–Sundberg strategy was successfully transformed into indoles with the help of catalytic mol% of the transition metals of Fe(II), Cu(II), and Pd(II), etc.^{143, 153, 154} by the combination of organic nitrogenous heterocyclic ligands and reducing silane reagents or surrogating equivalents of CO.^{155, 156} The combination of the approaches afforded to excellent yield of the product in all cases (Scheme 26a-d).¹⁵⁷ However, some other cobonylating catalysts such as Fe(CO)₅, Ru₃(CO)₁₂, and Rh₆(CO)₁₆ were also well-known for the indoles synthesis from the nitrostyrenes, but it led to the formation of the amines and other incredible products.¹⁵⁸ Thus, these procedures are less frequent for the synthesis of indoles.

In 2019, Driver and coworkers developed a multi-chambered reduction process for reductive transformations of the *ortho*-nitrostyrenes to indoles. The reaction conditions are slightly chaotic with a tedious protocol.¹⁵⁹ However, the reaction conditions are well tolerated for most of the functional groups, and chemoselectively nitro group, after the reduction process, forms the heterocyclic compound indoles with excellent yields (Scheme 27).



Scheme 26. Indoles Synthesis with the Help of Reducing Reagents or CO-Surrogating Reagents.



Scheme 27. Single-Pot Three-Chambered Reductive-Annulation of *ortho*-Nitrostyrenes to Indoles.

In further approaches, the oldest mode of the Cadogan annulation approach using TEP was replaced by triphenylphosphine at mild to severe conditions also explored.^{160a-163} However, many other approaches were reported for the improvement of the indoles in the presence of the additive of the salts/complexes of the transition metals, etc., with the PPh₃. Herein, the indoles were synthesized in two

steps, such as hydroarylation and Mo-catalyzed Cadogan annulation. In both of the approaches, the indoles were synthesized with good to excellent yields (Scheme 28a).^{160a, 134} Herein, we followed a stereospecific *syn*-addition approach similar to the hydroboration reaction as established by Yamamoto et al. (Scheme 28b).^{160b} The reaction proceeds in the presence of a copper catalyst and undergoes an exchange




reaction to form complex I. This is followed by coordination with an aryl boronic acid to form complex III via intermediate complex II. Subsequently, a stereospecific *syn*-addition with the alkyne yields compound V. Rapid regeneration of the catalytic complex occurs in the presence of methanol, affording the major product VI. In the subsequent sequence (Scheme 28c),^{160c} MoO₂Cl₂(DMF)₂ undergoes an exchange reaction with PPh₃, followed by coordination with a nitrostyrene analogue through intermediates VII and VIII, to yield complex IX. The nitro group facilitates the formation of a strained four-membered intermediate X by displacing *N,N*-dimethylformamide (DMF). This intermediate is then transformed into complex XI, which subsequently generates the catalyst precursor XII and finally the active catalytic species VII. Throughout this sequence, the formation of a nitro-reduced product XIII is observed, arising via a cascade of exchange reactions involving DMF and complexes

XI and XII. Ultimately, these transformations conclude in an intramolecular annulation reaction to furnish the corresponding indole derivatives, consistent with earlier transformations described in Schemes 19–20.


In 2015, Nelson and coworkers modified the Cadogan-Sundberg annulation with the help of stoichiometric amounts of triphenyl phosphine and the catalytic mol% of molybdenum dichloride dioxide bis(*N,N*-dimethylformamide) [MoO₂Cl₂(DMF)₂] in the microwave (MW) condition during the synthetic exploration of eumelanin-inspired polymer from vanillin.^{164, 165} The reported reaction offered a good yield (Scheme 29).^{166, 167} The mechanistic approach of the reaction is similar to Scheme 28c, and the applied microwave condition reduces the time tenure of the reaction with improved yield of the desired product.

a) Reaction methodologies




b) Stereospecific addition

ref. *Chem. Commun.* 2008, **44**, 2010–2012



c) Mechanism of Indoles synthesis

ref. *Adv. Synth. Catal.* 2023, **365**, 3155–3161




Scheme 28. Molybdenum Catalysts-Assisted Indoles Synthesis.



REVIEW

Organic & Biomolecular Chemistry



View Article Online
DOI: 10.1039/D5OB00338E


Scheme 29. Microwave-Assisted Cadogan–Sundberg Cyclization.

In 1869, Baeyer and Emmerling reported a novel approach for indole synthesis using iron metal from *ortho*-nitrocinnamic acid in strongly alkaline conditions, illustrated in Scheme 30a.¹⁶⁸ The reaction proceeds *via* the reductive annulation, followed by the decarboxylation process is commonly known as Baeyer–Emmerling indole synthesis.¹⁶⁹ During the exploration of the reaction, the nitro functionality is converted into the corresponding nitroso group, which plays a crucial role in the formation of the five-membered ring (Scheme 30b).¹⁷⁰

In the 19th century, Arnold Reiesert developed two-step indole synthesis from *ortho*-nitrotoluenes, and it was a hot topic for chemists. In this procedure, various types of *ortho*-nitrotoluenes are subjected to indole syntheses with the help of diethyl oxalate in the presence of sodium metal followed by reduction with Zinc/Acetic acid leads to the formation of an amino-phenylpyruvic acid intermediate. Furthermore, this intermediate undergoes a cyclization reaction followed by the decarboxylation process that produces an indole carboxylic acid derivative. This procedure afforded a good to excellent yield of the indole product, and this procedure is commonly known as the Reiesert indole synthesis.^{171–174} The other alternative


method for the hydrogenation methods of the nitrophenylpyruvic acid intermediate was also explored with the help of ferrous sulphate/ammonium hydroxide, stannous chloride/dehydrate, and Pd/C that also gives the desired product as indoles. The mechanistic approach of these reactions is also shown in Scheme 31.^{175, 176}

In the following examples, the potassium metal was dissolved into the absolute ethanol (without stirring) to prepare the potassium ethoxide base, and then it was mixed with the solution of diethyl oxalate in the inert atmosphere at room temperature. Further, the nitrotoluene was also added for the synthesis of the potassium salt of ethyl *o*-nitrophenylpyruvate, which was later subjected to the hydrogenation reaction at high pressure using hydrogen gas in the presence of the catalytic amount of platinum(IV) oxide in acetic acid. The yield of the product varied based on the purification methods through several washes. The synthesis of indoles *via* the Arnold Reiesert method is a challenging and hectic procedure; thus, it was replaced by many other alternative synthetic methods (Scheme 32).¹⁷²




Scheme 30. Synthesis of Indoles from the 2-Nitrocinnamic Acid.






Scheme 31. Synthesis of Indole *via* Reissert Indole Synthesis.



Scheme 32. Modified Synthesis of Indole *via* Reissert Indole Synthesis.

Independently, Nagashima group and Chung group reported the indole synthesis *via* the hydrogenation reaction of 2-(2-nitrophenyl)acetonitrile using the [Pt]@SiC₆ and Co-Rh heterobimetallic nanoparticles under hydrogen of 10 atm and 1 atm pressures in ethyl acetate and wet methanol, respectively (Scheme 33).¹⁷⁷⁻¹⁷⁹

In the next reaction, low-valent titanium(III) chloride was used to look into *ortho*-nitrostyrene analogues. Herein, the titanium acts as a reducing reagent for the nitro group and transforms it into the corresponding nitroso derivatives.¹⁸⁰ Further, it cyclized into the corresponding product hydroxyindoles *via* path a, and later it reduced to indole derivatives due to the presence of an excessive amount of valent titanium(III) chloride (Scheme 34a). However, the trisubstituted olefins of *ortho*-nitrostyrene analogues also proceed with the same mode as path a and the 1,2-shift *via* path b, which further undergo rearrangement followed by aromatization and deoxygenation reactions that



Scheme 33. Nitrile Hydrogenations in Indoles Synthesis.




finish 2,3-disubstituted indoles as the major product (Scheme 34b). The mechanistic detailed information is shown in Scheme 34c.

In 2019, Song et al. established the development of 3-amino/aminoalkyl-2-keto-indole analogs from *ortho*-nitrochalcones.¹⁸¹ They treated the corresponding chalcones with ammonia and Hantzsch ester in the basic medium leading to the formation of the 3-amino-indoles (Scheme 35a). Interestingly, the methodology was also employed for the synthesis of COX-2 and tubulin polymerization inhibitors. Furthermore, the ammonia was substituted by primary amines to perform the successful synthesis of 3-aminoalkyl-indoles with a yield of 27-86% (Scheme 35b).



REVIEW

Organic & Biomolecular Chemistry

View Article Online
DOI: 10.1039/D5OB00338E**Scheme 34.** Titanium(III)-Supported Intramolecular Reductive-Annulation of Indoles Synthesis.**Scheme 35.** Synthesis of Indoles from the Chalcones.**Scheme 36.** Synthesis of *N*-Hydroxyindoles from 3-(2-Nitrophenyl)acrylic Acid.

According to recent studies, it was reported that stannous chloride was utilized to reduce the 3-(2-nitrophenyl)acrylic acid to alkyl 3-(2-nitrosophenyl)acrylate intermediate, which further undergoes annulation with the double bond of acrylate, followed by an aromatization process that produces alkyl 1-hydroxyindole carboxylates as the major product. At the same time, annulation with the lone pair of nitro with the carbonyls of the ester gives a 2-hydroxyquinoline derivative as the side product (Scheme 36).¹⁸²

Song et al. reported a novel approach for the synthesis of indoles from 3-(2-nitrophenyl)acrylic acids and stilbenes with the help of bis(pinacolato)diboron (B₂Pin₂) and KF in an ethanolic solution (Scheme 37a-b).¹⁸³ Virtually, the bis(pinacolato)diboron reagent does the deoxygenation of nitro-group and shows it in the form of a nitroso derivative. Later, it again reacts with another mole of bis(pinacolato)diboron, and then annulation followed by the [1,5]-sigmatropic shift affording the indoles (Scheme 37c).




The following alkynes can easily be prepared by the most frequent reaction, i.e., Sonogashira coupling, and then the obtained starting materials were further treated with the metals (In and Zn) in activation with the additives of HI/Aliquat, dibromoethane.¹⁸⁴ Remarkably, both metals were known for the reduction of the selective functionality *via* the electron transfer mechanism. Herein, the nitro group is reduced by the metals, and the salts are formed during these reactions, activating the triple bond for the annulation reaction, which leads to the formation of the heterocyclic compound indoles (Scheme 38).¹⁸⁵

Oh and Shin have reported a tandem reaction for the synthesis of 1-hydroxyindoles from the *ortho*-iodonitrobenzene.¹⁸⁶ Herein, these reactions were performed *via* a cascade of reactions, i.e., Sonogashira coupling, selective partial reduction, followed by the Pd(II)-catalyzed electrophilic annulation reaction affording the hydroxyindole with an overall moderate to good yield of the reaction (Scheme 39).¹⁸⁷


In 2021, Ying et al. have vastly developed a novel approach for the 1-aryl-2-arylindoles through the coupling reaction between disubstituted *ortho*-nitroaryl acetylene with iodoarenes in the presence of Zn/ZnI₂/Co₂(CO)₈ and the catalytic amount of Ni(II) salt and 4,4'-Di-*tert*-butyl-2,2'-dipyridyl (dtbbpy) ligand into the DMF at 120 °C (Scheme 40a).¹⁸⁸ In 2022, the same group modified the previous protocol

by replacing the iodoarenes, ZnI₂, and Ni(dme)Cl₂ with aryl boronate esters, TMSCl, and Ni(OTf)₂ by changing their corresponding stoichiometric amounts and elevating the temperature, respectively (Scheme 40b).¹⁸⁹ The later improved protocol afforded better yields of the indoles than the previous one, and both reactions were well tolerated for the variety of the functionality of the substrates. They have proposed the mechanism of this coupling process by using a Ni(II) catalyst by generating the Ni(0) species by using dtbbp and Zn/ZnI₂. Next, the Ni(0) species undergo oxidative insertion with iodoarene, followed by carbonylation *via* in situ-generated Co₂(CO)₈ reagent that forms active species A to B. At the same time, chemoselective reduction of the nitroalkynes generates the aniline derivative due to the presence reducing agent Zn/ZnI₂ and then immediately reacts with in situ generated reactive species B and generating an intermediate product C. Finally, it undergoes an annulation reaction that produces the corresponding *N*-aryl indole with the aid of Co₂(CO)₈, zinc powder, and ZnI₂ (Scheme 40c).¹⁸⁸

Herein, the cobalt nitrogen-doped carbon-900 (Co-N-C-900) catalyst was accompanied for the in situ hydrogen transfer reaction.¹⁹⁰ The set of reactions proceeds with the redox process, in which the dihydroindoles are reduced with the nitro group as the reactions are involved in the Skraup reaction during the quinoline synthesis depicted in Scheme 41.



Scheme 37. Reductive Annulation of Nitrostyrenes to Indoles using B₂Pin₂ and KF Reagents.






Scheme 38. Synthesis of Indoles Using In and Zn metals for Intramolecular Reductive-Annulation Reaction.



REVIEW

Organic & Biomolecular Chemistry

Scheme 39. Synthesis of *N*-Hydroxyindoles via Pd(II)-Catalyzed Intramolecular Annulation Reaction.Scheme 40. Ni(II) Catalyzed Synthesis of *N*-Aroylindoles from the 2-Nitroalkynes.

Scheme 41. Oxidation of Dihydroindoles with Nitrobenzene.

Cycloaddition Approach.

In 2002, K. M. Nicholas and his group developed a novel approach for indole synthesis in reaction with nitroarenes and alkenes in the presence of $[\text{Cp}^*\text{Ru}(\text{CO})_2]_2$ and carbon monoxide atm. at 750 psi (Scheme 42a).¹⁹¹ This developed procedure also had some limitations, such as regioselectivity and solvent selection, which resulted in inferior yields. At the same time, his group also did the mechanistic investigation by additionally adding the corresponding nitrosoarene in the same reaction


mixture and reported a superior yield of the indole. Later, they explored indoles from nitrosoarenes with alkynes and found that the current procedure of the synthesis gives higher yields than the previous one.¹⁹² Ragaini et al. utilized the palladium-phenanthroline complexes to get the indole derivative similarly at 60 bar pressure and synthesized the library of the compounds with a yield of 8-45% (Scheme 42b).^{193a} However, the exact mechanism is unclear. While mechanistic studies were being attempted, it was also found that the nitro group converted into the nitroso derivative and then underwent an addition reaction




with alkynes to furnish the indoles as the major products. During the controlled experiment, they were observed that the absence of either catalyst or carbon monoxide does not favour the reaction in the forward direction. Based on their controlled experiments, they also assumed that the interaction between the nitrosoarene (it might be generated from the reduction of nitroarene with CO) and the alkyne takes place outside the metal's coordination sphere, as suggested by Nicholas and co-workers (Scheme 42c)¹⁹¹⁻¹⁹². The regioselective addition found with alkenes and nitrosoarenes and generating dipolar α -styryl cations (theoretical support^{193b}), which also can generate the α -styryl radicals (EPR-support^{193c}). Further, it affords the indoles via intramolecular annulation followed by carbon monoxide deoxygenation (Scheme 42c, path A). At the same an alternative path was also found that leads to the formation of azoxyarenes

via self-dimerization in the presence of carbon monoxide. We assume that this can be an insightful and selective synthetic approach for the indole analogues.

Mancini et al. introduce a different route for the synthesis of indoles via the cycloaddition approach than the previous reports without any kind of metal catalysts.¹⁹⁴ In this work, they elaborated on the reactions between the different types of 1-tosyl-nitropyrrole (electrophilic dienophile) with 1-methoxy-3-trimethylsilyloxy-1,3-butadiene (nucleophilic Danishefsky's diene) in the non-polar benzene at 120-140 °C (Schemes 43a-b). Noticeably, the 1-tosyl-2-nitropyrroles on cycloaddition with Danishefsky's diene, followed by the nitrous acid elimination



Scheme 42. Cycloaddition Approach for the Indoles Synthesis.



Scheme 43. Synthesis of Indoles from Electrophilic Dienophiles with Danishefsky's Dienes




REVIEW

and desilylation, produced corresponding 5-hydroxyindoles, whereas 6-hydroxyindoles were formed by the 1-tosyl-3-nitropyrroles due to the cycloaddition of the electron-demanding rules.¹⁹⁵ However, this work was performed with the isoprene and *N*-(buta-1,3-dien-1-yl)-*N*-propylacetamide, which also produces the annulation product with poor yields. Furthermore, the reaction was explored by the same group as indole products in microwave-assisted with and without solvent conditions, where they found a higher yield than the previous reaction protocols (Scheme 43c-d). Similarly, the reactions were also performed with two more types of nucleophilic dienes, such as isoprene and 1-trimethylsilyl-1,3-butadiene, which also afforded poor yields of *N*-tosylindoles (for the preparation of the [HMIM][BF₄], [HMIM][PF₆], and [NEA] solvents according to the previously reported methods.¹⁹⁶

Peters and his coworkers reported an electrochemical reductive approach¹⁹⁷ for indole synthesis at carbon cathodes in deoxygenated *N,N*-dimethylformamide using electrolyte as tetramethylammonium tetrafluoroborate and an excessive amount of proton donor source as phenol or methyl 3-oxobutanoate at room temperature with yields of 35-82% (Scheme 44).¹⁹⁸ For the justification of the indoles, it was believed that the reaction proceeded with the nitrene intermediate as formed in the case of the Soderberg approach. In the following reaction, the synthesis of 2-phenyl benzindole phenol offered a poor yield of 35%. Hence, it was replaced by the same equivalents of a strong acidic proton source as methyl 3-oxobutanoate then it furnishing a superior yield of 65% than the prior protocol.¹⁹⁹

Electrochemical Approach.



Scheme 44. Electrochemical Approach for the Synthesis of Indoles



Photochemical Approach.

Wang et al. did the synthesis of indoles from the *ortho*-nitrohomobenzyl alcohols by employing the diborane and DIPEA at 400 nm in blue LEDs (Scheme 45a). Herein, the nitro- and hydroxyl-groups undergo intramolecular redox reactions in the presence of blue LEDs to form 2-(2-nitrosophenyl)acetaldehyde, and then it is reduced with the diborane derivative to form 2-(2-(hydroxyamino)phenyl)acetaldehyde.²⁰⁰ Further, it cyclized to form the indoline-1,2-diol intermediate, which later proceeded with base-mediated dehydration and diborane-mediated deoxygenation and aromatization reactions as the indoles (Scheme 45b).²⁰¹

Waheed and his coworkers successfully attempted the photoinduced synthesis of indoles from nitrostyrenes with the help of the Knölker catalyst and the 4-tolylmethanol in a basic medium at rt.^{202, 203} This synthetic procedure was employed for the exploration of the variety of indoles in good to excellent yields (Scheme 46a). They have proposed the mechanism of reaction, where the Knölker catalyst plays a significant role in the reduction of the nitro group to nitroso and then hydroxyindoles to indole derivatives. During these reactions, 4-tolylmethanol acts as a reducing agent of the Knölker catalyst during the complete set of reactions. The formation of the intermediate nitroso and hydroxyindoles was confirmed by the HRMS data analysis (Scheme 46b).



REVIEW

**Scheme 45.** Synthesis of Indoles from *ortho*-Nitrohomobenzyl Alcohols.**Scheme 46.** Photochemical Synthesis of Indoles from Nitrostyrenes Using Knölker Catalyst.

Dayong Shi and coworkers explored the novel photochemical approach of Bartoli-Indole synthesis on the reaction between nitroarenes with gem-methylstyrene in the mild acidic medium with inert atm.²⁰⁴ by applying 390 nm of the intensity of light in

ethyl acetate. The reaction worked well for a wide range of substrates having halogens, nitriles, acids, esters, etc. (Scheme 47a). They have proposed the mechanism that the nitroarene produces diradical species *via* the ISC process employing the




REVIEW




Organic & Biomolecular Chemistry

390 nm wavelength of light, which forms a [3+2]-cycloadduct with methylstyrene, and then it produces a dipolar intermediate *via* a favoured polar ring fragmentation reaction. Next, this dipolar species forms the 2-(2-aminophenyl)acetophenone intermediate *via* the cascade of hydride and [3,3]-sigmatropic shifts. Finally, 2-(2-aminophenyl)acetophenone forms the desired product on the dehydration reaction in the presence of acetic acid (Scheme 47b).

In 2024, Studer and his group developed an excellent approach for the synthesis of *N*-alkylated indoles *via* photochemical [3+2]-cycloaddition reaction.²⁰⁵ To demonstrate [3+2]-annulation, they were chosen as the nitroarenes and vinyl ketones/esters as the 3- and 2-skeleton constituents, respectively. The nitroarenes are activated in combination with vinyl ketones/esters using the iridium-based photocatalyst (PC) to initiate a chain reaction of the radical annulation process while



Scheme 47. Photoinduced Bartoli-Indoles Synthesis.



Scheme 48. Photocatalyst-Induced Reaction of Nitroarenes with Vinylketones/esters Indoles Synthesis.



REVIEW

retaining the light intensity of 455 nm with the help of a 5-watt light-emitting diode (LED) at the ambient temperature. The developed reaction condition was well tolerated for the synthesis of myriads of indoles and afforded moderate to good yields of 21 to 80%. In some cases, the reaction suffered from regiochemical annulation, which dropped the formation of desired outcomes. This work also had good enough novelty in the direct synthesis of pharmacophores and natural product components with a combination of indoles such as estrone, L-(-)-menthol, diacetone-D-glucose, cholesterol, and β -citronellol with yields of 52-66% (Scheme 48).

Conclusions

This review describes the classical to modern approach of the indoles synthesis using the nitroarenes and through its precursor routes. The chemists are putting their interests due to the high bioactivity potential of indoles. Importantly, the various synthons were utilized with commercial nitroarenes for the VNS reaction, which were treated as precursor intermediates of indoles. Also, added the synthetic approach of hydroxyindoles, to further transformed into the desired indoles by easily accessible reduction methods. During the classical approach, most of the indole analogues were synthesized from *ortho*-substituted nitroarenes, such as Cadogan, Bartoli, Knochel, Batcho–Leimgruber, etc., and also without *ortho*-substituted arene as Bartoli indole synthesis. Furthermore, several other recent approaches have been developed, where the indoles synthesis are conducted using transition metal catalysts in diverse modified advanced protocols to make the convenient, and easily available source to develop cost-effective synthesis such as palladium, molybdenum, nickel, zinc, etc. Furthermore, the novel photochemical and electrochemical approaches were also making it promising candidates in the progression of indole synthesis. We assume that this study can provide a better understanding in the synthesis of indole derived natural products, drugs, and complex heterocyclic molecules. Thus, our study can play a significant role in both academic and research.

Future Aspects. Herein, various approaches based on the cascade, tandem, and multistep reactions have been studied. We assume that this study can provide ample knowledge for the construction of indoles and can provide a better platform in its development.

1. Additionally, the VNS approach has been added to develop the precursor of the indoles, which can play an important role in the synthesis substituted desired indole derivatives.

2. The N-hydroxyindoles are also added to get an alternative way, which can easily be helpful in its via selective reduction methods. In contrast, classical indole syntheses such as the Fischer, Bartoli, or Gassman approaches offer more predictable regioselectivity and often proceed in fewer steps, albeit sometimes with less functional group tolerance.

3. Despite the vast progress in indole chemistry, the chiral approach remains underexplored, highlighting the need for the development of catalytic as well as enantioselective methods.

4. These methods will not only allow the creation of indoles, but can also provide the platform for the easy synthesis of indole derivatives like carbazoles, indolocarbazoles, diphenyl indoles, 1,2-disubstituted indoles, etc.

5. This study provides the approaches of single-pot one-shot as well as multistep reactions and tandem synthetic approaches for various indoles.

6. This study affords the direct and sustainable routes from nitroarenes/*ortho*-substituted nitroarenes to indoles by leveraging photochemical, electrochemical, and metal-free catalysis. Additionally, expanding the substrate scope to electron-deficient and sterically hindered nitroarenes, improving the step economy, and integrating continuous flow processes may address current limitations and establish nitroarene-based approaches as mainstream strategies for indole synthesis.

7. Recent advancements of photochemical strategies enabling more direct access to indoles from nitroarenes, often bypassing the intermediate aniline and focusing on such emerging methodologies to provide a critical comparison with traditional multistep routes. Recent advancements have introduced one-pot, tandem, and photochemical strategies enabling more direct access, often bypassing the aniline intermediate.

8. The nitroarenes/*ortho*-substituted nitroarenes have moderate to excellent tolerance against the acid as well as base during the synthesis of target molecules. Hence, these methodologies can provide a great baseline in the development of several bioactive indoles in pharma and chemical industries in future aspects.

Conflicts of interest

All the authors declared that there are no conflicts to declaration.

Acknowledgments

Individually, we all appreciate and thank our respective departments of the current institutions for giving us the great



REVIEW

opportunity as the funding and infrastructures to work as the higher post to explore the field of education and research.

Dr. Ajeet Chandra has completed his doctoral degree in oxidative transformation using hypervalent iodine (V) reagents. He completed his BSc and MSc programs from the TDPG Collage, Jaunpur, in affiliation with VBSPU Jaunpur and the University of Allahabad, Prayagraj, Uttar Pradesh (UP), respectively. In pursuit of his research interests in research, he was enrolled in his doctoral program at IIT Delhi. Due to his interest in pure organic synthesis, he came to IIT Kanpur and completed his doctoral program under the supervision of Prof. J. N. Moorthy. Later, he joined as an institute postdoctoral fellow at IIT Bombay and explored the sulfonyl phthalides and asymmetric amines. Recently, he has been working as a research professor in the department of information display at Kyung Hee University, Seoul, South Korea. With his interest in organic synthesis, he is exploring organic materials for device applications in the orientation of commercialization purposes for real display applications. My sincere, gratitude to my thesis supervisor for providing me a significant background in research to explore my knowledge in education and research.



Dr. Suresh C. Yadav is working as an assistant professor in the Department of Chemistry at Satish Chandra College, Ballia, India. He did his master's and doctoral programs from the Department of Chemistry, TDPG College Jaunpur, and the University of Allahabad, respectively. He defended his thesis in the area of oxidative reaction kinetics and synthetic transformations with the help of transition metals and an excessive amount of oxidizing reagents under the supervision of Prof. PK Tandon. Dr. Yadav is highly dedicated to teaching and research education and did the reaction kinetics as well as the mechanistic investigations of the transition metal-catalysed reactions.



Dr. Subba Rao Cheekatla, born in Bobbillanka village, Andhra Pradesh, India, earned his master's degree in organic chemistry from Andhra University. He completed his PhD in synthetic organic chemistry at IIT Bombay, Mumbai, under Prof. Sambasivarao Kotha. Subba worked as a research associate at IIT Palakkad before joining Korea University College of Medicine as a postdoctoral fellow under the supervision of Prof. Jun-Seok Lee. His research focuses on diverse heterocycles, cage hydrocarbons, aryl/heteroaryl fused macrocycles, and designing fluorescent probes for chemical proteomics and biological applications



Dr. Abhijeet Kumar obtained his B.Sc. (Hons.) and M.Sc. from Banaras Hindu University (BHU), India. Later he moved to R&D (API) division to get the industrial experience at Dr. Reddy's Laboratory, Hyderabad. After obtaining industrial exposure and experience, he moved to IIT Kanpur to do the doctoral program under the supervision of Prof. M. L. N. Rao. Where, he developed metal catalysed various efficient protocol for the cross-coupling reactions to perform the synthesis of important heterocyclic molecules. Further, he joined the faculty position for a short tenure at the National Institute of Technology, Raipur, and then shifted to the Department of Chemistry, Mahatma Gandhi Central University in 2016 as an assistant professor. Presently, he has been involved in both teaching and research along with non-professional work. Currently, he is teaching of common chemistry and green chemistry as elective papers. In the past, he was published various research and review articles in the reputed national and international journals.



References

1. E. Kabir and M. Uzzaman, A review on biological and medicinal impact of heterocyclic compounds, *Results Chem.*, 2022, **4**, 100606.
2. B. Yu and X. Yang, Why are heterocycles so special in medicinal chemistry?, *Chem. Biol. Drug Des.*, 2022, **100**, 763-764.
3. I. Pibiri, Recent Advances: Heterocycles in Drugs and Drug Discovery, *Int. J. Mol. Sci.*, 2024, **25**, 9503.
4. P. H. M, Zabiulla and S. A. Khanum, An Overview of the Synthetic Routes and Pharmacological Aspects of Pyridine, Isoxazole, Thiazole, and Indole Derivatives, *Polycycl. Aromat. Compd.*, 2024, **44**, 7168-7196.
5. S. Kotha and S. R. Cheekatla, DESIGN AND SYNTHESIS OF PENTACYCLOUNDECANE CAGE COMPOUND CONTAINING OXAZOLE MOIETY, *Heterocycles : an international journal for reviews and communications in heterocyclic chemistry*, 2020, **100**, 1623-1632.
6. S. Kotha and S. R. Cheekatla, Synthesis of Bisoxazole and Bromo-substituted Aryloxazoles, *Molbank*, 2022, **2022**, M1440.
7. É. Frank and G. Szöllösi, Nitrogen-Containing Heterocycles as Significant Molecular Scaffolds for Medicinal and Other Applications, *Molecules*, 2021, **26**, 4617.
8. S. Kotha, M. Salman and S. R. Cheekatla, Synthesis of Indenoindole Derivatives under Deep Eutectic Solvent Conditions, *ChemistrySelect*, 2024, **9**, e202402530.
9. D. Monica, N-heterocycles: Recent Advances in Biological Applications, *Mini-Rev. Org. Chem.*, 2023, **20**, 735-747.
10. C. Prandi and E. G. Occhiato, From synthetic control to natural products: a focus on N-heterocycles, *Pest Manag. Sci.*, 2019, **75**, 2385-2402.
11. N. Kerru, L. Gummidi, S. Maddila, K. K. Gangu and S. B. Jonnalagadda, A Review on Recent Advances in Nitrogen-Containing Molecules and Their Biological Applications, *Molecules*, 2020, **25**, 1909.
12. A. Amin, T. Qadir, P. K. Sharma, I. Jeelani and H. Abe, A Review on The Medicinal And Industrial Applications of N-Containing Heterocycles, *Open J. Med. Chem.*, 2022, **16**, e187410452209010.



13. A. Rusu, I.-M. Moga, L. Uncu and G. Hancu, The Role of Five-Membered Heterocycles in the Molecular Structure of Antibacterial Drugs Used in Therapy, *Pharm.*, 2023, **15**, 2554.
14. S. Pramod, S. Jitendra, J. P. Geeta and S. M. R. Mohan, 2-Pyrazolines as Biologically Active and Fluorescent Agents, An Overview, *Anti-Cancer Agents iMed. Chem.*, 2018, **18**, 1366-1385.
15. Y.-J. Wu, in *Prog. Heterocycl. Chem.*, eds. G. W. Gribble and J. A. Joule, Elsevier, 2021, vol. 33, pp. 277-292.
16. M. M. Heravi and V. Zadsirjan, Prescribed drugs containing nitrogen heterocycles: an overview, *RSC Adv.*, 2020, **10**, 44247-44311.
17. A. Ansari, A. Ali, M. Asif and Shamsuzzaman, Review: biologically active pyrazole derivatives, *New J. Chem.*, 2017, **41**, 16-41.
18. R. Khajuria, S. Dham and K. K. Kapoor, Active methylenes in the synthesis of a pyrrole motif: an imperative structural unit of pharmaceuticals, natural products and optoelectronic materials, *RSC Adv.*, 2016, **6**, 37039-37066.
19. P. K. Sharma, A. Amin and M. Kumar, A Review: Medicinally Important Nitrogen Sulphur Containing Heterocycles, *Open J. Med. Chem.*, 2020, **14**, 49-64.
20. E. Vitaku, D. T. Smith and J. T. Njardarson, Analysis of the Structural Diversity, Substitution Patterns, and Frequency of Nitrogen Heterocycles among U.S. FDA Approved Pharmaceuticals, *J. Med. Chem.*, 2014, **57**, 10257-10274.
21. S. Rao Cheekatla, D. P. Murale, L. Gopala and J.-S. Lee, Sensing and Imaging Agents for Cyclooxygenase Enzyme, *ChemMedChem*, 2024, **2024**, e202400636.
22. J. Dhuguru and R. Skouta, Role of Indole Scaffolds as Pharmacophores in the Development of Anti-Lung Cancer Agents, *Molecules*, 2020, **25**, 1615.
23. J. N. Marcelo and K. L. Hannah, Indole and Indoline Scaffolds in Antimicrobials: Overview, Synthesis and Recent Advances in Antimicrobial Research, *Curr. Med. Chem.*, 2021, **28**, 4828-4844.
24. Y. Wan, Y. Li, C. Yan, M. Yan and Z. Tang, Indole: A privileged scaffold for the design of anti-cancer agents, *Eur. J. Med. Chem.*, 2019, **183**, 111691.
25. A. Kumari and R. K. Singh, Medicinal chemistry of indole derivatives: Current to future therapeutic prospectives, *Bioorg. Chem.*, 2019, **89**, 103021.
26. S. Dadashpour and S. Emami, Indole in the target-based design of anticancer agents: A versatile scaffold with diverse mechanisms, *Eur. J. Med. Chem.*, 2018, **150**, 9-29.
27. T. V. Sravanthi and S. L. Manju, Indoles — A promising scaffold for drug development, *Eur. J. Pharm. Sci.*, 2016, **91**, 1-10.
28. A. Singh, C. Bhutani, P. Khanna, S. Talwar, S. K. Singh and L. Khanna, Recent report on indoles as a privileged anti-viral scaffold in drug discovery, *Eur. J. Med. Chem.*, 2025, **281**, 117017.
29. E. Barresi, E. Baglini, V. Poggetti, J. Castagnoli, D. Giorgini, S. Salerno, S. Taliani and F. Da Settimo, Indole-Based Compounds in the Development of Anti-Neurodegenerative Agents, *Molecules*, 2024, **29**, 2127.
30. T. Nishith, A. Khushbu, M. P. Tarun, P. Archita, P. Samir, S. Umang, S. Shaileshkumar, R. Khushman and P. Krupa, A Review of the Therapeutic Importance of Indole Scaffold in Drug Discovery, *Curr. Drug Discov. Technol.*, 2023, **20**, 9-37.
31. A. Fernando Rodrigues de Sa, J. B. Eliezer and F. Carlos Alberto Manssour, From Nature to Drug Discovery: The Indole Scaffold as a Privileged Structure, *Mini-Rev. Med. Chem.*, 2009, **9**, 782-793. [DOI: 10.1039/D5OB00338E](https://doi.org/10.1039/D5OB00338E)
32. R. Pino-Rios and M. Solà, The Relative Stability of Indole Isomers Is a Consequence of the Glidewell-Lloyd Rule, *J. Phys. Chem. A*, 2021, **125**, 230-234.
33. M. Bandini, Electrophilicity: the “dark-side” of indole chemistry, *Org. Biomol. Chem.*, 2013, **11**, 5206-5212.
34. C. C. J. Loh and D. Enders, Exploiting the Electrophilic Properties of Indole Intermediates: New Options in Designing Asymmetric Reactions, *Angew. Chem. Int. Ed.*, 2012, **51**, 46-48.
35. C. T. Walsh, Biological Matching of Chemical Reactivity: Pairing Indole Nucleophilicity with Electrophilic Isoprenoids, *ACS Chem. Biol.*, 2014, **9**, 2718-2728.
36. A. M. Janeiro and C. S. Marques, Biological Profile of Synthetic and Natural Indole Derivatives: Paving New Paths in Cancer Treatment, *Drugs Drug Candidates*, 2024, **3**, 488-511.
37. S. Kumar and Ritika, A brief review of the biological potential of indole derivatives, *Futur. J. Pharm. Sci.*, 2020, **6**, 121.
38. L. Tianze and X. Hui, Recent Progress of Bioactivities, Mechanisms of Action, Total Synthesis, Structural Modifications and Structure-activity Relationships of Indole Derivatives: A Review, *Mini-Rev. Med. Chem.*, 2022, **22**, 2702-2725.
39. P. V. Thanikachalam, R. K. Maurya, V. Garg and V. Monga, An insight into the medicinal perspective of synthetic analogs of indole: A review, *Eur. J. Med. Chem.*, 2019, **180**, 562-612.
40. J. W. Huffman, in *The Cannabinoid Receptors*, ed. P. H. Reggio, Humana Press, Totowa, NJ, 2009, DOI: 10.1007/978-1-59745-503-9_3, pp. 49-94.
41. Z. Dvořák, K. Pouliková and S. Mani, Indole scaffolds as a promising class of the aryl hydrocarbon receptor ligands, *Eur. J. Med. Chem.*, 2021, **215**, 113231.
42. C. T. Walsh, S. Garneau-Tsodikova and A. R. Howard-Jones, Biological formation of pyrroles: Nature's logic and enzymatic machinery, *Nat. Prod. Rep.*, 2006, **23**, 517-531.
43. S. Taliani, F. Da Settimo, C. Martini, S. Laneri, E. Novellino and G. Greco, Exploiting the Indole Scaffold to Design Compounds Binding to Different Pharmacological Targets, *Molecules*, 2020, **25**, 2331.
44. K. Himani, T. Abhishek and T. Varsha, Indole Derivatives: Versatile Scaffolds in Drug Development and Cancer Therapeutics, *Lett. Org. Chem.*, 2025, **22**, 1-39.
45. N. J. Pravin, R. S. Kavalapure, S. G. Alegaon, S. Gharge and S. D. Ranade, Indoles as promising Therapeutics: A review of recent drug discovery efforts, *Bioorg. Chem.*, 2025, **154**, 108092.
46. D. Paul and J. John, Recent Advances towards the Synthesis and Material Applications of Indoloindoles, *Chem. Asian J.*, 2022, **17**, e202200460.
47. O. Younis, M. Sayed, A. A. K. Mohammed, M. S. Tolba, R. Hassanien, A. M. Kamal El-Dean, O. Tsutsumi and M. Ahmed, Solid-State Luminescent Materials Containing Both Indole and Pyrimidine Moieties: Design, Synthesis, and Density Functional Theory Calculations, *ACS Omega*, 2022, **7**, 15016-15026.
48. A. V. Terry, J. J. Buccafusco and C. Wilson, Cognitive dysfunction in neuropsychiatric disorders: Selected serotonin receptor subtypes as therapeutic targets, *Behav. Brain Res.*, 2008, **195**, 30-38.



REVIEW

Organic & Biomolecular Chemistry

49. C. Kriegebaum, L. Gutknecht, A. Schmitt, K. P. Lesch and A. Reif, Serotonin Kompakt – Teil 1, *Neurobiologische und entwicklungsgenetische Grundlagen*, 2010, **78**, 319-331.
50. B. Alka and S. Shalu, Serotonergic 5-HT₆ Receptor Antagonists: Heterocyclic Chemistry and Potential Therapeutic Significance, *Curr. Top. Med. Chem.*, 2015, **15**, 1643-1662.
51. X. Li and A. Polter, Glycogen Synthase Kinase-3 is an Intermediate Modulator of Serotonin Neurotransmission, *Front. mol. neurosci.*, 2011, **4**, 1-14.
52. J. Opacka-Juffry, The role of serotonin as a neurotransmitter in health and illness: A review, *British J. Neurosci. Nurs.*, 2008, **4**, 272-277.
53. A. S. Girgis, S. S. Panda, B. M. Kariuki, M. S. Bekheit, R. F. Barghash and D. R. Aboshouk, Indole-Based Compounds as Potential Drug Candidates for SARS-CoV-2, *Molecules*, 2023, **28**, 6603.
54. H. S. Soor, S. D. Appavoo and A. K. Yudin, Heterocycles: Versatile control elements in bioactive macrocycles, *Biorg. Med. Chem.*, 2018, **26**, 2774-2779.
55. D. Garcia Jimenez, V. Poongavanam and J. Kihlberg, Macrocycles in Drug Discovery—Learning from the Past for the Future, *J. Med. Chem.*, 2023, **66**, 5377-5396.
56. S. R. Cheekatla, D. Barik, G. Anand, R. Mol K. M. and M. Porel, Indole-Based Macrocyclization by Metal-Catalyzed Approaches, *Organics*, 2023, **4**, 333-363.
57. S. R. Cheekatla, L. Thurakkal, A. Jose, D. Barik and M. Porel, Aza-Oxa-Triazole Based Macrocycles with Tunable Properties: Design, Synthesis, and Bioactivity, *Molecules*, 2022, **27**, 3409.
58. M. D. Cummings and S. Sekharan, Structure-Based Macrocycle Design in Small-Molecule Drug Discovery and Simple Metrics To Identify Opportunities for Macrocyclization of Small-Molecule Ligands, *J. Med. Chem.*, 2019, **62**, 6843-6853.
59. S. A. Patil, S. A. Patil and R. Patil, Medicinal applications of (benz)imidazole- and indole-based macrocycles, *Chem. Biol. Drug Des.*, 2017, **89**, 639-649.
60. C. G. Neochoritis, M. Kazemi Miraki, E. M. M. Abdelraheem, E. Surmiak, T. Zarganes-Tzitzikas, B. Łabuzek, T. A. Holak and A. Dömling, Design of indole- and MCR-based macrocycles as p53-MDM2 antagonists, *Beilstein J. Org. Chem.*, 2019, **15**, 513-520.
61. R. N. Das, M. Andréasson, R. Kumar and E. Chorell, Macrocyclization of bis-indole quinolines for selective stabilization of G-quadruplex DNA structures, *Chem. Sci.*, 2020, **11**, 10529-10537.
62. K. Devendra, S. Sahil, K. Sourav, S. Gurpreet, M. Vikramdeep and K. Bhupinder, Medicinal Perspective of Indole Derivatives: Recent Developments and Structure-Activity Relationship Studies, *Curr. Drug Targets*, 2020, **21**, 864-891.
63. Y. Chen, H. Li, J. Liu, R. Zhong, H. Li, S. Fang, S. Liu and S. Lin, Synthesis and biological evaluation of indole-based peptidomimetics as antibacterial agents against Gram-positive bacteria, *Eur. J. Med. Chem.*, 2021, **226**, 113813.
64. M. Tsuda, Y. Takahashi, J. Fromont, Y. Mikami and J. i. Kobayashi, Dendridine A, a Bis-indole Alkaloid from a Marine Sponge Dictyodendrilla Species, *J. Nat. Prod.*, 2005, **68**, 1277-1278.
65. X. Mo, D. P. Rao, K. Kaur, R. Hassan, A. S. Abdel-Samea, S. M. Farhan, S. Bräse and H. Hashem, Indole Derivatives: A Versatile Scaffold in Modern Drug Discovery—An Updated Review on Their Multifaceted Therapeutic Applications (2020–2024), *Molecules*, 2024, **29**, 4770. [10.1039/D5OB00338E](https://doi.org/10.1039/D5OB00338E)
66. D. R. Aboshouk, M. A. Youssef, M. S. Bekheit, A. R. Hamed and A. S. Girgis, Antineoplastic indole-containing compounds with potential VEGFR inhibitory properties, *RSC Adv.*, 2024, **14**, 5690-5728.
67. N. Chadha and O. Silakari, Indoles as therapeutics of interest in medicinal chemistry: Bird's eye view, *Eur. J. Med. Chem.*, 2017, **134**, 159-184.
68. P. A. Kumar, Md Iftekhar; Singh, Sangam; Pratam, Mohammad Rizki Fadhil; and Mishra, Arun K., Recent advancements on biological activity of indole and their derivatives: A review, *Thai J. Pharm. Sci.*, 2022, **46**, 233-250.
69. L. Ferrer, M. Mindt, V. F. Wendisch and K. Cankar, Indoles and the advances in their biotechnological production for industrial applications, *Syst. Microbiol. Biomanufacturing.*, 2024, **4**, 511-527.
70. S. M. Hassan, A. Farid, S. S. Panda, M. S. Bekheit, H. Dinkins, W. Fayad and A. S. Girgis, Indole Compounds in Oncology: Therapeutic Potential and Mechanistic Insights, *Pharm.*, 2024, **17**, 922.
71. L. K. Golightly, Pindolol: A Review of Its Pharmacology, Pharmacokinetics, Clinical Uses, and Adverse Effects, *Pharmacotherapy: J. Hum. Pharmacol. Drug Ther.*, 1982, **2**, 134-147.
72. R. R. Ruffolo and G. Z. Feuerstein, Pharmacology of Carvedilol: Rationale for Use in Hypertension, Coronary Artery Disease, and Congestive Heart Failure, *Cardiovasc. Drugs Ther.*, 1997, **11**, 247-256.
73. K. A. Shaw, J. Turner and C. Del Mar, Tryptophan and 5-Hydroxytryptophan for depression, *Cochrane Database Syst. Rev.*, 2002, DOI: 10.1002/14651858.CD003198.
74. M. D. Ferrari and P. R. S. Saxena, Clinical effects and mechanism of action of sumatriptan in migraine, *Clin. Neurol. Neurosurg.*, 1992, **94**, 73-77.
75. H. D. Langtry and S. P. Clissold, Pergolide, *Drugs*, 1990, **39**, 491-506.
76. C. P. Rains, H. M. Bryson and A. Fitton, Cabergoline, *Drugs*, 1995, **49**, 255-279.
77. M. P. Yavropoulou, P. Makras and A. D. Anastasilakis, Bazedoxifene for the treatment of osteoporosis, *Expert Opin. Pharmacother.*, 2019, **20**, 1201-1210.
78. S. Nalamachu and R. Wortmann, Role of Indomethacin in Acute Pain and Inflammation Management: A Review of the Literature, *Postgrad. Med.*, 2014, **126**, 92-97.
79. P. G. Casali, C. R. Garrett, M. E. Blackstein, M. Shah, J. Verweij, G. McArthur, I. Judson, J. Li, C. M. Baum and G. D. Demetri, Updated results from a phase III trial of sunitinib in GIST patients (pts) for whom imatinib (IM) therapy has failed due to resistance or intolerance, *J. Clin. Oncol.* 2006, **24**, 9513-9513.
80. S. Kotha, S. R. Cheekatla, A. K. Chinnam and T. Jain, Design and synthesis of polycyclic bisindoles via Fischer indolization and ring-closing metathesis as key steps, *Tetrahedron Lett.*, 2016, **57**, 5605-5607.
81. S. Kotha and C. Chakrapalli, Application of Fischer Indolization under Green Conditions using Deep Eutectic Solvents, *Chem. Rec.*, 2017, **17**, 1039-1058.
82. K. Hayakawa, T. Yasukouchi and K. Kanematsu, A new approach to the efficient indole synthesis by allene intramolecular cycloaddition, *Tetrahedron Lett.*, 1986, **27**, 1837-1840.



83. J. J. Li, in *Name Reactions: A Collection of Detailed Mechanisms and Synthetic Applications Fifth Edition*, ed. J. J. Li, Springer International Publishing, Cham, 2014, DOI: 10.1007/978-3-319-03979-4_182, pp. 415-416.
84. (a) A. R. Akanksha Porwal, Parwz Alam, Himanshu Singh, Kratika Singh, Ayush Dubey, Indole Moiety in Organic Synthesis: A Comprehensive Review of Methods and Mechanisms, *Int J. Pharm. Investigation*, 2024, **14**, 1052-1060; (b) D. I. Bugaenko, A. V. Karchava and M. A. Yurovskaya, Synthesis of indoles: recent advances, *Russian Chem. Rev.*, 2019, **88**, 99.
85. (a) D. F. Taber and P. K. Tirunahari, Indole Synthesis: A Review and Proposed Classification, *Tetrahedron*, 2011, **67**, 7195-7210; (b) R. Vicente, Recent Advances in Indole Syntheses: New Routes for a Classic Target, *Org. Biomol. Chem.*, 2011, **9**, 6469-6480; (c) J. S. S. Neto and G. Zeni, Recent Advances in the Synthesis of Indoles from Alkynes And Nitrogen Sources, *Org. Chem. Front.*, 2020, **7**, 155-210; (d) B. S. Mathada and N. G. Yernale Current Advances in Transition Metal-Free Access to Indoles. *Org. Prep. Proced. Int.* 2023, **55**, 299-316; (e) F. A. Almalki and A. O. Baryyan, Recent advances in the green synthesis of indole and its derivatives using microwave irradiation and the role of indole moiety in cancer, *Green Chem. Lett. Rev.* 2024, **17**, 2362925, DOI: <https://doi.org/10.1080/17518253.2024.2362925>; (f) G. W. Gribble, Indole Ring Synthesis: From Natural Products to Drug Discovery, John Wiley & Sons Ltd. 2016, DOI: 10.1002/9781118695692.
86. M. Makosza and J. Winiarski, Vicarious nucleophilic substitution of hydrogen, *Acc. Chem. Res.*, 1987, **20**, 282-289.
87. G. W. Gribble, in *Indole Ring Synthesis*, 2016, DOI: <https://doi.org/10.1002/9781118695692.ch42>, pp. 349-353.
88. A. Frischmuth, A. Unsinn, K. Groll, H. Stadtmüller and P. Knochel, Preparations and Reactions of SF₅-Substituted Aryl and Heteroaryl Derivatives via Mg and Zn Organometallics, *Chem. Eur. J.* 2012, **18**, 10234-10238.
89. T. Eicher, Hauptmann, Siegfried, in *The Chemistry of Heterocycles*, 2003, DOI: <https://doi.org/10.1002/352760183X.ch5a>, pp. 52-121.
90. L. Marion, Heterocyclic Compounds with Indole and Carbazole Systems. The Chemistry of Heterocyclic Compounds, *J. Am. Chem. Soc.*, 1955, **77**, 2917-2917.
91. S. Cacchi and G. Fabrizi, Synthesis and Functionalization of Indoles Through Palladium-catalyzed Reactions, *Chem. Rev.*, 2005, **105**, 2873-2920.
92. M. Bandini and A. Eichholzer, Catalytic Functionalization of Indoles in a New Dimension, *Angew. Chem. Int. Ed.*, 2009, **48**, 9608-9644.
93. V. Sharma, P. Kumar and D. Pathak, Biological importance of the indole nucleus in recent years: A comprehensive review, *J. Heterocycl. Chem.*, 2010, **47**, 491-502.
94. T. Kawasaki and K. Higuchi, Simple indole alkaloids and those with a nonrearranged monoterpenoid unit, *Nat. Prod. Rep.*, 2005, **22**, 761-793.
95. P. Beier, T. Pastýřková and G. Iakobson, Preparation of SF₅ Aromatics by Vicarious Nucleophilic Substitution Reactions of Nitro(pentafluorosulfanyl)benzenes with Carbanions, *J. Org. Chem.*, 2011, **76**, 4781-4786.
96. G. Iakobson, M. Pošta and P. Beier, Synthesis of Pentafluorosulfanyl-Containing Indoles and Oxindoles, *Synlett*, 2013, **24**, 855-859.
97. G. D. Heffernan, R. D. Coghlan, E. S. Manas, R. E. McDevitt, Y. Li, P. E. Mahaney, A. J. Robichaud, C. H. Huelston, P. Alifan, A. Bray, S. A. Cosmi, G. H. Johnston, T. Kenney, E. Koury, R. C. Winneker, D. C. Decher and E. J. Trybulski, Dual acting norepinephrine reuptake inhibitors and 5-HT_{2A} receptor antagonists: Identification, synthesis and activity of novel 4-aminoethyl-3-(phenylsulfonyl)-1H-indoles, *Biorg. Med. Chem.*, 2009, **17**, 7802-7815.
98. M. Makosza and A. Tyrala, Alkylation and the Knoevenagel Condensation of Nitrobenzyl Sulfones and Nitriles.1, *Synth. Commun.*, 1986, **16**, 419-423.
99. Z. Wróbel and M. Mąkosza, Synthesis of 1-hydroxyindoles and indoles from ortho-nitroarylethanes, *Tetrahedron*, 1997, **53**, 5501-5514.
100. Z. Wróbel and M. Mąkosza, New Simple Synthesis of N-Hydroxy 2-Vinylindoles, *Synlett*, 1993, **1993**, 597-598.
101. Z. Wróbel and M. Mąkosza, Transformations of o-nitroarylallyl carbanions. Synthesis of quinoline N-oxides and N-hydroxyindoles, *Tetrahedron*, 1993, **49**, 5315-5326.
102. J. Stalewski, The vicarious nucleophilic substitution of hydrogen proceeding via a cyclopropane ring-opening, *Tetrahedron Lett.*, 1998, **39**, 9523-9526.
103. M. Mkosza, J. Stalewski and O. S. Maslennikova, Synthesis of 7,8-Dimethoxy-2-oxo-1,3,4,5-tetrahydropyrrolo[4,3,2-de]quinoline: A Key Intermediate en Route to Makaluvamines, Discorhabdin C and Other Marine Alkaloids of this Group via Vicarious Nucleophilic Substitution of Hydrogen, *Synthesis*, 1997, **1997**, 1131-1133.
104. J. P. Hénichart, S. Chackal, F. Dudouit and R. Houssin, On the synthesis of two dimethoxy-1,3,4,5-tetrahydropyrrolo[4,3,2-de]quinoline regioisomers, *Heterocycles*, 2003, **60**, 615-622.
105. R. Bujok, Z. Wróbel and K. Wojciechowski, Expedient Synthesis of 1-Hydroxy-4- and 1-Hydroxy-6-nitroindoles, *Synlett*, 2012, **23**, 1315-1320.
106. J. K. Sutherland, Reaction of 1,8-diazabicyclo[5.4.0]undec-8-ene with methyl 3,5-dinitrobenzoate and 1,3,5-trinitrobenzene, *Chem. Commun.*, 1997, DOI: 10.1039/A607723D, 325-326.
107. B.-C. H. Chen, Jr., John; R. Pandit, Chennagiri; Zhao, Rulin; P. Skoumbourdis, Amanda; Wu, Hong; E. Sundeen, Joseph; Leftheris, Katerina . A GENERAL LARGE SCALE SYNTHESIS OF 2-ALKYL-7-METHOXYINDOLES, *Heterocycl.*, 2001, **55**, 951-960.
108. M. S. Islam, C. Brennan, Q. Wang and M. M. Hossain, Convenient Method of Synthesizing 3-Ethoxycarbonyl Indoles, *J. Org. Chem.*, 2006, **71**, 4675-4677.
109. B. T. McClay, K. E. Lambson, S. R. Banini, T. N. Akhmedov and B. C. G. Söderberg, A base-mediated synthesis of N-hydroxy- and N-alkoxyindoles from 2-nitrostyrenes, *Tetrahedron*, 2023, **144**, 133578.
110. Y. Maeyama, Y. Nakashima, H. Kato, Y. Hitora, K. Maki, N. Inada, S. Murakami, T. Inazumi, Y. Ise, Y. Sugimoto, H. Ishikawa and S. Tsukamoto, Amakusamine from a Psammocinia sp. Sponge: Isolation, Synthesis, and SAR Study on the Inhibition of RANKL-Induced Formation of Multinuclear Osteoclasts, *J. Nat. Prod.*, 2021, **84**, 2738-2743.
111. G. W. Gribble, in *Indole Ring Synthesis*, 2016, DOI: <https://doi.org/10.1002/9781118695692.ch2>, pp. 41-115.
112. L. Novellino, M. d'Ischia and G. Protà, Expedient Synthesis of 5,6-Dihydroxyindole and Derivatives via an Improved Zn(II)-



REVIEW

Organic & Biomolecular Chemistry

- Assisted 2,β-Dinitrostyrene Approach, *Synthesis*, 1999, **1999**, 793-796.
113. F. He, Y. Bo, J. D. Altom and E. J. Corey, Enantioselective Total Synthesis of Aspidophytine, *J. Am. Chem. Soc.*, 1999, **121**, 6771-6772.
 114. A. K. Sinhababu and R. T. Borchardt, Silica gel assisted reductive cyclization of alkoxy-2, .beta.-dinitrostyrenes to alkoxyindoles, *J. Org. Chem.*, 1983, **48**, 3347-3349.
 115. Y. Fukuyama, C. Iwatsuki, M. Kodama, M. Ochi, K. Kataoka and K. Shibata, Antimicrobial indolequinones from the mid-intestinal gland of the muricid gastropod *Drupella fragum*, *Tetrahedron*, 1998, **54**, 10007-10016.
 116. L.-M. Yang, C.-F. Chen and K.-H. Lee, Synthesis of rutaecarpine and cytotoxic analogues, *Bioorg. Med. Chem. Lett.*, 1995, **5**, 465-468.
 117. G. W. Gribble, in *Indole Ring Synthesis*, 2016, DOI: <https://doi.org/10.1002/9781118695692.ch4>, pp. 121-130.
 118. G. Bartoli, G. Palmieri, M. Bosco and R. Dalpozzo, The reaction of vinyl grignard reagents with 2-substituted nitroarenes: A new approach to the synthesis of 7-substituted indoles, *Tetrahedron Lett.*, 1989, **30**, 2129-2132.
 119. G. Bartoli, M. Bosco, R. Dalpozzo, G. Palmieri and E. Marcantoni, Reactivity of nitro- and nitroso-arenes with vinyl grignard reagents: synthesis of 2-(trimethylsilyl)indoles, *J. Chem. Soc., Perkin Trans. 1*, 1991, DOI: 10.1039/P19910002757, 2757-2761.
 120. D. Dobson, A. Todd and J. Gilmore, The Synthesis of 7-Alkoxyindoles, *Synth. Commun.*, 1991, **21**, 611-617.
 121. K. R. Buszek, N. Brown and D. Luo, Concise Total Synthesis of (±)-cis-Trikentrin A and (±)-Herbindole A via Intermolecular Indole Aryne Cycloaddition, *Org. Lett.*, 2009, **11**, 201-204.
 122. J. J. Li, in *Gassman indole synthesis; Name Reactions: A Collection of Detailed Mechanisms and Synthetic Applications Fifth Edition*; , ed. J. J. Li, Springer International Publishing, Cham, 2014, DOI: 10.1007/978-3-319-03979-4_116, pp. 276-277.
 123. P. G. Gassman, T. J. Van Bergen, D. P. Gilbert and B. W. Cue, Jr., General method for the synthesis of indoles, *J. Am. Chem.*, 1974, **96**, 5495-5508.
 124. T. D. Lash, C. Wijesinghe, A. T. Osuma and J. R. Patel, Synthesis of novel porphyrin chromophores from nitroarenes: Further applications of the Barton-Zard pyrrole condensation, *Tetrahedron Lett.*, 1997, **38**, 2031-2034.
 125. H.-J. Xu, Z. Shen, T. Okujima, N. Ono and X.-Z. You, Synthesis and spectroscopic characterization of meso-tetraarylporphyrins with fused phenanthrene rings, *Tetrahedron Lett.*, 2006, **47**, 931-934.
 126. C. S. B. Gomes, P. T. Gomes, M. T. Duarte, R. E. Di Paolo, A. L. Maçanita and M. J. Calhorda, Synthesis, Structure, and Photophysical Characterization of Blue-Green Luminescent Zinc Complexes Containing 2-Iminophenanthropyrrolyl Ligands, *Inorg. Chem.*, 2009, **48**, 11176-11186.
 127. S. Sahoo, M. H. Mahadavaiah, D. Usharani and H. Rath, NIR Absorbing Aromatic E-Ethylene Bridged Hexaphyrins (2.1.1.2.1.1): Synthesis, Characterization, and Protonation Studies, *Org. Lett.*, 2023, **25**, 1491-1496.
 128. W. Dohle, A. Staubitz and P. Knochel, Mild Synthesis of Polyfunctional Benzimidazoles and Indoles by the Reduction of Functionalized Nitroarenes with Phenylmagnesium Chloride, *Chem. Eur. J.*, 2003, **9**, 5323-5331.
 129. R. D. Clark, D. B. Repke,, The Leimgruber-Batcho Indole Synthesis, *Heterocycl.*, 1984, **22**, 195-221.
 130. A. D. L. Batcho, W. , INDOLES FROM 2-METHYLNITROBENZENES BY CONDENSATION WITH FORMAMIDE ACETALS FOLLOWED BY REDUCTION: 4-BENZYLOXYINDOLE, *Org. Synth.* 1985, **63**, 214-220.
 131. J. Siu, I. R. Baxendale and S. V. Ley, Microwave assisted Leimgruber-Batcho reaction for the preparation of indoles, azaindoles and pyrrolylquinolines, *Org. Biomol. Chem.*, 2004, **2**, 160-167.
 132. A. P. Kozikowski, H. Ishida and Y.-Y. Chen, New synthesis and some selected reactions of the potential ergot alkaloid precursor indole-4-carboxaldehyde, *J. Org. Chem.*, 1980, **45**, 3350-3352.
 133. G. W. Gribble, in *Indole Ring Synthesis*, 2016, DOI: <https://doi.org/10.1002/9781118695692.ch26>, pp. 266-277.
 134. J. I. G. Cadogan and R. K. Mackie, Tervalent phosphorus compounds in organic synthesis, *Chem. Soc. Rev.*, 1974, **3**, 87-137.
 135. J. I. G. Cadogan, M. Cameron-Wood, R. K. Mackie and R. J. G. Searle, 896. The reactivity of organophosphorus compounds. Part XIX. Reduction of nitro-compounds by triethyl phosphite: a convenient new route to carbazoles, indoles, indazoles, triazoles, and related compounds, *J. Chem. Soc. (Resumed)*, 1965, DOI: 10.1039/JR9650004831, 4831-4837.
 136. Z. Qu, X. Chen, S. Zhong, G.-J. Deng and H. Huang, NaI/PPh₃-Mediated Photochemical Reduction and Amination of Nitroarenes, *Org. Lett.*, 2021, **23**, 5349-5353.
 137. J. I. G. Cadogan, B. S. Tait and N. J. Tweddle, Formation of carbazoles by photolytic extrusion of dimethyl phenylphosphonate from oxazaphosphoranes. A phosphorus analogue of the Graebe-Ullmann reaction, *J. Chem. Soc., Chem. Commun.*, 1975, DOI: 10.1039/C39750000847, 847-848.
 138. R. J. Sundberg, Deoxygenation of Nitro Groups by Trivalent Phosphorus. Indoles from o-Nitrostyrenes, *J. Org. Chem.*, 1965, **30**, 3604-3610.
 139. R. J. Sundberg, A Study of the Deoxygenation of Some o-Alkylnitro- and o-Alkylnitrosobenzenes in Triethyl Phosphite, *J. Am. Chem. Soc.*, 1966, **88**, 3781-3789.
 140. R. J. Sundberg and T. Yamazaki, Rearrangements and ring expansions during the deoxygenation of .beta.,.beta.-disubstituted o-nitrostyrenes, *J. Org. Chem.*, 1967, **32**, 290-294.
 141. H. Majgier-Baranowska, J. D. Williams, B. Li and N. P. Peet, Studies on the mechanism of the Cadogan-Sundberg indole synthesis, *Tetrahedron Lett.*, 2012, **53**, 4785-4788.
 142. N. H. Ansari and B. C. G. Söderberg, Short syntheses of the indole alkaloids alocasin A, scalaridine A, and hyrtinadine A-B, *Tetrahedron*, 2016, **72**, 4214-4221.
 143. F. Ferretti, D. R. Ramadan and F. Ragaini, Transition Metal Catalyzed Reductive Cyclization Reactions of Nitroarenes and Nitroalkenes, *ChemCatChem*, 2019, **11**, 4450-4488.
 144. G. Ghimire and B. C. G. Söderberg, Short syntheses of the tricyclic indole alkaloids cimitrypazepine and fargesine, *Tetrahedron Lett.*, 2016, **57**, 3873-3876.
 145. J. T. Kuethe, A. Wong and I. W. Davies, Effective Strategy for the Preparation of Indolocarbazole Aglycons and Glycosides: Total Synthesis of Tjipanazoles B, D, E, and I, *Org. Lett.*, 2003, **5**, 3721-3723.
 146. R. W. Clawson, C. A. Dacko, R. E. Deavers, N. G. Akhmedov and B. C. G. Söderberg, Attempted synthesis of 3-hydroxy-2-octadecylindole. Proposed structural revision of previously



- prepared 3-hydroxy-2-octadecylindole and a proposed structure of fistulosin, *Tetrahedron*, 2009, **65**, 8786-8793.
147. T. Bzeih, K. Zhang, A. Khalaf, A. Hachem, M. Alami and A. Hamze, One-Pot Reaction between N-Tosylhydrazones and 2-Nitrobenzyl Bromide: Route to NH-Free C2-Arylindoles, *J. Org. Chem.*, 2019, **84**, 228-238.
 148. A. Kasahara, T. Izumi, S. Murakami, K. Miyamoto and T. Hino, A regiocontrolled synthesis of substituted indoles by palladium-catalyzed coupling of 2-bromonitrobenzenes and 2-bromoacetanilides, *J. Heterocycl. Chem.*, 1989, **26**, 1405-1413.
 149. M. Akazome, T. Kondo and Y. Watanabe, Novel Synthesis of Indoles via Palladium-Catalyzed Reductive N-Heterocyclization of o-Nitrostyrene Derivatives, *Chem. Lett.*, 1992, **21**, 769-772.
 150. M. Akazome, T. Kondo and Y. Watanabe, Palladium complex-catalysed reductive N-heterocyclization of N-(2-nitrobenzylidene)amines into 2H-indazole derivatives, *J. Chem. Soc., Chem. Commun.*, 1991, DOI: 10.1039/C39910001466, 1466-1467.
 151. M. Akazome, T. Kondo and Y. Watanabe, Palladium Complex-Catalyzed Reductive N-Heterocyclization of Nitroarenes: Novel Synthesis of Indole and 2H-Indazole Derivatives, *J. Org. Chem.*, 1994, **59**, 3375-3380.
 152. B. C. Söderberg and J. A. Shriver, Palladium-Catalyzed Synthesis of Indoles by Reductive N-Heteroannulation of 2-Nitrostyrenes, *J. Org. Chem.*, 1997, **62**, 5838-5845.
 153. M. Shevlin, X. Guan and T. G. Driver, Iron-Catalyzed Reductive Cyclization of o-Nitrostyrenes Using Phenylsilane as the Terminal Reductant, *ACS Catal.*, 2017, **7**, 5518-5522.
 154. J. G. Taylor and C. R. D. Correia, Stereoselective Synthesis of Unsymmetrical β,β -Diarylacrylates by a Heck-Matsuda Reaction: Versatile Building Blocks for Asymmetric Synthesis of β,β -Diphenylpropanoates, 3-Aryl-indole, and 4-Aryl-3,4-dihydro-quinolin-2-one and Formal Synthesis of (-)-Indatraline, *J. Org. Chem.*, 2011, **76**, 857-869.
 155. F. Ragaini, F. Ferretti and M. A. Fouad, Phenyl Formate as a CO Surrogate for the Reductive Cyclization of Organic Nitro Compounds to Yield Different N-Heterocycles: No Need for Autoclaves and Pressurized Carbon Monoxide, *Catal.* 2023, **13**, 224; DOI: <https://doi.org/10.3390/catal13020224>.
 156. F. Zhou, D.-S. Wang and T. G. Driver, Palladium-Catalyzed Formation of N-Heteroarenes from Nitroarenes using Molybdenum Hexacarbonyl as the Source of Carbon Monoxide, *Adv. Synth. Catal.*, 2015, **357**, 3463-3468.
 157. Y. Dong, X. Wang, M. Cal, M. Kaiser and J. L. Vennerstrom, Activity of diimidazoline amides against African trypanosomiasis, *Bioorg. Med. Chem. Lett.*, 2014, **24**, 944-948.
 158. C. Crotti, S. Cenini, B. Rindone, S. Tollari and F. Demartin, Deoxygenation reactions of ortho-nitrostyrenes with carbon monoxide catalysed by metal carbonyls: a new route to indoles, *J. Chem. Soc., Chem. Commun.*, 1986, DOI: 10.1039/C39860000784, 784-786.
 159. X. Guan, H. Zhu, Y. Zhao and T. G. Driver, Pd-Catalyzed Reductive Cyclization of Nitroarenes with CO₂ as the CO Source, *Eur. J. Org. Chem.*, 2020, **2020**, 57-60.
 160. (a) Y. Yamamoto, E. Ohkubo and M. Shibuya, Synthesis of 3-Aryl-2-(trifluoromethyl)indoles via Copper-Catalyzed Hydroarylation and Subsequent Cadogan Cyclization, *Adv. Synth. Catal.*, 2017, **359**, 1747-1751.; (b) Y. Yamamoto, N. Kirai, Y. Harada, Cu-catalyzed stereoselective conjugate addition of arylboronic acids to alkynoates, *Chem. Commun.*, 2008, **44**, 2010-2012, DOI: 10.1039/B802231G/(cc)Xbdu30.
 161. J. I. G. Cadogan, Phosphite-Reduction of Aromatic Nitro-Compounds as a Route to Heterocycles, *Synthesis*, 1969, **1969**, 11-17.
 162. I. W. Davies, V. A. Guner and K. N. Houk, Theoretical Evidence for Oxygenated Intermediates in the Reductive Cyclization of Nitrobenzenes, *Org. Lett.*, 2004, **6**, 743-746.
 163. A. W. Freeman, M. Urvoy and M. E. Criswell, Triphenylphosphine-Mediated Reductive Cyclization of 2-Nitrobiphenyls: A Practical and Convenient Synthesis of Carbazoles, *J. Org. Chem.*, 2005, **70**, 5014-5019.
 164. S. Selvaraju, K. A. Niradha Sachinthan, R. A. Hopson, F. M. McFarland, S. Guo, A. L. Rheingold and T. L. Nelson, Eumelanin-inspired core derived from vanillin: a new building block for organic semiconductors, *Chem. Commun.*, 2015, **51**, 2957-2959.
 165. R. Sanz, J. Escribano, M. R. Pedrosa, R. Aguado and F. J. Arnáiz, Dioxomolybdenum(VI)-Catalyzed Reductive Cyclization of Nitroaromatics. Synthesis of Carbazoles and Indoles, *Adv. Synth. Catal.*, 2007, **349**, 713-718.
 166. H. Gao, Q.-L. Xu, M. Yousufuddin, D. H. Ess and L. Kürti, Rapid Synthesis of Fused N-Heterocycles by Transition-Metal-Free Electrophilic Amination of Arene C-H Bonds, *Angew. Chem. Int. Ed.*, 2014, **53**, 2701-2705.
 167. T. N. Poudel and Y. R. Lee, Construction of highly functionalized carbazoles via condensation of an enolate to a nitro group, *Chem. Sci.*, 2015, **6**, 7028-7033.
 168. A. Baeyer and A. Emmerling, Synthese des Indols, *Berichte der deutschen chemischen Gesellschaft*, 1869, **2**, 679-682.
 169. B. Robinson, The Fischer Indole Synthesis, *Chem. Rev.*, 1963, **63**, 373-401.
 170. H. E. R. Indigo and its Artificial Production, *Nature*, 1881, **24**, 227-231.
 171. A. Reissert, Einwirkung von Oxalester und Natriumäthylat auf Nitrotoluole. Synthese nitrirter Phenylbrenztraubensäuren, *Berichte der deutschen chemischen Gesellschaft*, 1897, **30**, 1030-1053.
 172. W. E. N. a. F. J. Baude, ETHYL INDOLE-2-CARBOXYLATE, *Org. Synth.*, 1963, **43**, 40.
 173. J. R. Johnson, R. B. Hasbrouck, J. D. Dutcher and W. F. Bruce, Gliotoxin. V. The Structure of Certain Indole Derivatives Related to Gliotoxin1,2, *J. Am. Chem. Soc.*, 1945, **67**, 423-430.
 174. S. Katayama, N. Ae and R. Nagata, Synthesis of Tricyclic Indole-2-carboxylic Acids as Potent NMDA-Glycine Antagonists, *J. Org. Chem.*, 2001, **66**, 3474-3483.
 175. A. V. Butin, T. y. A. Stroganova, I. V. Lodina and G. D. Krapivin, Furan ring opening—indole ring closure: a new modification of the Reissert reaction for indole synthesis, *Tetrahedron Lett.*, 2001, **42**, 2031-2033.
 176. G. W. Gribble, in *Indole Ring Synthesis*, 2016, DOI: <https://doi.org/10.1002/9781118695692.ch40>, pp. 332-337.
 177. Y. Motoyama, K. Kamo and H. Nagashima, Catalysis in Polysiloxane Gels: Platinum-Catalyzed Hydrosilylation of Polymethylhydrosiloxane Leading to Reusable Catalysts for Reduction of Nitroarenes, *Org. Lett.*, 2009, **11**, 1345-1348.



REVIEW

Organic & Biomolecular Chemistry

178. I. Choi, H. Chung, J. W. Park and Y. K. Chung, Active and Recyclable Catalytic Synthesis of Indoles by Reductive Cyclization of 2-(2-Nitroaryl)acetonitriles in the Presence of Co–Rh Heterobimetallic Nanoparticles with Atmospheric Hydrogen under Mild Conditions, *Org. Lett.*, 2016, **18**, 5508–5511.
179. E. Ucciani and A. Bonfand, Catalytic synthesis of 3-substituted indoles using CO as building block and supported rhodium as catalyst, *J. Chem. Soc., Chem. Commun.*, 1981, DOI: 10.1039/C39810000082, 82–83.
180. S. Tong, Z. Xu, M. Mamboury, Q. Wang and J. Zhu, Aqueous Titanium Trichloride Promoted Reductive Cyclization of o-Nitrostyrenes to Indoles: Development and Application to the Synthesis of Rizatriptan and Aspidospermidine, *Angew. Chem. Int. Ed.*, 2015, **54**, 11809–11812.
181. G. Zhang, L. Lin, K. Yang, S. Wang, Q. Feng, J. Zhu and Q. Song, 3-Aminoindole Synthesis from 2-Nitrochalcones and Ammonia or Primary Amines, *Adv. Synth. Catal.*, 2019, **361**, 3718–3722.
182. Y. Hakmaoui, A. Eşme, R. E. Ajlaoui, M. E. Belghiti, A. Zeroual and E. M. Rakib, Efficient One-Pot Synthesis of Indolhydroxy Derivatives Catalyzed by SnCl₂, DFT Calculations and Docking Study, *Chem. Africa*, 2022, **5**, 827–839.
183. K. Yang, F. Zhou, Z. Kuang, G. Gao, T. G. Driver and Q. Song, Diborane-Mediated Deoxygenation of o-Nitrostyrenes To Form Indoles, *Org. Lett.*, 2016, **18**, 4088–4091.
184. J. S. Kim, J. H. Han, J. J. Lee, Y. M. Jun, B. M. Lee and B. H. Kim, Indium-HI-mediated one-pot reaction of 1-(2-arylethynyl)-2-nitroarenes to 2-arylindoles, *Tetrahedron Lett.*, 2008, **49**, 3733–3738.
185. K. Okuma, J.-i. Seto, K.-i. Sakaguchi, S. Ozaki, N. Nagahora and K. Shioji, Palladium-free zinc-mediated hydroamination of alkynes: efficient synthesis of indoles from 2-alkynylaniline derivatives, *Tetrahedron Lett.*, 2009, **50**, 2943–2945.
186. S. M. Oh and S. Shin, Synthesis of -Hydroxyindole Derivatives via Pd-Catalyzed Electrophilic Cyclization, *Bull. Korean Chem. Soc.*, 2021, **42**, 925–928.
187. M. S. Lokolkar, P. A. Mane, S. Dey and B. M. Bhanage, Synthesis of 2-Substituted Indoles by Pd-Catalyzed Reductive Cyclization of 1-Halo-2-nitrobenzene with Alkynes, *Eur. J. Org. Chem.*, 2022, **2022**, e202101505.
188. L. Yao, J. Ying and X.-F. Wu, Nickel-catalyzed cascade carbonylative synthesis of N-benzoyl indoles from 2-nitroalkynes and aryl iodides, *Org. Chem. Front.*, 2021, **8**, 6541–6545.
189. L. Yao, P. Wei, J. Ying and X.-F. Wu, Nickel-catalyzed carbonylative domino cyclization of arylboronic acid pinacol esters with 2-alkynyl nitroarenes toward N-aryl indoles, *Org. Chem. Front.*, 2022, **9**, 2685–2689.
190. D. Xu, R. Liu, J. Li, H. Zhao, J. Ma and Z. Dong, Atomically dispersed Co-N₄ sites anchored on N-doped carbon for aqueous phase transfer hydrogenation between nitroarenes and saturated N-heterocycles, *Appl. Catal. B: Environ.*, 2021, **299**, 120681.
191. A. Penoni and K. M. Nicholas, A novel and direct synthesis of indoles via catalytic reductive annulation of nitroaromatics with alkynes, *Chem. Commun.*, 2002, DOI: 10.1039/B110370A, 484–485.
192. A. Penoni, J. Volkmann and K. M. Nicholas, Regioselective Synthesis of Indoles via Reductive Annulation of Nitrosoaromatics with Alkynes, *Org. Lett.*, 2002, **4**, 699–701.
193. (a) F. Ragaini, A. Rapetti, E. Visentin, M. Monzani, A. Caselli and S. Cenini, Synthesis of Indoles by Intramolecular Cyclization of Unfunctionalized Nitroarenes and Alkynes, Catalyzed by Palladium–Phenanthroline Complexes, *J. Org. Chem.*, 2006, **71**, 3748–3753; (b) C. Galli, P. Gentili, A. Guarnieri, S. Kobayashi, Z. Rappoport, Competition of Mechanisms in the Photochemical Cleavage of the C–X Bond of Aryl-Substituted Vinyl Halides, *J. Org. Chem.*, 1998, **63**, 9292–9299. (c) B. Brocklehurst, J.S. Robinson, D.N. Tawn, The fluorescence spectrum of the α-styryl radical, C₆H₅–CH₂, *Chem. Phys. Lett.*, 1972, **12**, 610–611. [https://doi.org/10.1016/0009-2614\(72\)80019-8](https://doi.org/10.1016/0009-2614(72)80019-8).
194. C. Della Rosa, M. Kneeteman and P. Mancini, Comparison of the reactivity between 2- and 3-nitropyrroles in cycloaddition reactions. A simple indole synthesis, *Tetrahedron Lett.*, 2007, **48**, 1435–1438.
195. P. M. E. Mancini, M. N. Kneeteman, M. Cainelli, C. M. Ormachea and L. R. Domingo, Nitropyrroles, Diels-Alder reactions assisted by microwave irradiation and solvent effect. An experimental and theoretical study, *J. Mol. Struct.*, 2017, **1147**, 155–160.
196. P. Geerlings, F. De Proft and W. Langenaeker, Conceptual Density Functional Theory, *Chem. Rev.*, 2003, **103**, 1793–1874.
197. P. Du, J. L. Brosmer and D. G. Peters, Electrosynthesis of Substituted 1H-Indoles from o-Nitrostyrenes, *Org. Lett.*, 2011, **13**, 4072–4075.
198. T. Wirtanen, E. Rodrigo and S. R. Waldvogel, Recent Advances in the Electrochemical Reduction of Substrates Involving N–O Bonds, *Adv. Synth. Catal.*, 2020, **362**, 2088–2101.
199. C. G. S. Bjorn, Synthesis of Heterocycles via Intramolecular Annulation of Nitrene Intermediates, *Curr. Org. Chem.*, 2000, **4**, 727–764.
200. B. Wang, H. Ren, H.-J. Cao, C. Lu and H. Yan, A switchable redox annulation of 2-nitroarylethanol affording N-heterocycles: photoexcited nitro as a multifunctional handle, *Chem. Sci.*, 2022, **13**, 11074–11082.
201. K. Teng, Q. Liu, J. Lv and T. Li, The application of nitroarenes in catalyst-free photo-driven reactions, *ChemPhotoChem*, 2024, **8**, e202300236.
202. M. Waheed, M. A. Alsharif, M. I. Alahmdi, S. Mukhtar and H. Parveen, Iron-catalyzed intramolecular reductive cyclization of o-nitroarenes to indoles under visible light irradiation, *Tetrahedron Lett.*, 2023, **123**, 154543.
203. A. Berkessel, S. Reichau, A. von der Höh, N. Leconte and J.-M. Neudörfl, Light-Induced Enantioselective Hydrogenation Using Chiral Derivatives of Casey's Iron–Cyclopentadienone Catalyst, *Organomet.*, 2011, **30**, 3880–3887.
204. H. Qin, R. Liu, Z. Wang, F. Xu, X. Li, C. Shi, J. Chen, W. Shan, C. Liu, P. Xing, J. Zhu, X. Li and D. Shi, Photoinduced Bartoli Indole Synthesis by the Oxidative Cleavage of Alkenes with Nitro(hetero)arenes, *Angew. Chem. Int. Ed.*, 2025, **64**, e202416923.
205. J. Zhang, C. Mück-Lichtenfeld, M.-A. Wiethoff and A. Studer, Photocatalytic PPh₃-Mediated Synthesis of C3-Functionalized Indoles via Radical Annulation of Nitroarenes and Alkenes, *Angew. Chem. Int. Ed.*, 2024, **63**, e202416726.



Open Access Article. Published on 30 May 2025. Downloaded on 7/5/2025 12:59:24 PM.
This article is licensed under a Creative Commons Attribution-NonCommercial 3.0 Unported Licence.



Data availability statement

No primary research results, software or code have been included, and no new data were generated or analyzed as part of this review.

