

Cite this: *Org. Biomol. Chem.*, 2025, **23**, 10143

Digold photosensitizer for visible light [2 + 2] cycloaddition and *E/Z* isomerization reactions

Tommaso Ruggiero,^a James Fortwengler,^{b,c} Kristof Van Hecke,^d Mark E. Thompson,^{b,c} Thomas S. A. Heugebaert^d and Steven P. Nolan^{*,a}

Energy transfer photocatalysis has emerged as a powerful platform for a wide variety of transformations, yet advances in this area are often limited by the visible-light absorption of catalysts and the very high cost of state-of-the-art iridium-based sensitizers. We report the synthesis, characterization, and photocatalytic use of a novel digold(i) complex, {[Au(IPr)]₂(DHIC)} (**PhotAu3**), designed to overcome these limitations by absorbing visible light in the blue region and proving catalytically effective under 450 nm LED irradiation. The complex was successfully applied to a range of EnT-mediated photocatalytic reactions, including intramolecular [2 + 2] cycloadditions and *E/Z* isomerizations, achieving high conversions at low catalyst loadings in short reaction times.

Received 18th September 2025,
Accepted 16th October 2025

DOI: 10.1039/d5ob01503k

rsc.li/obc


Introduction

Over the past fifteen years, the area of photocatalysis has experienced remarkable growth, becoming a powerful synthetic tool enabling efficient late-stage functionalization and the construction of very diverse molecular frameworks.^{1–4} While photoredox catalysis *via* electron transfer (ET) has largely dominated the field,^{5–7} energy transfer (EnT) catalysis, particularly through a triplet-triplet energy transfer (TTEnt) mechanism,⁸ has recently gained attention as a promising strategy for molecular assembly.^{9–12} Although a wide variety of organosensitizers,^{13–17} including ruthenium,^{18–20} and other transition metal-based photocatalysts^{21–24} have been explored in energy transfer (EnT) photocatalysis, iridium complexes remain the state-of-the-art^{9,25} due to their exceptionally long excited triplet-state lifetimes, high quantum yields, tunable ligand frameworks, and, most notably, their high triplet energy (E_t) values that permit excitation of a broad range of organic substrates.²⁶ Despite recent advances where E_t values of up to 73 kcal mol⁻¹ have been achieved,^{27,28} the majority of Ir-based photosensitizers still fall below the 64 kcal mol⁻¹ threshold, thereby oftentimes limiting substrate scope. Furthermore, their complex multi-step syntheses,²⁹ high

cost,³⁰ limited compatibility with green solvents due to their ionic nature, and poor recyclability^{31,32} collectively highlight the pressing need for more sustainable, cost-effective, and efficient alternatives. As a more cost-effective and synthetically accessible alternative to iridium-based photosensitizers, Nolan and co-workers have developed and made commercially available³³ the gold carbazolyl complexes [Au(SIPr)(Cbz)] (**PhotAu1**) (SIPr = [*N,N*-bis(2,6-diisopropylphenyl)imidazolin-2-ylidene]; Cbz = carbazolyl) and [Au(IPr)(Cbz)] (**PhotAu2**) (IPr = [*N,N*-bis(2,6-diisopropylphenyl)imidazol-2-ylidene]) which exhibit high triplet energies (67.9 and 67.6 kcal mol⁻¹, respectively), remarkably long lifetimes in solution (266 and 335 μ s respectively), and intense energy emission, enabling previously inaccessible energy transfer (EnT) transformations such as the intramolecular [2 + 2] cycloaddition of unsubstituted indoles in notably short reaction times.^{34,35} The successful use of these organogold complexes in additional EnT photocatalytic processes further demonstrates their potential and versatility. These have shown effective in the intermolecular [2 + 2] cycloaddition of coumarins,³⁶ isooxazolines³⁷ and benzothio-phenes³⁸ with various alkenes, the cyclization of acrylamides³⁹ and in the synthesis of cyclohepta-indoles.⁴⁰

Dinuclear gold(i) complexes have only recently been studied in catalysis,^{41,42} materials science^{43–45} and as promising anti-cancer agents.^{46,47} Dinuclear amido-bearing NHC complexes for photocatalysis have few precedents, and this synthetic challenge represents an opportunity to explore the fundamental question of synergistic bimetallic interactions and their impact on photocatalytic behavior and reactivity. In this regard, novel dinuclear [(bisNHC){Au(Cbz)}₂] complexes have been described.⁴⁸ Investigation of their photocatalytic activity and photophysical properties revealed that, while their pro-

^aDepartment of Chemistry and Center for Sustainable Chemistry, Ghent University, Krijgslaan 289, building S-3, 9000 Ghent, Belgium. E-mail: steven.nolan@ugent.be^bMork Family Department of Chemical Engineering and Materials Science, University of Southern California, Los Angeles, California 90089, USA^cDepartment of Chemistry, University of Southern California, Los Angeles, California 90089, USA. E-mail: met@usc.edu^dDepartment of Green Chemistry and Technology, Ghent University, Coupure Links 653, 9000 Ghent, Belgium. E-mail: thomas.heugebaert@ugent.be




Scheme 1 State-of-the-art Ir and gold photosensitizers and novel $[\text{Au}(\text{IPr})]_2(\text{DHIC})$ (**PhotAu3**).

properties resemble those of $[\text{Au}(\text{SI}Pr)(\text{Cbz})]$, the NHC-Au-Cbz fragments in the binuclear complexes appear to operate independently from one another in solution without the apparent existence of cooperative interaction. Additionally, binuclear Janus-type and tandem-NHC gold complexes reported by Thompson and coworkers exhibit thermally activated delayed fluorescence (TADF), although they have not been employed as sensitizers, and their synthesis remains challenging.^{49,50} A major limitation of the new gold complexes **PhotAu1** and **PhotAu2** is the position of their absorbance in the UV region. Irradiation in this section of the spectrum can lead to photochemical initiation of reactions without the need for a sensitizer.³⁴ Thus far, only light sources emitting at 365 nm have been successfully used in experiments conducted by the Nolan group with gold sensitizers. In contrast, iridium-based photocatalysts can operate with 450 nm irradiation, utilizing visible blue light, which is generally considered a milder and more selective condition than UV light irradiation. The use of visible light is preferred due to its lower energy consumption, safer handling, and reduced potential for degradation of both the starting material and products, as well as a lower likelihood of side reactions and byproduct formation.^{51–53} To overcome these issues, we sought to design a gold photocatalyst that could operate (like iridium complexes) in this region of the visible light spectrum. Here, we report the facile synthesis of a novel gold sensitizer $[\text{Au}(\text{IPr})]_2(\text{DHIC})$ (**PhotAu3**) and its catalytic use in $[2 + 2]$ cycloaddition and *E/Z* isomerization reactions (Scheme 1).

Results and discussion

Inspired by the catalytic behavior of amido-complexes and the efficiency of carbazole in these key photocatalytically active architectures, 5,11-dihydroindolo[3,2-*b*]carbazole (**2**) was employed as a new nitrogen donor. The initial synthetic approach involved the weak base route with $[\text{Au}(\text{IPr})(\text{Cl})]$, K_2CO_3 and **2** (see SI). Unfortunately, this first attempt proved unsuccessful. We reasoned that an alternative gold nucleophilic synthon could be deployed to form the amido-gold bond.

To this end, $[\text{Au}(\text{IPr})(\text{OH})]$ (**1**)⁵⁴ was employed and cleanly led to the formation of **PhotAu3** (**3**) in ethyl acetate at room temperature, according to Scheme 2. The excellent solubility of **3** in THF permitted full spectroscopic characterization as well as the growth of single crystals, suitable for X-ray diffraction study. Photoluminescent spectroscopy was also possible in this solvent. The crystal structure of **PhotAu3** exhibits an asymmetric unit, consisting of two halves of symmetry-independent molecules, with different spatial disposition of the NHC and amido fragments (Fig. 1). One molecule (Fig. 1, left) presents the NHC moiety and the donor ligand on the same plane, while the other shows the two fragments lying perpendicular to one another. The second molecular conformation is similar to the one exhibited by **PhotAu1**.³⁵ Interestingly, other similar molecular structures bearing IPr show conformations with the two fragments set on the same plane.^{55–58} Absorption spectra of the free ligand **2** with that of **PhotAu3**, measured in THF solution, show that the last absorption band of 5,11-dihydroindolo[3,2-*b*]carbazole ends at around 420 nm, with the last peak at 405 nm (see SI). By adding the IPr-Au moiety to the donor ligand, the number of π -conjugated orbitals increases, thus red-shifting the absorption band of the complex: for **PhotAu3**



Scheme 2 Synthesis of **PhotAu3**.




Fig. 1 Crystal structure of **PhotAu3**, containing two different conformations of the **PhotAu3** molecule. Thermal displacement ellipsoids are shown at the 50% probability level. Selected bond distances (Å): Au1–N1 = 1.978(4), Au1–C1 = 2.003(4), Au2–N2 = 1.976(4), Au2–C2 = 2.005(3). CCDC number: 2465569.



two new bands are present with maximum absorption at 400, 440, and 470 nm, still absorbing up to 480 nm, indicating the ability to absorb from a standard blue LED at 450 nm.

The PL emission spectrum shows a peak of $\lambda_{\max} = 562$ nm, which indicates that **PhotAu3** has an E_T value of *ca.* 50.9 kcal mol⁻¹. This value is in line with reactivity found using the catalyst, which proved unreactive with substrates possessing higher values than 60 kcal mol⁻¹. The PLQY reported for the catalyst is 3% and the lifetime of the photoluminescence is 39 μ s in THF (Table 1).

Both of those values are inferior compared to those reported for **PhotAu1** and **PhotAu2**. The low k_r is most likely a result of emission from a ligand locally excited triplet state (³LE). The vibronic fine structure seen in the emission band is characteristic of LC-based emission, whereas CT-based emission is typically broad and featureless. To further support the ³LC assignment, the phosphorescence spectrum of **PhotAu3** in 2Me-THF was collected at 77 K. The 77 K spectrum and the room temperature spectra are the same, consistent with the emission at both temperatures coming from the ³LE state. We have found in previous (carbene)M(amido) complexes that cooling the sample in a polar solvent destabilizes the CT and often gives rise to ³LC emission.^{57,59–62} Here the ³LC emission is observed at both RT and 77 K. The gold is facilitating rapid intersystem crossing from the ¹ICT to the ³ICT, which internally converts to the ³LE state. The involvement of the ³LE state leads to the long-measured lifetime, which is beneficial for photocatalysis requiring diffusion of the substrate to the excited catalyst (Fig. 2).

Table 1 Key photophysical data of **PhotAu3** at room temperature in THF

λ_{\max}	Φ PLQY	τ (μ s)	k_r (10^3 s ⁻¹)	k_{nr} (10^5 s ⁻¹)
562	3%	39	7.7	2.5




Fig. 2 Molar absorptivity of the DHIC ligand and **PhotAu3** in THF, as well as photoluminescence spectra at room temperature of **PhotAu3** in THF and at 77 K in 2-Me-THF are shown.

Our new photosensitizer was initially tested in the intramolecular [2 + 2] cycloaddition of substrates that have previously been investigated using iridium catalysts and our gold photosensitizers. Cinnamyl ether **4** was initially selected as the model substrate.^{34,35,48} Compound **4** has previously been tested by Yoon and coworkers as a benchmark for EnT catalysis.⁶³ In their seminal work, 1 mol% of the [Ir(dF(CF₃)ppy)₂(dtbbpy)](PF₆) sensitizer was used and the reaction proceeded in very dilute DMSO solution (0.01 M). Ruthenium sensitizers were unable to mediate this reaction as triplet energies of styrenes are estimated to be *ca.* 60 kcal mol⁻¹.⁶⁴

Optimization focused mainly on decreasing the catalyst loading and reaction time while confirming that visible light could facilitate the reaction. We employed 450 nm irradiation wavelength at 100 W. Surprisingly, full conversion and a yield of 96% were obtained after 30 minutes with 0.5 mol% of the catalyst in both THF and 2Me-THF (entries 2 and 3, Table 2). By decreasing the catalyst loading to 0.25 mol%, a conversion of 72% was reached after 1 h and 83% after 2 h (entries 4 and 5, Table 2). Control experiments confirmed both the need for light and for photosensitizer (entries 6 and 7, Table 2). To confirm the need for gold, compound **2** (the carbazole ligand alone) was used as photocatalyst in a 10 mol% amount (entry 8, Table 2). At the 450 nm irradiation for 6 hours, a poor 23% conversion was obtained. These catalytic data agree with expected reactivity based on the absorption spectra of both catalyst and ligand 5,11-dihydroindolo[3,2-*b*]carbazole.

An additional family of spiroindoline precursor substrates was examined. This series has previously been examined by

Table 2 Intramolecular [2 + 2] cycloaddition of cinnamyl ether **4** with **PhotAu3** as photocatalyst^a

Entry	Mol%	Solvent	Time (h)	Conv. (%)	Yield (%)
1	0.5	THF	1	100	98
2	0.5	THF	0.5	100	96
3	0.5	2Me-THF	0.5	100	96 (90) ^b
4	0.25	THF	1	72	68
5	0.25	THF	2	83	73
6	0	THF	6	0	0
7	0.5 ^c	THF	6	0	0
8	10 ^d	THF	6	23	17
9	4 ^e	THF	1	0	0
10	4 ^f	THF	1	15	9


^a General reaction conditions: **4** (0.1 mmol, 1 eq.), **PhotAu3** (0.5–0.25 mol%), THF or 2Me-THF, room temperature, 0.5–6 h, under nitrogen, under 450 nm light irradiation at 100 W. NMR conversion and yield are calculated with 1,3,5-trimethoxybenzene as internal standard as average of two runs. ^b Isolated yield. ^c Reaction performed in the dark. ^d 5,11-Dihydroindolo[3,2-*b*]carbazole used as catalyst. ^e Benzophenone used as catalyst. ^f Thioxantone used as catalyst.



You,^{65–69} Fu,⁷⁰ Koenig¹⁶ and Oderinde.⁷¹ Polycyclic spiroindolines are privileged scaffolds found in numerous naturally occurring alkaloids.⁷² In order to generate the products by photocatalysis, it is often necessary to pre-functionalize the precursors, as branched indoles, with the introduction of an electron-withdrawing group on the C2 of the indole, since this functionalization has been shown to significantly lower the excitation threshold of the precursor molecule, thereby facilitating the photochemical transformation.⁶⁵ Keeping in mind the exciting results brought by compound **4**, the optimization of intramolecular [2 + 2] cycloaddition of tethered indoles at low catalyst loading and short reaction times was investigated (Table 2).


Both substrates **6** and **8** reached full conversion after 1 hour with 0.5 mol% of catalyst (1 mol% gold). Once again THF and 2Me-THF were selected as solvents to fully solubilize the sensitizer. Interestingly but not surprisingly, compound **10**, without the phenyl group at C2, did not react under these conditions as it possesses an E_T value of 65 kcal mol⁻¹ (ref. 61) which is well above the value at which the photocatalyst can operate (Scheme 3, see SI for reaction conditions).

The *E/Z* isomerization of double bonds *via* EnT photocatalysis was next explored. The synthesis of *Z*-cinnamyl ethers, their corresponding alcohols and, more generally, *Z*-alkenes present a significant challenge. Various conceptually distinct methodologies have been developed to tackle this challenge including Wittig reactions,⁷³ cross-coupling of alkenyl halides,⁷⁴ stereoselective semi-hydrogenation of alkynes,⁷⁵ *Z*-selective olefin metathesis,⁷⁶ and Peterson olefination reactions.⁷⁷ However, these reported transformations generally suffer from limited selectivity for the *Z*-configuration and often depend on energy-intensive reagents or costly transition metal-based catalysts.



Scheme 3 Intramolecular [2 + 2] cycloadditions of various branched indoles.

Therefore, an alternative strategy that enables the synthesis of the *Z*-isomer from the more accessible *E*-isomer, while allowing diversification from a common and easily available intermediate, is highly desirable. Various types of photosensitizers have been used recently for this purpose, obtaining the *Z* isomer from its *E* counterpart through *E/Z* isomerization. Those photocatalysts include iridium-based triplet photosensitizers,^{78–80} organic dyes such as riboflavin,⁸¹ aromatic ketones like 2-iodo-9-fluorenone,⁸² and covalent organic frameworks (COFs).^{83,84} Our first substrate candidate was cinnamyl alcohol (**12**) (Scheme 4), which has been reported by Wang and coworkers to yield the corresponding *Z* isomer with a ratio of *Z/E* = 81/19 in a system with state-of-the-art *fac*-Ir(ppy)₃ iridium sensitizer at 1 mol% in THF with 24 h of blue LED irradiation.⁸⁵ The experiments, reported in Table 3, show that reaction times longer than 2 hours were unnecessary to reach high *Z/E* ratio (entries 1, 2 and 3). A 0.1 M solution showed the same result as that of 0.2 M (entry 4). A catalyst



Scheme 4 *E/Z* isomerization reactions of cinnamyl alcohol and stilbene.

Table 3 Optimization of *E/Z* isomerization of cinnamyl alcohol **12**^a

Entry	Mol%	Solvent	Time (h)	Conc. (M)	<i>Z/E</i>
1	1	THF	16	0.2	74/26
2	1	THF	6	0.2	74/26
3	1	THF	2	0.2	74/26
4	1	THF	2	0.1	74/26
5	1	THF	1	0.2	70/30
6	2	THF	1	0.2	70/30
7	0.5	THF	2	0.2	74/26
8	0.5	2Me-THF	2	0.2	56/44
9	0.25	THF	2	0.2	27/73
10	0	THF	2	0.2	0/100
11	0.5 ^b	THF	2	0.2	0/100
12	4 ^c	THF	2	0.2	0/100
13	4 ^d	THF	2	0.2	23/77

^a General reaction conditions: 15 (0.2 mmol, 1 eq.), PhotAu3 (0.25–2.0 mol%), THF or 2Me-THF, room temperature, 1–16 h, under nitrogen, under 450 nm light irradiation at 100 W. *Z/E* ratios were confirmed by NMR spectroscopy, average of two runs. ^b Reaction performed in the dark. ^c Benzophenone used as catalyst. ^d Thioxanthone used as catalyst.



loading of 0.5 mol% allowed *Z/E* ratios of 74/26 (entry 7). These findings suggest an equilibrium-based reaction. Similar results were observed for the stilbene substrate (**14**) (Scheme 4), which reached a *Z/E* ratio of 79/21 with 0.5 mol% catalyst in just 1 hour. Reactions performed in THF proved better than those conducted in Me-THF in both cases.

A recent report disclosed the application of a new organic sensitizer that is able to yield *Z*-stilbene with *Z/E* = 74/26,⁸⁶ while with state-of-the-art iridium catalysts it is possible to reach *Z/E* ratios of 99/1.⁸⁷ Control experiments confirmed that for the two substrates examined with **PhotAu3**, both 450 nm irradiation and the sensitizer are necessary for the reaction to occur (entries 10 and 11, Table 3). To complete this section of the study, the isomerization reactions were carried out with common and commercially available photosensitizers such as benzophenone⁸⁸ and thioxanthone,⁸⁹ which have been used extensively in a variety of photochemical transformations.⁹⁰

These two photocatalysts proved completely ineffective in the isomerization reactions. This can be easily understood as the absorbance band of benzophenone does not go above *ca.* 380 nm while thioxanthone absorbs around 410 nm.⁹¹ The results shown in Tables 2 and 3 confirm the inefficacy of benzophenone in reactions involving **4** and **12**. Since thioxanthone can absorb a portion of visible light emitted, we were able to reach 15% conversion with **4** as substrate and 23/77 *Z/E* ratio for cinnamyl alcohol with 4 mol% of catalyst loading. **PhotAu3** proves significantly more effective.

Conclusions

A simple synthetic protocol to access a new digold photosensitizer, which has been fully characterized by spectroscopic and X-ray diffraction studies on a single crystal, has been established. This new photocatalyst is the first gold complex to be employed successfully for energy transfer photocatalysis under visible light irradiation and proves a viable alternative to blue light absorbing state-of-the-art iridium sensitizers. **PhotAu3** has been deployed as catalyst in three different intramolecular [2 + 2] cycloadditions, and two *E/Z* isomerizations at low catalyst loading and proved effective in short reaction times. Further studies involving extensive reaction scope in various transformations using **PhotAu3** are ongoing. The evolution of the synthetic design of gold complexes for photocatalysis is being further explored in our laboratories.

Author contributions

T. R.: conceptualization, investigation and synthesis and catalysis experiments; characterization NMR and/or SC-XRD; writing and editing of the manuscript. J. F.: photophysical analysis and characterization of the catalyst and writing part of the manuscript on the topic. K. V. H.: review and editing of manuscript; secured funding. M. E. T.: supervision of the work; writing and editing of the manuscript. T. S. A. H.: super-

vision of the work; writing and editing of the manuscript; secured funding. S. P. N.: conceptualization, supervision of the project and work; writing and editing of the manuscript; secured funding.

Conflicts of interest

There are no conflicts to declare.

Data availability

Additional data supporting this article have been included as part of the supplementary information (SI). Supplementary information is available. See DOI: <https://doi.org/10.1039/d5ob01503k>.

CCDC 2465569 contains the supplementary crystallographic data for this paper.⁹²

Acknowledgements

We gratefully acknowledge support from the Special Research Fund (BOF/24J/2023/084 to K. V. H.) of Ghent University, the Special Research Fund (BOF/STA/202102/003 to T. S. A. H.) of Ghent University and The Research Foundation – Flanders (FWO) (G0A6823N to S. P. N.). Umicore AG is thanked for generous gifts of materials.

References

- 1 L. Marzo, S. K. Pagire, O. Reiser and B. König, *Angew. Chem., Int. Ed.*, 2018, **57**, 10034–10072.
- 2 B. König, *Eur. J. Org. Chem.*, 2017, 1979–1981.
- 3 N. Corrigan, S. Shanmugam, J. Xu and C. Boyer, *Chem. Soc. Rev.*, 2016, **45**, 6165–6212.
- 4 T. P. Yoon, M. A. Ischay and J. Du, *Nat. Chem.*, 2010, **2**, 527–532.
- 5 N. A. Romero and D. A. Nicewicz, *Chem. Rev.*, 2016, **116**, 10075–10166.
- 6 S. P. Pitre and L. E. Overman, *Chem. Rev.*, 2022, **122**, 1717–1751.
- 7 M. Reckenthäler and A. G. Griesbeck, *Adv. Synth. Catal.*, 2013, **355**, 2727–2744.
- 8 F. Strieth-Kalthoff and F. Glorius, *Chem*, 2020, **6**, 1888–1903.
- 9 Q.-Q. Zhou, Y.-Q. Zou, L.-Q. Lu and W.-J. Xiao, *Angew. Chem., Int. Ed.*, 2019, **58**, 1586–1604.
- 10 J. Großkopf, T. Kratz, T. Rigotti and T. Bach, *Chem. Rev.*, 2022, **122**, 1626–1653.
- 11 F. Strieth-Kalthoff, M. J. James, M. Teders, L. Pitzer and F. Glorius, *Chem. Soc. Rev.*, 2018, **47**, 7190–7202.
- 12 S. Dutta, J. E. Erchinger, F. Strieth-Kalthoff, R. Kleinmans and F. Glorius, *Chem. Soc. Rev.*, 2024, **53**, 1068–1089.



- 13 W. C. de Souza, B. T. Matsuo, P. M. Matos, J. T. M. Correia, M. S. Santos, B. König and M. W. Paixão, *Chem. – Eur. J.*, 2021, **27**, 3722–3728.
- 14 X. Li, R. J. Kutta, C. Jandl, A. Bauer, P. Nuernberger and T. Bach, *Angew. Chem., Int. Ed.*, 2020, **59**, 21640–21647.
- 15 A. Hölzl-Hobmeier, A. Bauer, A. V. Silva, S. M. Huber, C. Bannwarth and T. Bach, *Nature*, 2018, **564**, 240–243.
- 16 A. B. Rolka and B. Koenig, *Org. Lett.*, 2020, **22**, 5035–5040.
- 17 E. Pylova, B. Lasorne, N. D. McClenaghan, G. Jonusauskas, M. Taillefer, S. N. Konchenko, A. Prieto and F. Jaroschik, *Chem. – Eur. J.*, 2024, **30**, e202401851.
- 18 T. R. Blum, Z. D. Miller, D. M. Bates, I. A. Guzei and T. P. Yoon, *Science*, 2016, **354**, 1391–1395.
- 19 F. M. Hörmann, C. Kerzig, T. S. Chung, A. Bauer, O. S. Wenger and T. Bach, *Angew. Chem.*, 2020, **132**, 9746–9755.
- 20 Y.-Q. Zou, S.-W. Duan, X.-G. Meng, X.-Q. Hu, S. Gao, J.-R. Chen and W.-J. Xiao, *Tetrahedron*, 2012, **68**, 6914–6919.
- 21 P.-K. Chow, G. Cheng, G. S. M. Tong, C. Ma, W.-M. Kwok, W.-H. Ang, C. Y.-S. Chung, C. Yang, F. Wang and C.-M. Che, *Chem. Sci.*, 2016, **7**, 6083–6098.
- 22 N. Hu, H. Jung, Y. Zheng, J. Lee, L. Zhang, Z. Ullah, X. Xie, K. Harms, M.-H. Baik and E. Meggers, *Angew. Chem., Int. Ed.*, 2018, **57**, 6242–6246.
- 23 M. Beliaeva, A. Belyaev, E. V. Grachova, A. Steffen and I. O. Koshevoy, *J. Am. Chem. Soc.*, 2021, **143**, 15045–15055.
- 24 C. Cruché, W. Neiderer and S. K. Collins, *ACS Catal.*, 2021, **11**, 8829–8836.
- 25 M. Bera, H. S. Hwang, T.-W. Um, S. M. Oh, S. Shin and E. J. Cho, *Org. Lett.*, 2022, **24**, 1774–1779.
- 26 D. Jespersen, B. Keen, J. I. Day, A. Singh, J. Briles, D. Mullins and J. D. I. Weaver, *Org. Process Res. Dev.*, 2019, **23**, 1087–1095.
- 27 L. Schmid, F. Glaser, R. Schaer and O. S. Wenger, *J. Am. Chem. Soc.*, 2022, **144**, 963–976.
- 28 T. O. Paulisch, L. A. Mai, F. Strieth-Kalthoff, M. J. James, C. Henkel, D. M. Guldi and F. Glorius, *Angew. Chem., Int. Ed.*, 2022, **61**, e202112695.
- 29 M. S. Lowry, J. I. Goldsmith, J. D. Slinker, R. Rohl, R. A. Pascal, G. G. Malliaras and S. Bernhard, *Chem. Mater.*, 2005, **17**, 5712–5719.
- 30 In February 2023, Umicore reports an average price of 29.6 € per mmol for Ir (<https://pmm.umicore.com/en/prices/iridium/>), and 12.1 € per mmol for gold (<https://pmm.umicore.com/en/prices/gold/>).
- 31 Z.-Y. Xu, Y. Luo, D.-W. Zhang, H. Wang, X.-W. Sun and Z.-T. Li, *Green Chem.*, 2020, **22**, 136–143.
- 32 J. Ma, F. Strieth-Kalthoff, T. Dalton, M. Freitag, J. L. Schwarz, K. Bergander, C. Daniliuc and F. Glorius, *Chem*, 2019, **5**, 2854–2864.
- 33 Strem catalogue numbers: 79-0000 [Au(IPr)(Cbz)]; 79-0002 [Au(SIPr)(Cbz)].
- 34 E. A. Martynova, V. A. Voloshkin, S. G. Guillet, F. Bru, M. Beliš, K. V. Hecke, C. S. J. Cazin and S. P. Nolan, *Chem. Sci.*, 2022, **13**, 6852–6857.
- 35 N. V. Tzouras, E. A. Martynova, X. Ma, T. Scattolin, B. Hupp, H. Busen, M. Saab, Z. Zhang, L. Falivene, G. Pisanò, K. Van Hecke, L. Cavallo, C. S. J. Cazin, A. Steffen and S. P. Nolan, *Chem. – Eur. J.*, 2021, **27**, 11904–11911.
- 36 V. A. Voloshkin, M. Villa, E. A. Martynova, M. Beliš, K. V. Hecke, P. Ceroni and S. P. Nolan, *Chem. Sci.*, 2024, **15**, 4571–4580.
- 37 S. G. Guillet, A. A. Logvinov, V. A. Voloshkin, E. A. Martynova and S. P. Nolan, *Org. Lett.*, 2023, **25**, 1403–1408.
- 38 E. A. Martynova, V. V. Voloshkin, M. Villa, A. Fiorentino, M. Beliš, K. V. Hecke, P. Ceroni and S. P. Nolan, *J. Catal.*, 2025, **442**, 115850.
- 39 Y. Zhao, S. G. Guillet, V. A. Voloshkin and S. P. Nolan, *Org. Lett.*, 2024, **26**, 10322–10327.
- 40 Y. Zhao, V. A. Voloshkin, E. A. Martynova, B. Maity, L. Cavallo and S. P. Nolan, *Chem. Commun.*, 2024, **60**, 3174–3177.
- 41 T. A. C. A. Bayrakdar, F. Nahra, D. Ormerod and S. P. Nolan, *J. Chem. Technol. Biotechnol.*, 2021, **96**, 3371–3377.
- 42 T. A. C. A. Bayrakdar, F. Nahra, J. V. Davis, M. M. Gamage, B. Captain, M. Temprado, M. Marazzi, M. Saab, K. V. Hecke, D. Ormerod, C. D. Hoff and S. P. Nolan, *Organometallics*, 2020, **39**, 2907–2916.
- 43 R. W. Y. Man, C.-H. Li, M. W. A. MacLean, O. V. Zenkina, M. T. Zamora, L. N. Saunders, A. Rousina-Webb, M. Nambo and C. M. Crudden, *J. Am. Chem. Soc.*, 2018, **140**, 1576–1579.
- 44 H. Shen, S. Xiang, Z. Xu, C. Liu, X. Li, C. Sun, S. Lin, B. K. Teo and N. Zheng, *Nano Res.*, 2020, **13**, 1908–1911.
- 45 M. Jin, R. Ando, M. J. Jellen, M. A. Garcia-Garibay and H. Ito, *J. Am. Chem. Soc.*, 2021, **143**, 1144–1153.
- 46 N. Mirzadeh, M. A. Bennett and S. K. Bhargava, *Coord. Chem. Rev.*, 2013, **257**, 2250–2273.
- 47 N. Mirzadeh, T. S. Reddy and S. K. Bhargava, *Coord. Chem. Rev.*, 2019, **388**, 343–359.
- 48 X. Ma, V. A. Voloshkin, E. A. Martynova, M. Beliš, M. Peng, M. Villa, N. V. Tzouras, W. Janssens, K. V. Hecke, P. Ceroni and S. P. Nolan, *Catal. Sci. Technol.*, 2023, **13**, 4168–4175.
- 49 J. Ma, J. Schaab, S. Paul, S. R. Forrest, P. I. Djurovich and M. E. Thompson, *J. Am. Chem. Soc.*, 2023, **145**, 20097–20108.
- 50 T. Li, D. G. Shlian, P. I. Djurovich and M. E. Thompson, *Chem. – Eur. J.*, 2021, **27**, 6191–6197.
- 51 S. Poplata, A. Tröster, Y.-Q. Zou and T. Bach, *Chem. Rev.*, 2016, **116**, 9748–9815.
- 52 N. Hoffmann, *Synthesis*, 2004, 481–495.
- 53 M. T. Crimmins and T. L. Reinhold, in *Organic Reactions*, John Wiley & Sons, Ltd, 2004, pp. 297–588.
- 54 S. Gaillard, A. M. Z. Slawin and S. P. Nolan, *Chem. Commun.*, 2010, **46**, 2742–2744.
- 55 S. Heo, B. H. Jhun, S. Woo, H. S. Kim, I. Kim, J. Kim, W. Son, Y. Jung, K. Kim, J. Cho and Y. You, *Adv. Opt. Mater.*, 2024, **12**, 2302148.



- 56 I. Ibni Hashim, N. V. Tzouras, W. Janssens, T. Scattolin, L. Bourda, S. Bhandary, K. Van Hecke, S. P. Nolan and C. S. J. Cazin, *Chem. – Eur. J.*, 2022, **28**, e202201224.
- 57 R. Hamze, M. Idris, D. S. Muthiah Ravinson, M. C. Jung, R. Haiges, P. I. Djurovich and M. E. Thompson, *Front. Chem.*, 2020, **8**, 401.
- 58 J. Ma, J. Schaab, P. Djurovich and M. E. Thompson, chemrxiv, 2023, preprint, chemrxiv:2023-03j7h, DOI: [10.26434/chemrxiv-2023-03j7h](https://doi.org/10.26434/chemrxiv-2023-03j7h).
- 59 R. Hamze, J. L. Peltier, D. Sylvinson, M. Jung, J. Cardenas, R. Haiges, M. Soleilhavoup, R. Jazzar, P. I. Djurovich, G. Bertrand and M. E. Thompson, *Science*, 2019, **363**, 601–606.
- 60 R. Hamze, S. Shi, S. C. Kapper, D. S. Muthiah Ravinson, L. Estergreen, M.-C. Jung, A. C. Tadler, R. Haiges, P. I. Djurovich, J. L. Peltier, R. Jazzar, G. Bertrand, S. E. Bradforth and M. E. Thompson, *J. Am. Chem. Soc.*, 2019, **141**, 8616–8626.
- 61 S. Shi, M. C. Jung, C. Coburn, A. Tadler, M. R. Daniel Sylvinson, P. I. Djurovich, S. R. Forrest and M. E. Thompson, *J. Am. Chem. Soc.*, 2019, **141**, 3576–3588.
- 62 T.-Y. Li, S.-J. Zheng, P. I. Djurovich and M. E. Thompson, *Chem. Rev.*, 2024, **124**, 4332–4392.
- 63 Z. Lu and T. P. Yoon, *Angew. Chem., Int. Ed.*, 2012, **51**, 10329–10332.
- 64 T. Ni, R. A. Caldwell and L. A. Melton, *J. Am. Chem. Soc.*, 1989, **111**, 457–464.
- 65 M. Zhu, C. Zheng, X. Zhang and S.-L. You, *J. Am. Chem. Soc.*, 2019, **141**, 2636–2644.
- 66 M. Zhu, X. Zhang, C. Zheng and S.-L. You, *ACS Catal.*, 2020, **10**, 12618–12626.
- 67 M. Zhu, H. Xu, X. Zhang, C. Zheng and S.-L. You, *Angew. Chem., Int. Ed.*, 2021, **60**, 7036–7040.
- 68 M. Zhu, X.-L. Huang, S. Sun, C. Zheng and S.-L. You, *J. Am. Chem. Soc.*, 2021, **143**, 13441–13449.
- 69 M. Zhu, X.-L. Huang, H. Xu, X. Zhang, C. Zheng and S.-L. You, *CCS Chem.*, 2020, **3**, 652–664.
- 70 Z. Zhang, D. Yi, M. Zhang, J. Wei, J. Lu, L. Yang, J. Wang, N. Hao, X. Pan, S. Zhang, S. Wei and Q. Fu, *ACS Catal.*, 2020, **10**, 10149–10156.
- 71 M. S. Oderinde, E. Mao, A. Ramirez, J. Pawluczyk, C. Jorge, L. A. M. Cornelius, J. Kempson, M. Vetrichelvan, M. Pitchai, A. Gupta, A. K. Gupta, N. A. Meanwell, A. Mathur and T. G. M. Dhar, *J. Am. Chem. Soc.*, 2020, **142**, 3094–3103.
- 72 M. Ishikura, K. Yamada and T. Abe, *Nat. Prod. Rep.*, 2010, **27**, 1630.
- 73 A. Pelter, E. Buss and E. Colclough, *J. Chem. Soc., Chem. Commun.*, 1987, 297–299.
- 74 E. Negishi, Z. Huang, G. Wang, S. Mohan, C. Wang and H. Hattori, *Acc. Chem. Res.*, 2008, **41**, 1474–1485.
- 75 H. Lindlar, *Helv. Chim. Acta*, 1952, **35**, 446–450.
- 76 M. J. Koh, R. K. M. Khan, S. Torker and A. H. Hoveyda, *Angew. Chem., Int. Ed.*, 2014, **53**, 1968–1972.
- 77 L. F. van Staden, D. Gravestock and D. J. Ager, *Chem. Soc. Rev.*, 2002, **31**, 195–200.
- 78 K. Nakajima, X. Guo and Y. Nishibayashi, *Chem. – Asian J.*, 2018, **13**, 3653–3657.
- 79 J. J. Molloy, J. B. Metternich, C. G. Daniliuc, A. J. B. Watson and R. Gilmour, *Angew. Chem., Int. Ed.*, 2018, **57**, 3168–3172.
- 80 H. Zhang, Q. Xu, L. Yu and S. Yu, *Eur. J. Org. Chem.*, 2020, 1472–1477.
- 81 K. Livingstone, M. Tenberge, F. Pape, C. G. Daniliuc, C. Jamieson and R. Gilmour, *Org. Lett.*, 2019, **21**, 9677–9680.
- 82 W. Cai, H. Fan, D. Ding, Y. Zhang and W. Wang, *Chem. Commun.*, 2017, **53**, 12918–12921.
- 83 M. Bhadra, S. Kandambeth, M. K. Sahoo, M. Addicoat, E. Balaraman and R. Banerjee, *J. Am. Chem. Soc.*, 2019, **141**, 6152–6156.
- 84 A. López-Magano, X. Solans-Monfort, N. Salaverri, L. Marzo, R. Mas-Ballesté and J. Alemán, *Sol. RRL*, 2024, **8**, 2300768.
- 85 H. Li, H. Chen, Y. Zhou, J. Huang, J. Yi, H. Zhao, W. Wang and L. Jing, *Chem. – Asian J.*, 2020, **15**, 555–559.
- 86 C.-X. Liu, Y. Yao, Z.-W. Zhou, S. Qin, Z.-P. Yu, F.-Y. Tao, W.-D. Li, X.-Q. Yu and N. Wang, *ChemSusChem*, 2025, **18**, e202401387.
- 87 D. C. Fabry, M. A. Ronge and M. Rueping, *Chem. – Eur. J.*, 2015, **21**, 5350–5354.
- 88 S. I. Faßbender, J. J. Molloy, C. Mück-Lichtenfeld and R. Gilmour, *Angew. Chem., Int. Ed.*, 2019, **58**, 18619–18626.
- 89 N. F. Nikitas, P. L. Gkizis and C. G. Kokotos, *Org. Biomol. Chem.*, 2021, **19**, 5237–5253.
- 90 I. K. Sideri, E. Voutyritsa and C. G. Kokotos, *Org. Biomol. Chem.*, 2018, **16**, 4596–4614.
- 91 J. Corpas, P. Mauleón, R. Gómez Arrayás and J. C. Carretero, *Adv. Synth. Catal.*, 2022, **364**, 1348–1370.
- 92 CCDC 2465569: Experimental Crystal Structure Determination, 2025, DOI: [10.5517/ccdc.csd.cc2nrmhk](https://doi.org/10.5517/ccdc.csd.cc2nrmhk).

