INORGANIC CHEMISTRY







FRONTIERS

RESEARCH ARTICLE

View Article Online
View Journal | View Issue



Cite this: *Inorg. Chem. Front.*, 2025, **12**, 3166

Organophosphonate- and dimethylarsinatefunctionalized hexamolybdates(v) and their solution and gas phase properties†

Here we report a comprehensive study on the synthesis and characterization of dimethylarsinate-functionalized phosphomolybdates(v), $[RPMo_6^VO_{15}(OH)_3\{AsO_2(CH_3)_2\}_3]^{2-}$ (R = H, HO, CH₃, HO₂CCH₂, HO₂CC₂H₄, C₆H₅, 4-FC₆H₄, 4-F₃COC₆H₄) and the monoanionic mixed-valent heptamolybdate $[HOMo^{VI}Mo_6^VO_{15}(OH)_3\{AsO_2(CH_3)_2\}_3]^{-}$. These nine novel polyanions feature a reduced, cyclic hexanuclear $\{Mo_6^VO_{24}\}$ core, decorated peripherally by three dimethylarsinate ligands and centrally by six different organophosphonate groups including fluorinated ones, as well as phosphate and phosphite. The polyanions were synthesized under simple one-pot aqueous solution conditions and characterized (i) in the solid state by single-crystal and powder XRD, TGA, elemental analysis, (ii) in solution by multinuclear (1H , ^{31}P , ^{19}F , ^{13}C) NMR, and (iii) in the gas phase by ESI mass spectrometry.

Received 14th December 2024, Accepted 17th February 2025 DOI: 10.1039/d4qi03215b

rsc.li/frontiers-inorganic

1. Introduction

In materials science, the burgeoning domain of hybrid organic-inorganic materials carries profound implications, driving innovation in the development of multifunctional materials. The synergistic combination of the structural intricacies and electronic characteristics inherent to polyoxometalates (POMs), 1a,b which are discrete, well-defined, anionic metal-oxo clusters composed of early transition metal ions (M = Mo^{VI}, W^{VI}, V^V, Nb^V and Ta^V) in high oxidation states, with the flexibility, tunability and functionality of organic or organometallic components, 1c,d represent a rapidly growing field of research with unique, potential applications in areas such as catalysis, 1ef molecular nanosciences, 1g,h photo (electro-) chemistry, 1i,j biotechnology, and medicine. 1k-m New hybrid architectures could be generated through interaction of classical lacunary POMs with organic or organometallic groups, or via in situ self-assembly between metal building blocks and O-donating organic ligands. These hybrid POMs, featuring reactive sites, offer multifaceted platforms for subsequent post-functionalization.²

The conventional domains of polyoxomolybdates and -tung-states constitute the primary subsets within POM frameworks. The formation of polyoxomolybdate-organic hybrids is highly desirable due to the attractive redox properties and stability of multielectron-reduced polyoxomolybdates. Nevertheless, it poses a notable hurdle due to the increased lability and the consequent instability associated with polyoxomolybdates, resulting in limited documentation on the synthesis of molecular hybrids involving lacunary polyoxomolybdates and multidentate organic ligands. The investigation of *in situ* hybrid POMs gathered significant impetus after Pope and coworkers structurally characterized the organoarsenic functionalized polyoxomolybdates, $[(RAs)_2Mo_6O_{24}]^{4-}$ ($R = CH_3$, C_6H_5 , C_6H_5 , 4c,d initially proposed by Rosenheim and co-workers.

The first organophosphonate derivatized heteropolyanions were the pentamolybdobisphosphonate complexes, 5a [(RP)₂Mo $_5^{VI}$ O₂₁]⁴⁻ (R = H, CH₃, C₂H₅, C₆H₅, C₂H₄NH₃⁺, p-CH₂C₆H₄NH₃⁺), reported by Pope's group in 1975. Subsequently, in 1976, Stalick and Quicksall elucidated the structures of [(CH₃P)₂Mo₅O₂₁]⁴⁻ and [(NH₃C₂H₄P)₂Mo₅O₂₁]^{2-,5b} The crystal structure of [(HP)₂Mo₅O₂₁]⁴⁻ was published by Sasaki's group in 1988, 5c while Lyxell and Strandberg concurrently reported crystal structure of [(C₆H₅P)₂Mo₅O₂₁]^{4-,5d} Subsequently, Kortz's group has published a series of [(RP)₂Mo $_5^{VI}$ O₂₁]⁴⁻ derivatives, featuring aliphatic phosphonocarboxylate groups (R = O₂CCH₂, O₂CC₂H₄)^{6a} and aryl carboxyphenyl phosphonate groups (R = 4-O₂CC₆H₄, 4-O₂CC₆H₄CH₂). 6b The

^aSchool of Science, Constructor University, Campus Ring 1, 28759 Bremen, Germany. E-mail: ukortz@constructor.university

^bDepartment of Chemistry, BITS Pilani K. K. Birla Goa Campus, 403726 Goa, India † Electronic supplementary information (ESI) available. CCDC 2394921–2394926 and 2406188–2406190. For ESI and crystallographic data in CIF or other electronic format see DOI: https://doi.org/10.1039/d4qi03215b

incorporation of enantiopure aminoalkylphosphonates (R = CH₃CH(NH₃), CH₃CH(CH₃)CH(NH₃)) induced the formation of self-assembled fibrillar nanostructures with heightened optically activity.6c The hexamolybdic wheel structural $[RPMo_6^{VI}O_{21}(O_2CCH_2NH_3)_3]^{2-}$ (R = OH, CH₃, C₂H₅, H) underwent covalent modification at two distinct sites: peripherally by glycine molecules and centrally by organophosphonate group, vielding molybdenum oxo frameworks characterized by their flexibility in both steric and electronic aspects.^{6d} Additionally, similar structural archetype has been reported earlier, $[XMo_6O_{21}(O_2CRNH_3)_3]^{n-}$ (n = 2, X = Se^{IV}, Te^{IV}; n = 3, X = As^{III}, Sb^{III} , Bi^{III} ; $R = CH_2$, C_2H_4 , C_3H_6 , $CHCH_3$, $CH(CH_2)_4NH_2$), where lone-pair containing heteroatoms occupy the central position and are surrounded by amino acids bound peripherally via their carboxylate functions.6e

In recent years Kortz's group has reported a number of polyoxo-noble-metalate clusters functionalized by dimethylarsinate (also known as cacodylate) ligands. The inclusion of this functional moiety in reduced polyoxomolybdate chemistry led to the discovery of an organofunctionalized mixed-valent polyoxomolybdate wheel $\left[Mo_{18}^{VI}Mo_{12}^{V}O_{84}\left\{AsO_{2}\left(CH_{3}\right)_{2}\right\}_{18}\right]^{18-}$ (Mo₃₀) and a neutral, reduced tetramolybdate cubane cluster [Mo₄^VO₈{AsO₂(CH₃)₂}₄] (Mo₄).⁸ The introduction of phosphorous into the same system, at elevated pH levels, resulted in a series of reduced hexanuclear, dimethylarsinate- functionalized phosphomolybdenum clusters, with general formula $[RPMo_6^VO_{15}(OH)_3\{AsO_2(CH_3)_2\}_3]^{n-}$ reminiscent Anderson archetype, where the polyoxoanion core can be centrally modified systematically and rationally through incorporation of phosphite, phosphate and hydrolytically stable alkyl and aryl phosphonates, represented as RPO₃²⁻. This particular type of hexanuclear molybdic wheel was initially isolated by Haushalter's group as the reduced molybdenum phosphate, {P₄Mo^V₆}, where peripheral phosphates and hydroxo-bridges mutually connect {MoV2O4} dimeric structural units.9 Subsequent studies demonstrated the versatility of this prototype, where PO₄³⁻ could be substituted with C₆H₅PO₃²⁻, ^{10a-c} $C_6H_5AsO_3^{2-,10d}CO_3^{2-,10e}$ and $SO_3^{2-,10f}$ Thiomolybdate analogues, $[X_4Mo_6S_6O_6(OH)_3]^{5-}$ $(X = HPO_4, HAsO_4 as assem$ bling groups) with dimolybdenum pairs bridged by six sulfur atoms, have also been documented, 11a with a study showcasing regioselective substitution of sulfur atoms in oxothio analogue {P₄Mo₆S₆} to obtain {P₄Mo₆S₃O₃} illustrated by NMR reported later. 11b The mixed-valent MoV/MoVI molecular anion, [Mo₇O₁₆(O₃PCH₂PO₃)₃]⁸⁻{P₆Mo₇}, anchored by peripheral diphosphonate groups could accommodate Mo^{VI}O₄ tetrahedron in the center, due to the larger aperture created from corner-sharing arrangement of the molybdenum dimers differing from the edge-sharing mode observed in $\{P_4Mo_6\}$.¹²

Now we decided to investigate the system of molybdate in the presence of cacodylate and phosphonates in aqueous solution under reducing conditions in detail, and we isolated a new family of eight functionalized polyoxomolybdates(v) with interesting solid state, solution and gas phase properties.

Experimental details

2.1 Synthesis of $Na_{1.8}(NH_4)_{0.2}[HOPMo_6^VO_{15}(OH)_3\{AsO_2(CH_3)_2\}_3]$ 0.8NH₄Cl·6H₂O (NaNH₄-PMo₆)

 $(NH_4)_6Mo_7O_{24}\cdot 4H_2O$ (0.265 g, 0.215 mmol) and $N_2H_4\cdot 2HCl$ (0.052 g, 0.5 mmol) were dissolved completely in 10 mL 0.5 M sodium dimethylarsinate buffer (prepared as follows: 0.5 M aqueous solution of dimethylarsinic (cacodylic) acid (pH 4) adjusted to pH 7 by adding NaOH pellets). To this mixture, sodium phosphate, Na₃PO₄ (0.041 g, 0.25 mmol) was added followed by pH adjustment to 6.8-6.9 by dropwise addition of 6M NaOH solution. The initial dark green solution, which progressively turned dark red, was heated at 80 °C for 1 hour in closed vial. Dark-red block-shaped crystals of NaNH4-PMO6 started to form within 3-4 days and were collected after 1 week and air dried. Isolated yield: 0.140 g (37% based on P). Elemental analysis (%): calcd (found) for NaNH₄-PMo₆: Na, 2.70 (2.68); Mo, 37.6 (38.2); As, 14.7 (14.7); P, 2.0 (2.3); C, 4.7 (4.8); H, 2.55 (2.62); N, 1.10 (1.33). FT-IR data (KBr pellet) in cm⁻¹: 3410–3232 (s) $[\nu(O-H)]$, 3025 (w), 2926 (w) $[\nu(C-H)]$, 2091 (w), 1632 (m) $[\delta(O-H)]$, 1407 (m) $[\delta(C-H)]$, 1278 (m), 1257 (sh) $[\nu(As-C)]$, 1089 (m), 1010 (w) $[\nu(P-O)]$, 957–920 (s) $[\nu(Mo=O)]$, 857 (sh), 825 (s) $[\nu(As-O)]$, 751 (s), 655 (w), 608, 491 (s), 459 (s) $[\delta(Mo-O(Mo))].$

2.2 Synthesis of $Na_{1,2}H_{0,8}[HPMO_6^VO_{15}(OH)_3\{AsO_2(CH_3)_2\}_3]$. 11H₂O (Na-HPMo₆)

The polyanion HPMo₆ was prepared using the same method as that for PMo₆, except that sodium phosphite, Na₂HPO₃·5H₂O (0.054 g, 0.25 mmol) was used instead of sodium phosphate. The pH of the reaction mixture was adjusted to 6.8 with 6M NaOH before heating at 80 °C. Dark-red block crystals were isolated after 1 week and air dried. Isolated yield: 0.145 g (38% based on P). Elemental analysis (%): calcd (found) for Na-HPMo₆: Na, 1.80 (1.80); Mo, 37.5 (36.8); As, 14.6 (14.3); P, 2.0 (2.1); C, 4.7 (4.8); H, 2.8 (2.6). FT-IR data (KBr pellet) in cm⁻¹: 3400 (s) $[\nu(O-H)]$, 3019(w), 2928 (w) $[\nu(C-H)]$, 2427 (w) $[\nu(P-H)]$, 2088 (w), 1632 (m) $[\delta(O-H)]$, 1404 (m) $[\delta(C-H)]$, 1274 (w), 1257 (sh) $[\nu(As-C)]$, 1143 (sh), 1072(s) $[\nu(P-O)]$, 972 (s), 944 (s) $[\nu(Mo=O)]$, 852 (sh), 820 (s) $[\nu(As-O)]$, 752 (s), 656 (m), 615 (w), 540 (m), 496 (s), 460 (m) $[\delta(Mo-O(Mo))]$.

2.3 Synthesis of NaNH₄[CH₃PMo $_6^V$ O₁₅(OH)₃{AsO₂(CH₃)₂}₃]. 11H₂O (NaNH₄-CH₃PMo₆)

The polyanion CH₃PMo₆ was prepared similarly to PMo₆, substituting methyl phosphonic acid (CH₃PO₃H₂) (0.024 g, 0.25 mmol) for sodium phosphate. pH of the reaction mixture was adjusted to 6.8 with 6 M NaOH prior to stirring and heating at 80 °C for 1 hour. As the reaction mixture cooled to room temperature, dark-red block crystals suitable for single crystal XRD emerged within a day at the bottom of the vial. After a week of crystallization in open air, the compound was collected and characterized by IR spectroscopy, resulting in a total yield of 0.180 g (46% based on phosphorus). Elemental analysis (%): calcd (found) for NaNH₄-CH₃PMo₆: Na, 1.5 (1.5); Mo, 36.8 (36.7); As, 14.4 (14.7); P, 2.0 (1.9); C, 5.38 (5.43); H,

3.22 (3.39); N, 0.90 (1.06). FT-IR data (KBr pellet) in cm⁻¹: 3488 (s) $[\nu(O-H)]$, 3017 (w), 2926 (w) $[\nu(C-H)]$, 1629 (m) $[\delta(O-H)]$, 1404 (m) $[\delta(C-H)]$, 1311–1274 (w) $[\nu(P-C)]$, 1275 (m), 1257 (sh) $[\nu(As-C)]$, 1116(sh), 1051 (m) $[\nu(P-O)]$, 999 (w), 967 (s), 939 (m) $[\nu(Mo=O)]$, 854 (sh), 822 (s) $[\nu(As-O)]$, 778 (w), 750 (m), 655 (w), 616 (w), 552 (m), 493 (s), 461 (s) $[\delta(Mo-O(MO))]$.

2.4 Synthesis of $Na_{1.6}(NH_4)_{0.4}[HO_2CCH_2PMO_6^VO_{15}(OH)_3 \{AsO_2(CH_3)_2\}_3]\cdot 0.5NH_4Cl\cdot 9H_2O\ (NaNH_4-HO_2CCH_2PMO_6)$

The polyanion HO₂CCH₂PMo₆ was prepared similarly to PMo₆, substituting phosphonoacetic acid (H₂O₃PCH₂COOH) (0.035 g, 0.25 mmol) for sodium phosphate. pH of the reaction mixture was adjusted to 6.8 with 6 M NaOH and the dark red solution is heated at 80 °C for 1 hour. The dark-red block-shaped crystals of NaNH4-HO2CCH2PMo6 that formed within 2 days in the vial kept closed at room temperature, was later characterized by single crystal XRD. After 2 weeks of open-air crystallization, the compound was collected and characterized by IR. Total isolated yield: 0.125 g (31% based on P). Elemental analysis (%): calcd (found) for NaNH₄-HO₂CCH₂PMo₆: Na, 2.33 (2.43); Mo, 36.5 (37.0); As, 14.3 (14.2); P, 2.0 (2.2); C, 6.10 (5.97); H, 2.82 (3.08); N, 0.80 (0.84). FT-IR data (KBr pellet) in cm⁻¹: 3404 (s), 3020 (w) $[\nu(O-H)]$, 2927 (w), 2085 (w) $[\nu(C-H)]$, 1635 (m) $[\delta(O-H)]$, 1581 (m) $[\nu(C=O)]$, 1405 (m) $[\delta(C-H)]$, 1378 (m) $[\nu(P-C)]$, 1275 (w), 1259 (sh) $[\nu(As-C)]$, 1215 (w), 1132 (w), 1056 (m) $[\nu(P-O)]$, 967 (s), 942 (m) $[\nu(Mo=O)]$, 851 (sh), 822 (s) $[\nu(As-O)]$, 754 (s), 656 (w), 612 (w), 553 (m), 495 (s), 465 (s) $[\delta(Mo-O(Mo))].$

2.5 Synthesis of NaNH₄[$HO_2CC_2H_4PMo_6^VO_{15}(OH)_3$ { $AsO_2(CH_3)_2$ }₃]·10H₂O (NaNH₄- $HO_2CC_2H_4PMo_6$)

The synthesis of HO₂CC₂H₄PMo₆ was accomplished by using a procedure identical to that of HO2CCH2PMo6, except for substituting phosphonoacetic acid with phosphonopropionic acid (H₂O₃PC₂H₄COOH) (0.039 g, 0.25 mmol). The pH of the reaction mixture was adjusted to 6.8 with 6M NaOH before heating at 80 °C for 1 hour. Dark-red block-shaped crystals started forming at the bottom of the vial within two days. Total isolated yield after 2 weeks: 0.120 g (30% based on P). Elemental analysis (%): calcd (found) for NaNH₄-HO₂CC₂H₄PMo₆: Na, 1.43 (1.72); Mo, 35.9 (36.6); As, 14.0 (14.5); P, 1.9 (1.9); C, 6.74 (6.57); H, 3.14 (3.40); N, 0.87 (1.15). FT-IR data (KBr pellet) in cm⁻¹: 3400 (s), 3247 (s) $[\nu(O-H)]$, 3008 (w), 2923 (w) $[\nu(C-H)]$, 2094 (w), 1631 (m) $[\delta(O-H)]$, 1559 (m) $[\nu(C=O)]$, 1444 (sh) $[\nu(P-D)]$ C)], 1403 (m) $[\delta(C-H)]$, 1320 (w), 1273 (w), 1254 (sh) $[\nu(As-C)]$, 1258 (w), 1213 (w), 1050 (m) $[\nu(P-O)]$, 990 (sh), 966 (s), 938 (s) $[\nu(Mo=O)]$, 852 (sh), 822 (s) $[\nu(As-O)]$, 748 (s), 656 (w), 612 (w), 549 (m), 494 (s), 462 (s) $[\delta(Mo-O(Mo))]$.

2.6 Synthesis of Na $_{0.7}$ (NH $_4$) $_{1.3}$ [C $_6$ H $_5$ PMO $_6^V$ O $_{15}$ (OH) $_3$ {AsO $_2$ (CH $_3$) $_2$ } $_3$]·9H $_2$ O (NaNH $_4$ -C $_6$ H $_5$ PMO $_6$)

The polyanion $C_6H_5PMo_6$ was synthesized using a method analogous to PMo_6 , replacing sodium phosphate with phenyl phosphonic acid ($C_6H_5PO_3H_2$) (0.040 g, 0.25 mmol). The pH of the solution is adjusted to 6.8 with 6M NaOH, followed by stirring and heating at 80 °C for 1 hour in a closed vial. The reac-

tion mixture is later kept for cooling at room temperature in the closed vial for two days to procure good quality red block crystals. After isolation of the initial product, the dark red solution is kept for crystallization in open air for 2 weeks. The combined isolated yield after two weeks is 0.160 g (39% based on P). IR was used to characterize the products separated at different time intervals. Elemental analysis (%): calcd (found) for NaNH₄-C₆H₅PMo₆: Na, 1.01 (1.15); Mo, 36.3 (36.8); As, 14.2 (14.4); P, 2.0 (2.0); C, 9.08 (9.01); H, 3.12 (3.37); N, 1.15 (1.19). FT-IR data (KBr pellet) in cm⁻¹: 3416 (s), 3218 (s) $[\nu(O-H)]$, 3017 (w), 2928 (w) $[\nu(C-H)]$, 2071 (w), 1623 (m) $[\delta(O-H)]$, 1441 (sh) $[\nu(C=C), \nu(P-C)]$, 1404 (m) $[\delta(C-H)]$, 1275 (w), 1255 (sh) $[\nu(As-C)]$, 1142 (m), 1079 (sh), 1058 (m) $[\nu(P-O)]$, 965 (s), 940 (m) $[\nu(Mo=O)]$, 854 (sh), 821 (s) $[\nu(As-O)]$, 748 (s), 655 (w), 611(w), 578 (w), 567 (sh), 557 (m), 493 (s), 461 (s) [δ(Mo-O (Mo))].

2.7 Synthesis of (4-fluorophenyl) phosphonic acid (L_{PF})

The reported synthetic procedure¹³ was slightly modified and optimized as follows: 1-bromo-4-fluoro benzene (9.0 mmol), triethyl phosphite (1.89 mL, 10.8 mmol), and nickel chloride catalyst (0.117 g, 0.90 mmol) were accurately weighed and transferred into a standard microwave reaction vessel equipped with a pressure controller. The assembly was placed in a CEM Discover MW reactor, and the reaction mixture was irradiated at 140 °C for 5 hours under continuous stirring at a power setting of 200 W. After the reaction, the crude product was purified by vacuum distillation to obtain pure diethyl (4-fluorophenyl) phosphonate. ¹H NMR (CDCl₃, 400 MHz, ppm): δ 1.18 (t, 6H, CH₃), 3.99 (m, 4H, CH₂CH₃), 7.01 (m, 2H, Ar), 7.68 (m, 2H, Ar); ¹⁹F NMR (CDCl₃, 376.5 MHz, ppm): δ –106; ³¹P{¹H} NMR (CDCl₃, 161.83 MHz, ppm): δ 18.3; ¹³C{¹H} NMR (CDCl₃, 100 MHz, ppm): δ 16.21, 62.12, 115.5, 124.4 (d, ${}^{1}J_{CP}$ = 189.4 Hz), 134.4, 165.5 (d, ${}^{1}J_{CF}$ = 252.1 Hz).

The phosphonic ester was then refluxed at 110 °C for 48 hours in 20 mL of concentrated hydrochloric acid (12 M). The resulting mixture was fully evaporated using a rotavapor, and the resulting white solid was washed with n-hexane and dried, yielding (4-fluorophenyl) phosphonic acid ($\mathbf{L_{PF}}$) as a fine white powder. ¹H NMR (D₂O, 400 MHz, ppm): δ 7.04 (m, 2H, Ar), 7.57 (m, 2H, Ar); ¹⁹F NMR (D₂O, 376.5 MHz, ppm): δ -107.9; ³¹P{¹H} NMR (D₂O, 161.83 MHz, ppm): δ 15.4; ¹³C{¹H} NMR (D₂O, 100 MHz, ppm): δ 115.7, 127.2 (d, ¹ J_{CP} = 183.5 Hz), 132.8, 164.4 (d, ¹ J_{CF} = 248.4 Hz).

2.8 Synthesis of NaNH₄[4-FC₆H₄PMo $_6^V$ O₁₅(OH)₃{AsO₂(CH₃)₂}₃]-8H₂O (NaNH₄-FC₆H₄PMo₆)

 $(NH_4)_6Mo_7O_{24}\cdot 4H_2O$ (0.053 g, 0.04 mmol), $N_2H_4\cdot 2HCl$ (0.011 g, 0.1 mmol) and (4-fluorophenyl) phosphonic acid (L_{PF}), $FC_6H_4PO_3H_2$ (0.009 g, 0.05 mmol) is added to 3 mL 0.5 M sodium dimethylarsinate buffer (pH 7) and the resulting mixture is stirred at room temperature for 1 hour. Dark-red block-shaped crystals start forming after one to two days. Total isolated yield after one week is 0.035 g (42% based on P). Elemental analysis (%): calcd (found) for $NaNH_4$ - $FC_6H_4PMo_6$: Na, 1.45 (1.63); Mo, 36.2 (38.4); As, 14.2 (14.7); P, 1.95 (1.82); C,

9.07 (8.14); H, 2.85 (2.95); N, 0.88 (1.10). FT-IR data (KBr pellet) in cm⁻¹: 3406 (s) [ν (O-H)], 3027 (w), 2929 (w) [ν (C-H)], 2079 (w), 1631 (m) [δ (O-H)], 1499 (w) [ν (C-F)], 1441 (sh) [ν (C=C), ν (P-C)], 1401 (m) [δ (C-H)], 1281 (w), 1223 (w) [ν (As-C)], 1166 (sh), 1137 (w), 1062 (m) [ν (P-O)], 965(s), 937(sh) [ν (Mo=O)], 827 (s) [ν (As-O)], 747 (s), 661 (w), 609 (w), 574 (w), 528 (sh), 488 (s), 466 (s) [δ (Mo-O(Mo))].

2.9 Synthesis of (4- trifluoromethoxyphenyl) phosphonic acid $\left(L_{POCF3}\right)$

The reported synthetic procedure ¹³ was modified with the reaction mixture containing 1-bromo-4-(trifluoromethoxy) benzene (1.34 mL, 9.0 mmol), triethyl phosphite (1.89 mL, 10.8 mmol), and NiCl₂ (0.117 g, 0.90 mmol) being heated up to 130 °C for 5 hours with a power input of 150 W in a standard microwave reaction pressure vial. This reaction was repeated 5 times to obtain approximately 15 mL of crude product which was further purified through vacuum distillation yielding the desired product diethyl (4-trifluoromethoxyphenyl) phosphonate ester. ¹H NMR (CDCl₃, 400 MHz, ppm): δ 1.28 (t, 6H, CH₃), 4.07 (m, 4H, CH₂CH₃), 7.24 (m, 2H, Ar), 7.82 (m, 2H, Ar); ¹⁹F NMR (CDCl₃, 376.5 MHz, ppm): δ –57.7; ³¹P{¹H} NMR (CDCl₃, 161.83 MHz, ppm): δ 17.6; ¹³C{¹H} NMR (CDCl₃, 100 MHz, ppm): δ 16.3, 62.3, 120.3 (q, ¹ $J_{\rm CF}$ = 256.7 Hz), 127.2 (d, ¹ $J_{\rm CP}$ = 188.9 Hz), 120.5 (d, ² $J_{\rm CP}$ = 15.4 Hz), 133.8 (d, ³ $J_{\rm CP}$ = 10.6 Hz), 152.2.

(4-Trifluoromethoxyphenyl) phosphonic acid (L_{POCF3}) was isolated as a fine white powder after refluxing the phosphonic ester in a similar procedure used for L_{PF} . ¹H NMR (D₂O, 400 MHz, ppm): δ 7.39 (d, 2H, Ar), 7.74 (m, 2H, Ar); ¹⁹F NMR (D₂O, 376.5 MHz, ppm): δ -56.6; ³¹P{¹H} NMR (D₂O, 161.83 MHz, ppm): δ 11.4; ¹³C{¹H} NMR (D₂O, 100 MHz, ppm): δ 120.5 (q, ¹ J_{CF} = 255 Hz), 121.1 (d, ² J_{CP} = 13.4 Hz), 133.4 (d, ³ J_{CP} = 9.9 Hz), 133.9 (d, ¹ J_{CP} = 173.8 Hz), 150.6.

2.10 Synthesis of Na_{0.5}(NH₄)_{1.5}[4-F₃COC₆H₄PMo₆^VO₁₅(OH)₃ {AsO₂(CH₃)₂}₃]·0.3NaO₂As(CH₃)₂·9H₂O (NaNH₄-F₃COC₆H₄PMo₆)

The polyanion F₃COC₆H₄PMo₆ was synthesized in a similar manner to FC₆H₄PMo₆ by replacing (4-fluorophenyl) phosphonic acid with (4-trifluoromethoxyphenyl) phosphonic acid (L_{POCE3}) , $F_3COC_6H_4PO_3H_2$, (0.012 g, 0.05 mmol). The reaction mixture (pH 7) was continuously heated and stirred at 80 °C for 1 hour and dark-red block-shaped crystals started forming within a day after cooling to room temperature. The total isolated yield of the compound after one week is 0.030 g (38% based on P). Elemental analysis (%): calcd (found) for NaNH₄- $F_3COC_6H_4PMo_6$: Na, 1.07 (1.08); Mo, 33.5 (35.0); As, 14.4 (15.2); P, 1.80 (1.32); C, 9.50 (8.66); H, 2.98 (3.20); N, 1.22 (1.36). FT-IR data (KBr pellet) in cm⁻¹: 3398 (s), 3247 (s) $[\nu(O-$ H)], 3022 (w), 2931 (w) $[\nu(C-H)]$, 2077 (w), 1628 (m) $[\delta(O-H)]$, 1496 (w) $[\nu(C-F)]$, 1456 (sh) $[\nu(C-C)]$, $\nu(P-C)$, 1401 (m) $[\delta(C-C)]$ H)], 1261 (m), 1226 (w) [ν(As-C)], 1166 (w), 1140 (sh), 1068(m) $[\nu(P-O)]$, 966 (s), 939 (sh) $[\nu(Mo=O)]$, 829 (s) $[\nu(As-O)]$, 751 (s), 657 (w), 618 (sh), 580 (m), 495 (s), 465 (s) δ (Mo-O(Mo))].

2.11 Synthesis of Na[HOMo VI Mo $_{6}^{V}$ O₁₅(OH)₃{AsO₂(CH₃)₂}₃]·NH₄O₂As(CH₃)₂·11H₂O (Na-Mo₇)

The mixed-valent polyanion \mathbf{Mo}_7 was synthesized by dissolving $(\mathrm{NH}_4)_6\mathrm{Mo}_7\mathrm{O}_{24}\cdot 4\mathrm{H}_2\mathrm{O}$ (0.265 g, 0.215 mmol) in 5 mL 0.5 M sodium cacodylate buffer (pH 7) followed by addition of $\mathrm{N}_2\mathrm{H}_4\cdot 2\mathrm{HCl}$ (0.52 g, 0.5 mmol) with continuous stirring. pH of the dark red solution is adjusted to 6.9–7.0 with 6 M NaOH followed by heating at 50 °C for 1 hour with continuous stirring. The isolated crystal yield after 2 weeks is around 0.040 g (11% based on Mo). Elemental analysis (%): calcd (found) for Na-Mo₇: Na, 1.30 (1.69); Mo, 38.0 (37.7); As, 17.1 (18.3); C, 5.44 (5.12); H, 3.08 (3.20); N, 0.79 (0.83). FT-IR data (KBr pellet) in cm⁻¹: 3403 (s) [ν (O-H)], 3010 (w), 2922 (w) [ν (C-H)], 2054 (w), 1633 (m) [δ (O-H)], 1401 (m) [δ (C-H)], 1274 (w), 1255 (sh) [ν (As-C)], 969 (s) [ν (Mo=O)], 849 (sh), 820 (s) [ν (As-O)], 747 (s), 653 (w), 613 (w), 555 (sh), 524 (sh), 495 (s), 455 (s) [δ (Mo-O(Mo))].

3. Results and discussion

The polyanions $[HPMo_6^VO_{15}(OH)_3\{AsO_2(CH_3)_2\}_3]^{2-}$ $(HPMo_6)_3$ $[HOPMo_6^VO_{15}(OH)_3\{AsO_2(CH_3)_2\}_3]^{2-}$ $(PMo_6)_{,}$ $[CH_3PMo_6^VO_{15}(OH)_3$ $\{AsO_2(CH_3)_2\}_3^{2-}$ (CH_3PMo_6) [HO₂CCH₂PMo₆VO₁₅(OH)₃{AsO₂ $(CH_3)_2\}_3^{2-}$ $(HO_2CCH_2PMO_6)$ $[HO_2CC_2H_4PMO_6^VO_{15}(OH)_3\{AsO_2\}]$ $(CH_3)_2\}_3]^{2-}$ (HO₂CC₂H₄PMo₆) $[C_6H_5PMo_6^VO_{15}(OH)_3\{AsO_2\}]$ $(CH_3)_2\}_3^{2-}$ $(C_6H_5PMo_6)$, $[FC_6H_4PMo_6^VO_{15}(OH)_3\{AsO_2(CH_3)_2\}_3]^{2-}$ $[F_3COC_6H_4PMo_6^VO_{15}(OH)_3\{AsO_2(CH_3)_2\}_3]^{2-}$ (FC₆H₄PMo₆), $(F_3COC_6H_4PMo_6)$ and $[Mo^{VI}Mo_6^VO_{15}(OH)_4\{AsO_2(CH_3)_2\}_3]^ (Mo_7)$, were synthesized in a one-pot aqueous reaction by stirring a solution containing ammonium heptamolybdate/hydrazine hydrochloride/heterogroup/sodium cacodylate buffer in the molar ratio of 1/2/1/20. The molecular structures of the polyanions $[RPMO_6^VO_{15}(OH)_3\{AsO_2(CH_3)_2\}_3]^{2-}$ (R = H, HO, CH₃, HO₂CCH₂, $HO_2CC_2H_4$, C_6H_5 , $4-FC_6H_4$, $4-F_3COC_6H_4$) and $[HOMo^{VI}Mo_6^VO_{15}]$ $(OH)_3[AsO_2(CH_3)_2]_3$ feature three edge-shared $\{Mo_2^VO_{10}\}$ fragments interconnected via edges by hydroxo bridges (confirmed by bond valence sum (BVS) calculations, Tables S2-S10†) forming a six-membered molybdenum(v)-oxo ring surrounding the RPO3 or MoO4 hetero group, and being capped by three peripheral dimethylarsinate ligands, creating a flowerpot-like structure (Fig. 1a and b). BVS calculations for the crystallographically independent molybdenum atoms resulted in values close to 5 (Tables S2-S10†), further confirming the Mo^V oxidation state. The reduced Mo₂^V pairs exhibit a severely distorted octahedral coordination geometry, and the coordination of each molybdenum center is mediated by two μ_2 -oxo, one μ_2 -hydroxo, one terminal oxo group, one oxo-donor from the terminal dimethylarsinate and one oxodonor from the central heterogroup.

As a result, the cyclic hexanuclear $\{Mo_6O_{24}\}$ core comprises exclusively edge-shared octahedra with alternating short Mo-Mo bonded contacts (*ca.* 2.6 Å) and longer Mo···Mo nonbonding contacts (*ca.* 3.6 Å).

In the solid state, the individual polyanions form Na^+ counter cation-mediated dimers via six $\mathrm{Na}\cdots\mathrm{O}$ interactions ($ca.\ 2.3\ \mathrm{\mathring{A}}$) with oxo groups from the bonded $\mathrm{Mo^V}_2$ pairs opposite to the side where the dimethylarsinates are bound (Fig. 1c). It appears that




Fig. 1 (a) Combined polyhedral/ball-and-stick representation of polyanion $RPMo_6$ [$RPMo_6^VO_{15}(OH)_3\{AsO_2(CH_3)_2\}_3]^2$ (R=H, HO, CH_3 , HO $_2CCH_2$, HO $_2CC_2H_4$, C_6H_5 , $4-FC_6H_4$, $4-F_3COC_6H_4$) and (b) the mixed-valent heptamolybdate [HOMO $^VIMo_6^VO_{15}(OH)_3\{AsO_2(CH_3)_2\}_3]^-$. (c) In the solid state, all nine novel polyanions form sodium ion-mediated dimers. Color code: MoO $_6$, purple octahedra; MoO $_4$, violet tetrahedron; As, green; C, yellow; P, navy blue; O, red; H, black; Na, light turquoise.

the formation of the Na $^+$ -mediated polyanion sandwich in the solid state is pivotal for the crystallization of the polyanions, as we could never isolate any polyanion without such motif. 9a,b

The novel polyanions can be obtained in the pH range 6.6–8.2 at room temperature as well as 50 or 80 °C, with optimal yields at pH 6.8–6.9. In all cases, the solutions turned dark red due to the reduction of the molybdenum centres from 6+ to 5+, with good quality crystals being formed within a week in decent yields. However, if the solutions are left standing open to the air during crystallization for a long time such as a month then $[Mo_{18}^{VI}Mo_{12}^{V}O_{84}\{AsO_2(CH_3)_2\}_{18}]^{18-}$ $\{Mo_{30}\}$ crystals started forming alongside the target compounds, probably due to a drop in pH and partial air-oxidation of the Mo^V centers (Fig. 2). This reemphasizes the fact that in POM chemistry pH plays a vital role in the reaction outcomes.

When using sodium molybdate (Na₂MoO₄·2H₂O) instead of ammonium heptamolydate ((NH₄)₆Mo₇O₂₄·4H₂O) as a reagent the target compounds could be obtained, but we observed extended crystallization periods and lower yields. Above pH 6.5, the same reaction mixture without addition of the phosphorous-containing heterogroups produced the heptanuclear $[Mo^{VI}Mo_6^VO_{15}(OH)_4\{AsO_2$ mixed-valent polyoxomolybdate $(CH_3)_2$ ₃]¹⁻ (Mo_7) with a central tetrahedral molybdate ion surrounded by a reduced MoV6-oxo ring. Varying the amount of reductant or the order of adding the reactants did not alter the outcome, suggesting that the suitable pH range and counter cations facilitated the crystal formation. However, the use of hydrazine hemisulphate (N₂H₄·0.5 H₂SO₄) as reducing agent for the Mo₇ reaction resulted in a disordered Mo atom in the center of the structure, likely due to its partial substitution by sulphate.




Fig. 2 Schematic representation of the pH importance in the synthesis of polyoxomolybdates in aqueous solution; (bottom and right) the novel polyanions described here. (Left) The Mo₃₀ wheel $[\text{Mo}_{18}^{\text{YM}} \text{Mo}_{12}^{\text{Y}} \text{O}_{84} \{\text{AsO}_2(\text{CH}_3)_2\}_{18}]^{18^-}$ and (top) the neutral $[\text{Mo}_4^{\text{Y}} \text{O}_8 \{\text{AsO}_2(\text{CH}_3)_2\}_4]$. Color code: $[\text{Mo}_{06}^{\text{Y}}, \text{purple octahedra; Mo}_{06}^{\text{Y}}, \text{oldet tetrahedron; P, navy blue; As, green; C, yellow; O, red; H, black.}$

The central hetero group and the three dimethylarsinate ligands are all grafted on the same side of the polyanions. Within each polyanion, the central hetero elements are slightly elevated above the plane of the six molybdenum atoms and are bound via three µ3-oxo groups resulting in a distorted tetrahedral coordination geometry. The central heterogroup can be systematically substituted by incorporation of phosphate (HOPO₃²⁻), phosphite (HPO₃²⁻), methyl phosphonate (CH₃PO₃²⁻), phosphonoacetate (HO₂CCH₂PO₃²⁻), phosphono- $(HO_2CC_2H_4PO_3^{2-}),$ propionate phenyl phosphonate $(C_6H_5PO_3^{2-})$, (4-fluorophenyl)phosphonate (4-FC $_6H_4PO_3^{2-})$ and (4-trifluoromethoxyphenyl)phosphonate $(4-F_3COC_6H_4PO_3^{2-}),$ wherein the overall charge of the polyanion mirrors with that of heterogroup.⁶ The mixed-valent heptamolybdate $[\text{Mo}^{\text{VI}}\text{Mo}_{6}^{\text{V}}\text{O}_{15}[\text{OH}]_{4}[\text{AsO}_{2}(\text{CH}_{3})_{2}]_{3}]^{1-} \quad (\textbf{Mo}_{7}) \quad \text{comprises} \quad \text{molyb-}$ denum atoms in both tetrahedral and octahedral coordination modes. The diphosphonate derivative [Mo₇O₁₆(O₃PCH₂PO₃)₃]⁸⁻ also contains one MoVI and six MoV centers, but the latter do not form an edge-shared six-membered ring, as the three Mo^V₂ units are bridged by a diphosphonate group.¹²

The eight novel phosphomolybdates(v) [RPMo $_{6}^{V}O_{15}(OH)_{3}$ {AsO $_{2}(CH_{3})_{2}$ } $_{3}$] $^{2-}$ (R = H, HO, CH $_{3}$, HO $_{2}$ CCH $_{2}$, HO $_{2}$ CC $_{2}$ H $_{4}$, C $_{6}$ H $_{5}$, 4-FC $_{6}$ H $_{4}$, 4-F $_{3}$ COC $_{6}$ H $_{4}$) and the mixed-valent heptamolybdate [HOMo $_{6}^{VI}$ Mo $_{6}^{VO}$ O1 $_{5}$ (OH) $_{3}$ {AsO $_{2}$ (CH $_{3}$) $_{2}$ } $_{3}$] $^{-}$ can be structurally described as follows: (a) terminal Mo $_{6}$ Oterm bonds of 1.675–1.708 Å; (b) Mo $_{6}$ O(As) bonds of 2.066–2.104 Å and Mo $_{6}$ O (Mo) bonds of 2.107–2.128 Å; and (c) Mo $_{6}$ O(P/Mo) bonds of 2.274–2.355 Å. The dimethylarsinate ((CH $_{3}$) $_{2}$ AsO $_{2}$) capping groups act as bidentate ligands and the As center is tetrahed-rally coordinated with average As $_{6}$ O(Mo) distances from

1.68–1.70 Å.^{4d} The terminal P–O bond (1.52 Å) in **PMo**₆ and the terminal Mo^{VI}–O bond (1.82 Å) of the central tetrahedral molybdenum atom in \mathbf{Mo}_7 are both monoprotonated corresponding to a hydroxo group, as confirmed by BVS calculations (Tables S2 and S10†).

The crystalline nature and phase purity of the synthesized polyanions were assessed through powder-XRD (PXRD) measurements on crystals directly taken from the mother liquor, see Fig. S18.† The diffractograms showed a good agreement between the experimental patterns and simulated patterns derived from the respective single-crystal X-ray diffraction analyses. Differences in diffraction peak intensities are attributed to various crystal orientations. The PXRD patterns of all compounds complement each other confirming their isostructural nature.

Upon closer examination of the diffraction patterns for the phosphite derivative Na-HPMo6 and the phosphate derivative NaNH₄-PMo₆ (Fig. S19†), an additional peak was observed at low 2θ angles 7.5–10°. Further studies revealed that polyanions with short groups on the central P hetero atom lose their crystal waters rapidly upon exposure to atmospheric air, causing the clear red crystals to transition into orange powder within one hour.14 Different degrees of hydration for Na-HPMo6 result in modifications of the packing arrangements in the crystalline lattice, thereby influencing the positions and intensities of the diffraction peaks observed in the air-dried sample. The rehydration of Na-HPMo6 in the presence of water vapor at room temperature overnight brings back the compound to the freshly synthesized states as shown by PXRD and TGA, indicating a reversible dehydrationrehydration process (Fig. 3). Conversely, compounds with longer groups attached to the hetero atom such as NaNH₄-HO₂CC₂H₄PMo₆ maintained their structural integrity and crystalline nature even after extended drying periods (Fig. S19†).

3.1 NMR spectroscopy

To investigate the solution properties of the P-containing polyanions $[RPMo_6^VO_{15}(OH)_3\{AsO_2(CH_3)_2\}_3]^{2-}$ (R = H, HO, CH₃, HO₂CCH₂, HO₂CC₂H₄, C₆H₅, 4-FC₆H₄, 4-F₃COC₆H₄), we conducted multinuclear NMR spectra (^{31}P , ^{14}H , ^{19}F , $^{13}C\{^{1}H\}$) on the

salts dissolved in water. The ¹³C{¹H} NMR spectra required overnight collection due to the low solubility of the polyanions. The ¹H solution NMR spectra in H₂O/D₂O of the reference compounds cacodylic acid (H-Cac) (pH 4.1) and sodium cacodylate buffer (Na-Cac) (H-Cac adjusted to pH 7.0 with NaOH pellets) exhibit a distinct peak at 1.8 and 1.7 ppm, respectively, alongside the solvent water peak at 4.7 ppm. Their corresponding ¹³C{¹H} NMR spectra show a sharp singlet at 17.0 and 17.5 ppm, respectively, consistent with the expected signal from the two equivalent methyl groups on the As^V atom. ⁷a

The ${}^{31}P\{^{1}H\}$ NMR spectrum of **NaNH₄-PMo₆** displays three singlets at 2.80, 2.65 and 2.50 ppm, with an intensity ratio of 20:5:1. Notably, the peak at 2.80 ppm could be attributed to **PMo₆**, whereas the signals at 2.65 and 2.50 ppm likely correspond to protonated phosphate species (Fig. 4a). The ${}^{1}H$ NMR spectrum of **NaNH₄-PMo₆** dissolved in H_2O/D_2O at room temp-





Fig. 4 Multinuclear NMR study ((a) $^{31}P\{^1H\}$, (b) 1H , (c) $^{13}C\{^1H\}$) of NaNH₄-PMo₆ (pink spectra) dissolved in H₂O/D₂O at room temperature (pH 7). The spectra of the reference compounds are shown in blue and black.






Fig. 3 (left) Powder XRD patterns for freshly prepared, air-dried and rehydrated Na-HPMo₆ and (right) thermograms for air dried and rehydrated Na-HPMo₆ from room temperature to 400 °C under N₂ atmosphere.

erature exhibits two distinct singlets at δ = 2.3 and 2.0 ppm (intensity ratio 1:1) (Fig. 4b), whereas the ¹³C{¹H} NMR spectrum shows two sharp resonances at 19.0 and 18.0 ppm, clearly reflecting the inequivalent nature of the two methyl groups on each As atom (Fig. 4c). The ¹H NMR peak at 1.7 ppm is likely from a minute amount of cocrystallized or decomposed free cacodylate, whereas the minor peaks at 2.2 and 1.9 ppm could not be assigned. The solutions were monitored by NMR over a period of 1 week, during which the spectra remained unchanged, indicating no significant decomposition of PMo₆ (Fig. S20†). This observation suggests that in **PMo₆** and the other derivatives, the cacodylate groups are strongly bound to the polyanion, which is supported by Mo-O(As) bond lengths of 2.076-2.099 Å. For comparison, the Mo-O(P) bonds in Haushalter's [(H₃O)₂NaMo₆P₄O₂₄(OH)₇]²⁻ $\{P_4Mo_6^V\}$ are 2.1073–2.1075 Å, ^{9b} and for both compounds are much shorter than the Mo-O(C) bonds (2.260-2.322 Å) in Kortz's amino acid-decorated [Se^{IV}Mo₆O₂₁(O₂CCH₂NH₃)₃]²⁻ (which is one of 18 derivatives with different amino acids bound, including chiral ones).6e

Considering that the novel polyanions are formed easily in a one-pot procedure we performed in situ 31P{1H} NMR studies, in order to optimize the synthetic procedures. For the synthesis of PMo₆, a reaction mixture containing ammonium heptamolybdate, sodium phosphate, and hydrazine hydrochloride in sodium cacodylate buffer (pH 7) with a molar ratio of Mo:P at 36:1 was stirred and monitored at two different temperatures. This six-fold molybdenum excess was initially used to ensure optimal product yield. During the synthesis, we observed that at room temperature PMo6 did not form during the first hour (Fig. S21a†), whereas at 80 °C a gradual conversion from Na₃PO₄ at pH 7 (δ = 2.5 ppm) to **PMo₆** (δ = 3.0 ppm) occurred over the course of one hour, ultimately resulting in a clean singlet at 3.0 ppm, indicative of complete conversion (Fig. S21b†). To further understand the effect of molar ratios, reaction progress was monitored at 80 °C across various Mo: P ratios. Notably, with a stoichiometric Mo: P ratio of 6:1, only 35% conversion to the target product was observed after one hour (Fig. S21†), underscoring the importance of molybdenum excess in achieving higher conversion rates.

Regarding the NMR characteristics of dissolved crystalline material for the other compounds, the phosphite-analogue **HPMo**₆ displayed the expected NMR resonances in H₂O/D₂O with prominent sharp singlets at δ = 2.4 and 1.9 ppm for ¹H NMR (Fig. S22a†) as well as $\delta = 19.4$ and 17.9 ppm in ${}^{13}\text{C}_{1}^{1}\text{H}$ NMR (Fig. S22b†) attributed to the cacodylate methyl groups. The ³¹P ¹H} spectrum showed a significant downfield shift at 12.9 ppm, compared to the reference shift at 2.90 ppm for Na₂HPO₃ (adjusted from pH 9.2 to pH 7 with 4M HCl). Additionally, the ³¹P NMR spectra displayed doublet peaks at 15.12 and 10.84 ppm $\binom{1}{J_{P-H}}$ = 693 Hz) due to the strong coupling of the phosphorus atom with the directly attached proton (Fig. S23†).^{6d}

The ¹H NMR spectrum of the methylphosphonate analogue CH₃PMo₆ exhibits two sharp equal-intensity resonances at 2.4 and 2.0 ppm (for the inequivalent methyl protons of the coordinated cacodylates), a peak at 1.8 ppm for free cacodylate

(possibly some minor polyanion decomposition due to heating at 40 °C to increase solubility in H2O) and a doublet centered at 0.015 ppm $(^2J_{P-H} = 18 \text{ Hz})$ due to coupling of the central methylphosphonate protons with the phosphorous nucleus, significantly upfield shifted from the doublet at 1.06 ppm $\binom{2}{I_{P-H}}$ = 16.4 Hz) for the reference CH₃PO₃H₂ dissolved in water (adjusted to pH 6 with 6M NaOH) (Fig. S24a†). The ¹³C{¹H} NMR of CH₃PMo₆ shows the anticipated two peaks at 19.6 and 17.5 ppm for the two inequivalent methyl groups of each bound cacodylate, along with a sharp peak at 16.9 ppm (corresponding to the free cacodylates) and a doublet at 13.85 ppm $\binom{1}{J_{P-C}}$ = 150.8 Hz) (originating from the methyl group on the P heteroatom), shifted slightly downfield from the doublet of the reference centered at 13.3 ppm $\binom{1}{I_{P-C}}$ = 140.7 Hz) (Fig. S24b†). The $^{31}P\{^{1}H\}$ NMR of CH_3PMo_6 shows a sharp singlet at 37.2 ppm, shifted downfield as compared to the 24.0 ppm signal for the reference methylphosphonate (Fig. S25a†), whereas the non-decoupled ³¹P NMR spectrum of CH_3PMo_6 depicts a clear quartet centered at $\delta = 37.2$ ppm $(^2J_{\rm P-H}$ = 18.5 Hz), and more upfield at δ = 24 ppm $(^2J_{\rm P-H}$ = 16.2 Hz) the quartet for the reference (Fig. S25b†). 6d

salts of the polyanions HO₂CCH₂PMo₆, HO₂CC₂H₄PMo₆ and C₆H₅PMo₆ exhibited low solubility in H₂O and so we used d₆-DMSO as solvent to acquire clear and reliable NMR spectra. In ³¹P{¹H} NMR the polyanion $HO_2CCH_2PMo_6$ displays a major peak $\delta = 20.6$ ppm, as compared to 15.9 ppm for the reference and a minor peak at 16.8 ppm, which could not be assigned (Fig. S26a†). 6a It was difficult to obtain a good 13C(1H) NMR spectrum due to the low solubility of the compound. On the other hand, the ³¹P {1H} spectrum of HO₂CC₂H₄PMo₆ in d₆-DMSO exhibits a sharp signal at 31.5 ppm, whereas the reference signal is at 25.5 ppm (Fig. S26b†). 6a The 13 C 1 H 1 NMR of $HO_{2}CC_{2}H_{4}PMo_{6}$ redissolved in d₆-DMSO (Fig. S27†) displays two signals at 18.2 and 20.1 ppm (for the inequivalent methyl carbons of coordinated cacodylates), along with signals at $\delta = 24.4$ (23.4), 27.9 (28.2), 174.4 (174.4) ppm from the 2-carboxyethyl phosphonate hetero group (values in parentheses are the chemical shifts for the corresponding C atoms of free 2-carboxyethyl phosphonic acid dissolved in d₆-DMSO). The peak at 24.4 ppm is a doublet ${}^{1}J_{P-C}$ = 150.8 Hz), arising from C-P coupling.

The ¹³C{¹H} NMR spectrum of C₆H₅PMo₆ in d₆-DMSO shows resonances at $\delta = 126.7$ (128.6), 129.2 (131.5), 131.4 (131.0) and 137.0 (134.5) ppm for the phenyl group (values in parentheses are for the corresponding C atoms of free phenylphosphonic acid dissolved in d₆-DMSO). Two further signals at 20.0 and 17.8 ppm were observed originating from the inequivalent methyl carbons of the cacodylate groups (Fig. 5). The $^{13}\text{C}\{^{1}\text{H}\}$ peaks in the aromatic region at 137.0 ppm $(^{1}J_{\text{P-C}} = 191$ Hz) and 126.7 ppm (${}^2J_{P-C}$ = 20.1 Hz) could be ascribed to the ipso and ortho carbon atoms of the phenyl ring, whereas the peak at 131.4 ppm ${}^{3}J_{P-C}$ = 10.1 Hz) is due to the *meta* carbon atom. The ${}^{31}P{}^{1}H{}$ spectrum of $C_6H_5PMo_6$ exhibited a major peak at 20.0 for the hetero element, as compared to a signal at 13.6 ppm for the neat phenylphosphonate. The minor peak at 19.6 ppm we cannot assign with confidence (Fig. S28a†).




Fig. 5 The 13 C 14 H) NMR spectra of NaNH $_4$ -C $_6$ H $_5$ PMo $_6$ (upper left in blue) and the reference C $_6$ H $_5$ PO $_3$ H $_2$ (lower left in black) dissolved separately in d $_6$ -DMSO at room temperature. The right panel showcases the enhanced spectra for the aromatic region only for NaNH $_4$ -C $_6$ H $_5$ PMo $_6$ (blue) and C $_6$ H $_5$ PO $_3$ H $_2$ (black). The peaks corresponding to structurally and hence magnetically inequivalent carbon atoms are labelled accordingly.

The $^{31}P\{^{1}H\}$ NMR spectra for the two fluorinated polyanions $\mathbf{4\text{-FC}}_{\mathbf{6}}\mathbf{H_{4}PMo_{6}}$ and $\mathbf{4\text{-F}}_{\mathbf{3}}\mathbf{COC_{6}}\mathbf{H_{4}PMo_{6}}$ were recorded in $\mathbf{H_{2}O/D_{2}O}$ at pH 6 and displayed singlets at 22.8 ppm (Fig. S28b†) and 22.0 ppm (Fig. S28c†), respectively, indicating structural integrity. The chemical shifts of the free hetero groups (4-fluorophenyl) phosphonic acid ($\mathbf{L_{PF}}$) and (4-trifluoromethoxyphenyl) phosphonic acid ($\mathbf{L_{POCF3}}$) in the same solvent at the same pH appear at 12.1 and 11.8 ppm, respectively (Fig. S28†). The 19 F NMR spectra for 4 -FC₆H₄PMo₆ and 4 -F₃COC₆H₄PMo₆ in 4 -QO/D₂O appear at $^{-1}$ 08.8 and $^{-5}$ 7.7 ppm, respectively (Fig. S29†).

Overall, it can be concluded that all our compounds which were easily water-soluble (HPMo6, 4-FC6H4PMo6 and 4-F₃COC₆H₄PMo₆) were shown to be solution-stable by multinuclear NMR. On the other hand, those compounds which needed heating and/or sonication to accomplish dissolution in water, showed some extra peaks due to possible minor transformation/decomposition of the polyanion (PMo6 and CH_3PMo_6). Those compounds (HO₂CCH₂PMo₆, HO₂CC₂H₄PMo₆ and C₆H₅PMo₆) which were less soluble in water could be dissolved in DMSO and exhibited clean NMR spectra. It is to be noted that, the mixed-valent heptamolybdate $[Mo^{VI}Mo_6^VO_{15}(OH)_4\{AsO_2(CH_3)_2\}_3]^{1-}$ (Mo_7) lacks sufficient solubilty in all tested solvents to obtain clear spectra required for detailed analysis.

3.2 ESI-mass spectrometry

ESI mass spectra of the polyanions were acquired from aqueous solution in both positive and negative ion mode. In positive ion mode only \mathbf{Mo}_7 resulted in a high-quality spectrum. In the negative ion mode all polyanions analyzed showed high-quality ESI-MS spectra with systematic patterns

(Fig. S30–S36†). All compounds showed two strong cluster of ions, a first between m/z 680–820 and a second between m/z 1320 and 1410 depending on the central moiety. For a detailed discussion we selected as two representative examples $\mathbf{C_6H_5PMo_6}$ and $\mathbf{Mo_7}$. All mass spectrometry data are summarized in Table S11.†




Fig. 6 Negative mode ESI mass spectra of polyanion $C_6H_5PMo_6$. (A) Full scan full range view with two main signal clusters; (B) expanded region around m/z 690 with experimental spectra in upper panel and simulated isotope pattern in bottom panel; (C) expanded region around m/z 1406 with experimental spectra in upper panel (please note signals of dimeric species at 0.5 Da mass difference at lower intensity in between main signals) and simulated isotope pattern in bottom panel.




Fig. 7 Negative mode ESI mass spectra of polyanion Mo_7 . (A) Full scan full range view with two main signal clusters; (B) expanded region around m/z 680 with experimental spectra in upper panel and simulated isotope pattern in bottom panel; (C) expanded region around m/z 1390 with experimental spectra in upper panel (please note: signals of dimeric species at lower intensity in between main signals) and simulated isotope pattern in bottom panel.

The polyanion $C_6H_5PMo_6$ showed a first cluster of signals at m/z 692.54 and a second cluster at m/z 1406.07 (see Fig. 6). The first cluster was assigned to a doubly-charged negative ion with an elemental composition of $[Mo_6PAs_3C_{12}H_{27}O_{24}]^{2-}$. The observed isotope pattern is in full agreement with the structure as demonstrated by comparison to a simulation of the spectrum (see Fig. 6).

The second signal cluster at m/z 1406.07 was assigned to originate from two individual species in the gas phase. Firstly, a monomeric, singly-charged species with an elemental composition of [Mo₆PAs₃C₁₂H₂₇O₂₄Na] with a mass difference of 1 Da between isotope peaks. Secondly, in between these signals a second species with isotope peak distances of 0.5 Da was observed, which was assigned to a dimeric species with an elemental composition of Na[Mo₆PAs₃C₁₂H₂₇O₂₄Na]₂²⁻. Again, simulated spectra are in full agreement with experimental spectra. As demonstrated from single-crystal XRD data, the polyanion C₆H₅PMo₆ dimerizes also in the gas phase mediated by a central Na⁺ ion, reminiscent of a 15-crown-5 crown ether interacting with a sodium cation. From the relative ion intensities of the two monomeric species and the single dimeric species a ratio of dimer/monomer of 0.57 in the gas phase was determined.




Fig. 8 Negative mode ESI ion mobility mass spectra of compound CH_3PMo_6 and corresponding mobilograms. (A) Expanded view of ESI mass spectrum with two overlapping species; (B) extracted ion mobilogram of m/z 1346.7 showing single dimeric species $[2M-2H]^{2-}$. (C) ESI mass spectrum corresponding to dimeric species $[2M-2H]^{2-}$. (D) Extracted ion mobilogram of m/z 1346.2 showing single monomeric species $[M-H]^{2-}$. (E) ESI mass spectrum corresponding to monomeric species $[M-H]^{2-}$.

The polyanion Mo_7 showed a first cluster of signals at m/z683.54 and a second cluster at m/z 1390.08 (see Fig. 7). The first cluster was assigned to a doubly-charged negative ion with an elemental composition of $[Mo_7As_3C_6H_{18}O_{24}]^{2-}$. The observed isotope pattern is in full agreement with the structure as demonstrated by comparison to a simulation of the spectrum (see Fig. 7). The second cluster at m/z 1390.08 was assigned to originate from two individual species in the gas phase. Firstly, a monomeric, singly-charged species with an elemental composition of [Mo₇As₃C₆H₂₁O₂₅] with a mass difference of 1 Da between isotope peaks. Secondly in between these signals a second species with isotope differences of 0.5 Da was observed, which was assigned to a dimeric species with an elemental composition of Na[Mo₇As₃C₆H₂₂O₂₅]₂²⁻. Again, simulated spectra are in full agreement with experimental spectra. In this case the ratio of dimer to monomer in the gas phase was determined to be 0.24.

Additionally, the polyanion \mathbf{Mo}_7 allowed for acquisition of a spectrum in the positive ion mode with two clusters observed at m/z 760.57 for a doubly positive ion and a second cluster centered around m/z 1498.17 corresponding to a singly positively charged ion (see Fig. S37†). It is worth mentioning that for the polyanion $\mathbf{HO}_2\mathbf{CC}_2\mathbf{H}_4\mathbf{PMO}_6$ exclusively the signals at higher m/z values at 1403.22 were observed with the monomeric dianion signals being absent (see Fig. S34†).

3.3 Ion mobility mass spectrometry

With a view to obtain further evidence for the assignment of the higher m/z signal envelope as overlapping monomeric and dimeric species we decided to attempt separation of the ions using ion mobility techniques. Ion mobility separation of POMs has a precedent in early work by Cronin's group on Keplerates. ^{15a} Further ion mobility separation of complex inorganic species have been achieved by Clever's group on Pd cages, ^{15b} and summarised in a recent review by Geue *et al.* ^{15c}

The two polyanions Mo₇ and CH₃PMo₆ were selected for investigations with the latter described in detail and the former shown in the ESI (see Fig. S38†). For CH₃PMo₆ quadrupole selection isolated the peaks of the isotope envelope centred at m/z 1346 with an isolation width of 35 Da (see Fig. 8). The multiple isotopes were subjected to ion mobility separation in a cyclic ion mobility mass spectrometer. Extracted ion mobilograms were obtained at m/z 1346.7 corresponding to the dimeric doubly-charged species and m/z 1346.2 corresponding to a mixture of monomeric and dimeric species. For the dimeric species a single species was observed in the mobilogram, whereas for the ions at m/z 1346.2 two species were detected at distinct drift times of 18.4 ms and 24.9 ms. ESI spectra were recorded separately for both species confirming the identity of a dimeric $[2M - 2H]^{2-}$ ion at 18.4 ms and a monomeric $[M - H]^-$ ion at 24.9 ms. We believe this to be the first example of ion mobility separation of isobaric oligomeric POMs in the gas phase. The term isobaric refers to the fact that half of the monoisotopic peaks of the dimeric

species are isobaric to the monoisotopic peaks of the monomeric species.

4. Conclusions

We have performed a comprehensive study on the synthesis and characterization of organically functionalized, water soluble and stable reduced polyoxomolybdates. As a result, we have isolated a family of eight dimethylarsinate-functionalized phosphomolybdates(v), $[RPMo_6^VO_{15}(OH)_3\{AsO_2(CH_3)_2\}_3]^{2-}$ (R = H, HO, CH₃, HO₂CCH₂, HO₂CC₂H₄, C₆H₅, 4-FC₆H₄, 4-F₃COC₆H₄) and the mixed-valent heptamolybdate $[HOMo^{VI}Mo_6^VO_{15}(OH)_3\{AsO_2(CH_3)_2\}_3]^-$. All nine polyanions were synthesized via facile one-pot open beaker conditions in aqueous solution, by careful adjustment of the reagent ratios, ionic strength, and most prominently pH. The incorporation of various organophosphonates including fluorinated derivatives as well as phosphate and phosphite within a stable hexanuclear molybdenum(v)-oxo ring provided a nice platform to investigate solubility and stability in aqueous medium at around pH 6-7. We performed extensive multinuclear (1H, 31P, 19F, 13C) NMR studies to investigate the solution properties of the novel polyanions. Due to the systematic modification of the organic function on the polyanions their interaction with biomolecules is of interest, which is currently in progress and will be reported separately. Furthermore, we are in the process of expanding the current family of compounds by making other derivatives. Interestingly, the polyanions crystallized as a sodium-ion mediated dimer, which appears to be in equilibrium with the free polyanion after dissolution of the solid salts in water. ESI-MS allowed to prove the coexistence of both monomer and dimer, and it was even possible to separate both entities and then obtain clean spectra for each by using ion mobility mass spectrometry, indicating the power of this technique when it comes to separating two components during the measurement.

Data availability

- The data supporting this article have been included as part of the ESI.†
- Electronic supplementary information (ESI) available: single-crystal XRD data, bond valence sum calculations, ESI-MS data and spectra, multinuclear NMR spectra, FT-IR spectra, thermograms, and powder XRD patterns.
- CCDC 2394921–2394926 and 2406188–2406190, respectively.
- Crystallographic data for nine compounds have been deposited at the CCDC under the accession numbers 2394921 (NaNH₄-PMo₆), 2394922 (Na-Mo₇), 2394923 (Na-HPMo₆), 2394924 (NaNH₄-CH₃PMo₆), 2394925 (NaNH₄-C₆H₅PMo₆), 2394926 (HO₂CC₂H₄PMo₆·dmso), 2406188 (NaNH₄-FC₆H₄PMo₆), 2406189 (NaNH₄-HO₂CCH₂PMo₆) and 2406190 (NaNH₄-F₃COC₆H₄PMo₆).

Author contributions

U. K. supervised the project and guided the research direction. V. S. synthesized the compounds, conducted material characterization, and drafted this paper. A. P. was responsible for synthesizing and characterizing the fluorinated organophosphonate groups and the corresponding polyanions. S. B. discovered the parent compound with initial studies performed by A. S. B. S. B. contributed to the single - XRD measurements, data analysis discussions. J. H. and D. S. performed the mass spectrometry experiments, with N. K. analysing the data and writing the mass spectrometry section. J. O., D. K., R. G., A. K. assisted with the reproduction and characterization of the compounds during their bachelor's thesis. All authors reviewed and contributed to the final version of the manuscript.

Conflicts of interest

There is no conflict of interest to declare.

Acknowledgements

U. K. thanks the German Research Council (DFG KO-2288/31-1) and Constructor University (formerly Jacobs University) for research support. The polyanion structures in Fig. 1, 2, 4 and 5 were generated with Diamond, version 3.2 (Crystal Impact GbR). We acknowledge ion mobility mass spectrometry measurements carried out by Dr Susan Slade and Dr Malcolm Anderson from Waters (Wilmslow, UK).

References

1 (a) M. T. Pope, Heteropoly and Isopoly Oxometalates, Springer, Berlin, 1983; (b) M. T. Pope and A. Müller, Polyoxometalate Chemistry: An Old Field with New Dimensions in Several Disciplines, Angew. Chem., Int. Ed. Engl., 1991, 30, 34-48; (c) A. Proust, R. Thouvenot and P. Gouzerh, Functionalization of polyoxometalates: towards advanced applications in catalysis and materials science, Chem. Commun., 2008, 1837–1852; (d) A. Dolbecq, E. Dumas, C. R. Mayer and P. Mialane, Hybrid Organic-Inorganic Polyoxometalate Compounds: From Structural Diversity to Applications, Chem. Rev., 2010, 110, 6009-6048; (e) S.-S. Wang and G.-Y. Yang, Recent Advances in Polyoxometalate-Catalyzed Reactions, Chem. Rev., 2015, 115, 4893-4962; (f) N. Lotfian, M. M. Heravi, M. Mirzaei and B. Heidari, Applications of inorganic-organic hybrid architectures based on polyoxometalates in catalyzed and photocatalyzed chemical transformations, Appl. Organomet. Chem., 2019, 33, e4808; (g) A. Proust, B. Matt, R. Villanneau, G. Guillemot, P. Gouzerh and G. Izzet, Functionalization and post-functionalization: a step towards polyoxometalate-based materials, Chem. Soc. Rev.,

- 2012, 41, 7605-7622; (h) M.-P. Santoni, G. S. Hanan and B. Hasenknopf, Covalent multi-component systems of polyoxometalates and metal complexes: Toward multi-functional organic-inorganic hybrids in molecular and material sciences, Coord. Chem. Rev., 2014, 281, 64-85; (i) J. J. Walsh, A. M. Bond, R. J. Forster and T. E. Keves, Hybrid polyoxometalate materials for photo(electro-) chemical applications, Coord. Chem. Rev., 2016, 306, 217-234; (j) K. Xia, K. Yamaguchi and K. Suzuki, Recent advances in hybrid materials of metal nanoparticles and polyoxometalates, Angew. Chem., Int. Ed., 2023, 62, e202214506; (k) J. T. Rhule, C. L. Hill, D. A. Judd and R. F. Schinazi, Polyoxometalates in medicine, Chem. Rev., 1998, **98**, 327–358; (l) M. B. Čolović, M. Lacković, J. Lalatović, A. S. Mougharbel, U. Kortz and D. Z. Krstić, Polyoxometalates in biomedicine: Update and overview, Curr. Med. Chem., 2020, 27, 362–379; (m) H. Soria-Carrera, E. Atrián-Blasco, R. Martín-Rapún and S. G. Mitchell, Polyoxometalate-peptide hybrid materials: from structureproperty relationships to applications, Chem. Sci., 2023, 14, 10-28.
- 2 (a) A. V. Anyushin, A. Kondinski and T. N. Parac-Vogt, Hybrid polyoxometalates as post-functionalization platforms: from fundamentals to emerging applications, *Chem. Soc. Rev.*, 2020, 49, 382–432; (b) J. M. Cameron, G. Guillemot, T. Galambos, S. S. Amin, E. Hampson, K. M. Haidaraly, G. N. Newton and G. Izzet, Supramolecular assemblies of organo-functionalised hybrid polyoxometalates: from functional building blocks to hierarchical nanomaterials, *Chem. Soc. Rev.*, 2022, 51, 293–328.
- 3 (a) C. Rinfray, S. Renaudineau, G. Izzet and A. Proust, A covalent polyoxomolybdate-based hybrid with remarkable electron reservoir properties, Chem. Commun., 2014, 50, 8575-8577; (b) C. Li, N. Mizuno, K. Yamaguchi and K. Suzuki, Self-assembly of anionic polyoxometalateorganic architectures based on lacunary phosphomolybdates and pyridyl ligands, J. Am. Chem. Soc., 2019, 141, 7687-7692; (c) C. Li, K. Yamaguchi and K. Suzuki, Ligand-Directed Polyoxometalate Approach in Formation of a New Divacant Lacunary Polyoxomolybdate $[\gamma - PMo_{10}O_{36}]^{7-}$, Angew. Chem., Int. Ed., 2021, **60**, 6960– 6964; (d) A. Jimbo, C. Li, K. Yonesato, T. Ushiyama, K. Yamaguchi and K. Suzuki, Molecular hybrids of trivacant lacunary polyoxomolybdate and multidentate organic ligands, Chem. Sci., 2023, 14, 10280-10284.
- 4 (a) A. Rosenheim and R. Bilecki, Über Molybdänsäure–alkylarsinate. (Zur Kenntnis der Iso– Und Heteropolysalze. VIII. Mitteilung.), *Ber. Dtsch. Chem. Ges.*, 1913, 46, 539–557; (b) W. Kwak, L. M. Rajković, J. K. Stalick, M. T. Pope and C. O. Quicksall, Synthesis and Structure of Hexamolybdobis(organoarsonates), *Inorg. Chem.*, 1976, 15, 2778–2783; (c) K. M. Barkigia, L. M. Rajković, M. T. Pope and C. O. Quicksall, New type of heteropoly anion. Tetramolybdo complexes of dialkyl- and diarylarsinates, *J. Am. Chem. Soc.*, 1975, 97, 4146–4147; (d) K. M. Barkigia,

- L. M. Rajković-Blazer, M. T. Pope, E. Prince and C. O. Quicksall, Molybdoarsinate Heteropoly Complexes. Structure of the Hydrogen Tetramolybdodimethylarsinate (2-) Anion by X-Ray and Neutron Diffraction, *Inorg. Chem.*, 1980, 19, 2531–2537.
- 5 (a) W. Kwak, M. T. Pope and T. F. Scully, Stable Organic Derivatives of Heteropoly Anions, Pentamolybdobisphosphonates, J. Am. Chem. Soc., 1975, 97, 5735-5738; (b) J. K. Stalick and C. O. Quicksall, Two Heteropoly Anions Containing Organic Groups. Crystal and Molecular Structures of Ammonium Pentamolybdobis(methylphosphonate) Pentahydrate, $(NH_4)_4[(CH_{3P})_2Mo_5O_{21}]\cdot 5H_2O$, and Sodium Tetramethylammonium Pentamolybdobis(ethylammoniumphosphonate) Pentahydrate, $Na[N(CH_3)_4]$ $[(NH_3C_2H_4P)_2Mo_5O_{21}] \cdot 5H_2O$, Inorg. Chem., 1976, 15, 1577-1584; (c) T. Ozeki, H. Ichida, H. Miyamae and Y. Sasaki, Crystal Structure of K₄[H₂P₂Mo₅O₂₁]·2H₂O, Bull. Chem. Soc. Jpn., 1988, **61**, 4455–4457; (d) D. G. Lyxell and R. Strandberg, The structure of tetraguanidinium pentamolybdodiphenylphosphonate, Acta Crystallogr., Sect. C: Cryst. Struct. Commun., 1988, 44, 1535-1538.
- 6 (a) U. Kortz, C. Marquer, R. Thouvenot and M. Nierlich, Polyoxomolybdates Functionalized with Phosphonocarboxylates, Inorg. Chem., 2003, 42, 1158-1162; (b) P. Manna, S. Bhattacharya and U. Kortz, Arylarsonate- and Phosphonate-Capped Polyoxomolybdates, [(RC₆H₄As)₂Mo₆O₂₄]ⁿ⁻ and [(R' $C_6H_4P_{2}Mo_5O_{21}^{n-}$, Inorg. Chem., 2021, **60**, 7161–7167; (c) M. Carraro, A. Sartorel, G. Scorrano, C. Maccato, M. H. Dickman, U. Kortz and M. Bonchio, Chiral Strandberg-Type Molybdates [(RPO₃)₂Mo₅O₁₅]²⁻ as Molecular Gelators: Self-Assembled Fibrillar Nanostructures with Enhanced Optical Activity, Angew. Chem., Int. Ed., 2008, 47, 7275-7279; (d) U. Kortz, J. Vaissermann, R. Thouvenot and P. Gouzerh, Heteropolymolybdates of Phosphate, Phosphonate, and Phosphite Functionalized by Glycine, Inorg. Chem., 2003, 42, 1135-1139; (e) U. Kortz, M. G. Savelieff, F. Y. A. Ghali, Maalouf and D. I. Sinno, M. Khalil, S. A. Heteropolymolybdates of As^{III} , Sb^{III} , Bi^{III} , Se^{IV} , and Te^{IV} Functionalized by Amino Acids, Angew. Chem., Int. Ed., 2002, 41, 4070-4073.
- 7 (a) S. Bhattacharya, X. Ma, A. S. Mougharbel, M. Haouas, P. Su, M. F. Espenship, D. H. Taffa, H. Jaensch, A.-J. Bons, T. Stuerzer, M. Wark, J. Laskin, E. Cadot and U. Kortz, Discovery of a Neutral 40-PdII-Oxo Molecular Disk, $[Pd_{40}O_{24}(OH)_{16}\{(CH_3)_2AsO_2\}_{16}]$: Synthesis, Structural Characterization, and Catalytic Studies, Inorg. Chem., 2021, **60**, 17339–17347; (b) S. Bhattacharya, A. Barba-Bon, T. A. Zewdie, A. B. Müller, T. Nisar, A. Chmielnicka, I. A. Rutkowska, C. J. Schürmann, V. Wagner, N. Kuhnert, P. J. Kulesza, W. M. Nau and U. Kortz, Discrete, Cationic Palladium(II)-Oxo Clusters via f-Metal Ion Incorporation and their Macrocyclic Host-Guest Interactions with Sulfonatocalixarenes, Angew. Chem., Int. Ed., 2022, 61, e202203114; (c) X. Ma, S. Bhattacharya, T. Nisar, A. B. Müller, V. Wagner, N. Kuhnert and U. Kortz, Mixed-valent palladium $[Pd^{IV}O_6Pd^{II}_6((CH_3)_2AsO_2)_6]^{2-}$ (IV/II)-oxoanion,

- Commun., 2023, **59**, 904–907; (*d*) J. Zhang, S. Bhattacharya, A. B. Müller, L. Kiss, C. Silvestru, N. Kuhnert and U. Kortz, Mixed Noble Metal–Oxo Clusters: Platinum(IV)–Gold(III) Oxoanion [Pt^{IV}₂Au^{III}₃O₆((CH₃)₂AsO₂)₆]⁻, Chem. Commun., 2023, **59**, 5918–5921; (*e*) J. Zhang, S. Bhattacharya, B. E. Khsara, T. Nisar, A. B. Müller, M. Besora, J. M. Poblet, V. Wagner, N. Kuhnert and U. Kortz, Pt^{IV}-Containing Hexaplatinate(II) [Pt^{IV}Pt^{II}₆O₆(AsO₂(CH₃)₂)₆]²⁻ and Hexapalladate(II) [Pt^{IV}Pd^{II}₆O₆(AsO₂(CH₃)₂)₆]²⁻, Inorg. Chem., 2023, **62**, 13184–13194.
- 8 A. Sundar, S. Bhattacharya, J. Oberstein, X. Ma, B. S. Bassil, T. Nisar, D. H. Taffa, M. Wark, V. Wagner and U. Kortz, Organically Functionalized Mixed-Valent Polyoxo-30-molybdate Wheel and a Neutral Tetramolybdenum(V) Oxo Cluster, *Inorg. Chem.*, 2022, **61**, 11524–11528.
- 9 (a) R. C. Haushalter and F. W. Lai, $[Et_4N]_6[Na_{14}Mo_{24}P_{17}O_{97}(OH)_{31}]\cdot xH_2O:$ A Hollow Cluster Filled with 12 Na[⊕] Ions and a H₃PO₄ Molecule, Angew. Int. Engl., 1989, Chem., Ed.28, (b) R. C. Haushalter and F. W. Lai, Synthesis of a New One-Dimensional Sodium Molybdenum Phosphate Polymer: Structure of $[(H_3O)_2NaMo_6P_4O_{24}(OH)_7]^{2-}$, Inorg. Chem., 1989, 28, 2904–2905; (c) R. C. Haushalter and L. A. Mundi, Reduced Molybdenum Phosphates: Octahedral-Tetrahedral Framework Solids with Tunnels, Cages, and Micropores, Chem. Mater., 1992, 4, 31-48.
- (a) G. Cao, R. C. Haushalter and K. G. Strohmaier, A Novel Polyoxo Molybdenum(V) Organophosphonate Anion Having a Sandwich Structure: Synthesis and Crystal Structure of $[N(C_2H_5)_4]_2Na_3(H_3O)_4\{Na[Mo_6O_{15}(O_3PC_6H_5)]$ $(HO_3PC_6H_5)_3]_2$ }·14H₂O, Inorg. Chem., 1993, 32, 127–128; (b) M. I. Khan, Q. Chen and J. Zubieta, Hydrothermal synthesis and crystal structure of (NH₄)₅Na₄{Na[Mo₆O₁₅- $(HO_3PC_6H_5)_3(O_3PC_6H_5)]_2$ $(6H_2O_1, Inorg. Chim. Acta, 1993_1)$ 206, 131-133; (c) M. I. Khan, Q. Chen and J. Zubieta, Oxomolybdenum(V) polyanion clusters. Hydrothermal syntheses structures of (NH₄)₅Na₄{Na $[Mo_6O_{12}(OH)_3(O_3PC_6H_5)_4)]_2$ \:6H_2O and $(C_6H_5CH_2NMe_3)_4$ $K_4\{K_2[Mo_6O_{12}(OH)_3(O_3PC_6H_5)_4]_2\}\cdot 10H_2O$ and their relationship to the binuclear (Et₄N)[Mo₂O₄Cl₃(H₂O)₃]·5H₂O, Inorg. Chim. Acta, 1995, 235, 135-145; (d) Y. D. Chang and J. Zubieta, Investigations into the syntheses and structures of clusters of the Mo-O-REO₃²⁻ systems (E = P and As), *Inorg.* Chim. Acta, 1996, 245, 177-198; (e) M. J. Manos, A. D. Keramidas, J. D. Woollins, A. M. Z. Slawin and T. A. Kabanos, The first polyoxomolybdenum carbonate compound: Synthesis and crystal structure of $(NH_4)_5[(Mo_2^VO_4)_3(\mu_6-\mu_6)]$ CO_3 (μ - CO_3)₃(μ -OH)₃]·0.5CH₃OH, J. Chem. Soc., Dalton Trans., 2001, 3419-3420; (f) M. J. Manos, J. D. Woollins, A. D. Keramidas, A. M. Z. Slawin and T. A. Kabanos, Polyoxomolybdenum(v) Sulfite Complexes: Structural, and Physical Studies, Angew. Chem., Int. Ed., 2002, 41, 2801-2805.
- 11 (a) E. Cadot, A. Dolbecq, B. Salignac and F. Sécheresse, Self-Condensation of [Mo^V₂O₂S₂]²⁺ with Phosphate or Arsenate Ions by Acid-Base Processes in Aqueous Solution:

Syntheses, Crystal Structures, and Reactivity $[(HXO_4)_4Mo_6S_6O_6(OH)_3]^{5-}$, X= P, As, Chem. - Eur. J., 1999, 5, 2396-2403; (b) A. Dolbecq, E. Cadot, D. Eisner and F. Sécheresse, Regioselective S/O substitutions in heteropolyoxothioanions: 31P NMR study and X-ray crystal structure of the half-substituted anion [(HPO₄)₄Mo₆S₃O₉(OH)₃]⁵⁻, Inorg. Chim. Acta, 2000, 300-302, 151-157.

Research Article

- 12 (a) E. Dumas, C. Sassove, K. D. Smith and S. C. Sevov, Synthesis and Characterization $[Mo_7O_{16}(O_3PCH_2PO_3)_3]^{8-}$: Mixed-Valent Polyoxomolybdenum Diphosphonate Anion with Octahedrally and Tetrahedrally Coordinated Molybdenum, Inorg. Chem., 2002, 41, 4029-4032; (b) C. du Peloux, A. Dolbecq, P. Mialane, J. Marrot and F. Sécheresse, Template synthesis of $\{(Mo_2^VO_4)(O_3PCH_2PO_3)\}_n$ clusters (n = 3, 4, 10): solid state and solution studies, Dalton Trans., 2004, 1259-1263.
- 13 G. Keglevich, A. Grün, A. Bölcskei, L. Drahos, M. Kraszni and G. T. Balogh, Synthesis and Proton Dissociation Properties of Arylphosphonates: A Microwave-Assisted Catalytic Arbuzov Reaction with Aryl Bromides, Heteroat. Chem., 2012, 23, 574-582.

- 14 (a) D. Barats-Damatov, L. J. W. Shimon, Y. Feldman, T. Bendikov and R. Neumann, Solid-State Crystal-to-Crystal Phase Transitions and Reversible Structure-Temperature Behavior of Phosphovanadomolybdic Acid, H₅PV₂Mo₁₀O₄₀, Inorg. Chem., 2015, 54, 628-634; (b) B. Herzog, W. Bensch, T. Ilkenhans, R. Schlögl and N. Deutsch, Single crystal and powder diffraction studies of the structure of heteropolymolybdophosphoric acid catalysts, Catal. Lett., 1993, 20, 203-219.
- 15 (a) P. J. Robbins, A. J. Surman, J. Thiel, D.-L. Long and L. Cronin, Use of ion-mobility mass spectrometry (IMS-MS) to map polyoxometalate Keplerate clusters and their supramolecular assemblies, Chem. Commun., 2013, 49, 1909-1911; (b) K. E. Ebbert, L. Schneider, A. Platzek, C. Drechsler, B. Chen, R. Rudolf and G. H. Clever, Resolution of minor size differences in a family of heteroleptic coordination cages by trapped ion mobility ESI-MS, Dalton Trans., 2019, 48, 11070-11075; (c) N. Geue, R. E. P. Winpenny and P. E. Barran, Ion Mobility Mass Spectrometry for Large Synthetic Molecules: Expanding the Analytical Toolbox, J. Am. Chem. Soc., 2024, 146, 8800-8819.