

RESEARCH ARTICLE

View Article Online
View Journal | View IssueCite this: *Org. Chem. Front.*, 2025, 12, 2409Received 17th January 2025,
Accepted 11th February 2025

DOI: 10.1039/d5qo00100e

rsc.li/frontiers-organic

Recyclable g-C₃N₄ catalyzed decarboxylative alkenylation of *N*-aryl glycines with vinyl sulfones under visible-light irradiation†Chengjie Guo,^a Guozhi Zhao,^a Yabiao Feng,^a Dong Chen,^b Teck-Peng Loh,^b Dongping Wang^{*a} and Zhenhua Jia^{*a,c}

Allylamines are a versatile class of compounds with significant applications in pharmaceuticals and as building blocks in organic synthesis. Herein we present a straightforward protocol for visible-light driven g-C₃N₄-catalyzed decarboxylative alkenylation of *N*-aryl glycines with vinyl sulfones to access allylamines in moderate to excellent yields (up to 91%), and it demonstrated broad substrate compatibility, including primary, secondary, and tertiary *N*-aryl glycines and diverse vinyl sulfones. Notably, the g-C₃N₄ catalyst is recyclable up to five times without obvious loss of catalytic performance. Preliminary mechanistic studies indicated that visible light is essential to achieve the desired transformation efficiently.

Introduction


Allylamines are a class of compounds with diverse pharmacological effects, abundant in the pharmaceutical field as anti-fungals, antihistamines, antidepressants and more (Scheme 1A).¹ Over the past few decades, numerous methods for synthesizing allylamine derivatives have been developed.² However, most of these strategies rely on the use of expensive or toxic transition metal catalysts, high temperatures, and harsh reaction conditions, resulting in limited substrate compatibility. Recently, visible-light promoted photoredox catalysis has been widely used in organic synthesis, leveraging visible light as an energy source to drive chemical reactions in a clean, efficient, and sustainable manner.³ Homogeneous photoredox catalysts, such as transition metal complexes and organic dyes, have been widely utilized in organic synthesis.⁴ However, resource scarcity, high price and metal contamination in products limited their potential practical application. To address these challenges, the development of practical heterogeneous catalysts has become a promising alternative.

g-C₃N₄, also known as graphitic carbon nitride, is a semiconductor material composed of carbon and nitrogen elements and exhibits excellent photocatalytic and photothermal properties, making it applicable in various fields.⁵ Moreover, owing to its wide availability and remarkable stability, g-C₃N₄ has also been employed as a photoredox catalyst in organic synthesis.⁶

N-Aryl glycines, a class of natural amino acid derivatives, are widely utilized as pharmaceutical intermediates and a key raw material for generating α -aminoalkyl radicals *via* de-

^aHenan Linker Technology Key Laboratory, College of Advanced Interdisciplinary Science and Technology (CAIST), Henan University of Technology, Zhengzhou 450001, China.

E-mail: teckpeng@ntu.edu.sg, caist_wdp@haut.edu.cn, caist_zhjia@haut.edu.cn

^bCollege of Material Engineering, Henan International Joint Laboratory of Rare Earth Composite Materials, Henan University of Engineering, Zhengzhou 450001, Henan, China^cDivision of Chemistry and Biological Chemistry, School of Physical and Mathematical Sciences, Nanyang Technological University, Singapore 637371, Singapore†Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d5qo00100e>

Scheme 1 (A) Allylamines in pharmaceutical molecules. (B) *N*-Aryl glycines as radical precursors for decarboxylative functionalization. (C) This work: g-C₃N₄ as the recyclable photoredox catalyst for decarboxylative alkenylation.




carboxylation, enabling the efficient synthesis of high value-added products (Scheme 1B).^{7–17} Recently, silver-catalyzed decarboxylation of carboxylic acids has also emerged as a promising strategy for the synthesis of diverse functional molecules.¹⁸ Since Rueping *et al.* pioneered the first visible light-induced g-C₃N₄-catalyzed decarboxylative alkylation of *N*-aryl glycines in 2018, the recent advances in this field have opened novel avenues for organic transformations and the synthesis of complex molecules.^{19,20} Nevertheless, to the best of our knowledge, visible-light-induced g-C₃N₄-catalyzed decarboxylative alkenylation of *N*-aryl glycines has not yet been reported. In this study, we have reported g-C₃N₄ as a recyclable photoredox catalyst which induced the generation of α -aminoalkyl radicals from *N*-aryl glycines to undergo radical addition with vinyl sulfones for the efficient synthesis of allylamines under visible-light irradiation (Scheme 1C).

Results and discussion

Initially, we employed phenyl vinyl sulfones (**1a**) and *N*-phenyl glycine (**2a**) as substrates to optimize the reaction conditions (Table 1). When 10 mg g-C₃N₄ was used as a photoredox catalyst, NaHCO₃ (2.0 equiv.) as the base in DCM (2 mL), under 15 W 460 nm LED irradiation in air, the desired alkylation product **3a** was detected in a trace amount (entry 1). Then,

Table 1 Optimization of the reaction conditions^a




Entry	2a (equiv.)	Base	Solvent (mL)	T (°C)	Yield ^b (%)
1	1.5	NaHCO ₃	DCM (2)	25	Trace
2	1.5	CH ₃ COOK	DCM (2)	25	Trace
3	1.5	AgF	DCM (2)	25	0
4	1.5	NH ₄ F	DCM (2)	25	34
5	1.5	InF ₃	DCM (2)	25	Trace
6	1.5	KF	DCM (2)	25	36
7	1.5	NaF	DCM (2)	25	Trace
8	1.5	CsF	DCM (2)	25	53
9	1.5	CsF	DCE (2)	25	41
10	1.5	CsF	MeOH (2)	25	47
11	1.5	CsF	Toluene (2)	25	50
12	1.5	CsF	EA (2)	25	44
13	1.5	CsF	1,4-Dioxane (2)	25	43
14	1.5	CsF	MeCN (2)	25	47
15	1.5	CsF	THF (2)	25	48
16 ^c	1.5	CsF	DCM (2)	25	37
17	2.0	CsF	DCM (2)	25	60
18	2.0	CsF	DCM (1)	25	51
19	2.0	CsF	DCM (3)	25	72
20	2.0	CsF	DCM (4)	25	57
21	2.0	CsF	DCM (3)	35	85
22 ^d	2.0	CsF	DCM (3)	35	Trace
23 ^e	2.0	CsF	DCM (3)	35	0
24	2.0	—	DCM (3)	35	28

^a Reaction conditions: **1a** (0.2 mmol, 1.0 equiv.), **2a**, g-C₃N₄ (10 mg), base (2.0 equiv.), solvents, under 15 W 460 nm LED irradiation in air for 12 h. ^b Isolated yield. ^c g-C₃N₄ (15 mg). ^d In the absence of g-C₃N₄. ^e In the dark.

various bases were screened and it was found that CsF was the most effective, enhancing the yield to 53% (entries 2–8). Further screening of solvents indicated that DCM was the optimal medium (entries 9–15). When the amount of g-C₃N₄ was adjusted to 15 mg, the efficiency of the reaction was not improved obviously (entry 16). However, **3a** was obtained in 60% yield with the use of 2 equivalents of **2a** (entry 17). Screening of the reaction concentrations demonstrated that 3 mL of DCM resulted in 72% yield (entries 18–20). Finally, the product **3a** was obtained in 85% yield by increasing the reaction temperature to 35 °C (entry 21). Controlled experiments confirmed that visible light, g-C₃N₄ and base were all essential for achieving the desired transformation (entries 22–24).

Under the optimal conditions, we next explored the substrate scope of vinyl sulfones as shown in Scheme 2. Generally, aryl vinyl sulfones bearing diverse electron-donating and electron-withdrawing groups, such as -F, -Cl, -Br, -CN, -CF₃, -Me, -^tBu, -OPh, and -OAc on benzene rings at diverse positions, underwent the reaction smoothly, yielding the corresponding products in 70–86% yields (**3b–3n**). Multi-substituted aromatic vinyl sulfones were also compatible, delivering the products **3o–3q** in moderate to excellent yields. Moreover, naphthyl vinyl sulfone was also tolerated to afford **3r** in 78% yield. Furthermore, ester-substituted vinyl sulfone was also employed to achieve the target product **3u** in 34% yield. However, the unexpected dimer product **3v** was formed when 4-iodophenyl



Scheme 2 Substrate scope of vinyl sulfones. Reaction conditions: **1** (0.2 mmol, 1.0 equiv.), **2a** (2.0 equiv.), g-C₃N₄ (10 mg), CsF (2.0 equiv.), DCM (3 mL) at 35 °C under 15 W 460 nm LED irradiation in air for 12 h.




vinyl sulfone was used as the starting material under standard conditions.


Subsequently, a range of diverse *N*-aryl glycines were explored as shown in Scheme 3. Notably, we varied the steric hindrance or electronic properties of the aryl ring to evaluate the efficiencies, producing the desired products in 21–84% yields (**4a–4l**). Substituents such as –F (**4a**), –Cl (**4b**, **4k**), –Br (**4c**), –I (**4d**), –CN (**4e**) and –COMe (**4h**) at diverse positions of the benzene ring were compatible, allowing the synthesis. *N*-Aryl glycines with di- or tri-substituents were also accommodated under the optimal conditions to generate **4m–4p** in 38–65% yields. In addition, α -methyl or dimethyl substituted *N*-aryl glycines were effective acceptors, affording the corresponding products in 90% and 47% yields, respectively (**4q**, **4r**). Moreover, *N*-methyl-*N*-phenyl glycine was also suitable for the transformation, providing the product **4s** in 47% yield. Unfortunately, when *N*-phenyl- α -phenyl glycine was employed as the substrate, only a trace amount of the product was detected (**4t**).

To showcase the practical application, we scaled up the model reaction to 5 mmol, and the product **3a** was isolated in 61% yield (Scheme 4A). Furthermore, **3a** was efficiently transformed into the diene product **5** in 82% yield, underscoring its potential for practical synthetic applications (Scheme 4B). To evaluate the stability and recyclability of $g\text{-C}_3\text{N}_4$, a series of recycling experiments were conducted to assess its performance in multiple reaction cycles. As shown in Scheme 4C, $g\text{-C}_3\text{N}_4$ was successfully recycled and reused in five consecutive reaction cycles without any obvious loss in catalytic performance, highlighting the potential for a sustainable process.

Next, we performed radical trapping and light on/off experiments to elucidate the possible mechanism. As shown in




Scheme 4 (A) Scale up model reaction. (B) Further modification of **3a**. (C) Catalyst recycle experiments of the model reaction.



Scheme 5 (A) Radical trapping experiment. (B) Light on/off experiments. (C) Proposed mechanism.

Scheme 5, the addition of a radical scavenger, such as 2,6-*tert*-butyl-4-methylphenol (BHT), to the standard reaction conditions completely inhibited the formation of product **3a**. Moreover, the aminomethyl-BHT adduct was detected by



Scheme 3 Substrate scope of *N*-aryl glycines. Reaction conditions: **1a** (0.2 mmol, 1.0 equiv.), **2** (2.0 equiv.), $g\text{-C}_3\text{N}_4$ (10 mg), CsF (2.0 equiv.), DCM (3 mL) at 35 °C under 15 W 460 nm LED irradiation in air for 12 h.



high-resolution mass spectrometry, indicating the probable involvement of a radical species (Scheme 5A). In addition, the results of light on/off experiments suggested that a radical chain pathway was excluded, and product formation was sustained under light irradiation but completely ceased in the absence of light (Scheme 5B).

Based on the experimental results and precedent literature,^{18–21} the proposed reaction mechanism is illustrated in Scheme 5C. Initially, under visible-light irradiation, *g*-C₃N₄ absorbed photons to generate holes in the valence band (VB) and electrons in the conduction band (CB). Then, oxygen in air was reduced by electrons in the CB to produce the superoxide radical anion. Simultaneously, the deprotonation of **2a** proceeded in the presence of base to generate carboxylate anion **I**, which was oxidized by holes in the VB to form carboxyl radical **II**. Following the release of CO₂ from **II**, α -aminoalkyl radical **III** was generated, which underwent radical addition with **1a** to form the intermediate **IV**. The intermediate **IV** then underwent β -elimination, leading to the desired alkenylation product **3a** and phenylsulfone radical **V**. Finally, the phenylsulfone radical was reduced to the corresponding phenylsulfone anion **VI** by the superoxide radical anion.

Conclusions

In summary, we have developed an efficient protocol for the decarboxylative alkenylation of *N*-aryl glycines with vinyl sulfones using recyclable *g*-C₃N₄ as a heterogeneous photoredox catalyst. This strategy accommodated a wide variety of *N*-aryl glycines, demonstrating excellent functional group tolerance and facilitating the rapid synthesis of allylamines. Additionally, this method was scalable to the gram-scale and also provided direct access to valuable diene compounds through straightforward product derivatization, underscoring its practicality and potential application in organic synthesis.

Data availability

The data supporting this article have been included as part of the ESI.†

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgements

We gratefully acknowledge the grants from the National Natural Science Foundation of China (22401081), the Natural Science Foundation of Henan Province (242300420546, 242300421349) and the Start-up Grant of Henan University of Technology (2023BS007, 2024PYJH064). We gratefully acknowl-

edge the financial support from the Distinguished University Professor grant (Nanyang Technological University) and the Agency for Science, Technology, and Research (A*STAR) under its MTC Individual Research Grant (M21K2c0114) and RIE2025 MTC Programmatic Fund (M22K9b0049) for T.-P. L.

References

- (a) A. Stuetz and G. Petranyi, Synthesis and antifungal activity of (*E*)-*N*-(6,6-dimethyl-2-hepten-4-ynyl)-*N*-methyl-1-naphthalenemethanamine (SF 86–327) and related allylamine derivatives with enhanced oral activity, *J. Med. Chem.*, 1984, **27**, 1539; (b) L. Mohammadkhani and M. M. Heravi, Applications of Transition-Metal-Catalyzed Asymmetric Allylic Substitution in Total Synthesis of Natural Products: An Update, *Chem. Rev.*, 2021, **21**, 29; (c) R. Hairani, R. Mongkol and W. Chavasiri, Allyl and prenyl ethers of mansonone G, new potential semisynthetic antibacterial agents, *Bioorg. Med. Chem. Lett.*, 2016, **26**, 5300.
- (a) M. Johannsen and K. A. Jørgensen, Allylic Amination, *Chem. Rev.*, 1998, **98**, 1689; (b) P. Evans, R. Grange and E. Clizbe, Recent Developments in Asymmetric Allylic Amination Reactions, *Synthesis*, 2016, **48**, 2911; (c) R. A. Fernandes, P. Kattanguru, S. P. Gholap and D. A. Chaudhari, Recent advances in the Overman rearrangement: synthesis of natural products and valuable compounds, *Org. Biomol. Chem.*, 2017, **15**, 2672.
- (a) T. P. Yoon, M. A. Ischay and J. Du, Visible light photocatalysis as a greener approach to photochemical synthesis, *Nat. Chem.*, 2010, **2**, 527; (b) J. M. R. Narayanam and C. R. J. Stephenson, Visible light photoredox catalysis: applications in organic synthesis, *Chem. Soc. Rev.*, 2011, **40**, 102; (c) C. K. Prier, D. A. Rankic and D. W. C. MacMillan, Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis, *Chem. Rev.*, 2013, **113**, 5322.
- (a) N. A. Romero and D. A. Nicewicz, Organic Photoredox Catalysis, *Chem. Rev.*, 2016, **116**, 10075; (b) T. Y. Shang, L. H. Lu, Z. Cao, Y. Liu, W. M. He and B. Yu, Recent advances of 1,2,3,5-tetrakis(carbazol-9-yl)-4,6-dicyanobenzene (4CzIN) in photocatalytic transformations, *Chem. Commun.*, 2019, **55**, 5408; (c) W.-M. Cheng and R. Shang, Transition Metal-Catalyzed Organic Reactions under Visible Light: Recent Developments and Future Perspectives, *ACS Catal.*, 2020, **10**, 9170.
- (a) S. C. Yan, Z. S. Li and Z. G. Zou, Photodegradation of Rhodamine B and Methyl Orange over Boron-Doped *g*-C₃N₄ under Visible Light Irradiation, *Langmuir*, 2010, **26**, 3894; (b) M. Ismael and Y. Wu, A mini-review on the synthesis and structural modification of *g*-C₃N₄-based materials, and their applications in solar energy conversion and environmental remediation, *Sustainable Energy Fuels*, 2019, **3**, 2907; (c) N. Kumar, M. Kumari, M. Ismael, M. Tahir, R. K. Sharma, K. Kumari, J. R. Koduru and P. Singh, Graphitic carbon nitride (*g*-C₃N₄)-assisted materials for the



- detection and remediation of hazardous gases and VOCs, *Environ. Res.*, 2023, **231**, 116149.
- 6 (a) H. Ji, F. Chang, X. Hu, W. Qin and J. Shen, Photocatalytic degradation of 2,4,6-trichlorophenol over *g*-C₃N₄ under visible light irradiation, *Chem. Eng. J.*, 2013, **218**, 183; (b) F. Su, S. C. Mathew, L. Möhlmann, M. Antonietti, X. Wang and S. Blechert, Aerobic Oxidative Coupling of Amines by Carbon Nitride Photocatalysis with Visible Light, *Angew. Chem., Int. Ed.*, 2011, **50**, 657; (c) J. Guo, Y. Wang, Y. Li, K. Lu, S. Liu, W. Wang and Y. Zhang, Graphitic Carbon Nitride Polymer as a Recyclable Photoredox Catalyst for Decarboxylative Alkynylation of Carboxylic Acids, *Adv. Synth. Catal.*, 2020, **362**, 3898; (d) J. Wang, B. Ni, T. Niu and F. Ji, C₃N₄-Photocatalyzed aerobic oxidative cleavage of C≡C bonds in alkynes with diazonium salts leading to two different aldehydes or esters in one pot, *Catal. Sci. Technol.*, 2020, **10**, 8458; (e) J.-Q. Di, M. Zhang, Y.-X. Chen, J.-X. Wang, S.-S. Geng, J.-Q. Tang and Z.-H. Zhang, Copper anchored on phosphorus *g*-C₃N₄ as a highly efficient photocatalyst for the synthesis of *N*-arylpyridin-2-amines, *Green Chem.*, 2021, **23**, 1041; (f) B. Ni, B. Zhang, J. Han, B. Peng, Y. Shan and T. Niu, Heterogeneous Carbon Nitrides Photocatalysis Multicomponent Hydrosulfonylation of Alkynes To Access β -Keto Sulfones with the Insertion of Sulfur Dioxide in Aerobic Aqueous Medium, *Org. Lett.*, 2020, **22**, 670.
- 7 L. Chen, C. S. Chao, Y. Pan, S. Dong, Y. C. Teo, J. Wang and C.-H. Tan, Amphiphilic methyleneamino synthon through organic dye catalyzed-decarboxylative aminoalkylation, *Org. Biomol. Chem.*, 2013, **11**, 5922.
- 8 (a) T. Shao, Y. Yin, R. Lee, X. Zhao, G. Chai and Z. Jiang, Sequential Photoredox Catalysis for Cascade Aerobic Decarboxylative Povarov and Oxidative Dehydrogenation Reactions of *N*-Aryl α -Amino Acids, *Adv. Synth. Catal.*, 2018, **360**, 1754; (b) Y. Liu, X. Liu, J. Li, X. Zhao, B. Qiao and Z. Jiang, Catalytic enantioselective radical coupling of activated ketones with *N*-aryl glycines, *Chem. Sci.*, 2018, **9**, 8094; (c) G. Zeng, Y. Li, B. Qiao, X. Zhao and Z. Jiang, Photoredox asymmetric catalytic enantioconvergent substitution of 3-chlorooxindoles, *Chem. Commun.*, 2019, **55**, 11362; (d) X. Liu, Y. Yin and Z. Jiang, Photoredox-catalysed formal [3 + 2] cycloaddition of *N*-aryl α -amino acids with isoquinoline *N*-oxides, *Chem. Commun.*, 2019, **55**, 11527; (e) H. Yang, G. Wei and Z. Jiang, Access to Isoxazolidines through Visible-Light-Induced Difunctionalization of Alkenes, *ACS Catal.*, 2019, **9**, 9599; (f) W. Hu, Q. Zhan, H. Zhou, S. Cao and Z. Jiang, Radical-based functionalization-oriented construction: rapid assembly of azaarene-substituted highly functionalized pyrroles, *Chem. Sci.*, 2021, **12**, 6543; (g) X. Bai, J. Yao, W. Li, X. Zhao, Y. Yin, S. Yu and Z. Jiang, Enantioselective Hydroaminoalkylation of Azaaryl Ketones through Asymmetric Photoredox Catalysis, *Org. Lett.*, 2024, **26**, 5037.
- 9 (a) T. Shi, K. Sun, X.-L. Chen, Z.-X. Zhang, X.-Q. Huang, Y.-Y. Peng, L.-B. Qu and B. Yu, Recyclable Perovskite as Heterogeneous Photocatalyst for Aminomethylation of Imidazo-Fused Heterocycles, *Adv. Synth. Catal.*, 2020, **362**, 2143; (b) Y.-F. Si, K. Sun, X.-L. Chen, X.-Y. Fu, Y. Liu, F.-L. Zeng, T. Shi, L.-B. Qu and B. Yu, Arylaminoethyl Radical-Initiated Cascade Annulation Reaction of Quinoxalin-2(1*H*)-ones Catalyzed by Recyclable Photocatalyst Perovskite, *Org. Lett.*, 2020, **22**, 6960.
- 10 Y. Chen, X. Ye, F. He and X. Yang, Asymmetric synthesis of oxazolines bearing α -stereocenters through radical addition–enantioselective protonation enabled by cooperative catalysis, *Org. Chem. Front.*, 2021, **8**, 5804.
- 11 (a) W.-T. Ouyang, H.-T. Ji, J. Jiang, C. Wu, J.-C. Hou, M.-H. Zhou, Y.-H. Lu, L.-J. Ou and W.-M. He, Ferrocene/air double-mediated FeTiO₃-photocatalyzed semi-heterogeneous annulation of quinoxalin-2(1*H*)-ones in EtOH/H₂O, *Chem. Commun.*, 2023, **59**, 14029; (b) Y.-H. Lu, S.-Y. Mu, H.-X. Li, J. Jiang, C. Wu, M.-H. Zhou, W.-T. Ouyang and W.-M. He, EtOH-catalyzed electro-synthesis of imidazolidine-fused sulfamidates from *N*-sulfonyl ketimines, *N*-arylglycines and formaldehyde, *Green Chem.*, 2023, **25**, 5539; (c) Y.-H. Lu, Z.-T. Zhang, H.-Y. Wu, M.-H. Zhou, H.-Y. Song, H.-T. Ji, J. Jiang, J.-Y. Chen and W.-M. He, TBAI/H₂O-cooperative electrocatalytic decarboxylation coupling-annulation of quinoxalin-2(1*H*)-ones with *N*-arylglycines, *Chin. Chem. Lett.*, 2023, **34**, 108036.
- 12 H.-H. Li, J.-Q. Li, X. Zheng and P.-Q. Huang, Photoredox-Catalyzed Decarboxylative Cross-Coupling of α -Amino Acids with Nitrones, *Org. Lett.*, 2021, **23**, 876.
- 13 S. Pan, M. Jiang, J. Hu, R. Xu, X. Zeng and G. Zhong, Synthesis of 1,2-amino alcohols by decarboxylative coupling of amino acid derived α -amino radicals to carbonyl compounds via visible-light photocatalyst in water, *Green Chem.*, 2020, **22**, 336.
- 14 W. Zeng, W. Li, H. Chen and L. Zhou, Relay Photocatalytic Reaction of *N*-Aryl Amino Acids and 2-Bromo-3,3,3-trifluoropropene: Synthesis of 4-(Difluoromethylidene)-tetrahydroquinolines, *Org. Lett.*, 2022, **24**, 3265.
- 15 L. Dai, Q. Zhu, J. Zeng, Y. Liu, G. Zhong, X. Han and X. Zeng, Asymmetric synthesis of chiral imidazolidines by merging copper and visible light-induced photoredox catalysis, *Org. Chem. Front.*, 2022, **9**, 2994.
- 16 J.-F. Cui, W.-Q. Zhong and J.-M. Huang, Annulation Reaction of Quinoxalin-2(1*H*)-ones Initiated by Electrochemical Decarboxylation of *N*-Arylglycines, *J. Org. Chem.*, 2023, **88**, 1147.
- 17 K.-H. Lv, L. Chen, K.-H. Zhao, J.-M. Yang and S.-J. Yan, Cu-Catalyzed Decarboxylative Annulation of *N*-Phenylglycines with Maleimides: Synthesis of 1*H*-Pyrrolo[3,4-*c*]quinoline-1,3(2*H*)-diones, *J. Org. Chem.*, 2023, **88**, 2358.
- 18 Seminal studies on silver-catalyzed decarboxylation. (a) Z. Wang, L. Zhu, F. Yin, Z. Su, Z. Li and C. Li, Silver-Catalyzed Decarboxylative Chlorination of Aliphatic Carboxylic Acids, *J. Am. Chem. Soc.*, 2012, **134**, 4258; (b) F. Yin, Z. Wang, Z. Li and C. Li, Silver-Catalyzed Decarboxylative Fluorination of Aliphatic Carboxylic Acids in Aqueous Solution, *J. Am. Chem. Soc.*, 2012, **134**, 10401;



- (c) X. Liu, Z. Wang, X. Cheng and C. Li, Silver-Catalyzed Decarboxylative Alkynylation of Aliphatic Carboxylic Acids in Aqueous Solution, *J. Am. Chem. Soc.*, 2012, **134**, 14330; (d) C. Liu, X. Wang, Z. Li, L. Cui and C. Li, Silver-Catalyzed Decarboxylative Radical Azidation of Aliphatic Carboxylic Acids in Aqueous Solution, *J. Am. Chem. Soc.*, 2015, **137**, 9820; (e) L. Cui, H. Chen, C. Liu and C. Li, Silver-Catalyzed Decarboxylative Allylation of Aliphatic Carboxylic Acids in Aqueous Solution, *Org. Lett.*, 2016, **18**, 2188; (f) Y. Dong, Z. Wang and C. Li, Controlled radical fluorination of poly (meth)acrylic acids in aqueous solution, *Nat. Commun.*, 2017, **8**, 277; (g) X. Tan, Z. Liu, H. Shen, P. Zhang, Z. Zhang and C. Li, Silver-Catalyzed Decarboxylative Trifluoromethylation of Aliphatic Carboxylic Acids, *J. Am. Chem. Soc.*, 2017, **139**, 12430.
- 19 Y. Cai, Y. Tang, L. Fan, Q. Lefebvre, H. Hou and M. Rueping, Heterogeneous Visible-Light Photoredox Catalysis with Graphitic Carbon Nitride for α -Aminoalkyl Radical Additions, Allylations, and Heteroarylations, *ACS Catal.*, 2018, **8**, 9471.
- 20 (a) Y.-F. Si, X.-L. Chen, X.-Y. Fu, K. Sun, Xi. Song, L.-B. Qu and B. Yu, Divergent g -C₃N₄-catalyzed Reactions of Quinoxalin-2(1H)-ones with *N*-Aryl Glycines under Visible Light: Solvent-Controlled Hydroaminomethylation and Annulation, *ACS Sustainable Chem. Eng.*, 2020, **8**, 10740; (b) A. Shi, K. Sun, Y. Wu, P. Xiang, I. B. Krylov, A. O. Terent'ev, X. Chen and B. Yu, Oxygen-doped carbon nitride for enhanced photocatalytic activity in visible-light-induced decarboxylative annulation reactions, *J. Catal.*, 2022, **415**, 28; (c) T. Shi, Y.-T. Liu, S.-S. Wang, Q.-Y. Lv and B. Yu, Recyclable Carbon Nitride Nanosheet-Photocatalyzed Aminomethylation of Imidazo[1,2-*a*]pyridines in Green Solvent, *Chin. J. Chem.*, 2022, **40**, 97.
- 21 J. Tian, L. Zhao, C. Yang, C. Yang, L. Guo and W. Xia, Four-Component Synthesis of Spiro-Imidazolidines Enabled by Carbon Nitride Photocatalysis, *ACS Catal.*, 2023, **13**, 866.

