

## RESEARCH ARTICLE

View Article Online  
View Journal | View IssueCite this: *Org. Chem. Front.*, 2025, 12, 6480Received 9th July 2025,  
Accepted 13th August 2025

DOI: 10.1039/d5qo01005e

rsc.li/frontiers-organic



# Palladium/photoredox catalysis for a decarboxylative radical C(sp<sup>3</sup>)-C(sp<sup>3</sup>) cross-coupling reaction of $\gamma$ -methylidene- $\delta$ -valerolactones with alkyl carboxylic acids

Ran Song,<sup>a</sup> Jiaying Li,<sup>a</sup> Lin Li,<sup>a</sup> Huaixi Mu,<sup>a</sup> Wei Feng,<sup>a</sup> Daoshan Yang,<sup>a,b</sup> Wen Si<sup>\*a</sup> and Jian Lv<sup>id</sup> <sup>\*a</sup>

Developing applicable methods to forge linkages between sp<sup>3</sup>- and sp<sup>3</sup>-hybridized carbons is of great significance in drug discovery. Here, a radical decarboxylative C(sp<sup>3</sup>)-C(sp<sup>3</sup>) cross-coupling of  $\gamma$ -methylidene- $\delta$ -valerolactones (GMDVs) and alkyl carboxylic acids was developed by combining photocatalysis and palladium catalysis, providing the target products in good to excellent yields (up to 97% yield). The reaction exhibited a broad substrate scope and wide functional group compatibility.

Transition metal-catalyzed decarboxylative cross-coupling has emerged as a strategically significant methodology in modern organic synthesis, which features inherent atom economy through liberation of CO<sub>2</sub> as the sole byproduct, aligning with green chemistry principles.<sup>1</sup> Parallel advancements in photoredox catalysis and synergistic transition metal/photoredox dual catalytic systems have revolutionized radical-mediated C-C bond formation, enabling mild, energy-efficient protocols that circumvent traditional harsh reaction conditions.<sup>2</sup> The integration of decarboxylative strategies with photoredox catalysis presents a compelling synergy, combining operational simplicity with environmental benignity.<sup>3</sup> However, current methodologies predominantly require preactivation of carboxylic acids into reactive intermediates such as *N*-hydroxyphthalimide esters or hypervalent iodine(III) species.<sup>4</sup> This activation barrier has rendered direct utilization of naturally occurring alkyl carboxylic acids in photoredox-mediated Csp<sup>3</sup>-Csp<sup>3</sup> cross-coupling reactions exceptionally rare,<sup>5</sup> despite their theoretical potential as ideal coupling partners (Scheme 1A). The development of such underdeveloped yet fundamentally important transformations therefore represents a significant frontier in catalytic methodology, offering opportunities to streamline synthetic sequences while accessing structurally diverse molecular architectures through operationally simple, sustainable protocols.

$\gamma$ -Methylidene- $\delta$ -valerolactones (GMDVs), first established as versatile C<sub>4</sub>-synthons through Shintani and Hayashi's seminal [4 + *n*] cycloaddition chemistry in 2007,<sup>6,7</sup> have remained underexplored as coupling partners compared to conventional organometallic reagents. Our prior investigations revealed the unique reactivity profile of these cyclic systems, demonstrating their capacity as *O*-allylation reagents for phenolic substrates through Csp<sup>3</sup>-O bond formation under mild conditions.<sup>8a</sup> Very recent advances in our group further expanded this paradigm through the development of a Pd<sup>0</sup>/Cu<sup>II</sup>/VB<sub>2</sub>-catalyzed *N*-allylation protocol with aniline derivatives under blue LED irradiation, achieving efficient construction of allylamine architectures.<sup>8b</sup> Despite these advances, the direct engagement of GMDVs in Csp<sup>3</sup>-Csp<sup>3</sup> bond-forming processes

A) Transition metal-catalyzed decarboxylative Csp<sup>3</sup>-Csp<sup>3</sup> cross-coupling reactionB) This work: Pd/photoredox dual-catalyzed decarboxylative Csp<sup>3</sup>-Csp<sup>3</sup> reaction of GMDVs**Scheme 1** Strategies for the decarboxylative Csp<sup>3</sup>-Csp<sup>3</sup> cross-coupling reaction.

<sup>a</sup>Key Laboratory of Optic-electric Sensing and Analytical Chemistry for Life Science, MOE, College of Chemistry and Molecular Engineering, Qingdao University of Science and Technology, Qingdao, 266042, P. R. China. E-mail: hjian@iccas.ac.cn, siwen@qust.edu.cn


<sup>b</sup>National Engineering Research Center of Low-Carbon Processing and Utilization of Forest Biomass, Nanjing Forestry University, Nanjing 210037, China



has remained conspicuously absent from the synthetic toolbox. To address this gap, we herein disclose a pioneering dual decarboxylative strategy that synergistically combines GMDVs with alkyl carboxylic acids under palladium/photo-redox dual catalysis (Scheme 1B).

To validate our hypothesis, we commenced the investigation by employing GMDVs **1a** and 9H-fluorene-9-carboxylic acid **2** as model substrates in a palladium-catalyzed system (Table 1). The initial reaction utilizing Pd(PPh<sub>3</sub>)<sub>4</sub> (5 mol%), Ir(dFMeppy)<sub>2</sub>(dtbbpy)(PF<sub>6</sub>) (**PC-1**, 2 mol%) as a photocatalyst, and K<sub>2</sub>CO<sub>3</sub> (4 equiv.) in DMSO under 370 nm UV irradiation (10 W LED) at ambient temperature under a nitrogen atmosphere afforded the Csp<sup>3</sup>-Csp<sup>3</sup> coupling product **3a** in 82% isolated yield (entry 1). Control experiments revealed critical dependencies: partial suppression of **3a** formation occurred in the absence of a photocatalyst (entry 2), while exclusive generation of the Csp<sup>3</sup>-O byproduct **3a'** was observed under dark conditions (entry 3). Subsequent systematic optimization identified DMA as the superior solvent (96% yield, entry 7 vs. entries 1, 5 and 6), outperforming DMF (78%) and maintaining solvent superiority over DMSO (82%). The reaction demonstrated strict base dependence, with Cs<sub>2</sub>CO<sub>3</sub> (64%), Na<sub>2</sub>CO<sub>3</sub> (78%), KH<sub>2</sub>PO<sub>4</sub> (34%), NaH<sub>2</sub>PO<sub>4</sub> (39%), KOAc (60%) and NaOAc (60%) proving inferior to K<sub>2</sub>CO<sub>3</sub> (entry 1 vs. Table S1, entries 2–7). Notably, stoichiometric analysis revealed the

**Table 1** Screening of reaction conditions




Entry <sup>a</sup>	Variation from standard conditions	Yield <sup>b</sup> (%)
1	None	82
2	No photocatalyst	32
3	In the dark	np <sup>c</sup>
4	Other bases instead of K <sub>2</sub> CO <sub>3</sub>	34–78
5	DMF instead of DMSO	62
6	THF	52
7	DMA	96
8	K[2] instead of 2/K <sub>2</sub> CO <sub>3</sub>	nr
9 <sup>d</sup>	2/K <sub>2</sub> CO <sub>3</sub> (1 : 1, 3.0 equiv.)	64
10 <sup>d</sup>	PC-2 instead of PC-1	62
11 <sup>d</sup>	PC-3	78
12 <sup>d</sup>	PC-4	38
13 <sup>d</sup>	390 nm instead of 370 nm	93
14 <sup>d</sup>	427 nm	75
15 <sup>d</sup>	456 nm	54
16 <sup>d</sup>	2 h	94
17 <sup>d</sup>	1 h	76

<sup>a</sup> Standard reaction conditions: under a nitrogen atmosphere, **1a** (0.1 mmol), **2** (0.4 mmol), Pd(PPh<sub>3</sub>)<sub>4</sub> (5 mol%), Ir(dFMeppy)<sub>2</sub>(dtbbpy)(PF<sub>6</sub>) (2 mol%), and K<sub>2</sub>CO<sub>3</sub> (0.4 mmol) in DMSO (1.0 mL), 370 nm (10 W) at room temperature under N<sub>2</sub> for 12 h. <sup>b</sup> Isolated yield. <sup>c</sup> np = no product; by-product **3a'** was obtained in 7% yield. <sup>d</sup> DMA (1.0 mL) as solvent. nr = no reaction.

necessity of excess **2** (4 equiv.) and K<sub>2</sub>CO<sub>3</sub> (4 equiv.), as reduced equivalents diminished yields (entry 9), while substitution of K<sub>2</sub>CO<sub>3</sub> and **2** with potassium carboxylate K[**2**] completely abolished reactivity (entry 8), strongly implicating *in situ*-generated H<sub>2</sub>O or free carboxylic acid as crucial proton donors. Photocatalyst screening confirmed **PC-1**'s superiority over alternative photocatalysts (entry 7 vs. entries 10–12), with 370 nm UV irradiation proving optimal among the tested wavelengths (entry 7 vs. entries 13–15). Temporal analysis revealed exceptional reaction efficiency, achieving 94% yield within 2 hours (entry 16), though incomplete conversion (76%) resulted from premature termination at 1 hour (entry 17).

With the optimized conditions in hand, we next investigated the substrate scope of the decarboxylative radical C(sp<sup>3</sup>)-C(sp<sup>3</sup>) cross-coupling reaction between GMDVs **1** and alkyl carboxylic acid **2**. At first, we varied the GMDVs **1** and the results are summarized in Table 2. We were delighted to find that a wide variety of structurally diverse GMDVs bearing either electron-donating groups (Me and OMe), electron-withdrawing groups (Cl and Br) or electron-neutral groups (H) at various positions of the aryl ring smoothly reacted with 9H-fluorene-9-carboxylic acid **2**, affording the corresponding cross-coupling products **3a–j** in moderate to excellent yields (entries 1–10). Aliphatic GMDV **3k** did not work well (entry 11), but heteroaromatic 3-thiophene was also tolerated in the reaction, affording the target product **3l** in good yield (entry 12). Gratifyingly, ethyl ester and cyano substrates were also well tolerated, affording the target product **3m** in 91% yield and product **3n** in 97% yield (entries 13 and 14), respectively. Unfortunately,

**Table 2** Substrate scope for GMDVs **1**<sup>a</sup>



Entry <sup>a</sup>	R <sup>1</sup>	R <sup>2</sup>	3	Yield <sup>b</sup> (%)
1	Ph	CO <sub>2</sub> Me	<b>3a</b>	94
2	2-ClC <sub>6</sub> H <sub>4</sub>	CO <sub>2</sub> Me	<b>3b</b>	86
3	2-BrC <sub>6</sub> H <sub>4</sub>	CO <sub>2</sub> Me	<b>3c</b>	40
4	2-MeC <sub>6</sub> H <sub>4</sub>	CO <sub>2</sub> Me	<b>3d</b>	81
5	2-MeOC <sub>6</sub> H <sub>4</sub>	CO <sub>2</sub> Me	<b>3e</b>	85
6	4-MeC <sub>6</sub> H <sub>4</sub>	CO <sub>2</sub> Me	<b>3f</b>	75
7	4-MeOC <sub>6</sub> H <sub>4</sub>	CO <sub>2</sub> Me	<b>3g</b>	78
8	3,4-Me <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	CO <sub>2</sub> Me	<b>3h</b>	72
9	3,4-(MeO) <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	CO <sub>2</sub> Me	<b>3i</b>	89
10	3,4-(OCH <sub>2</sub> O) <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	CO <sub>2</sub> Me	<b>3j</b>	70
11	Bn	CO <sub>2</sub> Me	<b>3k</b>	25
12	3-Thiophene	CO <sub>2</sub> Me	<b>3l</b>	90
13	Ph	CO <sub>2</sub> Et	<b>3m</b>	91
14	Ph	CN	<b>3n</b>	97
15	Ph	Ph	<b>3o</b>	nd <sup>c</sup>


<sup>a</sup> Standard reaction conditions: under a nitrogen atmosphere, **1** (0.1 mmol), **2** (0.4 mmol, 4.0 equiv.), Pd(PPh<sub>3</sub>)<sub>4</sub> (5 mol%), Ir(dFMeppy)<sub>2</sub>(dtbbpy)(PF<sub>6</sub>) (2 mol%), and K<sub>2</sub>CO<sub>3</sub> (0.4 mmol, 4.0 equiv.) in DMA (1.0 mL), blue LEDs (370 nm, 10 W) at room temperature for 2 h. <sup>b</sup> Isolated yield. <sup>c</sup> nd = not determined.



no desired product was observed when di-phenyl-substituted GMDV was introduced into the reaction (entry 15).

Furthermore, different alkyl carboxylic acids were examined. As depicted in Scheme 2, the reaction proceeded well when primary carboxylic acids bearing electron-withdrawing groups (F, Cl, Br, and CF<sub>3</sub>), electron-donating groups (OMe and Ph) or electron-neutral groups (H) were applied, affording the desired products **5a–h** in moderate yield and a byproduct **5'** was observed. Moreover, substrates bearing various heterocycles, such as indole **4l** or thiophene **4m**, could also work, affording the target products **5l** and **5m** in slightly lower yields. In addition, secondary benzylic-type carboxylic acids **4i** and **4j** also exhibited good reactivity. To our delight, commercially available and commonly-used anti-inflammatory drugs, such as naproxen **4k**, were tolerated in this radical cross-coupling to afford the corresponding products. Finally, tertiary carboxylic acids were also investigated. To our satisfaction, the reactions could work well when flurenol **4n** and 9-methylfluorene-9-carboxylic acid **4o** were used as substrates, and the target products could be obtained in up to 50% yield. Unfortunately, the reaction did not work with inactivated aliphatic carboxylic acid **3p**.

Under the standard conditions, the developed decarboxylative cross-coupling reaction of GMDVs **1a** (1.08 g, 4.4 mmol) could be performed on a gram scale, affording the product **3a** in 86% yield (Scheme 3A). In addition, a sunlight-accelerated experiment was carried out, affording the desired product **3a** in 78% yield (Scheme 3B). To further confirm the structures of products **3** and **5**, compound **3a** was hydrolyzed with LiOH in a mixture of methanol and water to obtain carboxylic acid **6** in 89% yield. The solid structure of compound **6** (CCDC 2358852) was unambiguously assigned by X-ray crystallography (Scheme 3C-a).<sup>10</sup>




**Scheme 3** Gram-scale reaction and synthetic transformation of **3a**.<sup>a</sup>

<sup>a</sup> Reaction conditions: (a) LiOH·H<sub>2</sub>O (2.0 equiv.), MeOH/H<sub>2</sub>O (10 : 1, 2.2 mL), rt, and 12 h. (b) DIBAL-H (1.0 mL), THF (0.1 M), rt, and 12 min. (c) PCC (1.5 equiv.), NaOAc (1.5 equiv.), DCM (0.1 M), rt, and 12 h. (d) 4-ClC<sub>6</sub>H<sub>4</sub>OH (0.8 equiv.), DMAP (10 mol%), DCC (0.8 equiv.), DCM (0.5 mL), N<sub>2</sub>, rt, and 15 h. <sup>b</sup> ORTEP drawing at 30% ellipsoid probability.


Treatment of compound **6** with DIBAL-H in THF at room temperature led to the reduction of the carboxyl group, providing alcohol **7** in 91% yield (Scheme 3C-b), which could be further oxidized with PCC in DCM to afford aldehyde **8** in 53% yield (Scheme 3C-c). Moreover, compound **3a** could also react with 4-ClC<sub>6</sub>H<sub>4</sub>OH to afford product **9** in 69% yield (Scheme 3C-d).

To investigate the mechanism of this transformation, several control experiments were performed. Initially, three radical trapping experiments were conducted. Under the standard conditions, the addition of (2,2,6,6-tetramethylpiperidin-1-yl)oxidanyl (TEMPO), 2,6-di-tertbutyl-4-methylphenol (BHT) and 1,1-diphenylethylene to the reaction solution significantly suppressed these transformations, and the radical trapping intermediates **10–13** were detected by HRMS, respectively (Scheme 4A). These results, as well as the formation of byproduct **5'**, indicated the presence of allyl and benzyl radicals in this reaction. Stern–Volmer quenching experiments indicate that a combination of K<sub>2</sub>CO<sub>3</sub> and **2** quenches the photocatalyst (Fig. S8). Interestingly, when the combination of K<sub>2</sub>CO<sub>3</sub> and **2** was replaced by fluorene-9-carboxylic acid **2** or its potassium salt, this reaction did not work, suggesting that K<sub>2</sub>CO<sub>3</sub> should be used as a proton shuttle between **2** and Pd(η-1,4-zwitterionic complexes). In addition, it was found that this reaction also worked in the absence of a photosensitizer, albeit in lower yield. This result indicates that in addition to Ir(III) photooxidants, Pd(I) species could also oxidize the carboxylate of **2** to generate benzyl radicals but with lower reactivity. Finally, we performed an on–off experiment. The reaction progressed smoothly upon irradiation with light (Fig. S9), while the reaction paused during the “light-off” periods. These results, as well as the quantum yield (1.73%), suggest that the present transformation was not a radical-chain process.



**Scheme 2** Substrate scope for alkyl carboxylic acids **4**.<sup>a</sup> Standard reaction conditions: under a nitrogen atmosphere, **1a** (0.1 mmol), **4** (0.4 mmol, 4.0 equiv.), Pd(PPh<sub>3</sub>)<sub>4</sub> (5 mol%), Ir(dFMeppy)<sub>2</sub>(dtbbpy)(PF<sub>6</sub>) (2 mol%), K<sub>2</sub>CO<sub>3</sub> (0.4 mmol, 4.0 equiv.) in DMA (1.0 mL), blue LEDs (370 nm, 10 W) at room temperature for 2 h, Isolated product yield.





**Scheme 4** (A) Control experiment and (B) proposed mechanism.

Based on the above mechanistic studies and previous reports,<sup>9</sup> a mechanism for this decarboxylative cross-coupling is proposed (Scheme 4B). This transformation starts with the palladium-catalyzed decarboxylation of GMDVs to generate Pd(II)-1,4-zwitterionic complexes **A**, which are easily protonated to form intermediate **B** by the combination of  $K_2CO_3$  and **2**. The allyl Pd(II) intermediate **B** undergoes homolysis to form hybrid allyl Pd(I) radical **C** by utilizing light energy,<sup>10</sup> followed by the generation of free allyl radical **D** and Pd(I). On the one hand, the generated Pd(I) complex slowly reacted with carboxylate **E** from fluorene-9-carboxylic acid **2** to regenerate the Pd(0) catalyst and benzyl radical **F**. On the other hand, a single electron transfer (SET) between carboxylate **E** and excited Ir(III) released radical **F**, along with the generation of Ir(II), which much faster reacted with Pd(I) to regenerate the Pd(0) catalyst and Ir(III) photosensitizer. Finally, the generated allyl radical **D** coupled with benzyl radical **F** to form the desired product **3a**.

## Conclusions

In summary, here, we have successfully demonstrated a mild and efficient decarboxylative radical  $C(sp^3)-C(sp^3)$  cross-coupling reaction of  $\gamma$ -methylidene- $\delta$ -valerolactones with various alkyl carboxylic acids by combining photocatalysis and palladium catalysis. Primary, secondary and tertiary carboxylic acids can all be well tolerated to afford the target products in moderate to excellent yields. Meanwhile, GMDVs were first employed in the photoredox-catalyzed procedure and exhibited a broad

substrate scope and outstanding functional group tolerance, which provides an alternative approach for  $C(sp^3)-C(sp^3)$  bond formation.

## Conflicts of interest

There are no conflicts to declare.

## Data availability

The data supporting this article have been included as part of the SI. The SI contains: general experimental information, optimization of reaction conditions, experimental procedure, characterization data for target products, mechanistic studies, X-ray crystallography data and copies of NMR spectra of compounds. See DOI: <https://doi.org/10.1039/d5qo01005e>.

CCDC 2358852 contains the supplementary crystallographic data for this paper.<sup>11</sup>

## Acknowledgements

We thank the Natural Science Foundation of Shandong Province for Distinguished Young Scholars (ZR2020JQ07), the Natural Science Foundation of Shandong Province, China (No. ZR2019PB020), and Qingdao University of Science and Technology for financial support.

## References

- (a) L. J. Gooßen, N. Rodríguez and K. Gooßen, Carboxylic Acids as Substrates in Homogeneous Catalysis, *Angew. Chem., Int. Ed.*, 2008, **47**, 3100; (b) P. Liu, G. H. Zhang and P. P. Sun, Transition metal-free decarboxylative alkylation reactions, *Org. Biomol. Chem.*, 2016, **14**, 10763; (c) T. Patra and D. Maiti, Decarboxylation as the Key Step in C–C Bond-Forming Reactions, *Chem. – Eur. J.*, 2017, **23**, 7382; (d) A. Varenikov, E. Shapiro and M. Gandelman, Decarboxylative Halogenation of Organic Compounds, *Chem. Rev.*, 2021, **121**, 412.
- (a) C. K. Prier, D. A. Rankic and D. W. C. MacMillan, Visible Light Photoredox Catalysis with Transition Metal Complexes: Applications in Organic Synthesis, *Chem. Rev.*, 2013, **113**, 5322; (b) M. H. Shaw, J. Twilton and D. W. C. MacMillan, Photoredox Catalysis in Organic Chemistry, *J. Org. Chem.*, 2016, **81**, 6898; (c) K. L. Skubi, T. R. Blum and T. P. Yoon, Dual Catalysis Strategies in Photochemical Synthesis, *Chem. Rev.*, 2016, **116**, 10035; (d) H. Yi, G. Zhang, H. Wang, Z. Huang, J. Wang, A. K. Singh and A. Lei, Recent Advances in Radical C–H Activation/Radical Cross-Coupling, *Chem. Rev.*, 2017, **117**, 9016; (e) Q.-Y. Huang and M. Shi, Recent advancements in Ni/photoredox dual catalysis for  $Csp^3-Csp^3$  cross-coupling reactions, *Org. Chem. Front.*, 2024, **11**, 4913.



- 3 (a) S. B. Beil, T. Q. Chen, N. E. Intermaggio and D. W. C. MacMillan, Carboxylic Acids as Adaptive Functional Groups in Metallaphotoredox Catalysis, *Acc. Chem. Res.*, 2022, **55**, 3481; (b) S. Karmakar, A. Silamkoti, N. A. Meanwell, A. Mathur and A. K. Gupta, Utilization of C(sp<sup>3</sup>)-Carboxylic Acids and Their Redox-Active Esters in Decarboxylative Carbon–Carbon Bond Formation, *Adv. Synth. Catal.*, 2021, **363**, 3693.
- 4 (a) S. K. Parida, T. Mandal, S. Das, S. K. Hota, S. D. Sarkar and S. Murarka, Single Electron Transfer-Induced Redox Processes Involving N-(Acyloxy)phthalimides, *ACS Catal.*, 2021, **11**, 1640; (b) T. Qin, J. Cornella, C. Li, L. R. Malins, J. T. Edwards, S. Kawamura, B. D. Maxwell, M. D. Eastgate and P. S. Baran, A general alkyl-alkyl cross-coupling enabled by redox-active esters and alkylzinc reagents, *Science*, 2016, **352**, 801; (c) X. J. Zeng, W. H. Yan, S. B. Zacate, T. H. Chao, X. D. Sun, Z. Cao, K. G. E. Bradford, M. Paeth, S. B. Tyndall, K. D. Yang, T. C. Kuo, M. J. Cheng and W. Liu, Copper-Catalyzed Decarboxylative Difluoromethylation, *J. Am. Chem. Soc.*, 2019, **141**, 11398; (d) W. Liu, M. N. Lavagnino, C. A. Gould, J. Alcázar and D. W. C. MacMillan, A biomimetic SH<sub>2</sub> cross-coupling mechanism for quaternary sp<sup>3</sup>-carbon formation, *Science*, 2021, **374**, 1258; (e) C. Jiang, P. Chen and G. Liu, Cu/photoredox-catalyzed decarboxylative radical C(sp<sup>3</sup>)-C(sp<sup>3</sup>) cross-coupling reactions, *Sci. China: Chem.*, 2023, **66**, 2858; (f) M. Huang, H. Sun, F. Seufert, A. Friedrich, T. B. Marder and J. Hu, Photoredox/Cu-Catalyzed Decarboxylative C(sp<sup>3</sup>)-C(sp<sup>3</sup>) Coupling to Access C(sp<sup>3</sup>)-Rich gem-Diborylalkanes, *Angew. Chem., Int. Ed.*, 2024, **63**, e202401782; (g) Z. Huang, M. E. Akana, K. M. Sanders and D. J. Weix, A decarbonylative approach to alkylnickel intermediates and C(sp<sup>3</sup>)-C(sp<sup>3</sup>) bond formation, *Science*, 2024, **385**, 1331; (h) X. Zhao, C. Wang, L. Yin and W. Liu, Highly Enantioselective Decarboxylative Difluoromethylation, *J. Am. Chem. Soc.*, 2024, **146**, 29297; (i) L. Liang, Y.-H. Wang, C.-X. Cui, X.-S. Deng, S.-L. Wang, H.-M. Guo, Y. Li, H.-Y. Niu and R. Mao, NADH Analogues Enable Metal- and Light-Free Decarboxylative Functionalization, *Angew. Chem., Int. Ed.*, 2024, e202415131.
- 5 (a) C. P. Johnston, R. T. Smith, S. Allmendinger and D. W. C. MacMillan, Metallaphotoredox-catalysed sp<sup>3</sup>-sp<sup>3</sup> cross-coupling of carboxylic acids with alkyl halides, *Nature*, 2016, **536**, 322; (b) Y. Duan, M. Zhang, R. Ruzi, Z. Wu and C. Zhu, The direct decarboxylative allylation of N-arylglycine derivatives by photoredox catalysis, *Org. Chem. Front.*, 2017, **4**, 525; (c) K. C. Cartwright and J. A. Tunge, Organophotoredox/palladium dual catalytic decarboxylative Csp<sup>3</sup>-Csp<sup>3</sup> coupling of carboxylic acids and  $\pi$ -electrophiles, *Chem. Sci.*, 2020, **11**, 8167; (d) H. A. Sakai and D. W. C. MacMillan, Nontraditional Fragment Couplings of Alcohols and Carboxylic Acids: C(sp<sup>3</sup>)-C(sp<sup>3</sup>) Cross-Coupling via Radical Sorting, *J. Am. Chem. Soc.*, 2022, **144**, 6185; (e) A. V. Tsymbal, L. D. Bizzini and D. W. C. MacMillan, Nickel Catalysis via SH<sub>2</sub> Homolytic Substitution: The Double Decarboxylative Cross-Coupling of Aliphatic Acids, *J. Am. Chem. Soc.*, 2022, **144**, 21278; (f) N. Xiong, Y. Li and R. Zeng, Merging Photoinduced Iron-Catalyzed Decarboxylation with Copper Catalysis for C–N and C–C Couplings, *ACS Catal.*, 2023, **13**, 1678.
- 6 (a) R. Shintani, M. Murakami and T. Hayashi,  $\gamma$ -Methylidene- $\delta$ -valerolactones as a Coupling Partner for Cycloaddition: Palladium-Catalyzed [4 + 3] Cycloaddition with Nitrones, *J. Am. Chem. Soc.*, 2007, **129**, 12356; (b) R. Shintani, M. Murakami, T. Tsuji, H. Tanno and T. Hayashi, Palladium-Catalyzed Decarboxylative [4 + 3] Cyclization of  $\gamma$ -Methylidene- $\delta$ -valerolactones with 1,1-Dicyanocyclopropanes, *Org. Lett.*, 2009, **11**, 5642; (c) R. Shintani, T. Ito and T. Hayashi, Palladium-Catalyzed Synthesis of 4-Oxaspiro[2.4]heptanes via Central Attack of Oxygen Nucleophiles to  $\pi$ -Allylpalladium Intermediates, *Org. Lett.*, 2012, **14**, 2410; (d) S.-Y. Baek, J. Y. Lee, D. Ko, M.-H. Baik and E. J. Yoo, Rationally Designing Regiodivergent Dipolar Cycloadditions: Frontier Orbitals Show How To Switch between [5 + 3] and [4 + 2] Cycloadditions, *ACS Catal.*, 2018, **8**, 6353; (e) L.-C. Yang, Y.-N. Wang, R. Liu, Y. Luo, X.-Q. Ng, B. Yang, Z.-Q. Rong, Y. Lan, Z. Shao and Y. Zhao, Stereoselective access to [5.5.0] and [4.4.1] bicyclic compounds through Pd-catalysed divergent higher-order cycloadditions, *Nat. Chem.*, 2020, **12**, 860; (f) Z.-L. Jia, X.-T. An, Y.-H. Deng, L.-H. Pang, C.-F. Liu, L.-L. Meng, J.-K. Xue, X.-H. Zhao and C.-A. Fan, Palladium-Catalyzed Asymmetric (4 + 2) Annulation of  $\gamma$ -Methylidene- $\delta$ -valerolactones with Alkenes: Enantioselective Synthesis of Functionalized Chiral Cyclohexyl Spirooxindoles, *Org. Lett.*, 2021, **23**, 745; (g) C. Gao, X. Wang, J. Liu and X. Li, Highly Diastereo- and Enantioselective Synthesis of Tetrahydrobenzo[b]azocines via Palladium-Catalyzed [4 + 4] Cycloaddition, *ACS Catal.*, 2021, **11**, 2684; (h) C. Gao, T. Zhang, X. Li, J.-D. Wu and J. Liu, Asymmetric decarboxylative [3 + 2] cycloaddition for the diastereo- and enantioselective synthesis of spiro[2.4]heptanes via cyclopropanation, *Org. Chem. Front.*, 2022, **9**, 2121; (i) X. Wang, J. Yang, R. Lv, P. Song, D. Ye, J. Liu and X. Li, Palladium-catalyzed [4 + 4] cycloadditions for highly diastereo- and enantioselective synthesis of functionalized benzo[b]oxocines, *Org. Chem. Front.*, 2022, **9**, 3493; (j) Y. Cao, H. Wang, X. Chen, Y. Qiao and Z. Miao, Gold and Palladium Relay Catalytic [4 + 4] Cycloadditions of Enynamides and  $\gamma$ -Methylene- $\delta$ -valerolactones: Diastereoselective Construction of Furan-Fused Azacyclooctanes, *J. Org. Chem.*, 2023, **88**, 11822; (k) T.-X. Liu, Y. Ru, W. Guo, N. Ma, P. Yang, X. Li, P. Zhang, J. Bi and G. Zhang, Catalytic System-Controlled Regioselective 1,2- and 1,4-Carboannulations of [60] Fullerene, *Org. Lett.*, 2024, **26**, 2552.
- 7 For an example of GMDVs involving in cycloadditions reported by our group, see: X. Wu, W. Feng, J. Zhao, D. Yang, R. Song, W. Si and J. Lv, Neighboring  $\pi$ -Coordination in Palladium Catalyzed [4 + 4] Cycloaddition of  $\gamma$ -Methylidene- $\delta$ -Valerolactones with Divinylketones, *Adv. Synth. Catal.*, 2024, **366**, 4049.



- 8 (a) R. Song, Z. Lian, W. Feng, T. Guan, W. Si, D. Yang and J. Lv, Palladium-catalyzed decarboxylative O-allylation of phenols with  $\gamma$ -methylidene- $\delta$ -valerolactones, *Org. Chem. Front.*, 2022, **9**, 4365; (b) H. Yang, K. Pang, B. Xue, X. Li, R. Song, D. Yang, W. Si and J. Lv, Visible-Light-Accelerated Decarboxylative N-Allylation of  $\gamma$ -Methylidene- $\delta$ -valerolactones via Synergistic Pd(0)/Cu(I)/Riboflavin Catalysis, *Org. Lett.*, 2025, **27**, 4547.
- 9 (a) H.-H. Zhang, M. Tang, J.-J. Zhao, C. Song and S. Yu, Enantioselective Reductive Homocoupling of Allylic Acetates Enabled by Dual Photoredox/Palladium Catalysis: Access to C2-Symmetrical 1,5-Dienes, *J. Am. Chem. Soc.*, 2021, **143**, 12836; (b) B. Li, H.-H. Zhang, Y. Luo, S. Yu, W. A. Goddard III and Y. Dang, Interception of Transient Allyl Radicals with Low-Valent Allylpalladium Chemistry: Tandem Pd(0/II/I)–Pd(0/II/I/II) Cycles in Photoredox/Pd Dual-Catalytic Enantioselective C(sp<sup>3</sup>)–C(sp<sup>3</sup>) Homocoupling, *J. Am. Chem. Soc.*, 2024, **146**, 6377.
- 10 (a) P. M. Waddell, L. Tian, A. R. Scavuzzo, L. Venigalla, G. D. Scholes and B. P. Carrow, Visible light-induced palladium–carbon bond weakening in catalytically relevant T-shaped complexes, *Chem. Sci.*, 2023, **14**, 14217; (b) H. Dau, A. Keyes, H. E. B. Alhan, E. Ordonez, E. Tsogtgerel, A. P. Gies, E. Auyeung, Z. Zhou, A. Maity, A. Das, D. C. Powers, D. B. Beezer and E. Harth, Dual Polymerization Pathway for Polyolefin-Polar Block Copolymer Synthesis via MILRad: Mechanism and Scope, *J. Am. Chem. Soc.*, 2020, **142**, 21469; (c) W.-M. Chen and R. Shang, Transition Metal-Catalyzed Organic Reactions under Visible Light: Recent Developments and Future Perspectives, *ACS Catal.*, 2020, **10**, 9170; (d) K. P. S. Cheung, S. Sarkar and V. Gevorgyan, Visible Light-Induced Transition Metal Catalysis, *Chem. Rev.*, 2022, **122**, 1543.
- 11 W. Feng, CCDC 2358852 (6): Experimental Crystal Structure Determination, 2024, DOI: [10.5517/ccdc.csd.cc2k510c](https://doi.org/10.5517/ccdc.csd.cc2k510c).

