

CORRECTION

View Article Online

View Journal | View Issue

Cite this: *Org. Chem. Front.*, 2025, **12**, 3942

Correction: Twisted organic TADF triads based on a diindolocarbazole donor for efficient photoisomerization of stilbene and photo-arylation of heteroarenes

Sushil Sharma and Sanchita Sengupta*

DOI: 10.1039/d5qo90048d
rsc.li/frontiers-organicCorrection for 'Twisted organic TADF triads based on a diindolocarbazole donor for efficient photoisomerization of stilbene and photo-arylation of heteroarenes' by Sushil Sharma *et al.*, *Org. Chem. Front.*, 2023, **10**, 6087–6095, <https://doi.org/10.1039/D3QO01542D>.

The authors regret that the singlet and triplet energy values were incorrectly reported in Table 1. The corrected Table 1 appears below.

Table 1 Absorption wavelength (λ_{abs}), emission wavelength (λ_{em}), Stokes shift, molar extinction coefficient (ϵ) and singlet energy (S_1), triplet energy (T_1) and singlet–triplet energy gap (ΔE_{ST}) of DI-PF and DI-PI

Compounds	λ_{ex} (nm)	λ_{em} (nm)	Stokes shift (nm)	ϵ ($\text{M}^{-1} \text{cm}^{-1}$)	S_1 (eV)	T_1 (eV)	ΔE_{ST} (eV)
DI-PF	279, 373, 394	590	196	48 268	2.39	2.38	0.01
DI-PI	328, 363, 393	519	126	80 783	3.14	2.77	0.37

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

