


 Cite this: *RSC Adv.*, 2025, **15**, 3904

Metal contamination – a global environmental issue: sources, implications & advances in mitigation

Gabrijel Ondrasek, ^{*a} Jonti Shepherd,^a Santosh Rathod,^b Ramesh Dharavath,^c Muhammad Imtiaz Rashid,^d Martin Brtnicky,^e Muhammad Shafiq Shahid, ^f Jelena Horvatinec^a and Zed Rengel^{gh}

Metal contamination (MC) is a growing environmental issue, with metals altering biotic and metabolic pathways and entering the human body through contaminated food, water and inhalation. With continued population growth and industrialisation, MC poses an exacerbating risk to human health and ecosystems. Metal contamination in the environment is expected to continue to increase, requiring effective remediation approaches and harmonised monitoring programmes to significantly reduce the impact on health and the environment. Bio-based methods, such as enhanced phytoextraction and chemical stabilisation, are being used worldwide to remediate contaminated sites. A systematic plant screening of potential metallophytes can identify the most effective candidates for phytoremediation. However, the detection and prediction of MC is complex, non-linear and chaotic, and it frequently overlaps with various other constraints. Rapidly evolving artificial intelligence (AI) algorithms offer promising tools for the detection, growth and activity modelling and management of metallophytes, helping to fill knowledge gaps related to complex metal-environment interactions in different scenarios. By integrating AI with advanced sensor technologies and field-based trials, future research could revolutionize remediation strategies. This interdisciplinary approach holds immense potential in mitigating the detrimental impacts of metal contamination efficiently and sustainably.

Received 25th June 2024

Accepted 28th January 2025

DOI: 10.1039/d4ra04639k

rsc.li/rsc-advances

1. Introduction

The persistent and toxic nature of metals represents a significant threat to human health, the environment, and food production.^{1,2} In general, most toxic metals found in soil are adsorbed onto soil particles (with kinetics being rapid initially, followed by a slower phase), and then are (re)distributed into diverse chemical forms with different mobility, toxicity and bioavailability.² The mobility of metals in soils, and their transfer through other environmental niches, including

potential entry into the food chain, is significantly influenced by their chemical speciation.^{3,4} The distribution of metals in soil is controlled by a multitude of biogeochemical reactions and processes in the pedosphere, such as complex formation in the soil solution, ion exchange, adsorption/desorption, uptake by soil biota and their dissolution/precipitation.^{5,6} It was confirmed that high concentrations of metals in biological systems affect enzymatic processes and cell organelles and their components, including the nucleus, mitochondria, cell membrane, lysosomes, and endoplasmic reticulum, leading to DNA damage, changes in the cell cycle, carcinogenesis, and cell apoptosis.⁷ The high toxicity and carcinogenicity of arsenic (As) and metals such as lead (Pb), cadmium (Cd), mercury (Hg), chromium (Cr) are frequently the result of oxidative stress due to the formation of reactive oxygen species (ROS).⁸ These metal(lloid)s are systemic toxins capable of damaging various organs even at low doses, prompting leading environmental and public health organizations to classify them as highly toxic, carcinogenic, and a significant threat to all living organisms.⁹

The chemical forms of metals in contaminated soils are influenced by many factors, primarily the soil organic matter (SOM) content and pH, and metal interactions with other soil variables.¹⁰ For example, the mobility of metals can be reduced through organic complexation by increasing the SOM in the

^aFaculty of Agriculture, The University of Zagreb, 10000 Zagreb, Croatia. E-mail: gondrasek@agr.hr

^bICAR—Indian Institute of Rice Research, Hyderabad 500030, India

^cDepartment of Computer Science and Engineering, Indian Institute of Technology (ISM), Dhanbad, 826004, Jharkhand, India

^dCenter of Excellence in Environmental Studies, King Abdulaziz University, 22252, Jeddah, Saudi Arabia

^eDepartment of Agrochemistry, Soil Science, Microbiology and Plant Nutrition, Faculty of AgriSciences, Mendel University in Brno, 61300 Brno, Czech Republic

^fDepartment of Plant Sciences, College of Agricultural and Marine Sciences, Sultan Qaboos University, Al-Khoud 123, Muscat, Oman

^gUWA School of Agriculture and Environment, The University of Western Australia, Perth, WA 6009, Australia

^hInstitute for Adriatic Crops and Karst Reclamation, 21000 Split, Croatia



soil. The use of soil bio-based conditioners such as biochar influences soil pH and increases SOM, which expands soil surface area (interface for metal adsorption), soil porosity, microbial activities and plant growth, ultimately leading to relatively rapid remediation of metal-contaminated soils.¹¹ For instance, adsorption and desorption are the primary processes influencing soil metal accumulation, with studies typically indicating stronger accumulation of metal(lloid)s in fine soil particles (clay, silty clay), although some studies have reported greater accumulation in coarse particles under specific conditions.^{10,12} Additionally, it was shown (albeit over a relatively narrow pH range) that the free cationic forms (most mobile and bioavailable) dominate in acidic (pH < 5) conditions, whereas the poorly mobile and bioavailable forms (*e.g.* carbonates, phosphates) or crystalline/amorphous forms (malachite, otavite, smithsonite) dominate in alkaline (pH > 8) environments.¹³ More specifically, a pH increases from 4 to 7 can decrease the amount of the most bioavailable Cd²⁺ by >60% at the expense of less bioavailable organo-complexed forms of Cd (that increased 7-fold). Consequently, the effective pH management of metal-contaminated acidic soils, using the addition of alkaline matrices can significantly reduce metal mobility and bioavailability, thereby mitigating their adverse impacts on (agro) ecosystems (more in Section 8).

Several pedovariables (pH, salinity) may contribute to high Cd transfer between soils and plants, as confirmed in radish,¹⁴ maize¹³ and strawberry¹⁵ due to formation of soluble and more mobile Cd-complexes. Moreover, this concept has practical applications in chemical remediation of metal-contaminated soils. Chloride salts such as CaCl₂ and FeCl₃ have proven to be effective soil-washing agents, reducing metal concentrations in soil and crops by promoting proton release and forming soluble Cd complexes¹² (more in Section 7).

The extent of the toxicity of metals in the soil environment is determined by the chemical forms and the total concentration of the metals. It has been shown that metals of anthropogenic origin that accumulate in soils are more mobile and bioavailable than metals from lithogenic or pedogenic sources. Furthermore, simulation models show that anthropogenic (*vs.* natural) atmospheric emissions generate 3-to-7-fold greater quantities of toxic metals.¹⁶ However, it should be noted that the availability of metals is influenced by numerous abiotic/biotic factors and interactions, such as temperature, adsorption, phase association, sequestration, solubility and complexation kinetics, genotype, plant species, ecotype, *etc.*¹⁷ In metal-polluted areas with more than 300 years of Pb mining and smelting, indoor environmental conditions can vary significantly. For example, attic dust primarily consists of calcium sulfates and metal-containing particles, whereas house dust is mainly composed of carbon-containing particles.¹⁸ Additionally, attic dust in these areas can have 7 times more metal-containing particles and 13 times more metal species of geogenic or anthropogenic origin compared to outdoor snow deposits. Consequently, in such metal-polluted regions, uniform mitigation approaches may prove ineffective on a small scale.

Here, we discuss the most important sources of metal contamination, their pathways within the biosphere, and the

current remediation methods based on the recent scientific advancements. The objective is to disseminate awareness of the sustainable and efficient use of contemporary technologies, materials, and approaches in remediating metal-contaminated soils. This includes the integration of biorenewable technologies aligned with the medium-term green plans and policies. A critical discussion of specific sites facing significant challenges with the existing on-site or off-site remediation methods underscores gaps, limitations, and opportunities for improvement. Notably, we emphasize the need for (i) more stringent regulatory measures or comprehensive risk assessment protocols, and (ii) multidisciplinary approaches to effectively remediate metal contamination.

2. Soil metal contamination: a critical environmental threat

Metals enter the biosphere through a combination of natural and anthropogenic sources and processes (Fig. 1). Natural sources include weathering of parent rocks, volcanic activity, erosion, sediment resuspension, and metal corrosion, whereas agriculture emerges as the most prominent anthropogenic contributor to global metal emissions⁷ (Fig. 1). Since the industrial revolution in the 1760s, pollution of soils has been on the rise due to contamination by metal(lloid) emissions from rapidly expanding industrial sources, such as manufacturing plants, coal burning, petrochemical releases/spills, atmospheric deposition, mining activities, waste disposal, application of wastewater for irrigation, agrochemicals such as pesticides and fertilizers, and soil amendments (Fig. 1). Zinc, Pb, Cd, As, and Cr are frequently found in contaminated sites,¹⁹ with Cu, Hg and Ni also commonly present.²⁰ Based on the emission sources, two groups of metal(lloid)s can be distinguished: (i) the As–Cr–Ni group primarily originates from natural processes/resources, whereas the Pb–Zn–Cu–Cd–Hg group is largely attributed to human activities²¹ (Fig. 1).

The most exploited ores globally are those vital for the construction, manufacturing, technology, and energy sectors, often containing Pb, Zn or Cd as associated metals or impurities, which complicates extraction, separation, purification and environmental management. For instance, in Celje area (Slovenia), the 100 years anthropogenic emissions from the former Zn smelting plant facility, with an estimated amount of >1700 t of Zn (~0.3% of total Zn production) and >9 t of Cd, resulted in a heavily contaminated area with maximum concentrations of Zn (up to 5.6% w/w in attic dust and 0.85% w/w in the soil) and Cd (456 mg kg⁻¹ in attic dust and 59 mg kg⁻¹ in soil).²² Similarly, in the area around Kosovska Mitrovica (Kosovo) in the former mining area covering 302 km², the maximum concentrations of heavy metals in the topsoil layer were also many times higher (*e.g.* in mg kg⁻¹; Pb 35 000; Zn 12 000; Cu 1,600, Cd 47)²³ than in uncontaminated soils.⁹ In some heavily polluted areas of China, the chronic daily intake of metals by residents near Zn–Pb mining sites has been documented to surpass the safe reference dose by as much as 15-fold.²⁴






Fig. 1 Schematic representation of the most important natural (rectangles bordered by solid black lines) and anthropogenic (rectangles with dashed black lines) sources of metal contamination, with metal transfer pathways (ovals) in the environment as well as into the human food chain, and on-site and off-site approaches to remediate metal-contaminated soils.

With increasing industrialisation, urbanisation, energy consumption and intensive agricultural production, especially in developing countries, metal pollution has accelerated worldwide, posing an increasing challenge to soil quality, food security and human health.²⁵ The presence of Cd in critical concentrations in soil is associated with various harmful structural, physiological and chemical changes in plants. There are approaches to mitigate the toxic effects of Cd in plants, such as applying (in)organic amendments to reduce Cd mobility or using plants that can accumulate Cd from the soil without translocating it to edible parts.¹⁴

Zinc fertilization has been shown to decrease Cd uptake and oxidative stress while increasing the net photosynthetic rate,²⁶ making it a potential candidate for mitigating the toxic effects of Cd. However, the competition between Zn and Cd uptake due to their similar chemical properties can be affected by the concentration of Zn, because high concentrations of Zn can become toxic to plants by increasing the formation of ROS and reducing growth, respiration, and photosynthesis.²⁷ Therefore, further research is necessary to understand the role of Zn in phytoaccumulation of Cd, considering important factors (such as plant species, genotype, metal concentrations, and duration of exposure) to effectively mitigate Cd toxicity.¹⁴

In soil solution, a small proportion of Pb is phytoavailable because most of Pb forms various complexes with soil components.²⁸ Although it is not a phytonutrient, Pb is taken up *via* the

apoplastic pathway or Ca-permeable channels from the rhizosphere.²⁹ The Pb dynamics in the soil and uptake by plants are influenced by soil pH, ion speciation, soil particle size, root surface area, and cation exchange capacity.³⁰ When it enters the plant, it accumulates mainly in the root cells because it is blocked by Caspary strip in the root endodermis.³¹ Furthermore, it is retained by negative charges in the walls of the root cells.³² Damage to plant tissues and negative impacts on morphological, physiological and biochemical functions are the main problems resulting from the excessive Pb phytoaccumulation.²⁸ The Pb accumulation induces phytotoxicity by altering the permeability of the plasma membrane. This alteration is attributed to the interaction between Pb and various active enzyme groups, particularly phosphates, which play a crucial role in plant metabolism.³³ Lipid peroxidation and DNA damage due to excessive generation of ROS and inhibition of ATP production have been modelled in Pb toxicity.³⁴ Lead damaged chlorophyll production, transpiration, protein content, seed germination, root elongation, and seedling development. Lead toxicity negatively affects plant growth by inhibiting Calvin cycle enzymes, leading to a deficiency of CO₂ resulting in stomatal closure, reducing the uptake of essential macro- and micro-nutrients such as Mg and Fe, and hindering the electron transport system.³⁵ However, there are adaptive mechanisms in plants involving a number of components that reduce the uptake of Pb into cells through cross-functional actions and



provide resistance to Pb toxicity.³⁶ In particular, Pb is sequestered in vacuoles through the formation of complexes with phytochelates, glutathione and amino acids.³⁷ The activation of various antioxidants as a secondary type of defence mechanism serves to combat the Pb-induced increase in ROS production.³⁸

As a result of modern activities such as mining, the use of agrochemicals and waste management, contamination with arsenic (As) is becoming a growing problem. Arsenic is found throughout the earth crust, and is a highly toxic metalloid. In water and soil matrices, it occurs mainly in forms such as arsenite (AsIII) and arsenate (AsV), which are more toxic than organic As species.²

After drinking water, the consumption of rice is the second most common way of As exposure.² Arsenic contamination of rice agroecosystems is one of the greatest threats to safe and sustainable rice production. Due to the phytotoxicity of As, grain quality changes and yield decreases.³⁹ Plants take up AsV mainly *via* phosphate transporters, whereas AsIII is taken up *via* aquaporins. Therefore, lower phytoaccumulation of As in rice grains and higher food safety can be achieved by genetically modifying rice varieties with altered expression and/or activity of specific transporters (the concept is explained in more detail in the following sections). The quality of water resources used for irrigation of rice agroecosystems has a significant impact on crop safety and consumption, as rice cultivation generally requires water. A study by ref. 2 has shown that more than 300 million people are affected by As contamination of groundwater sources and that, in Bangladesh alone, chronic As exposure is responsible for an annual death of up to 43 000 people.

A very common component of volcanic rocks and dust matrices is Cr (Fig. 1), which is used in leather tanning, the metal and alloy industry, ceramics and glass manufacturing.⁴⁰ Chromium poses a significant threat to the environment by contaminating soil and water. Once Cr enters the food chain, it can pose a serious risk to human health. The oxidation states in which it occurs range from 0 to +6, with Cr(III) and Cr(V) being the most stable and toxic to humans, animals and plants. The bioaccumulation of Cr(III) in living organisms and accumulation in the environment is exacerbated by the occurrence of (micro)plastic pollution, as polyethylene terephthalate and polystyrene serve as a vector for the transfer of pollutants to the aquatic environment.⁴¹ Due to its high toxicity, mutagenicity, genotoxicity and carcinogenicity⁴² the remediation of Cr is an important and urgent area of research in environmental science and engineering.

3. Agrochemicals as a source of metal contamination

Agriculture contributes significantly to soil metal inputs because many metal(lloid)s are effective agrochemicals.⁴³ Thus, in order to ensure optimal phytonutrition, soil pH balance, SOM content, and effective pest and weed management, the application of agrochemicals in intensive conventional farming systems must be carefully managed and monitored. The long-term annual application of different agrochemicals (fertilizers,

soil amendments, plant protection agents, growth regulators) has been identified as a significant source of soil metal inputs/contamination^{44,45} (Fig. 1 and Table 1), even following the recommended dosage according to the specific agro-ecological conditions and the current national legislations.⁴⁶ For instance,⁴⁶ suggest for some of the agroecosystems in South Brazil that [excluding Fe and Mn due to their high soil background concentrations], the soil accumulation of metals through the application of agrochemicals increased in the following order: Hg > Pb > Co > Cd > As > Cr > Ni > Cu > Zn. Considering some related studies, the average annual input of metals through agrochemical applications ranges (in g ha⁻¹): 0.03–0.71 for Hg, 0.8–12.8 for Pb, 0.6–3.7 for Co, 0.9–2.4 for Cd, 1–6.8 for As, 5.6–28 for Cr, 3.6–23.4 for Ni, 8–122 for Cu and 40–230 for Zn.^{46,53,54} Recently²¹ found that even at sites with the same current crop and agrochemical applications (herbicides and fertilizers), metal accumulation patterns differed, suggesting that the total amount and type of chemicals applied over time plays a significant role in determining Fe, Zn, Mn, Cu and Al content. The study also highlights the potential impact of crop rotation on soil metal accumulation, relating such outcome with well-known phytoremediation capacities of many crops⁵⁵ (Fig. 1).

The regulations regarding the use of different agrochemicals vary significantly among countries. The maximum permitted limits of metal(lloid)s in fertilizers differ by several orders of magnitude for the same metal(lloid), *e.g.*, in Brazil, the limit for As is 10 mg kg⁻¹, but in Canada it is 775 mg kg⁻¹; for Cd, Brazil allows up to 20 mg kg⁻¹, whereas Canada permits 207 mg kg⁻¹; for Pb, the limit is 100 mg kg⁻¹ in Brazil compared to 5169 mg kg⁻¹ in Canada.⁴⁶ According to the same authors, differences among maximum permitted limits of metal(lloid)s in the growing (in)organic substrates are even higher. This inconsistency likely reflects the varying metal concentrations in raw materials (ores, rocks), limiting national exploitation options.⁴³ In addition,⁴⁵ many countries have imposed restrictions or prohibitions on a number of common pesticides that are used intensively in agri-/forest sectors, and are the source of significant concentrations of metals (Fig. 2 and Table 1). Pesticides are extensively used in conventional agriculture as an effective and economical approach to ensure stable crop yields, thus ensuring food security (Fig. 2 and ref. 58). Global cropland areas have expanded by 6% over the period from 1990 (1.48 billion ha) to 2022 (1.57 billion ha) (Fig. 2). Global agricultural pesticide use reached 3.63 Mt of active ingredients in 2022, with herbicides accounting for 55%, insecticides for 22%, fungicides and bactericides for 22%, and other categories for 3.6% of that amount (Fig. 2). This marked a 4% increase compared to 2021, a 13% rise over the past decade, and a doubling since 1990 (Fig. 2 and ref. 59). Between 1990 and 2022, the intensity of pesticide use grew at varying rates: application per unit cropland area surged by 94%, use per unit of agricultural production value rose by 5%, and use per capita increased by 35%.⁵⁶ However, pesticide use in Europe declined by 5% since 1990, with a 7% reduction in the last decade, largely attributed to stricter regulations under the European Common Agricultural Policy⁶⁰ which enforces rigorous pesticide monitoring programs



Table 1 Metal concentrations in major agrochemicals, growth substrates, and livestock feed additives in nano-forms (based on ref. 46–52)

Type	mg kg ⁻¹											
	As	Ba	Cd	Pb	Co	Cu	Cr	Fe	Mn	Hg	Ni	Zn
Herbicides												
Median	<3.9		<0.2	<2.7	<0.2	1.71	0.9	43.1	2.02	0.08	0.55	1.8
Minimum	<3.9		<0.2	<2.7	<0.2	1.32	0.75	8.29	1.45	<0.05	<0.2	0.87
Maximum	<3.9		<0.2	<2.7	<0.2	1.85	1.68	53	2.08	0.08	0.55	4.54
Insecticides												
Median	7.4		<0.2	3.7	0.91	3.86	2.66	31.1	5.93	0.11	3.46	6.88
Minimum	<3.9		<0.2	<2.7	<0.1	1.02	1.19	1.71	1.49	<0.025	<0.1	3.06
Maximum	7.4		<0.2	5.8	10.9	242	42	592	436	0.21	10.8	44.5
Fungicides												
Median	<3.9		2.95	22.3	77.1	10.3	11.2	417	11.3	0.17	12.2	5.86
Minimum	<3.9		<0.2	9.2	<0.2	7.4	5.47	188	1	<0.05	<0.2	1.52
Maximum	<3.9		3.2	367	159	307	50.5	529	92.6	0.73	84.1	11.3
Anti-sprouting agents												
Median	<3.9		<0.2	<2.7	<0.2	3.34	1.49	27.7	2.01	<0.05	0.95	5.67
Minimum	<3.9		<0.2	<2.7	<0.2	2.35	1.33	18.5	1.47	<0.05	<0.2	2.43
Maximum	<3.9		<0.2	<2.7	<0.2	3.9	1.52	80.1	98.6	<0.05	0.95	5.95
Phosphatic fertilizers												
Median			13	13		26	60			22	236	
Minimum	2	200	0.1	7	1	1	66		40	0.01	7	50
Maximum	1200		170	225	12	300	600		2000	1.2	38	1450
Nitrogen fertilizers												
Median			0.9	1.9		2	3.4			6	5	
Minimum	1		0.05	2	5	1	3			0.03	7	1
Maximum	120		8.5	1450	12	15	19			3	38	42
Lime fertilizers												
Median			0.2	8.2		5.6	6.5			6.3	22	
Minimum	0.1	120	0.04	20	0.4	2	10		40	0.05	10	10
Maximum	24	250	0.1	1250	3	125	15		1200		20	450
Manures												
Minimum	3	270	0.3		0.3	2	5.25			0.09	15	
Maximum	150		0.8		24	60	55			26	250	
Growing substrates												
Median	10.7		<0.2	<2.7	15.7	13	11.4	4.3	159	0.2	3.1	35.2
Minimum	<3.9		<0.2	<2.7	<0.22	10.9	5.8	1.4	70.3	<0.05	2.2	29.6
Maximum	11.2		<0.2	12.3	31.7	76.6	544	6.1	220	0.4	517	69.1
Ameliorative impact of nano-mineral additives on livestock growth performance or productivity												
Se					Cu					Zn		
0.1–1.2 ^a					50 ^a					100–200 ^a		
										500–3000 ^b		

^a Broiler chickens. ^b Piglets.

and control. In contrast, the Americas have been the leading pesticide consumer since the mid-1990s, recording a 210% increase in usage between 1990 and 2022, with a notable 31% rise only in the last decade⁵⁶ (Fig. 2).

While pesticides containing Hg, As, Cu and Pb, once widely used, have now been banned in many countries, the presence

of other metal(loid)s as impurities in these products may still pose a significant environmental risk, depending on the management practices adopted and the specific products used.⁴⁶ An example of this is Bordeaux mixture, a CuSO₄-based pesticide that has been banned in most EU countries and the UK due to excessive Cu concentrations in the environment.¹⁶






Fig. 2 Global distribution of pesticide use on croplands globally from 1990 to 2022 (based on data from ref. 56 and 57)

Application of foliar spraying with a Bordeaux mixture for plant protection¹⁷ would result in >64% of vineyards older than 40 years in NW Croatia having received between 80 and 200 kg Cu per ha, with most of the applied Cu being in the surface soil layer where decomposition of biomass occurred (Fig. 1).

The activities in the transport sector and forestry are still, or were until recently (e.g. tetraethyl lead), responsible for a significant emission of metals that could enter water bodies and other ecological environments through (sub)surface runoff and/or leaching (Fig. 1). In the UK, approximately 5% of currently authorised insecticides and fungicides are based on compounds containing metals such as Pb, Zn, Cu, Hg and Mn.⁶¹

4. Bio-based amendments as a source of metal contamination

Various types of biosolids, including compost, animal manure, and municipal sewage sludge, are persistent sources of metal contaminants such as Cd, Zn, Pb, Hg, Cr, Ni, Cu, Mo, Se, Tl, and Sb.⁶¹ Unlike fertilizers and pesticides, with targeted and direct applications, biosolids continually contribute to metal contamination despite careful consideration of their use⁶¹ (Fig. 3). Due to the favourable biochemical composition, the application of animal wastes such as cattle, pig, and poultry manure in the form of solids or slurry is encouraged on croplands, pastures and urban public areas. Indeed, the increased concentration of metals in most fertilisers and wastewater from livestock farms comes from feed additives (growth promoters) added to animal feed for their regulatory, structural and catalytic role⁶³ in maintaining animal health and productivity.⁶⁴ To

prevent excessive growth of pathogenic microorganisms and the occurrence of diarrhoea and to have a positive effect on production parameters (yield and feed conversion), it is a common practice to enrich the feed with Zn and Cu in excess of the nutritional requirements, which, especially after the ban on antibiotics as growth promoters,⁶⁵ has proven successful in practice, particularly in pigs.⁶⁶ In addition, mineral additives are widely used in animal feeds to supply essential nutrients like Cu, Zn, Fe, Cr, Mn, and Co, improving livestock growth and performance;⁴⁷ however, the low purity of these additives often introduces non-essential and toxic metal(lloid)s (Cd, Hg, As, Pb) into the food chain.²⁴ The risk to the (agricultural) environment lies in the increased excretion of faeces from the livestock and the accumulation of metals in the manures.⁶³ For these reasons, the use of some metal-based additives in livestock feed is limited (e.g. for Zn to 150 mg kg⁻¹ and for Cu to 170 mg kg⁻¹ for pigs up to 12 weeks⁶⁵) but these prescribed doses still exceed the nutritional requirements of pigs (50–100 mg per kg Zn and 3–20 mg per kg Cu).⁶⁷ An additional potential source of (agricultural) pollution comes from the liquid component of manure containing significant amounts of metals being used in fertilisation/fertigation.

From the municipal sector, wastewater treatment plant facilities generate sewage sludges (commonly called biosolids) that contain solid organo-mineral compounds⁶⁸ dependent on the wastewater source and the level of purification treatment.⁶⁹ Organic N and inorganic P and K represent the majority of the total nutrient content in biosolids, with approximately 4.7% total N, 2.3% total P and 0.3% total K (on a dry weight basis),⁷⁰ suggesting this matrix is a valuable source of phyto-nutrients.




Fig. 3 Transformation of bio-based resources into bio-based amendments (reproduced (adapted) with permission from ref. 62 Copyright © 2024, Royal Society of Chemistry).

As of 2025, global biosolids generation (on a wet weight basis) is estimated at approximately 150–200 Mt, reflecting a significant increase from the previous estimate of 100–125 Mt.⁷¹ It is a common practice to utilize biosolids for land applications as a way of beneficial recycling;⁷² however, the legislative framework for the land application of biosolids is not uniform (inter)nationally. At the EU level, the regulation of

heavy metals in sewage sludge for agricultural use has been in place for nearly 40 years, following the adoption of the Council Directive.⁷³ The Directive was established to safeguard biota, agricultural soils, and the broader environment by limiting the potential negative impacts of biosolids application due to elevated concentrations of heavy metals. However, the scope of the directive is limited because it only sets permissible levels

Table 2 Limit values for metals in (A) sewage sludge used in agriculture, (B) agricultural soil treated with sludge, and (C) that can be annually added to agricultural soil treated with sludge⁷³

Metal	A mg per kg dw	B mg per kg dw of soil with a pH 6–7	C kg per ha per year
Cd	20–40	1–3	0.15
Cu	1000–1750	50–140	12
Ni	300–400	30–75	3
Pb	750–1200	50–300	15
Zn	2500–4000	150–300	30
Hg	16–25	1–1.5	0.1

with relatively broad concentration ranges for only a few toxic metals (Table 2). Despite this regulatory effort, the framework does not account for emerging pollutants, variability in soil conditions, or the evolving understanding of long-term ecological impacts. Furthermore, inconsistencies in the national implementation of the directive result in varying levels of environmental protection, both across EU member states and globally.

For example, in the USA more than 50% of approximately 9.6 Mt of dry biosolids produced annually are applied on agricultural land,⁷⁴ and similarly in the EU more than 42% of biosolids are utilized as fertilizers/amendments in agriculture.⁷⁵ In addition, certain European countries (Norway, Sweden, Denmark) have not only prohibited the disposal of biosolids in landfills, but also implemented taxes on landfilling and waste incineration;⁷¹ by contrast, other EU member states like Croatia have strictly forbidden any application of biosolids on agricultural land for food production.⁷⁶ Such restrictions on use of sewage sludge is related to the presence of heavy metals (Table 2), and emergent contaminants (pharmaceuticals, drugs, micro/nano plastics); hence, additional treatments (*e.g.* drying, alkalinisation, composting, blending) are necessary prior to safe application as soil amendments (Fig. 3).

The recent trend of mixing composted sewage sludge with various organic materials such as food waste from urban areas, straw, sawdust, garden and municipal waste, and plant biomass from agriculture and forestry (Fig. 3) has led to the spread of metal contamination (Fig. 1 and ref. 77). Metals from sewage sludge (Cd, Pb, Cr, Zn, Ni, Cu) can be found in high concentrations in mixed composts,⁷⁸ depending on the processes used in the treatment of sewage sludge.⁷⁷ The application of sewage sludge to soil can result in metals leaching through the soil profile and into groundwater. Recent studies, particularly from developed countries, are focusing on methods to safely apply composted sewage sludge to soil without risking groundwater contamination,⁷⁹ with this issue reported in soils treated with sewage sludge in New Zealand.⁷⁷ Factors that influence soil contamination with metals include the feedstock materials used in compost production, soil depth and profile properties, temperature, moisture content, and the surrounding landscape.⁷⁹

5. Wastewaters as a source of metal contamination

The use of various wastewaters is often associated with high risks of metal toxicity⁴² and/or salinity.⁸⁰ Nevertheless, wastewaters can partially or fully meet the water and nutrient requirements of crops.⁸¹ It is estimated that 20–25 million hectares of agricultural land worldwide are irrigated with wastewater.⁷⁹ In Asia and Africa, farmers are generally more focused on maximizing vegetable yield and therefore profits, rather than environmental protection.¹⁶ Studies indicate that agricultural irrigation with wastewater accounts for approximately 50% of all vegetables supplied to urban markets.⁸²

In developed countries, strict regulations result in relatively low concentrations of metals in processed wastewater

compared to untreated wastewater.⁸³ However, an analysis of heavy metal content in wastewater discharged across regions in China during 2011 indicated a total discharge volume of ~66 billion tons, containing substantial quantities of hazardous metals and As, including Pb (155 t), Hg (2.8 t), Cd (36 t), Cr(vi) (106 t), Cr (293 t), and As (146 t).⁸⁴ These findings underscore the severe environmental challenges posed by industrial and municipal wastewater discharges in regions with less efficient and/or stringent wastewater treatment and management systems. Elevated metal concentrations in wastewaters pose a serious environmental challenge, particularly when wastewater is reused for agricultural irrigation. Over time, the application of metal-enriched wastewater can result in the accumulation of these metals in soil and their transfer to crops, thus highlighting potential risks to soil quality, food safety/security, and ecological health. Unfortunately, the primary source of metal accumulation in food¹ is the widespread use of untreated (unfiltered, unsterilized) wastewater for irrigation in (peri)urban and industrial areas adjacent to arable land.⁸⁵ Furthermore, hazardous metals discharged from the pulp and paper industries contaminate large areas of agricultural and freshwater environments, affecting aquatic biota.⁸⁶ In addition, nearly 80% of tanneries are involved in the Cr tanning, emitting up to 3200 tonnes of Cr per year to the environment.⁴²

Activities related to mining have caused the most widespread metal contamination in soil, especially in recent decades (Fig. 1). Since the late 1990s, rapid industrialization and urbanisation have driven significant growth in global metal consumption rates, with annual increases of 6% for Mn, 5% for Al, Cr and Ni, 4% for Zn, and 3% for Cu and steel.⁸⁷ Moreover, it is expected that a transition from fossil-based to renewable energy sources will significantly increase the demand for specific metals because C-neutral energy infrastructures require substantially more raw materials per megawatt of installed capacity compared to traditional fossil fuel-based facilities.^{87,88} Additionally, energy transition, increasing urbanization, and the high demand for ores and minerals in industry suggest that more metals will be mined by the middle of this century than in the entire previous century. Accordingly, transitioning to renewable energy as the sole energy source is expected to require approximately 330 Mt of Cu (a nearly 20-fold increase over current global annual production), 8 Mt of Li (a 190-fold increase), 66 Mt of Ni (a 30-fold increase), and 31 kt Pt (a 15-fold increase).⁸⁸ Mine tailings, consisting of heavier and larger particles that settle at the bottom of flotation cells during mining, are often discharged directly into natural depressions in the landscape or accumulated in tailings dams.¹⁶ The rapid extraction of ores and subsequent smelting processes have led to global soil pollution, posing significant risks to human health and the environment⁸⁹ (Fig. 1).

6. Airborne sources of metal contamination

Most metals emitted into the atmosphere are released as particles in the gas stream.⁹⁰ Fugitive emissions, which consist



of gases or vapours, are released from stacks, ducts, chemical storage facilities, or waste dumps containing soils contaminated with various (frequently non-characterized) substances that cause toxic events, thereby becoming sources of airborne metals⁹⁰ (Fig. 1). During high-temperature processing, some metals and metalloids (such as Pb, Cd, and As⁹¹) can vaporize and condense into fine dust particles if a reducing atmosphere is not maintained.⁹⁰ After certain time, dust particles in the atmosphere settle on land and water, and gaseous metal elements can dissolve on these surfaces, collectively increasing environmental metal pollution.⁸⁴ In general, emissions from stacks are dispersed over large areas by natural air currents but can be removed from the atmosphere by precipitation (rain, snow), thus polluting aquatic and terrestrial ecosystems (Fig. 1). In contrast, fugitive emissions are distributed over smaller areas and are released close to the ground.⁹² Notably, fugitive emissions resulting from incomplete combustion were high, indicating that total emissions from combustion are considerably underestimated if leakage is not accounted for.⁹³ In both types of emissions, different metals are emitted from different sources. The reasons for this are varied and range from what is produced or destroyed to the types of filters that may be used to capture the emissions.⁴

Since the industrial revolution widespread metal pollution has been triggered by fossil fuels, and high concentrations of Pb, Cd and Zn have been measured in plants, soils and waters near smelters.⁹⁴ Extremely high Pb concentrations in soils in urban areas and along major roads were previously associated with the combustion of petrol containing tetraethyl Pb.⁹⁰ However, the use of tetraethyl Pb has steadily decreased over the last 30 years; as a result, a major problem in soils near major roads presently is that they contain high concentrations of Cd and Zn used in the production of tyres and lubricating oils.⁹⁵

The legacy of metal pollution since industrial revolution extends into modern challenges posed by rapid industrial growth and urbanization. For instance, in China, waste gas emissions increased from 2003 (199 trillion m³) to 2010 (519 trillion m³), with an average annual emission of 359 billion m³.⁸⁴ This increase has contributed to a large increase in the release of particulate matter (PM), which is a significant carrier of atmospheric metals. Recent findings reveal that PM10 (≤ 10 μm in diameter), primarily originating from the Earth crust, road traffic, and fuel combustion, poses significant risks in both developed and developing countries, with a more severe impact in Asian nations (notably China and India) compared to Europe and the USA, where levels have declined over the past two decades.⁹⁶ Recent analyses of 118 full-scale industrial plant facilities revealed that the majority (~98%) of particulate matter (PM) had diameters < 2.5 μm , with 79% having diameters below 1 μm ;⁹⁷ it should be borne in mind that PM < 10 μm poses the greatest health problems. According to the same source, annual atmospheric releases of Fe, heavy metals (Cd, Cr, Cu, Ni, Pb, Zn), As and five crystalline metallic compounds (ZnO, PbSO₄, Mn₃O₄, Fe₃O₄, Fe₂O₃) contained in fine PM from these industrial activities are estimated globally to be 51 Mt, 70 Mt, and 78 Mt, respectively.

The persistence of airborne metal pollution highlights the ongoing environmental and health challenges posed by industrial emissions and urbanization, particularly in rapidly developing regions. Mitigating the significant risks associated with airborne fine particulate matter and heavy metals requires urgent attention not only to control emission and achieve sustainable industrial practices but also to refine strategies addressing global climate change. Meteorological conditions, such as storms, wind speed, air temperature, and relative humidity, have been shown to significantly influence the spatiotemporal variability in concentration of various particulate matter sizes, both locally and regionally.⁹⁶

7. Mechanisms of metal uptake and redistribution in plants

One of the main pedovariables driving the solubility and bioavailability of metals in the rhizosphere is pH reaction (Fig. 4).¹⁰⁰ Metals such as Zn, Cd, and Cu are among the most soluble and phytoavailable in the rhizosphere and exhibit relatively low selectivity for phyto-uptake. This trait is not limited to metallophyte plant species (more in next section), but is common to most cultivated plants, allowing these metals to relatively easily overcome numerous rhizosphere-plant barriers (Fig. 4). For instance, by using the high-resolution secondary ion mass spectrometry (nano-SIMS) as one the most advanced *in situ* approaches for metal mapping, it was documented that Cd and Zn, even after short-term exposure (24 hours) to very low equimolar concentration (2.2 μM) can rapidly cross root¹⁰¹ and shoot barriers and reach edible parts of widely consumed vegetable.¹⁴

The consummation of crops produced in metal-contaminated surroundings represents the main route of metals entering food and feed sources (Fig. 1). In general, concentration as well as distribution of most metals decrease following the reach of the upstream transpiration stream: roots > shoots > fruits > seeds¹⁰² (Fig. 3). Plants absorb metals from the rhizosphere through their roots, store a portion in the underground tissues, and translocate the rest to the shoots *via* transpiration stream (xylem sap), followed by redistribution *via* phloem sap and sequestration in different tissue and cell compartments⁹⁹ (Fig. 4).

Some toxic metals resemble essential metals in physical (*e.g.*, ionic radii: Cd²⁺ 97 pm, Ca²⁺ 99 pm, Mn²⁺ 80 pm, Fe²⁺ and Zn²⁺ 74 pm) and chemical properties (*e.g.*, redox activity, Lewis acidity), thereby likely entering the pathways specific to these essential elements.^{80,99} Metal uptake across the plasma membrane of root cells has been shown to occur *via* concentration-dependent mechanisms that exhibit saturable kinetics,^{103,104} with confirmed Cd-Zn; Cd-Mn and Cd-Cu competition^{103,105} or Fe-Cd complementarity.¹⁰⁶ In addition, the mobility and uptake of metals are highly dependent on: (i) the pH reaction of the xylem and phloem sap, and (ii) the presence of (in)organic ligands (*e.g.*, organic acid anions, chlorides, carbonates, sulfates) with a strong potential for creating metallo-complexes^{80,99} (Fig. 4). Metal complexation with ligands can




Fig. 4 Schematic presentation of biogeochemical reactions in the rhizosphere and plant vascular tissues that impacts metals mobility, uptake and deposition in plants. Based on the chemical composition of xylem and phloem sap in wild tobacco (*Nicotiana glauca* Grah.)^{99*}, and castor bean (*Ricinus communis* L.) at 10 μM Cd treatment^{99**}.

enhance metal desorption from the solid interfaces (aluminosilicates, hydroxyoxides, humics) in the rhizosphere, and impact the uptake and long-distance transport of some metals in plants (Fig. 4). For instance, the cationic metal forms may adsorb onto negatively charged cell walls and plasma membrane and can easily precipitate in alkaline media (pH > 8; Fig. 4) as observed in the phloem sap of rice.⁹⁹

Different biomolecules, channel protein (CP) and transport proteins (TP) incorporated in the plant membranes (e.g. Cu-DMA TPs, Zn-Fe permease, ATPases, cation diffusion facilitators, cation exchangers, etc.) embedded in the plant cell plasma membranes play crucial roles in the uptake and redistribution of metals^{107,108} (Fig. 4). For instance, uptake of metals from the rhizosphere is possible by specific transporters (for nutrients)

(Ca-CP; Zn-TP; Fe-TP; Cu-DMA-TP), whereas toxic metals enter by competing for nutrient transporters^{109,110} (Fig. 4).

Metal-binding proteins (e.g. Cu chaperone ATX1-like proteins, glutathione – Gtn, metallothioneins – Mts, phytochelatins – Pht), organic acid anions, amino acids (e.g. histidine, cysteine, glutamine, asparagine), and peptides are essential for binding, sequestering, and detoxifying toxic metals in plant tissues^{42,108,111} (Fig. 4). These activities primarily occur in the cuticle, epidermis, and trichomes, areas where the cellular damage poses a relatively minor risk to plant survival.^{107,112} For instance, in the root cytoplasm metals are likely to be complexed by Gtn, i.e. a precursor of metal-Pht complexation (Fig. 4). Both Gtn and Pht are low-molecular-weight cysteine-based peptides that keep metal cationic species in the cytoplasm at low concentrations.¹¹³ As a Pht-Cd complex, Cd can

cross the tonoplast *via* certain TPs and create Pht-based Cd complexes in the vacuole.^{103,104} In addition, free cytosolic metal forms can be anti-ported into the vacuole¹⁰³ and sequestered with organic acid anions; these metal from the vacuolar pool may be remobilised into the cytosol (*e.g.* by Nramp3 proteins) and chelated with sulfur-containing ligands (Fig. 4).

Due to diverse organo-mineral composition and acidic reaction in the xylem sap, it is likely that metals are complexed (Fig. 4), *e.g.* Fe-citrates and Fe-phytosiderophores in some Poaceae (maize, rice, barley) and non-graminaceous plants (tomato, soybean, castor bean).^{114,115} The examples are Ni-histidine complex in the xylem sap of Ni accumulator *Alyssum lebiacum*¹¹⁶ and Cu complexed with 2'-deoxymugineic acid in the xylem sap of rice.¹⁰⁸ It is shown that metal deposition in some plants is specifically targeted to particular leaf cells, such as trichomes (*i.e.* leaf hair or gland cells derived from a specialized epidermal layer on the leaf or stem surfaces) (Fig. 4). In the phloem sap that is alkaline, the metal forms are present mostly as organo-metallo-complexes, *e.g.* Cu-nicotianamine, Cu-histidine, and other Cu complexes (>3 kDa in size) in the rice phloem sap,¹⁰⁸ although the existence of inorganic complexes should not be disregarded (Fig. 4).

8. Remediation of metal-contaminated soils

Numerous remediation methods are currently available for metal-contaminated environments and can be broadly categorized into on-site and off-site approaches (Fig. 1). These include a variety of physical (*e.g.*, soil washing, excavation, solidification), chemical (*e.g.*, flushing, immobilisation), biological (*e.g.*, phytoremediation, bioremediation), electrical (*e.g.*, electrokinetics), and thermal (*e.g.*, vitrification) processes, tailored to target specific contaminants and site conditions (Fig. 1). We focus on three promising approaches: phytoremediation and the use of bio-based materials such as bioashes and biochars for chemical conditioning of contaminated sites. These methods offer sustainable, cost-effective solutions while minimizing secondary environmental impacts. Phytoremediation leverages the capacity of plants to extract, stabilize, and/or degrade contaminants, making it an eco-friendly and low-cost remediation technique. Similarly, biochars (derived from organic materials through pyrolysis) and bioashes (produced by the oxidation of organic residues) demonstrate significant potential for immobilizing heavy metals and enhancing soil health.

8.1. Phytoremediation of metal-contaminated soils

Plant tolerance to metals is a crucial requirement for metal accumulation and phytoremediation, and it is regulated by various biomolecules (Fig. 4,^{99,107}). The (hyper)accumulating plants (metalophytes) are used to extract or 'excavate' potentially toxic metals from contaminated soils.¹¹⁷ Metalophytes include zinc violet (*Viola calaminaria*), plantain (*Plantago lanceolata*), alpine pennycress (*Thlaspi caerulescens*), *Cochlearia* spp, common bent (*Agrostis capillaris*), spring sandwort

(*Minuartia verna*), sea thrift (*Armeria maritima*)¹¹⁷ and many others. Many members of the Brassicaceae family (90 species, about $\frac{1}{4}$ of the family) tend to hyperaccumulate metals from the soil.

Industrial hemp (*Cannabis sativa*) has a great potential for metal hyperaccumulation, which unfortunately has not been utilised for legal reasons.¹¹⁸ However, hemp grown on metal-contaminated soils offers a wide range of potential biomass uses,¹¹⁸ with the necessary further testing (*e.g.* fibre strength, chemical composition) to mitigate the restrictions on the use of *Cannabis sativa* biomass resulting from remediation practices, but also to ensure compliance with food and safety guidelines.¹¹⁹

In recent years, significant progress has been made in elucidating the mechanisms responsible for metal accumulation and detoxification in plants, including chemical and microbiological components, and in optimising field management practices to maximise the remediation potential by hyperaccumulation^{120,121} reported hyperaccumulation of metals in plant tissues, with concentrations reaching up to 3% by weight, without exhibiting phytotoxic symptoms. More than 500 plant species have been identified as hyperaccumulators of metals and metalloids,¹²² which represents approximately 0.006% of all angiosperms. Notably, about 75% of these hyperaccumulators have the ability to hyperaccumulate Ni.¹²³ The Brassicaceae family is the richest in hyperaccumulators, though hyperaccumulators are found in over 34 different plant families. Within Brassicaceae, both Zn and Cd hyperaccumulators are abundant, particularly in the genus *Noccaea* (*e.g.* *N. caerulescens*, formerly *Thlaspi caerulescens*). Some *Noccaea* species/populations hyperaccumulate Ni in their natural serpentine soil environments and are capable of hyperaccumulating Zn under controlled conditions. Interestingly, hyperaccumulators are restricted to the genera *Noccaea* (Zn and Cd) and *Odontarrhena* (Ni), suggesting a monophyletic origin of hyperaccumulation. The *Alyssum* species that hyperaccumulate Ni do not accumulate Zn.¹²³ These examples raise important questions about the evolution of hyperaccumulation mechanisms in plants.

During the last three decades, various selection factors were hypothesized to have led to evolution of hyperaccumulation,¹²⁴ such as: drought tolerance, allelopathy, increased metal tolerance, *etc.* Among the populations of hyperaccumulating species, due to a large variability in the capacity to tolerate and accumulate metals, there was an opportunity to analyse the underlying genetic determinants of the different variations, which opened a possibility of specifically selecting/breeding/genetically engineering a hyperaccumulating ideotype suitable for the remediation of specific metal-contaminated soils. A systematic study of metalophytes can identify priority candidates for the remediation of metal-contaminated soils and the development of environmentally friendly approaches based on the removal of contaminating metals from the soil (Fig. 1).

8.2. Bioashes and metal-contaminated soils

The oxidation of various biological residues (from forestry, agriculture or household) in different plant facilities (for



electrical or thermal energy production, dryers, *etc.*) produces bioash, a reactive inorganic material characterised by an extremely complex composition and alkalinity ($\text{pH} > 12$), a variety of minerals, phytonutrients and trace elements¹³ (Fig. 5). In contrast to biochar (see the next section), with which it shares the same origin, the carbon content in bioash is low (<1% w/w), but has different ecological effects, is more alkaline due to the specific production conditions and has unique physicochemical properties (Fig. 5). For instance, principal component analysis (PCA) of 37 different types of bioashes

explained 71% of the variation in their composition. The PC1 explained more than 41% of the variation, with high concentrations of pozzolanic oxides (Si, Al, Ti, Fe) being most influential. The PC2 explained almost 20% of the variation, predominantly with alkali oxides (Na, K), and PC3 explained almost 10% of the variation, predominantly with MgO .¹²⁶ One of the key roles of bottom ash type used as soil amendment is in controlling soil biochemistry, due to high Si content in ash matrix, leading to increased metal adsorption capacity and shortening the time to reach equilibrium (ref. 127 and Fig. 5).




Fig. 5 Characterization of bottom (A) and fly (B) bioash by scanning electron microscope and X-ray diffraction, with schematic representation of the most relevant biogeochemical reactions of metals in the presence of bioash matrix (C) (reproduced (adapted) with permission from ref. 100 Copyright© 2024, Elsevier and ref. 125 Copyright© 2019, American Chemical Society).



According to the studies of,¹²⁸ the mobility of metals and their uptake by the corresponding biota is influenced by the addition of alkaline bioash to clay soils, which would change the biogeochemistry of metals in the soil. For example, in soils contaminated with metals, the addition of fly ash (5% w/w) reduced the leaching of Cu and Pb by >91% and >87%, respectively, due to the increased number of chemisorption sites and altered soil pH from 4.1 to 6.8; hence, the uptake of both metals by plants and bacteria in the soil decreased.¹²⁹ The combined application of fly ash from wood and coal with peat was also more effective in terms of chemisorption of Cu and Pb than their separate application. Fly ash was effective in removing various metals from aqueous solutions (Fe > Cu > Zn > Mn)¹³⁰ due to an increase in the pH of the liquid (from 4.2 to 8.0), which shifted the biogeochemistry of the metals towards physical adsorption at the interfaces of the ash and/or chemical deposition.⁵

Recently, two types of bioashes, rice husk ash (RHA) and sugarcane bagasse ash (SBA), were approved as soil amendments to immobilize metals in contaminated soils, reducing the metal toxicity and health risk associated with metals in wheat.¹³¹ Specifically, SBA proved more effective, reducing Cr, Ni, Cu, Zn, and Cd in seeds by 13.5%, 33.8%, 17.6%, 7.8%, and 10.0%, respectively, compared to RHA reductions of 6.8%, 16.9%, 8.8%, 3.9%, and 5.0%, with metals accumulating most in roots and least in seeds. Absorption and dissolution processes could explain some of results of the aforementioned studies under strong pH influence. This is because the proportion of cationic metal forms increases at low pH values, whereas anionic forms dominate at high pH values. The addition of bioash to soil can significantly alter key physico-chemical pedovariables, influencing metal biogeochemistry, including shifts in soil pH, electrical conductivity, bulk density, and water-holding capacity,¹⁰⁰ along with increases in carbon and phytonutrient content.¹³¹ These modifications underscore the potential of bioash to impact soil functionality and nutrient cycling, making it a valuable tool for soil management in contaminated or degraded ecosystems.

8.3. Biochar and metal-contaminated soils

Biochar is an organic material produced by pyrolysis in which the temperature inside the container is gradually raised to 300–700 °C (Fig. 3), using filters to capture and store potential pollutants as well as C.¹³² Biochar is C rich (containing >50% w/w C), and one of the most efficient matrices to convert C into a stable form that can then be incorporated into the soil as a soil conditioner.¹¹⁸ Consequently, as a very stable and porous substance, biochar has gained significant attention due to its potential benefits in reducing the availability and leaching of metals (Fig. 6), and ultimately limiting the accumulation of metals in edible tissues.¹³⁵ For example, the addition of biochar at 5% w/w reduced the availability of Pb, Cd and Zn by around 50%.¹³⁶ In the study by ref. 137 biochar derived from rice straw and bamboo significantly reduced the extractable fraction of metals in the soil as well as the fractions of organic complexes, CaCl_2 - and DTPA. The capacity of biochar to stabilise metals in the soil has been demonstrated in

many studies in the field and under controlled conditions but stabilization has not been considered effective enough to be used as a practical solution to the problem of soil metal contamination. To achieve the desired level of toxic metal stabilisation, options include expanding the range of feedstocks, optimizing the pyrolysis conditions used to produce biochar, and increasing the rate of biochar application.¹³⁸ However, due to the costs associated with biochar production and application, the maximum biochar application rate tested at 20% w/w is still too high to be of practical use.⁷² Other soil remediation approaches, such as phytoremediation,¹³⁹ myco-remediation, or the removal of contaminated soil, may incur costs similar to or higher than biochar production and application, providing alternative strategies for addressing metal contamination.

The widespread use of biochar in agroecosystems is hindered by the fact that high biochar application rates are not practical for farmers or environmental engineers¹⁴⁰ because the application of high amounts of biochar or bioash would increase soil alkalinity/salinity, thus jeopardising plant development and growth.¹⁴¹ Extensive research is currently being conducted to identify and characterise an alternative method that can improve the efficiency of biochar in metal stabilisation at relatively low application rates, thus achieving the remediation of metal-contaminated soils at an acceptable price and without negative effects on plant growth.¹²⁰ To improve the stabilisation of metals in soil, many studies have focused on biochar modification¹⁴² by using organic solvents and acids, iron compounds and hydroxides.¹⁴³ Potentially viable options are *e.g.* increasing cation exchange capacity by increasing the number of functional groups on the surface of biochar, which can significantly improve its effectiveness and would obviate a need for high application rates of biochar to the soil.¹⁴⁴ Optimising the specific surface area of biochar is also crucial. By increasing the specific surface area, the sorption capacity for metals can be significantly improved, which would further increase the efficiency of biochar in removing heavy metals from the environment.¹⁴⁵ Biochar can be modified by adding other materials or chemicals that improve its adsorption properties. For example, impregnating biochar with metal oxides or nanoparticles can increase its capacity to bind pollutants. By combining these approaches, it is possible to develop biochar with several enhanced properties to underpin greater efficiency in remediation of soil and water while requiring lower rates of biochar. These advanced solutions would not only reduce costs and environmental impact, but also contribute to more sustainable resource management and environmental protection.

So far, most of the biochar modifications have relied on using one compound or solution (*e.g.* Fe compounds, HNO_3 , H_2SO_4), with very few studies testing multiple modifiers simultaneously.¹⁴⁶ Hence, a potential of using multiple additives to improve biochar properties, and/or applying biochar with multiple modifiers to improve a range of specific soil properties¹⁴⁷ is yet to be explored fully.

Due to the complexity of different soils combined with the limitations of currently available analytical techniques, it is still difficult to accurately determine the mechanisms by which






Fig. 6 Characterization of wood chips-derived biochar by scanning electron microscope (SEM) and energy-dispersive X-ray spectroscopy (A) and of sewage sludge-derived biochar by SEM,⁶² with solid-state ¹³C CP-MAS NMR (Cross-Polarization Magic Angle Spinning Nuclear Magnetic Resonance) spectra of fresh and aged biochar (reproduced (adapted) with permission from ref. 133, Copyright © 2015 Taylor & Francis), and schematic representation of the most relevant biogeochemical reactions and immobilisation mechanisms of metals/metalloids in the presence of biochar matrix and different pH conditions (D) (reproduced (adapted) with permission from ref. 134 Copyright © 2021, Elsevier).

biochar immobilise/removes metal(lloid)s.¹²⁰ Several relevant mechanisms have been proposed, such as oxidation [e.g. Hg⁰]¹⁴⁸, reduction (e.g. Se⁶⁺),¹⁴⁹ intra-lamellar and/or intra-pore positioning (e.g. Zn²⁺, Cu²⁺),¹⁵⁰ physical adsorption, ion exchange (e.g. Cr³⁺, Cd²⁺),^{151,152} co-precipitation (e.g. Pb²⁺),¹⁵³ electrostatic adsorption and surface complexation (e.g. Zn²⁺, Cd²⁺, Cu²⁺; As³⁺, As⁵⁺) with functional groups containing oxygen and an electron located in a pi shell or a double or triple bond, or in some cases in a conjugated pi shell^{134,154} (Fig. 6).

Dominance of a specific metal-immobilising mechanism and its efficiency are strongly dependent on type and aging of particular biochar,¹⁵⁵ pH conditions (e.g. the ratio of anionic and cationic metal forms), SOM and metals properties (e.g. ionic radius, valence)^{156,157} (Fig. 6). For instance, recently performed meta-analyses revealed that soil pH exerted the greatest influence on metal bioavailability in soils amended with biochar, with soil texture, aging time, biochar pyrolysis temperature, metal species and applied dosage following in

significance.¹⁵⁵ In soil amended with biochar under alkaline pH conditions, enhanced metal immobilisation can be explained by (i) deprotonation of acidic radicals and their preference for anionic metal forms, (ii) the release of Na^+ , K^+ , Cl^- , various phosphate forms and OH^- , promoting the formation of relatively stable inner-sphere metallo-complexes and/or (iii) metal precipitates (with CO_3^{2-} , H_2PO_4^- , HPO_4^{2-} or OH^-)¹⁵⁵ (Fig. 4).

The oxidation of biochar surfaces, expected to produce negatively charged organic functional groups, plays a significant role in determining CEC and pH of biochar, consequently impacting the leaching patterns of metals (Fig. 4). Moreover, compared to fresh biochar, aged biochar contains a higher concentration of carboxyl, aromatic, *O*-alkyl, and alkyl surface functional radicals, thereby enhancing its capacity for retaining metals¹⁵³ and reducing bioavailability of metals in soils amended with biochar¹⁵⁵ (Fig. 6).

Some studies on adding biochar to soils reported only a slight change in soil pH and cation exchange capacity, possibly related to the removal of alkaline components from the tested biochar during the modification process.¹⁵⁸ It can be inferred that ion exchange and co-precipitation of metals (controlled by alkaline groups and pH) may not be the main mechanisms for enhanced stabilisation of metals in contaminated soils.¹⁵⁹ Increasing the number of functional groups ($-\text{SH}$, $-\text{OH}$, $-\text{COOH}$) in biochar matrix (Fig. 6) has been shown to enhance its remediation potential by significantly increasing its effective surface area and total pore volume, making it an important approach for biochar modification.¹⁶⁰ Higher efficiency in metal stabilisation was achieved by modified (compared to unmodified) biochar,¹⁶¹ suggesting perhaps the most important mechanism for Cd stabilisation and remediation of Cd, Pb and Zn pollution in various soils by effectively reducing availability and leaching,¹²⁹ through surface complexation with functional groups.⁷² The harsh conditions in metal-polluted soils are at least partially improved by converting metals into more stable fractions.¹⁶² In addition to physico-chemical changes in metal binding and speciation (Fig. 4), biochar also improves a range of other pedovariables, including enhanced soil microbiomes and enzyme activity,¹⁶³ water retention, nutrient availability,^{118,156} and many others (see ref. 164).

9. Estimation and prediction of metal dynamics in soils using artificial intelligence

Artificial intelligence (AI) is a branch of computer science focused on developing algorithms that replicate human brain functions, including the ability to learn from specific patterns in datasets. Machine learning (ML), a subset of AI, enables computers to learn without explicit programming by using statistical methods that allow machines to improve through iterations.¹⁶⁵ Unlike traditional statistical models, ML algorithms are highly flexible as they do not require assumptions about the data distribution. Instead, they learn from the patterns in the training datasets. Among the most widely used

and adaptable ML models is the artificial neural network (ANN) that effectively handles varying levels of linearity and complexity in datasets.¹⁶⁶ Other popular ML models include support vector machines (SVM), random forests (RF), fuzzy techniques, and k-nearest neighbours (KNN) (Table 3). Given the complex, nonlinear, and chaotic nature of metal-soil-plant interactions, ML models often yield more accurate predictions compared to classical models. However, there is no one-size-fits-all model. Developing an ML model requires selecting the optimal combination of hyperparameters, which significantly influences model accuracy. Although limited literature exists on using ML algorithms for metal identification and prediction, there have been substantial advancements in simulating metal behaviour in soils over the past two decades.

Single- or multi-stage digestion/extraction using a wide range of elemental and isotopic analytical tools is typically needed to determine metal forms and dynamics in soils and the related (microbial, plant) matrices,² which requires time, resources and labour. Artificial intelligence techniques are proving to be important alternatives. They may be superior to standard chemical analyses for solving spatial and temporal dynamics of metals in soil. Furthermore, AI approaches can predict the efficiency of metal immobilization in wide range of scenarios, identifying optimal environmental conditions and a selection of amendment(s) considering their complex and reactive matrices (*e.g.* biochar *vs.* bioash) to maximize metal immobilization (Fig. 4). In addition, the use of AI can (i) explain how the prevailing atmospheric conditions can affect the dispersion of metals from source locations, (ii) suggest crops that may be suitable for cultivation in specific metal-contaminated soils (Fig. 1), (iii) take into account various spatio-temporal hydro-geo-pedogenic processes to predict the range of metal concentration in a given area (Table 3). Recently,¹⁵⁷ have successfully applied three AI techniques to predict metal immobilization in biochar-amended soils, revealing that N content in the biochar (ranging from 0.3% to 25.9%) and its application rate (ranging from 0.5% to 10%) as the most influential factors, with the causal analysis indicating the following hierarchy of empirical categories for metal immobilization efficiency: biochar properties *>* experimental conditions *>* soil properties *>* metal properties. In addition,¹⁶⁸ developed the ANN models to determine the dynamics of metals such as Fe, Mn and Zn as influenced by the concentrations of Ca, K and Mg in soil samples obtained from different altitudes. Recently,¹⁷² proposed and validated algorithms against different AI models (ANN, SVM, RF) for Pb prediction by using 13 input variables (*i.e.* total metal concentrations) in the sediments of two Australian bays, with Zn being the most effective predictor for Pb, followed by Ni and Cu.

The tuning of the hyperparameters in the AI predictive models can be achieved by an advanced AI optimization¹⁷² to make modelling performance more reliable and applicative, which is especially valid under conditions of limited technical and logistical resources, *e.g.* in developing countries. Namely, one of the main advantages of AI algorithms is a possibility of using different big data repositories, which in combination with an appropriate technical support (*e.g.* supercomputers) enables



Table 3 Performance of machine learning (ML) models in metal analysis and prediction^a

Description	Best model	Other models tested	Reference
Estimation of Cd and Pb in polluted soil, Gilan province, Iran	ANN	ANFIS	167
Estimation of Fe, Mn, and Zn in Mount Ida, Turkey	ANN		168
Estimation of Ni, Pb, Cr, Hg, Cd, As, Cu, and Zn in polluted soil, Huanghua City, China	Hybrid LASSO-GA-BPNN	SVR, RF RF, ordinary kriging	169
Estimation of Zn, Cu, Cr, and Pb in topsoil of the Dammam area, Saudi Arabia	ANN and SVR		170
Prediction of Pb and Cd in soils from mining areas	BPNN		171
Prediction of Pb from sediments of two bays in Queensland, Australia	XGBoost	ANN, SVM, RF	172
Analysis and calibration of trace elements (Pb, Cu, Ni, Zn, Co, Cd, As, Sc, Hg, Mn, Cr, Ti, Sb, Sr, V, Ba) in soil of a waste disposal site, Rajbandh, Khulna, Bangladesh	ANFIS	ANN, SVM	173
Multivariate calibration model prediction of Ag concentrations in different soils, Lyon, France	ANN	BPNN	174
Prediction of Cr concentration in subarctic soil, Novy Urengoy, Russia	MLP	GRNN	175
Prediction of Cd, As and Pb in soil of mining area, Jiangsu Province, China	RF	SVM, ELM, PLS	176
Prediction of immobilisation of metals (Cu, Zn, Pb, Cd, Fe, Ni and Mn) in biochar-amended soils from 15 different studies	RF	ANN, SVM	157
Identification of the source and spatial prediction of metals (Zn, Pb, Hg, Ni, Cu, Cr, Cd) and As in peri-urban soil, Hefei City, China	SVM and RF	SVM, ANN	177

^a ANFIS – adaptive neuro-fuzzy inference system; ANN – artificial neural network; BPNN – back-propagation neural network; ELM – extreme learning machine; GA – genetic algorithm; IDW – inverse distance weighting; LASSO – least absolute shrinkage and selection operator; MARS – multivariate adaptive regression spline; PLS – partial least squares; RF – random forest; SVM – support vector machine; SVR – support vector regression; XGBoost – extreme gradient boosting.

fast processing of numerus iterations and generation of relevant scenarios.

Among the various AI models used to predict metals in soils, ANN is the predominant one; however, some newly introduced hybrid machine learning models have been confirmed as superior for metals in specific pedospheres.¹⁷⁸ Consequently, derived AI approaches could assist and significantly accelerate metal detection in the soil as well assessing economic, health and environmental impacts in various conditions, thus improv policies and regulations to diminish metal contamination and secure the sites that may be sources of metals.

10 Conclusions and future perspectives

For restoration of metal-contaminated soils there are initial requirements to characterize the contamination type and source, as well as the spatial range and depth of soil contamination.¹⁷⁹ Contemporary legislation designed for the protection

of public and environmental health is based purely on the chemical characterization within each tested site. Particular attention should be focused on potentially toxic metals that may enter the food chain, water bodies or atmosphere.⁷ Given that soil characterization for legislative and regulatory purposes is done based on total metal concentrations, a substantial improvement to underpin successful remediation strategies is needed *via* determining metal speciation that governs bioavailability and mobility.¹⁸⁰ As a risk-minimization strategy, decision makers seek solutions that can guarantee the removal or *in situ* immobilisation of the metal contaminants in the most cost-effective manner while preserving both public and ecosystem health.¹⁸¹ The impact of military activities, including the manufacture and disposal of weapons, the utilization of ammunition during military exercises, and engagement in conflicts and wars, is frequently underestimated despite their significant and multiple contributions to metal pollution (Fig. 1). These activities include the releases high loads of toxic metals as common constituents of ammunition,¹⁸² coatings,



electronic devices, *etc.* The shells and bullets used in firing ranges can remain in the soil for long periods, resulting in persistent metal pollution.¹⁸³ A recent study by¹⁸⁴ found that military activities have led to the accumulation of Pb, Cd and Hg in soils around military bases. Similarly, a study by¹⁸⁵ found that military activities in Malaysia have resulted in elevated levels of Pb and Cu in soils around firing ranges. The Russian–Ukrainian war is expected to have a significant and long-lasting multipli-cative impact on the levels of metal contamination in various environmental niches, not only in the affected regions, then globally.¹⁸⁶ A massive destruction of infrastructure has already resulted in the release of metals and other pollutants into the environment, hindering efforts to achieve a metal-clean environment, as well as energy and food security.¹⁸⁶ The most recent studies confirm significant increases in metal concentrations (Pb, Cd, Hg, Cu, Zn, Ni, Co, Sn, Mn, Se, Al) in the soils of Ukrainian regions affected by the Russian invasion.^{187,188} This is particularly significant, as Ukraine's chernozems are among the most productive and high-quality soils on Earth, and their restoration and remediation after the war will be crucial.

Due to a greater awareness based on the scientific knowledge, governments are dealing with contaminated soils and their implications for human and ecosystem health, by funding the further development of remediation strategies while simultaneously assessing, mapping and classifying metal contamination within their borders. However, increasing environmental pressures due to various anthropogenic activities (*e.g.* plastic, persistent organics) exacerbate the problems with metal-contaminated soils due to synergistic negative repercussions on environmental health, food safety and security. For example, recent studies warn that addition of micro-plastic¹²⁰ or combined application of humates and chloride salts⁹ to Cd-contaminated rhizosphere markedly change biochemical reactions, metabolites and their pathways, increasing Cd availability and promoting its uptake by crops. Importantly, the interactions among various pedo-variables lead to multicollinearity, making specific and synergistic effects extremely complex to discern. However, due to possibility of fast processing of numerous predictors under different relevant scenarios, the fast-developing algorithms driven by AI appear to be a promising approach to optimising the management of metal-contaminated areas, addressing the numerous knowledge gaps in metal interactions.

Consent for publication

All authors agreed to publication.

Data availability

All data will be made available upon request.

Author contributions

GO conceptualisation, funding and project administration. GO, JS and JH graphical data presentation. GO, JS, SR, RD, MIR, MB, MSS, JH and ZR drafted the manuscript. ZR discussed, revised and amended the text. All authors finalised the review.

Conflicts of interest

The authors declare no competing interest.

Acknowledgements

This study was partially performed within the DOK-2021-02 and IP-2022-10-7906 projects founded by the Croatian Science Foundation.

Notes and references

- 1 M. U. Khan, R. N. Malik and S. Muhammad, Human health risk from Heavy metal via food crops consumption with wastewater irrigation practices in Pakistan, *Chemosphere*, 2013, **93**, 2230–2238.
- 2 J. S. Y. Preetha, M. Arun, N. Vidya, K. Kowsalya, J. Halka and G. Ondrasek, Biotechnology Advances in Bioremediation of Arsenic: A Review, *Molecules*, 2023, **28**, 1474.
- 3 M. Uchimiya, D. Bannon, H. Nakanishi, M. B. McBride, M. A. Williams, T. Yoshihara and C. Speciation, Plant Uptake, and Toxicity of Heavy Metals in Agricultural Soils, *J. Agric. Food Chem.*, 2020, **68**, 12856–12869.
- 4 M. I. Inyang, B. Gao, Y. Yao, Y. Xue, A. Zimmerman, A. Mosa, P. Pullammanappallil, Y. S. Ok and X. Cao, A review of biochar as a low-cost adsorbent for aqueous heavy metal removal, *Crit. Rev. Environ. Sci. Technol.*, 2015, **46**, 406–433, DOI: [10.1080/10643389.2015.1096880](https://doi.org/10.1080/10643389.2015.1096880).
- 5 M. Kumar, N. Sawhney and R. Lal, Chemistry of heavy metals in the environment, *Heavy Metals in the Environment: Impact, Assessment, and Remediation*, 2021, pp. 9–37.
- 6 Y.-J. Du, S.-Y. Liu, Z.-B. Liu, L. Chen, F. Zhang and F. Jin, An Overview of Stabilization/Solidification Technique for Heavy Metals Contaminated Soils, *Advances in Environmental Geotechnics*, 2010, pp. 760–766.
- 7 J. Briffa, E. Sinagra and R. Blundell, Heavy metal pollution in the environment and their toxicological effects on humans, *Helijon*, 6(9), e04691, DOI: [10.1016/j.heliyon.2020.e04691](https://doi.org/10.1016/j.heliyon.2020.e04691).
- 8 Q. Y. Chen, T. DesMarais and M. Costa, Metals and Mechanisms of Carcinogenesis, *Annu. Rev. Pharmacol. Toxicol.*, 2019, **59**, 537.
- 9 G. Ondrasek, I. Jelovica Badovinac, R. Peter, M. Petracović, J. Macan and Z. Rengel, Humates and Chlorides Synergistically Increase Cd Phytoaccumulation in Strawberry Fruits, Heightening Health Risk from Cd in Human Diet, *Exposure Health*, 2022, **14**, 393–410.
- 10 B. Huang, Z. Yuan, D. Li, M. Zheng, X. Nie and Y. Liao, Effects of soil particle size on the adsorption, distribution, and migration behaviors of heavy metal(lloid)s in soil: A review, *RSC Environ. Sci.: Processes Impacts*, 2020, **8**, 1596, DOI: [10.1039/d0em00189a](https://doi.org/10.1039/d0em00189a).
- 11 C. E. Brewer, V. J. Chuang, C. A. Masiello, H. Gonnermann, X. Gao, B. Dugan, L. E. Driver, P. Panzacchi, K. Zygourakis and C. A. Davies, New approaches to measuring biochar density and porosity, *Biomass Bioenergy*, 2014, **66**, 176–185.



12 J. Shi, J. Pang, Q. Liu, Y. Luo, J. Ye, Q. Xu, B. Long, B. Ye and X. Yuan, Simultaneous removal of multiple heavy metals from soil by washing with citric acid and ferric chloride, *RSC Adv.*, 2020, **10**, 7432–7442.

13 G. Ondrasek, M. Zovko, F. Kranjčec, R. Savić, D. Romić and Z. Rengel, Wood biomass fly ash ameliorates acidic, low-nutrient hydromorphic soil & reduces metal accumulation in maize, *J. Clean. Prod.*, 2021, **283**, 124650, DOI: [10.1016/j.jclepro.2020.124650](https://doi.org/10.1016/j.jclepro.2020.124650).

14 G. Ondrasek, P. L. Clode, M. R. Kilburn, P. Guagliardo, D. Romić and Z. Rengel, Zinc and cadmium mapping in the apical shoot and hypocotyl tissues of radish by high-resolution secondary ion mass spectrometry (NanoSIMS) after short-term exposure to metal contamination, *Int. J. Environ. Res. Publ. Health*, 2019, **16**, 3, DOI: [10.3390/ijerph16030373](https://doi.org/10.3390/ijerph16030373).

15 G. Ondrasek, D. Romić, V. Tanaskovik, R. Savić, S. Rathod, J. Horvatinec and Z. Rengel, Humates mitigate Cd uptake in the absence of NaCl salinity, but combined application of humates and NaCl enhances Cd mobility & phytoaccumulation, *Sci. Total Environ.*, 2022, **848**, 157649, DOI: [10.1016/j.scitotenv.2022.157649](https://doi.org/10.1016/j.scitotenv.2022.157649).

16 WHO. Ambient (outdoor) air pollution, 2021, [https://www.who.int/news-room/fact-sheets/detail/ambient-\(outdoor\)-air-quality-and-health](https://www.who.int/news-room/fact-sheets/detail/ambient-(outdoor)-air-quality-and-health), accessed 20 October 2022.

17 G. Ondrasek, H. Bakić Begić, M. Zovko, L. Filipović, C. Meriño-Gergichevich, R. Savić and Z. Rengel, Biogeochemistry of soil organic matter in agroecosystems & environmental implications, *Sci. Total Environ.*, 2019, **658**, 1559–1573.

18 M. Miler and M. Gosar, Assessment of contribution of metal pollution sources to attic and household dust in Pb-polluted area, *Indoor Air*, 2019, **29**, 487–498.

19 R. Lal, Restoring Soil Quality to Mitigate Soil Degradation, *Sustainability*, 2015, **7**, 5875–5895.

20 S. Kumari, A. Mishra, S. Kumari and A. Mishra, Heavy Metal Contamination, *Soil Contamination - Threats and Sustainable Solutions*, DOI: [10.5772/INTECHOPEN.93412](https://doi.org/10.5772/INTECHOPEN.93412).

21 A. Naccarato, A. Tassone, F. Cavaliere, R. Elliani, N. Pirrone, F. Sprovieri, A. Tagarelli and A. Giglio, Agrochemical treatments as a source of heavy metals and rare earth elements in agricultural soils and bioaccumulation in ground beetles, *Sci. Total Environ.*, 2020, **749**, 141438.

22 R. Sajn, Using attic dust and soil for the separation of anthropogenic and geogenic elemental distributions in an old metallurgic area (Celje, Slovenia), *Geochemistry*, 2005, **5**, 59–67.

23 R. Šajn, M. Aliu, T. Stafilov and J. Alijagić, Heavy metal contamination of topsoil around a lead and zinc smelter in Kosovska Mitrovica/Mitrovicë, Kosovo/Kosovë, *J. Geochem. Explor.*, 2013, **134**, 1–16.

24 H. Yu, C. Li, J. Yan, Y. Ma, X. Zhou, W. Yu, H. Kan, Q. Meng, R. Xie and P. Dong, A review on adsorption characteristics and influencing mechanism of heavy metals in farmland soil, *RSC Adv.*, 2023, **6**, 3505–3519.

25 L. Bouida, M. Rafatullah, A. Kerrouche, M. Qutob, A. M. Alosaimi, H. S. Alorfi and M. A. Hussein, A Review on Cadmium and Lead Contamination: Sources, Fate, Mechanism, Health Effects and Remediation Methods, *Water*, 2022, **14**, 21, DOI: [10.3390/w14213432](https://doi.org/10.3390/w14213432).

26 M. Rizwan, S. Ali, A. Hussain, Q. Ali, M. B. Shakoor, M. Ziaur-Rehman, M. Farid and M. Asma, Effect of zinc-lysine on growth, yield and cadmium uptake in wheat (*Triticum aestivum* L.) and health risk assessment, *Chemosphere*, 2017, **187**, 35–42.

27 H. Kaur and N. Garg, Zinc toxicity in plants: a review, *Planta*, 2021, **253**, 129.

28 X. Li, S. Huang and M. B. McBride, Rhizosphere effect on Pb solubility and phytoavailability in Pb-Contaminated soils, *Environ. Pollut.*, 2021, **268**, 115840.

29 B. Pourrut, M. Shahid, F. Douay, C. Dumat and E. Pinelli, Molecular mechanisms involved in lead uptake, toxicity and detoxification in higher plants, *Heavy Met. Stress Plants*, 2013, 121–147.

30 P. Sun, Y. Chen, J. Liu, S. Lu, J. Guo, Z. Zhang and X. Zheng, Quantitative evaluation of the synergistic effect of biochar and plants on immobilization of Pb, *J. Environ. Manage.*, 2022, **316**, 115200.

31 S. Nanda and J. Abraham, Impact of heavy metals on the rhizosphere microflora of *Jatropha multifida* and their effective remediation, *Afr. J. Biotechnol.*, 2013, **10**, 11948–11955.

32 M. O. Asare, J. Száková and P. Tlustoš, The fate of secondary metabolites in plants growing on Cd-, As-, and Pb-contaminated soils—a comprehensive review, *Environ. Sci. Pollut. Res.*, 2022, **30**, 11378–11398.

33 M. Ali and F. S. Nas, The effect of lead on plants in terms of growing and biochemical parameters: a review, *Moj Ecol. Environ. Sci.*, 2018, **3**, 265–268, DOI: [10.15406/mojes.2018.03.00098](https://doi.org/10.15406/mojes.2018.03.00098).

34 I. Herath, P. Kumarathilaka, A. Navaratne, N. Rajakaruna and M. Vithanage, Immobilization and phytotoxicity reduction of heavy metals in serpentine soil using biochar, *J. Soils Sediments*, 2015, **15**, 126–138.

35 J. Sun, Q. Fan, J. Ma, L. Cui, G. Quan, J. Yan, L. Wu, K. Hina, B. Abdul and H. Wang, Effects of biochar on cadmium (Cd) uptake in vegetables and its natural downward movement in saline-alkali soil, *Environ. Pollut. Bioavailability*, 2020, **32**, 36–46.

36 K. Usman, M. H. Abu-Dieyeh, N. Zouari and M. A. Al-Ghouti, Lead (Pb) bioaccumulation and antioxidative responses in *Tetraena qataranse*, *Sci. Rep.*, 2020, **10**, 1–10.

37 A. Haque, C. K. Tang, S. Islam, P. G. Ranjith and H. H. Bui, Biochar Sequestration in Lime-Slag Treated Synthetic Soils: A Green Approach to Ground Improvement, *J. Mater. Civ. Eng.*, 2014, **26**, 06014024.

38 J. Kováčik, G. Rotková, M. Bujdoš, P. Babula, V. Peterková and P. Matúš, Ascorbic acid protects *Coccomyxa subellipsoidea* against metal toxicity through modulation of ROS/NO balance and metal uptake, *J. Hazard. Mater.*, 2017, **339**, 200–207.

39 D. Moulick, S. Samanta, S. Sarkar, A. Mukherjee, B. K. Pattnaik, S. Saha, J. P. Awasthi, S. Bhowmick, D. Ghosh, A. C. Samal, S. Mahanta, M. K. Mazumder,



S. Choudhury, K. Bramhachari, J. K. Biswas and S. C. Santra, Arsenic contamination, impact and mitigation strategies in rice agro-environment: An inclusive insight, *Sci. Total Environ.*, 2021, **800**, 149477.

40 A. Sharma, D. Kapoor, J. Wang, B. Shahzad, V. Kumar, A. S. Bali, S. Jasrotia, B. Zheng, H. Yuan and D. Yan, Chromium bioaccumulation and its impacts on plants: An overview, *Plants*, 2020, **13**(9), 100, DOI: [10.3390/plants9010100](https://doi.org/10.3390/plants9010100).

41 S. Liu, J. H. Huang, W. Zhang, L. X. Shi, K. X. Yi, H. B. Yu, C. Y. Zhang, S. Z. Li and J. N. Li, Microplastics as a vehicle of heavy metals in aquatic environments: A review of adsorption factors, mechanisms, and biological effects, *J. Environ. Manage.*, 2022, **302**, 113995.

42 P. Sharma, A. K. Pandey, A. Udayan and S. Kumar, Role of microbial community and metal-binding proteins in phytoremediation of heavy metals from industrial wastewater, *Bioresour. Technol.*, 2021, **326**, 124750.

43 A. A. Zunaidi, L. H. Lim and F. Metali, Assessments of Heavy Metals in Commercially Available Fertilizers in Brunei Darussalam, *Agric. Res.*, 2021, **10**, 234–242.

44 I. Dror, B. Yaron and B. Berkowitz, Microchemical contaminants as forming agents of anthropogenic soils, *Ambio*, 2017, **46**, 109–120.

45 M. Gosar, R. Šajn and T. Teršič, Distribution pattern of mercury in the Slovenian soil: Geochemical mapping based on multiple geochemical datasets, *J. Geochem. Explor.*, 2016, **167**, 38–48.

46 H. J. O. Zoffoli, N. M. B. Do Amaral-Sobrinho, E. Zonta, M. V. Luisi, G. Marcon and A. Tolón-Becerra, Inputs of heavy metals due to agrochemical use in tobacco fields in Brazil's Southern Region, *Environ. Monit. Assess.*, 2013, **185**, 2423–2437.

47 I. Michalak, K. Dziergowska, M. Alagawany, M. R. Farag, N. A. El-Shall, H. S. Tuli, T. Bin Emran and K. Dhama, The effect of metal-containing nanoparticles on the health, performance and production of livestock animals and poultry, *Vet. Q.*, 2022, **42**, 68–94.

48 N. Mroczek-Sosnowska, E. Sawosz, K. P. Vadalasetty, M. Łukasiewicz, J. Niemiec, M. Wierzbicki, M. Kutwin, S. Jaworski and A. Chwalibog, Nanoparticles of copper stimulate angiogenesis at systemic and molecular level, *Int. J. Mol. Sci.*, 2015, **16**, 4838–4849.

49 Z. Ouyang, P. Ren, D. Zheng, L. Huang, T. Wei, C. Yang, X. Kong, Y. Yin, S. He and Q. He, Hydrothermal synthesis of a new porous zinc oxide and its antimicrobial evaluation in weanling piglets, *Livest. Sci.*, 2021, **248**, 104499.

50 C. H. Hu, Y. L. Li, L. Xiong, H. M. Zhang, J. Song and M. S. Xia, Comparative effects of nano elemental selenium and sodium selenite on selenium retention in broiler chickens, *Anim. Feed Sci. Technol.*, 2012, **177**, 204–210.

51 F. Mohammadi, F. Ahmadi and A. M. Andi, Effect of zinc oxide nanoparticles on carcass parameters, relative weight of digestive and lymphoid organs of broiler fed wet diet during the starter period, *Int. J. Biosci.*, 2015, **6**, 389–394.

52 B. J. Alloway, Sources of Heavy Metals and Metalloids in Soils, in *Heavy Metals in Soils. Trace Metals and Metalloids in Soils and Their Bioavailability*, ed. Alloway, B. J., Springer, Dordrecht, The Netherlands, 2013, pp. 11–50.

53 F. A. Nicholson, S. R. Smith, B. J. Alloway, C. Carlton-Smith and B. J. Chambers, An inventory of heavy metals inputs to agricultural soils in England and Wales, *Sci. Total Environ.*, 2003, **311**, 205–219.

54 L. Luo, Y. Ma, S. Zhang, D. Wei and Y. G. Zhu, An inventory of trace element inputs to agricultural soils in China, *J. Environ. Manage.*, 2009, **90**, 2524–2530.

55 Y. Yang, C. Xiao, F. Wang, L. Peng, Q. Zeng and S. Luo, Assessment of the potential for phytoremediation of cadmium polluted soils by various crop rotation patterns based on the annual input and output fluxes, *J. Hazard. Mater.*, 2022, **423**, 127183.

56 FAO, 2024, <https://www.fao.org/faostat/en/#data/RP>, accessed 13 January 2025.

57 A. Alengebawy, S. T. Abdelkhalek, S. R. Qureshi and M. Q. Wang, Heavy Metals and Pesticides Toxicity in Agricultural Soil and Plants: Ecological Risks and Human Health Implications, *Toxics*, 2021, **9**, 42.

58 G. Ondrasek, Z. Rengel, R. G. Taylor and L. Zhuo, Centers for optimizing water management in agroecosystems & global food security, *Front. Sustain. Food Syst.*, 2024, **8**, 1398454.

59 A. Sharma, V. Kumar, B. Shahzad, M. Tanveer, G. P. S. Sidhu, N. Handa, S. K. Kohli, P. Yadav, A. S. Bali, R. D. Parihar, O. I. Dar, K. Singh, S. Jasrotia, P. Bakshi, M. Ramakrishnan, S. Kumar, R. Bhardwaj and A. K. Thukral, Worldwide pesticide usage and its impacts on ecosystem, *SN Appl. Sci.*, 2019, **1**, 1–16.

60 A. P. Alexoaei, R. G. Robu, V. Cojanu, D. Miron and A. M. Holobiuc, Good practices in reforming the common agricultural policy to support the European green deal – a perspective on the consumption of pesticides and fertilizers, *Amfiteatru Econ.*, 2022, **24**, 525–545.

61 H. Wu, C. Lai, G. Zeng, J. Liang, J. Chen, J. Xu, J. Dai, X. Li, J. Liu, M. Chen, L. Lu, L. Hu and J. Wan, The interactions of composting and biochar and their implications for soil amendment and pollution remediation: a review, *Crit. Rev. Biotechnol.*, 2016, **37**, 754–764.

62 G. Ondrasek, C. Meriño-Gergichevich, C. Manterola-Barroso, A. Seguel Fuentealba, S. M. Romero, R. Savić, S. S. Cholin and J. Horvatinec, Bio-based resources: systemic & circular solutions for (agro)environmental services, *RSC Adv.*, 2024, **14**, 23466.

63 M. Hejna, D. Gottardo, A. Baldi, V. Dell'Orto, F. Cheli, M. Zaninelli and L. Rossi, Review: Nutritional ecology of heavy metals, *Animal*, 2018, **12**, 2156–2170.

64 A. T. Adekanmi and A. T. Adekanmi, Health Hazards of Toxic and Essential Heavy Metals from the Poultry Waste on Human and Aquatic Organisms, *Veterinary Medicine and Science*, IntechOpen, 2022, DOI: [10.5772/intechopen.99549](https://doi.org/10.5772/intechopen.99549).

65 Regulation (EC) 1831/2003, <https://eur-lex.europa.eu/eli/reg/2003/1831/oj>, accessed 21 February 2024.

66 C. L. Walk, P. Wilcock and E. Magowan, Evaluation of the effects of pharmacological zinc oxide and phosphorus source on weaned piglet growth performance, plasma minerals and mineral digestibility, *Animal*, 2015, **9**, 1145–1152.

67 National Research Council (NRC), *A Framework for K-12 Science Education Practices, Crosscutting Concepts, and Core Ideas*, National Academies Press. - References - Scientific Research Publishing, Washington DC The, 2012, <https://www.scirp.org/reference/referencespapers?referenceid=1302130>, accessed 22 February 2024.

68 L. Lamastra, N. A. Suciu and M. Trevisan, Sewage sludge for sustainable agriculture: Contaminants' contents and potential use as fertilizer, *Chem. Biol. Technol. Agric.*, 2018, **5**, 1–6.

69 S. Marchuk, S. Tait, P. Sinha, P. Harris, D. L. Antille and B. K. McCabe, Biosolids-derived fertilisers: A review of challenges and opportunities, *Sci. Total Environ.*, 2023, **875**, 162555.

70 A. Gianico, C. M. Braguglia, A. Gallipoli, D. Montecchio and G. Mininni, Land application of biosolids in europe: Possibilities, con-straints and future perspectives, *Water*, 2021, **13**, 1, DOI: [10.3390/w13010103](https://doi.org/10.3390/w13010103).

71 Q. Le and G. W. Price, A review of the influence of heat drying, alkaline treatment, and composting on biosolids characteristics and their impacts on nitrogen dynamics in biosolids-amended soils, *Waste Manage.*, 2024, **176**, 85–104.

72 B. Wang, B. Gao and J. Fang, Recent advances in engineered biochar productions and applications, *Crit. Rev. Environ. Sci. Technol.*, 2018, **47**, 2158–2207.

73 Council Directive 86/278/EEC, <https://eur-lex.europa.eu/eli/dir/1986/278/oj/eng>, accessed 13 January 2025.

74 M. K. Hossain, V. Strezov Vladimir, K. Y. Chan, A. Ziolkowski and P. F. Nelson, Influence of pyrolysis temperature on production and nutrient properties of wastewater sludge biochar, *J. Environ. Manage.*, 2011, **92**, 223–228.

75 M. C. Collivignarelli, A. Abbà, A. Frattarola, M. C. Miino, S. Padovani, I. Katsoyiannis and V. Torretta, Legislation for the reuse of biosolids on agricultural land in Europe: Overview, *Sustainability*, 2019, **11**, 6015, DOI: [10.3390/su11216015](https://doi.org/10.3390/su11216015).

76 NN 71/2019 (26.7.2019.), Pravilnik o zaštiti poljoprivrednog zemljišta od onečišćenja - Zakon.hr, <https://www.zakon.hr/cms.htm?id=39921>, accessed 19 June 2024.

77 S. Ye, G. Zeng, H. Wu, J. Liang, C. Zhang, J. Dai, W. Xiong, B. Song, S. Wu and J. Yu, The effects of activated biochar addition on remediation efficiency of co-composting with contaminated wetland soil, *Resour. Conserv. Recycl.*, 2019, **140**, 278–285.

78 S. Schimmelpfennig and B. Glaser, One Step Forward toward Characterization: Some Important Material Properties to Distinguish Biochars, *J. Environ. Qual.*, 2012, **41**, 1001–1013.

79 A. F. Salleh and H. Hettiarachchi, *Safe Use of Wastewater in Agriculture (SUWA): Reflecting on the Journey of Supporting the UN Sustainable Development Goals since*, UNU-FLORES, 2015, DOI: [10.53325/FCVG7887](https://doi.org/10.53325/FCVG7887).

80 G. Ondrasek, Water scarcity and water stress in agriculture, *Physiological Mechanisms and Adaptation Strategies in Plants under Changing Environment*, 2014, vol. 1, pp. 75–96.

81 C. E. Nedelciu, K. V. Ragnarsdóttir and I. Stjernquist, From waste to resource: A systems dynamics and stakeholder analysis of phosphorus recycling from municipal wastewater in Europe, *Ambio*, 2019, **48**, 741–751.

82 C. B. Chisanga, Use of wastewater for irrigation in vegetable growing in the Kaufe Lagoon areas and along Ngwerere river, *Report PMA*, 2004, vol. 24, DOI: [10.13140/RG.2.1.2641.8646](https://doi.org/10.13140/RG.2.1.2641.8646).

83 J. Rogowska, M. Cieszynska-Semenowicz, W. Ratajczyk and L. Wolska, Micropollutants in treated wastewater, *Ambio*, 2020, **49**, 487–503.

84 H. Hu, Q. Jin and P. Kavan, A study of heavy metal pollution in China: Current status, pollution-control policies and countermeasures, *Sustainability*, 2014, **6**, 5820–5838.

85 R. Savic, G. Ondrasek, R. Zemunac, M. Bubalo Kovacic, F. Kranjec, V. Nikolic Jokanovic and A. Bezdan, Longitudinal distribution of macronutrients in the sediments of Jegricka watercourse in Vojvodina, Serbia, *Sci. Total Environ.*, 2021, **754**, 142138, DOI: [10.1016/j.scitotenv.2020.142138](https://doi.org/10.1016/j.scitotenv.2020.142138).

86 R. Chandra, P. Sharma, S. Yadav and S. Tripathi, Biodegradation of endocrine-disrupting chemicals and residual organic pollutants of pulp and paper mill effluent by biostimulation, *Front. Microbiol.*, 2021, **9**, 960, DOI: [10.3389/fmicb.2018.00960](https://doi.org/10.3389/fmicb.2018.00960).

87 O. Vidal, F. Rostom, C. François and G. Giraud, Global trends in metal consumption and supply: The raw material-energy nexus, *Elements*, 2017, **13**, 319–324.

88 A. García-Olivares, J. Ballabrera-Poy, E. García-Ladona and A. Turiel, A global renewable mix with proven technologies and common materials, *Energy Policy*, 2012, **41**, 561–574.

89 W. P. Covre, S. J. Ramos, W. V. da S. Pereira, E. S. de Souza, G. C. Martins, O. M. M. Teixeira, C. B. do Amarante, Y. N. Dias and A. R. Fernandes, Impact of copper mining wastes in the Amazon: Properties and risks to environment and human health, *J. Hazard. Mater.*, 2022, **421**, 126688.

90 F. Li and F. Li, Heavy Metal in Urban Soil: Health Risk Assessment and Management, *Heavy Metals*, InTech, 2018, DOI: [10.5772/intechopen.73256](https://doi.org/10.5772/intechopen.73256).

91 L. Zhou, H. Guo, X. Wang, M. Chu, G. Zhang and L. Zhang, Effect of occurrence mode of heavy metal elements in a low rank coal on volatility during pyrolysis, *Int. J. Coal Sci. Technol.*, 2019, **6**, 235–246.

92 O. D. Pedrayes, D. G. Lema, R. Usamentiaga and D. F. García, Detection and localization of fugitive emissions in industrial plants using surveillance cameras, *Comput. Ind.*, 2022, **142**, 103731.

93 Y. Chen, W. Du, S. Zhuo, W. Liu, Y. Liu, G. Shen, S. Wu, J. Li, B. Zhou, G. Wang, E. Y. Zeng, H. Cheng, W. Liu and S. Tao, Stack and fugitive emissions of major air pollutants from



typical brick kilns in China, *Environ. Pollut.*, 2017, **224**, 421–429.

94 X. Xu, X. Cao and L. Zhao, Comparison of rice husk- and dairy manure-derived biochars for simultaneously removing heavy metals from aqueous solutions: Role of mineral components in biochars, *Chemosphere*, 2013, **92**, 955–961.

95 M. Skorbiłowicz, E. Skorbiłowicz and W. Rogowska, Heavy Metal Concentrations in Roadside Soils on the Białystok-Budzisko Route in Northeastern Poland, *Minerals*, 2021, **11**, 1290.

96 A. Mukherjee and M. Agrawal, World air particulate matter: sources, distribution and health effects, *Environ. Chem. Lett.*, 2017, **15**, 283–309, DOI: [10.1007/s10311-017-0611-9](https://doi.org/10.1007/s10311-017-0611-9).

97 Q. Yang, G. Liu, J. Falandysz, L. Yang, C. Zhao, C. Chen, Y. Sun, M. Zheng and G. Jiang, Atmospheric emissions of particulate matter-bound heavy metals from industrial sources, *Sci. Total Environ.*, 2024, **947**, 174467.

98 P. J. Hocking, The Composition of Phloem Exudate and Xylem Sap from Tree Tobacco (*Nicotiana glauca* Grah.), *Ann. Bot.*, 1980, **45**, 633–643.

99 K. Hazama, S. Nagata, T. Fujimori, S. Yanagisawa and T. Yoneyama, Concentrations of metals and potential metal-binding compounds and speciation of Cd, Zn and Cu in phloem and xylem saps from castor bean plants (*Ricinus communis*) treated with four levels of cadmium, *Physiol. Plant.*, 2015, **154**, 243–255.

100 J. Horvatinec, J. Buczny and G. Ondrasek, Fly ash application impacts master physicochemical pedovariables: A multilevel meta-analysis, *J. Environ. Manage.*, 2024, **368**, 122066, DOI: [10.1016/j.jenvman.2024.122066](https://doi.org/10.1016/j.jenvman.2024.122066).

101 G. Ondrasek, Z. Rengel, P. L. Clode, M. R. Kilburn, P. Guagliardo and D. Romic, Zinc and cadmium mapping by NanoSIMS within the root apex after short-term exposure to metal contamination, *Ecotoxicol. Environ. Saf.*, 2019, **171**, 571–578.

102 V. Ördög, Plant physiology, <https://www.slideshare.net/slideshow/plant-physiology-by-vince-ordog/75183793>, accessed 13 January 2025.

103 D. E. Salt, M. Blaylock, N. P. B. A. Kumar, V. Dushenkov, B. D. Ensley, I. Chet and I. Raskin, Phytoremediation: A Novel Strategy for the Removal of Toxic Metals from the Environment Using Plants, *Bio/Technology*, 1995, **13**, 468–474.

104 J. J. Hart, R. M. Welch, W. A. Norvell, L. A. Sullivan and L. V. Kochian, Characterization of Cadmium Binding, Uptake, and Translocation in Intact Seedlings of Bread and Durum Wheat Cultivars, *Plant Physiol.*, 1998, **116**, 1413–1420.

105 J. J. Hart, R. M. Welch, W. A. Norvell and L. V. Kochian, Transport interactions between cadmium and zinc in roots of bread and durum wheat seedlings, *Physiol. Plant.*, 2002, **116**, 73–78.

106 H. Nakanishi, I. Ogawa, Y. Ishimaru, S. Mori and N. K. Nishizawa, Iron deficiency enhances cadmium uptake and translocation mediated by the Fe²⁺ transporters OsIRT1 and OsIRT2 in rice, *Soil Sci. Plant Nutr.*, 2006, **52**, 464–469.

107 G. Saxena, D. Purchase, S. I. Mulla, G. D. Saratale and R. N. Bharagava, Phytoremediation of Heavy Metal-Contaminated Sites: Eco-environmental Concerns, Field Studies, Sustainability Issues, and Future Prospects, *Rev. Environ. Contam. Toxicol.*, 2019, **249**, 71–131.

108 Y. Ando, S. Nagata, S. Yanagisawa and T. Yoneyama, Copper in xylem and phloem saps from rice (*Oryza sativa*): the effect of moderate copper concentrations in the growth medium on the accumulation of five essential metals and a speciation analysis of copper-containing compounds, *Funct. Plant Biol.*, 2012, **40**, 89–100.

109 S. Clemens, Toxic metal accumulation, responses to exposure and mechanisms of tolerance in plants, *Biochimie*, 2006, **88**, 1707–1719.

110 K. Yoneyama, A. A. Awad, X. Xie, K. Yoneyama and Y. Takeuchi, Strigolactones as Germination Stimulants for Root Parasitic Plants, *Plant Cell Physiol.*, 2010, **51**, 1095–1103.

111 A. Álvarez-Fernández, P. Díaz-Benito, A. Abadía, A. F. López Millán and J. Abadía, Metal species involved in long distance metal transport in plants, *Front. Plant Sci.*, 2014, **5**, 82117.

112 N. Sarwar, M. Imran, M. R. Shaheen, W. Ishaque, M. A. Kamran, A. Matloob, A. Rehim and S. Hussain, Phytoremediation strategies for soils contaminated with heavy metals: Modifications and future perspectives, *Chemosphere*, 2017, **171**, 710–721.

113 J. Sreeshma and C. Sudandiradoss, Identification of metal binding motifs in protein frameworks to develop novel remediation strategies for Hg²⁺ and Cr(VI), *BioMetals*, 2021, **34**, 621–638.

114 R. Rellán-Álvarez, J. Giner-Martínez-Sierra, J. Orduna, I. Orera, J. Á. Rodríguez-Castrilln, J. I. García-Alonso, J. Abadía and A. Álvarez-Fernández, Identification of a Tri-Iron(III), Tri-Citrate Complex in the Xylem Sap of Iron-Deficient Tomato Resupplied with Iron: New Insights into Plant Iron Long-Distance Transport, *Plant Cell Physiol.*, 2010, **51**, 91–102.

115 T. Ariga, K. Hazama, S. Yanagisawa and T. Yoneyama, Chemical forms of iron in xylem sap from graminaceous and non-graminaceous plants, *Soil Sci. Plant Nutr.*, 2014, **60**, 460–469.

116 U. Krämer, J. D. Cotter-Howells, J. M. Charnock, A. J. M. Baker and J. A. C. Smith, Free histidine as a metal chelator in plants that accumulate nickel, *Nature*, 1996, **379**, 635–638.

117 O. Akhatar, H. Sharma, I. Zoomi, K. L. Chaudhary and M. Kumar, *Metallophytes, Heavy Metals in Plants Physiological to Molecular Approach*, 2022, pp. 354–368.

118 L. L. Kong, W. T. Liu and Q. X. Zhou, Biochar: An effective amendment for remediating contaminated soil, *Rev. Environ. Contam. Toxicol.*, 2014, **228**, 83–99.

119 A. Mead, Legal and regulatory issues governing cannabis and cannabis-derived products in the United States,



Front. Plant Sci., 2019, 14, 697, DOI: [10.3389/fpls.2019.00697](https://doi.org/10.3389/fpls.2019.00697).

120 F. Wang, X. Wang and N. Song, Polyethylene microplastics increase cadmium uptake in lettuce (*Lactuca sativa* L.) by altering the soil microenvironment, *Sci. Total Environ.*, 2021, **784**, 147133.

121 L. Skuza, I. Szućko-Kociuba, E. Filip and I. Bożek, Natural Molecular Mechanisms of Plant Hyperaccumulation and Hypertolerance towards Heavy Metals, *Int. J. Mol. Sci.*, 2022, **23**, 9335, DOI: [10.3390/ijms23169335](https://doi.org/10.3390/ijms23169335).

122 Z. Ghori, H. Iftikhar, M. F. Bhatti, N. Um-Minullah, I. Sharma, A. G. Kazi and P. Ahmad, Phytoextraction: The Use of Plants to Remove Heavy Metals from Soil, *Plant Metal Interaction: Emerging Remediation Techniques*, 2016, pp. 385–409.

123 M. J. M. Christenhusz and J. W. Byng, The number of known plants species in the world and its annual increase, *Phytotaxa*, 2016, **261**, 201–217.

124 J. J. Cappa and E. A. H. Pilon-Smits, Evolutionary aspects of elemental hyperaccumulation, *Planta*, 2014, **239**, 267–275.

125 N. Allen, C. Dai, Y. Hu, J. D. Kubicki and N. Kabengi, Adsorption Study of Al 3+, Cr 3+, and Mn 2+ onto Quartz and Corundum using Flow Microcalorimetry, Quartz Crystal Microbalance and Density Functional Theory, *ACS Earth Space Chem.*, 2019, **3**(3), DOI: [10.1021/acsearthspacechem.8b00148](https://doi.org/10.1021/acsearthspacechem.8b00148).

126 G. Ondrasek, M. Bubalo Kovačić, I. Carević, N. Štirmer, S. Stipičević, N. Udiković-Kolić, V. Filipović, D. Romic and Z. Rengel, Bioashes and their potential for reuse to sustain ecosystem services and underpin circular economy, *Renew. Sustain. Energy Rev.*, 2021, **151**, 111540.

127 J. Mathur, P. Goswami, A. Gupta, S. Srivastava, T. Minkina, S. Shan and V. D. Rajput, Nanomaterials for Water Remediation: An Efficient Strategy for Prevention of Metal(lod) Hazard, *Water*, 2022, **14**, 24, DOI: [10.3390/w14243998](https://doi.org/10.3390/w14243998).

128 A. Kicińska, R. Pomykała and M. Izquierdo-Diaz, Changes in soil pH and mobility of heavy metals in contaminated soils, *Eur. J. Soil Sci.*, 2021, **73**, e13203, DOI: [10.1111/EJSS.13203](https://doi.org/10.1111/EJSS.13203).

129 D. Kanakaraju, D. Norfadzila, H. Nori and R. Wahi, Heavy metal leachability in fly ash remediated soil, *J. Sustainability Sci. Manage.*, 2019, **14**, 37–48.

130 V. Asokbunyarat, P. N. L. Lens and A. P. Annachhatre, *Permeable Reactive Barriers for Heavy Metal Removal*, 2017, pp. 65–100.

131 U. Kumar, P. K. Singh, I. Kumar and R. K. Sharma, Heavy metal accumulation, yield and health risk assessment of wheat crop grown in contaminated soil amended with bioash for sustainable agriculture, *J. Food Compos. Anal.*, 2025, **139**, 107140.

132 A. Tomczyk, Z. Sokołowska and P. Boguta, Biochar physicochemical properties: pyrolysis temperature and feedstock kind effects, *Rev. Environ. Sci. Biotechnol.*, 2020, **19**, 191–215.

133 D. Laird and N. Rogovska, Biochar effects on nutrient leaching, in *Biochar for Environmental Management: Science, Technology, and Implementation*, Earthscan, 2nd edn, 2015, pp. 521–524.

134 Y. Li, H. Yu, L. Liu and H. Yu, Application of co-pyrolysis biochar for the adsorption and immobilization of heavy metals in contaminated environmental substrates, *J. Hazard. Mater.*, 2021, **420**, 126655.

135 Y. Li, J. Shao, X. Wang, Y. Deng, H. Yang and H. Chen, Characterization of Modified Biochars Derived from Bamboo Pyrolysis and Their Utilization for Target Component (Furfural) Adsorption, *Energy Fuel.*, 2014, **28**, 5119–5127.

136 A. P. Puga, C. A. Abreu, L. C. A. Melo and L. Beesley, Biochar application to a contaminated soil reduces the availability and plant uptake of zinc, lead and cadmium, *J. Environ. Manage.*, 2015, **159**, 86–93.

137 K. Lu, X. Yang, G. Gielen, N. Bolan, Y. S. Ok, N. K. Niazi, S. Xu, G. Yuan, X. Chen, X. Zhang, D. Liu, Z. Song, X. Liu and H. Wang, Effect of bamboo and rice straw biochars on the mobility and redistribution of heavy metals (Cd, Cu, Pb and Zn) in contaminated soil, *J. Environ. Manage.*, 2017, **186**, 285–292.

138 Y. Sun, B. Gao, Y. Yao, J. Fang, M. Zhang, Y. Zhou, H. Chen and L. Yang, Effects of feedstock type, production method, and pyrolysis temperature on biochar and hydrochar properties, *Chem. Eng. J.*, 2014, **240**, 574–578.

139 V. Shah and A. Daverey, Phytoremediation: A multidisciplinary approach to clean up heavy metal contaminated soil, *Environ. Technol. Innov.*, 2020, **18**, 100774.

140 M. Ahmad, A. U. Rajapaksha, J. E. Lim, M. Zhang, N. Bolan, D. Mohan, M. Vithanage, S. S. Lee and Y. S. Ok, Biochar as a sorbent for contaminant management in soil and water: A review, *Chemosphere*, 2014, **99**, 19–33.

141 S. Khan, C. Chao, M. Waqas, H. Peter, H. Arp and Y.-G. Zhu, Sewage Sludge Biochar Influence upon Rice (*Oryza sativa* L) Yield, Metal Bioaccumulation and Greenhouse Gas Emissions from Acidic Paddy Soil, *Environ. Sci. Technol.*, 2013, **47**(15), DOI: [10.1021/es400554x](https://doi.org/10.1021/es400554x).

142 A. U. Rajapaksha, S. S. Chen, D. C. W. Tsang, M. Zhang, M. Vithanage, S. Mandal, B. Gao, N. S. Bolan and Y. S. Ok, Engineered/designer biochar for contaminant removal/immobilization from soil and water: Potential and implication of biochar modification, *Chemosphere*, 2016, **148**, 276–291.

143 A. S. Giwa, J. M. Ndungutse, Y. Li, A. Mabi, X. Liu, M. Vakili, A. G. Memon, L. Ai, Z. Chenfeng and M. Sheng, Modification of biochar with Fe3O4 and humic acid-salt for removal of mercury from aqueous solutions: a review, *Environ. Pollut. Bioavailability*, 2022, **34**, 352–364, DOI: [10.1080/26395940.2022.2115402](https://doi.org/10.1080/26395940.2022.2115402).

144 H. M. Anawar, F. Akter, Z. M. Solaiman and V. Strezov, Biochar: An Emerging Panacea for Remediation of Soil Contaminants from Mining, Industry and Sewage Wastes, *Pedosphere*, 2015, **25**, 654–665.

145 R. Zhou, M. Zhang and S. Shao, Optimization of target biochar for the adsorption of target heavy metal ion, *Sci. Rep.*, 2022, **12**, 1–17.



146 B. Sajjadi, T. Zubatiuk, D. Leszczynska, J. Leszczynski and W. Y. Chen, Chemical activation of biochar for energy and environmental applications: a comprehensive review, *Rev. Chem. Eng.*, 2018, **35**, 777–815.

147 G. Ondrasek, F. Kranjčec, L. Filipović, V. Filipović, M. Bubalo Kovačić, I. J. Badovinac, R. Peter, M. Petravić, J. Macan and Z. Rengel, Biomass bottom ash & dolomite similarly ameliorate an acidic low-nutrient soil, improve phytonutrition and growth, but increase Cd accumulation in radish, *Sci. Total Environ.*, 2021, **753**, 141902.

148 Y. Xu, F. Deng, Q. Pang, S. He, Y. Xu, G. Luo and H. Yao, Development of waste-derived sorbents from biomass and brominated flame retarded plastic for elemental mercury removal from coal-fired flue gas, *Chem. Eng. J.*, 2018, **350**, 911–919.

149 X. Wei, X. Li, L. Tang, J. Yu, J. Deng, T. Luo, J. Liang, X. Chen and Y. Zhou, Exploring the role of Fe species from biochar-iron composites in the removal and long-term immobilization of SeO_4^{2-} against competing oxyanions, *J. Hazard. Mater.*, 2021, **418**, 126311.

150 J. Deng, X. Li, X. Wei, Y. Liu, J. Liang, B. Song, Y. Shao and W. Huang, Hybrid silicate-hydrochar composite for highly efficient removal of heavy metal and antibiotics: Co-adsorption and mechanism, *Chem. Eng. J.*, 2020, **387**, 124097.

151 A. Ngambia, J. Ifthikar, I. I. Shahib, A. Jawad, A. Shahzad, M. Zhao, J. Wang, Z. Chen and Z. Chen, Adsorptive purification of heavy metal contaminated wastewater with sewage sludge derived carbon-supported $\text{Mg}^{(II)}$ composite, *Sci. Total Environ.*, 2019, **691**, 306–321.

152 D. Dias, M. Bernardo, I. Matos, I. Fonseca, F. Pinto and N. Lapa, Activation of co-pyrolysis chars from rice wastes to improve the removal of Cr^{3+} from simulated and real industrial wastewaters, *J. Clean. Prod.*, 2020, **267**, 121993.

153 R. Gao, Q. Fu, H. Hu, Q. Wang, Y. Liu and J. Zhu, Highly-effective removal of Pb by co-pyrolysis biochar derived from rape straw and orthophosphate, *J. Hazard. Mater.*, 2019, **371**, 191–197.

154 N. Bolan, A. Kunhikrishnan, R. Thangarajan, J. Kumpiene, J. Park, T. Makino, M. B. Kirkham and K. Scheckel, Remediation of heavy metal(lloid)s contaminated soils – To mobilize or to immobilize?, *J. Hazard. Mater.*, 2014, **266**, 141–166.

155 C. Yuan, B. Gao, Y. Peng, X. Gao, B. Fan and Q. Chen, A meta-analysis of heavy metal bioavailability response to biochar aging: Importance of soil and biochar properties, *Sci. Total Environ.*, 2021, **756**, 144058.

156 M. I. Rashid, G. A. Shah, M. Sadiq, N. ul Amin, A. M. Ali, G. Ondrasek and K. Shahzad, Nanobiochar and Copper Oxide Nanoparticles Mixture Synergistically Increases Soil Nutrient Availability and Improves Wheat Production, *Plants*, 2023, **12**, 1312.

157 K. N. Palansooriya, J. Li, P. D. Dissanayake, M. Suvarna, L. Li, X. Yuan, B. Sarkar, D. C. W. Tsang, J. Rinklebe, X. Wang and Y. S. Ok, Prediction of Soil Heavy Metal Immobilization by Biochar Using Machine Learning, *Environ. Sci. Technol.*, 2022, **56**, 4187–4198.

158 L. Zheng, Y. Lin, J. Wu and M. Zheng, The Associations of Tobacco Use, Sexually Transmitted Infections, HPV Vaccination, and Screening With the Global Incidence of Cervical Cancer: An Ecological Time Series Modelling Study, *SSRN Electron. J.*, 2023, **45**, DOI: [10.2139/ssrn.4003222](https://doi.org/10.2139/ssrn.4003222).

159 W. Cui, X. Li, W. Duan, M. Xie and X. Dong, Heavy metal stabilization remediation in polluted soils with stabilizing materials: a review, *Environ. Geochem. Health*, 2023, 1–37.

160 Y. Zhao, M. Zhao, L. Qi, C. Zhao, W. Zhang, Y. Zhang, W. Wen and J. Yuan, Coupled Relationship between Soil Physicochemical Properties and Plant Diversity in the Process of Vegetation Restoration, *Forests*, 2022, **13**(5), 645.

161 Y. Wang, K. Zheng, Z. Jiao, W. Zhan, S. Ge, S. Ning, S. Fang and X. Ruan, Simultaneous Removal of Cu^{2+} , Cd^{2+} and Pb^{2+} by Modified Wheat Straw Biochar from Aqueous Solution: Preparation, Characterization and Adsorption Mechanism, *Toxics*, 2022, **10**(6), 316.

162 A. Fayiga and O. Nwoke, Metal (lloid)s in Farmland Soils and Strategies to Reduce Bioavailability, *Open J. Environ. Biol.*, 2017, **2**, 009–024.

163 F. U. Haider, J. A. Coulter, L. Cai, S. Hussain, S. A. Cheema, J. Wu and R. Zhang, An overview on biochar production, its implications, and mechanisms of biochar-induced amelioration of soil and plant characteristics, *Pedosphere*, 2022, **32**, 107–130.

164 J. O'Laughlin and K. McElligott, *Biochar for Environmental Management: Science and Technology*, ed. J. Lehmann and S. M. Joseph, Earthscan, London UK, 2009, vol. 11, pp. 535–536.

165 P. Ongsulee, *Artificial Intelligence, Machine Learning and Deep Learning*, IEEE, DOI: [10.1109/ICTKE.2017.8259629](https://doi.org/10.1109/ICTKE.2017.8259629).

166 R. Song, D. Li, A. Chang, M. Tao, Y. Qin, A. A. Keller and S. Suh, Accelerating the pace of ecotoxicological assessment using artificial intelligence, *Ambio*, 2022, **51**, 598–610.

167 A. Bazoobandi, S. Emamgholizadeh and H. Ghorbani, Estimating the amount of cadmium and lead in the polluted soil using artificial intelligence models, *Eur. J. Environ. Civ. Eng.*, 2022, **26**, 933–951.

168 M. Sari, T. Cosgun, I. E. Yalcin, M. Taner and I. I. Ozyigit, Deciding Heavy Metal Levels in Soil Based on Various Ecological Information through Artificial Intelligence Modeling, *Appl. Artif. Intell.*, 2014, **36**, 2014189, DOI: [10.1080/08839514.2021.2014189](https://doi.org/10.1080/08839514.2021.2014189).

169 S. Shi, M. Hou, Z. Gu, C. Jiang, W. Zhang, M. Hou, C. Li and Z. Xi, Estimation of Heavy Metal Content in Soil Based on Machine Learning Models, *Land*, 2022, **11**, 1037.

170 B. Tawabini, M. A. Yassin, M. Benaafi, J. A. Adetoro, A. Al-Shaibani and S. I. Abba, Spatiotemporal Variability Assessment of Trace Metals Based on Subsurface Water Quality Impact Integrated with Artificial Intelligence-Based Modeling, *Sustainability*, 2022, **14**, 2192.

171 N. Luo, Methods for controlling heavy metals in environmental soils based on artificial neural networks, *Sci. Rep.*, 2024, **14**, 1–13.



172 S. K. Bhagat, T. M. Tung and Z. M. Yaseen, Heavy metal contamination prediction using ensemble model: Case study of Bay sedimentation, Australia, *J. Hazard. Mater.*, 2021, **403**, 123492.

173 S. Kumar Sarkar and I. M. Rafizul, *Analysis of Heavy Metal Concentration in Soils of a Waste Disposal Site in Khulna Using Artificial Intelligence Technique*, ICCESD, 2020.

174 C. Sun, Y. Tian, L. Gao, Y. Niu, T. Zhang, H. Li, Y. Zhang, Z. Yue, N. Delepine-Gilon and J. Yu, Machine Learning Allows Calibration Models to Predict Trace Element Concentration in Soils with Generalized LIBS Spectra, *Sci. Rep.*, 2019, **9**, 1–18.

175 D. A. Tarasov, A. G. Buevich, A. P. Sergeev, A. V. Shichkin and E. M. Baglaeva, Topsoil pollution forecasting using artificial neural networks on the example of the abnormally distributed heavy metal at Russian subarctic, *AIP Conf. Proc.*, 2017, **1836**, 020024.

176 W. Ma, K. Tan and P. Du, *Predicting Soil Heavy Metal Based on Random Forest Model*, IGARSS, 2016, pp. 4331–4334.

177 H. Zhang, S. Yin, Y. Chen, S. Shao, J. Wu, M. Fan, F. Chen and C. Gao, Machine learning-based source identification and spatial prediction of heavy metals in soil in a rapid urbanization area, eastern China, *J. Clean. Prod.*, 2020, **273**, 122858.

178 Z. M. Yaseen, An insight into machine learning models era in simulating soil, water bodies and adsorption heavy metals: Review, challenges and solutions, *Chemosphere*, 2021, **277**, 130126.

179 D. Houben, L. Evrard and P. Sonnet, Mobility, bioavailability and pH-dependent leaching of cadmium, zinc and lead in a contaminated soil amended with biochar, *Chemosphere*, 2013, **92**, 1450–1457.

180 Q. Li, Y. Wang, Y. Li, L. Li, M. Tang, W. Hu, L. Chen and S. Ai, Speciation of heavy metals in soils and their immobilization at micro-scale interfaces among diverse soil components, *Sci. Total Environ.*, 2022, **825**, 153862.

181 B. C. Ball, P. R. Hargreaves and C. A. Watson, A framework of connections between soil and people can help improve sustainability of the food system and soil functions, *Ambio*, 2018, **47**, 269–283, DOI: [10.1007/s13280-017-0965-z](https://doi.org/10.1007/s13280-017-0965-z).

182 A. J. Barker, J. L. Clausen, T. A. Douglas, A. J. Bednar, C. S. Griggs and W. A. Martin, Environmental impact of metals resulting from military training activities: A review, *Chemosphere*, 2021, **265**, 129110.

183 A. O. Fayiga and U. K. Saha, Soil pollution at outdoor shooting ranges: Health effects, bioavailability and best management practices, *Environ. Pollut.*, 2016, **216**, 135–144, DOI: [10.1016/j.envpol.2016.05.062](https://doi.org/10.1016/j.envpol.2016.05.062).

184 S. Shukla, G. Mbingwa, S. Khanna, J. Dalal, D. Sankhyan, A. Malik and N. Badhwar, Environment and health hazards due to military metal pollution: A review, *Environ. Nanotechnol. Monit. Manag.*, 2023, **20**, 100857.

185 R. Othman, N. H. Mohd Latiff, Z. M. Baharuddin, K. S. H. Y. Hashim and L. H. Lukman Hakim Mahamod, Closed landfill heavy metal contamination distribution profiles at different soil depths and radiiuses, *Appl. Ecol. Environ. Res.*, 2019, **17**, 8059–8067.

186 O. Shumilova, K. Tockner, A. Sukhodolov, V. Khilchevskyi, L. De Meester, S. Stepanenko, G. Trokhymenko, J. A. Hernández-Agüero and P. Gleick, Impact of the Russia–Ukraine armed conflict on water resources and water infrastructure, *Nat Sustainability*, 2023, **6**, 578–586.

187 O. Shebanina, I. Kormyshkin, A. Bondar, I. Bulba and B. Ualkhanov, Ukrainian soil pollution before and after the Russian invasion, *Int. J. Environ. Stud.*, 2024, **81**, 208–215.

188 A. Yakymchuk, Assessment of soil contamination of Ukraine with heavy metals during the war, *Scientific Papers of Silesian University of Technology Organization and Management Series*, 2024, vol. 196, DOI: [10.29119/1641-3466.2024.196.45](https://doi.org/10.29119/1641-3466.2024.196.45).

