RSC Advances



PAPER

View Article Online
View Journal | View Issue



Cite this: RSC Adv., 2025, 15, 12042

Received 23rd January 2025 Accepted 27th March 2025

DOI: 10.1039/d5ra00100e

rsc.li/rsc-advances

Iodine-mediated thio-arylation under electrochemical conditions†

Jiajia Yu,^a Tong Li,^b Qi Sun*^a and Zhiyong Wang (1)**ab

An efficient iodine-catalyzed thio-arylation reaction of aniline was developed under electrochemical conditions. A variety of diaryl sulfide compounds can be obtained under metal-free and chemical oxidant-free conditions. The reaction features a broad substrate scope, regulation of product distribution, and scalable preparation.

Sulfur-containing compounds play an important role in medicines,¹ natural products,² and functional materials.³ For instance, vortioxetine is a substance that regulates neurotransmitters and acts on multiple receptors, effectively alleviating depression-related symptoms. Dapson antibiotics have antibacterial and anti-inflammatory effects (Fig. 1).⁴ Owing to the excellent biological activity of sulfur-containing compounds, numerous methods have been proposed for synthesizing sulfur-containing compounds.

At present, the conventional synthesis strategies for sulfurcontaining compounds are mainly divided into two categories. One is the cross-coupling synthesis of thiols/disulfides and aryl halides in the presence of transition metal catalysts such as Pd,⁵ and Rh.⁶ The another one is the formation of diaryl sulfides in the presence of an I₂/peroxide catalytic system.⁷ For example, the Schoenebeck group successfully constructed C–S bonds using a Pd catalyst.⁸ Wang's group reported the formation of diaryl sulfides using an I₂/DTBP catalytic system.⁹ However, these traditional methods require the use of expensive substrates, chemical oxidants, metal catalysts, and harsh reaction conditions. However, green chemistry principles demand the development of facile synthesis strategies with atomic



Fig. 1 Organosulfur compounds in natural products and drugs.

*Institute of Advanced Technology, University of Science and Technology of China, Hefei 230000, China. E-mail: sunqi924@ustc.edu.cn; zwang3@ustc.edu.cn

† Electronic supplementary information (ESI) available. See DOI: https://doi.org/10.1039/d5ra00100e

economy and environmental benignity. In this study, an electrochemical method was adopted, wherein only 1.0 equivalent of Et₄NI was needed as the electrolyte, and the construction of C–S bonds was completed in 4 h. Compared with the I₂/DTBP catalytic system, this reaction did not require the excess addition of peroxides (such as DTBP). Additionally, this electrochemical method could be carried out under air atmosphere, while the I₂/DTBP catalysis required inert gas protection.

Electrochemical synthesis is an environmentally benign method to prepare organic compounds as electricity can provide the electrons needed for redox reactions, thereby avoiding the use of redox reagents. Previously, Zhou's group reported the construction of C–S bonds between quinoxalinone and thiol under electrochemical conditions. Intrigued by this electrochemical construction of diaryl sulfur compounds, we developed a reaction to directly construct aromatic sulfur bonds under electrochemical conditions, without the involvement of any transition metal catalysts or chemical oxidants (Scheme 1).

Results and discussion

As a model reaction, the reaction of diphenyl disulfide 1a and aniline 2a was conducted in the presence of Et4NI as the electrolyte and dimethyl sulfoxide (DMSO) as the solvent at a constant current density of 10 mA cm⁻² for 4 h under air atmosphere. The desired product was obtained with a high yield of 74% (entry 1, Table 1). Based on this result, we further optimized the reaction conditions. Firstly, different solvents were screened. The experimental results showed that DMSO was the optimal solvent, as shown in Table 1. Subsequently, various bases and acids were added to the reaction mixture, and the results demonstrated that these additions had a negative influence on the reaction, as shown in entries 4-8 of Table 1. To our delight, when H2O was added, the reaction yield was enhanced from 74% to 80% (entry 10, Table 1). The amount of H₂O added was also investigated. We found that a ratio of DMSO: $H_2O = 10:1$ gave the optimal result, affording 3aa in 86% yield (entry 11, Table 1). Afterwards, a variety of electrolytes, such as n-Bu₄NBF₄, n-Bu₄NPF₆, NH₄BF₄, were examined in

^bHefei National Research Center for Physical Sciences at Microscale, Key Laboratory of Precision and Intelligent Chemistry, School of Chemistry and Materials Science, University of Science and Technology of China, Hefei 230026, China

Previous Work

[a] Ar¹-X + Ar²-SH Pd, Cu, NI, Au, Rh, Fe, Ag Ar¹ S Ar²

X=CI, Br, I, B(OH)₂

[b] NH₂ + R²SH I₂(15 mol%) DTBP(5 equiv), 120°C H₂N R¹

J. Org. Chem. 2015, 80, 6083-6092

[c] R²(1 N H R³SH electricity undivided cell R²(1 N SR³)

J. Org. Chem. 2020, 85, 4365-4372

This Work

[Ar₁] SH Ar₂ NH₂ undivided cell NH₂ NH₂ NH₂ NH₂ NH₂ NH₃ NH₄ NH₂ NH₄ NH₂ NH₄ NH₄

Scheme 1 Previous work and this work for the synthesis of diaryl sulfides.

this reaction. However, no better result was obtained (entries 13–15, Table 1). It was noted that the iodide ion in the electrolyte was essential for constructing the $C(sp^2)$ –S bond. In the

absence of iodide in the electrolyte, no desired product was observed (entries 13–15, Table 1). As a result, we optimized different iodine salts and found that Et₄NI was the best electrolyte for the reaction (entries 16–20, Table 1). Furthermore, the reaction was performed at different temperature. It was found that low yield of 40% was obtained when the reaction was carried out at 100 °C. At temperatures below 100 °C, it was challenging to obtain the desired product, as shown in entries 1–8 of Table S1 in the ESI.† In the end, the electrode was optimized. We attempted to use carbon electrodes to replace the platinum electrodes. However, carbon electrode gave poor results, as shown in entries 21–23 of Table 1.

After establishing the optimal conditions, the substrate scope of thiols and thioethers was studied. As show in Scheme 2, the reaction yields ranged from 81% to 90% (3ba-3ga) when the substituent was an electron-withdrawing group. The position of the methyl substituent had no significant effect on the reaction, and the yield was generally above 83% (3ia-3ka). When the substituent was a strong electron-donating methoxy group, the yield was significantly reduced (3la-3ma). More importantly, when 4-(trifluoromethyl)thiophenol (1d'), 2-bromothiophenol (1f'), 3-chlorothiophenol (1g'), 2-methylbenzenethiol (1j') and 3-methylbenzenethiol (1k') were used as

Table 1 Optimization of reaction conditions^a

Entry	Solvent	Additive	Electrolyte	Electrode	Yield ^b [%]
1	DMSO	_	$\mathrm{Et_{4}NI}$	Pt(+)/Pt(-)	74
2	DMF	_	$\mathrm{Et_{4}NI}$	Pt(+)/Pt(-)	nd^d
3	DMA	_	$\mathrm{Et_{4}NI}$	Pt(+)/Pt(-)	20
4	DMSO	DIPEA	$\mathrm{Et_{4}NI}$	Pt(+)/Pt(-)	nd^d
5	DMSO	Cs_2CO_3	$\mathrm{Et_{4}NI}$	Pt(+)/Pt(-)	nd^d
6	DMSO	NaOH	$\mathrm{Et_{4}NI}$	Pt(+)/Pt(-)	nd^d
7	DMSO	H_2SO_4	$\mathrm{Et_{4}NI}$	Pt(+)/Pt(-)	nd^d
8	DMSO	$H_2C_2O_4$	$\mathrm{Et_{4}NI}$	Pt(+)/Pt(-)	nd^d
9	DMSO	H_2O	$\mathrm{Et_{4}NI}$	Pt(+)/Pt(-)	80
10	DMSO: $H_2O = 7:1$	_	$\mathrm{Et_{4}NI}$	Pt(+)/Pt(-)	80
11	DMSO: $H_2O = 10:1$	_	$\mathrm{Et_{4}NI}$	Pt(+)/Pt(-)	86
12	DMSO: $H_2O = 15:1$	_	$\mathrm{Et_{4}NI}$	Pt(+)/Pt(-)	82
13	DMSO: $H_2O = 10:1$	_	$n ext{-Bu}_4 ext{NBF}_4$	Pt(+)/Pt(-)	nd^d
14	DMSO: $H_2O = 10:1$	_	n-Bu ₄ NPF ₆	Pt(+)/Pt(-)	nd^d
15	DMSO: $H_2O = 10:1$	_	$\mathrm{NH_4BF_4}$	Pt(+)/Pt(-)	nd^d
16	DMSO: $H_2O = 10:1$	_	$n ext{-Bu}_4 ext{NI}$	Pt(+)/Pt(-)	55
17	DMSO: $H_2O = 10:1$	_	Me_4NI	Pt(+)/Pt(-)	66
18	DMSO: $H_2O = 10:1$	_	$\mathrm{NH_{4}I}$	Pt(+)/Pt(-)	65
19	DMSO: $H_2O = 10:1$	_	KI	Pt(+)/Pt(-)	60
20	DMSO: $H_2O = 10:1$	_	NaI	Pt(+)/Pt(-)	40
21	DMSO: $H_2O = 10:1$	_	$\mathrm{Et_{4}NI}$	C(+)/C(-)	30
22	DMSO: $H_2O = 10:1$	_	$\mathrm{Et_{4}NI}$	Pt(+)/C(-)	65
23	$DMSO: H_2O = 10:1$	_	$\mathrm{Et_{4}NI}$	C(+)/Pt(-)	70
24^c	DMSO: $H_2O = 10:1$	_	Et ₄ NI		nd^d

^a Standard conditions: platinum plate (10 mm \times 10 mm \times 0.2 mm) as the anode, platinum plate (10 mm \times 10 mm \times 0.2 mm) as the cathode, undivided cell, **1a** (0.15 mmol), **2a** (0.9 mmol), Et₄NI (0.3 mmol), and DMSO (3 mL), air, 120 °C, 4 h. ^b Isolated yield. ^c Without electricity. ^d Not detected.

Scheme 2 Scope of substrates. Unless otherwise noted, all reactions were performed with 1a-1n (0.15 mmol), 2a (0.9 mmol), Et_4NI (0.3 mmol), DMSO (3.0 mL). The reaction was carried out at 120 °C for 4 h. *Reaction conditions: thiophenols as substrate 1d', 1f', 1g', 1j', 1k' (0.3 mmol), 2a (0.9 mmol), Et_4NI (0.3 mmol), DMSO (3.0 mL). The reaction was carried out at 120 °C for 4 h.

substrates, the reactions proceeded smoothly, and the desired products were obtained with yields of 82–90%. In addition, bis(2-methyl-3-furyl) disulfide was also well tolerated and generated the desired product with 55% yield (3na).

Afterwards, we examined the range of aniline substrates. As shown in Scheme 3, the reaction yields were above 81% (3ab, 3ac) when the single methyl substituent was present on the phenyl ring, regardless of the substituent position. Regarding other substituents, the electronic effect influenced the reaction. When the substituent was located at the *meta*-position, the electron-withdrawing group favored the reaction, while the electron-donating group disfavored it (3ad-3ag). In contrast, when the substituent was located at the *ortho*-position, the electron-donating group was superior to the electron-withdrawing group (3ah-3ak). Regarding di-substitution, the steric effect influenced the reaction (3al, 3am). When the hydrogen on the amino group of the aniline was replaced, the moderate yields of 65–70% were obtained (3an-3ap).

In addition, we compared this electrochemical method with the chemical method. For instance, some substrates, such as 3aa, 3da, 3fa, 3ga, 3ka, 3ha and 3af, worked poorly in the $I_2/DTBP$ catalysis while worked well in this electrochemical reaction (see the control experiments on page S5 of the ESI†). To demonstrate the utility of this reaction, we performed the gramscale of the model reaction. This scale-up reaction was carried out smoothly affording the desired product with a high yield of 85% (Scheme 4). Additionally, methyl *ortho*-aminobenzoate, an important pharmaceutical intermediate of drugs for

Scheme 3 Scope of substrates. Unless otherwise noted, all reactions were performed with 1a (0.15 mmol), 2a-2p (0.9 mmol), Et_4NI (0.3 mmol), DMSO (3.0 mL). The reaction was carried out at 120 °C for 4 h.

psychotropic therapy and antimicrobial activity¹² (3aq) was successfully prepared using our method in one step with a yield of 60%.

To gain more insights into the reaction mechanism, cyclic voltammetry (CV) experiments were conducted. The electrochemical properties of the starting materials were investigated, as shown in Fig. 2. The oxidation peaks from $\rm Et_4NI$ (0.58 V, 0.98 V) [ν s. Ag/AgCl] were observed, while the oxidation peaks of $\rm 1a$ (1.91 V) and $\rm 2a$ (1.22 V) were also detected. This observation verified the catalytic role of $\rm Et_4NI$ in the reaction, indicating that the iodide anion should be oxidized first under standard conditions. We hypothesized that the iodide oxidation could generate some active species to initiate the reaction.

To further investigate the reaction mechanism, we conducted a series of control experiments (Scheme 5). Initially, the reaction proceeded well in the presence of BHT (butylated hydroxytoluene) and 1,1-diphenylethylene (DPE), suggesting that the reaction may not proceed through a radical process.

Scheme 4 Gram-scale experiments, and their employment in the preparation of natural products.

Paper

IUPAC Convention 4.0x10" a. background b. Et₄NI 3.5x10⁻⁴ c. **1a** 3.0x10⁻⁴ 1.91 V 1.22 V 2.5x10⁻⁴ 2 0x10⁻⁴ 0 98 V Current (A) 1.5x10 1.0x10 5.0x10⁻⁵ 0.0 -5.0x10⁻⁵ -1.0x10⁻⁴ 2.0 Potential (V) vs. Ag/AgCI

Fig. 2 Cyclic voltammetry experiments: cyclic voltammograms of 1a, 2a and Et_4NI in 0.1 M n-Bu $_4$ NBF $_4$ /MeCN = 10 mL using a Pt disk as the working electrode, and a Pt wire and Ag/AgCl as the counter and reference electrodes, respectively, at a scan rate of 100 mV s $^{-1}$; background (curve a), Et_4NI (5 mmol Et_4), curve Et_4), Et_4 0 and Et_4 0 and Et_4 0 and Et_4 1 (5 mmol Et_4 1, curve Et_4 1) and Et_4 2 (5 mmol Et_4 1, curve Et_4 3).

Scheme 5 Control experiments.

During the electrolysis process under standard conditions, the complete conversion of thiophenol to diphenyl disulfide was detected.

Based on these control experiments and some previous literature reports, 7,10,13,14 a relatively reasonable reaction mechanism was proposed (Scheme 6). First, iodine anions are oxidized to I_2 at the anode, which can oxidize thiophenol to diphenyl disulfide. The diphenyl disulfide can be further converted into the cationic species 3 by molecular iodine. Subsequently, the cationic species 3 reacts with aniline to afford 3aa. Meanwhile, the water is reduced at the cathode.

Scheme 6 Proposed reaction mechanism.

Conclusion

In summary, we developed a method to synthesize diaryl sulfide derivatives from aniline and thioether/thiophenol via electrochemical catalysis in the presence of Et_4NI . This electrocatalytic C–H/S–H coupling reaction can be carried out smoothly without the use of metals or oxidants, under the air atmosphere. Furthermore, the reaction features a broad substrate scope, rapid reaction rate and high atomic economy, providing a facile avenue for the synthesis of aryl sulfides.

Data availability

The datasets generated and/or analyzed during the current study are available from the corresponding author on reasonable request. The data includes experimental procedures and compound characterizations using NMR and HRMS.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We are grateful for the financial support from the National Natural Science Foundation of China (21772185).

References

- 1 T. Yoneda, N. Kojima, T. Matsumoto, D. Imahori, T. Ohta, T. Yoshida, T. Watanabe, H. Matsuda and S. Nakamura, *Org. Biomol. Chem.*, 2022, **20**, 196.
- 2 X. Chen and B. Li, Curr. Opin. Chem. Biol., 2023, 76, 102377.
- (a) T. Mori, T. Nishimura, T. Yamamoto, E. Miyazaki,
 I. Osaka and K. Takimiya, J. Am. Chem. Soc., 2013, 135,
 13900–13913; (b) H. Mutlu, E. B. Ceper, X. Li, J. Yang,
 W. Dong, M. M. Ozmen and P. Theato, Macromol. Rapid Commun., 2019, 40, 1800650.
- 4 J. Watanabe, J. Shimamoto and K. Kotani, *Antibiotics*, 2021, **10**, 156.
- 5 (a) Y. Liu, X. Mo, I. Majeed, M. Zhang, H. Wang and Z. Zeng, Org. Biomol. Chem., 2022, 20, 1532; (b) M. Iwasaki, T. Fujii, A. Yamamoto, K. Nakajima and Y. Nishihara, Chem.-Asian J., 2014, 9, 58-62.
- 6 (a) M. Arisawa, T. Suzuki, T. Ishikawa and M. Yamaguchi, J. Am. Chem. Soc., 2008, 130, 12214–12215; (b) F. Zhu and X. Wu, J. Org. Chem., 2018, 83, 13612–13617.
- 7 (a) H. Wang, T. Shi, W. Gao, Y. Wang, J. Li, Y. Jiang, Y. Hou,
 C. Chen, X. Peng and Z. Wang, Chem.-Asian J., 2017, 12,
 2675-2679; (b) X. Jiang, Z. Shen, C. Zheng, L. Fang,
 K. Chen and C. Yu, Tetrahedron Lett., 2020, 61, 152141; (c)
 Y. Yang, L. Liu, K. Li, Z. Zha, Q. Sun and Z. Wang, RSC Adv., 2022, 12, 7347-7351; (d) X. Zhang, J. Jiao, X. Zhang,
 B. Hu and X. Zhang, J. Org. Chem., 2016, 81, 5710-5716.
- 8 T. Scattolin, E. Senol, G. Yin, Q. Guo and F. Schoenebeck, *Angew. Chem., Int. Ed.*, 2018, 57, 12425–12429.

- 9 D. Yang, K. Yan, W. Wei, J. Zhao, M. Zhang, X. Sheng, G. Li and H. Wang, *J. Org. Chem.*, 2015, **80**, 6083–6092.
- 10 (a) Y. Yuan, Y. Cao, J. Qiao, Y. Lin, X. Jiang, Y. Weng, S. Tang and A. Lei, *Chin. J. Chem.*, 2019, 37, 49–52; (b) X. Sun, Y. Zhang, T. Li, K. Li, Q. Sun and Z. Wang, *Org. Lett.*, 2024, 26, 1566–1572; (c) J. Hua, Z. Fang, J. Xu, M. Bian, C. Liu, W. He, N. Zhu, Z. Yang and K. Guo, *Green Chem.*, 2019, 21, 4706–4711; (d) Y. Yu, Y. Jiang, S. Wu, Z. Shi, J. Wu, Y. Yuan and K. Ye, *Chin. Chem. Lett.*, 2022, 33, 2009–2014.
- 11 J. Zhou, Z. Li, Z. Sun, Q. Ren, Q. Zhang, H. Li and J. Li, *J. Org. Chem.*, 2020, **85**, 4365–4372.
- 12 S. Gao, Y. Shen, S. Yuan, Y. Quan, X. Li, Y. Wang, L. Yi and Y. Wang, *Int. J. Antimicrob. Agents*, 2023, **62**, 106996.
- 13 W. Zhang, Q. Zou, Q. Wang, D. Jin, S. Jiang and P. Qian, *J. Org. Chem.*, 2024, **89**, 5434–5441.
- 14 S. B. Annes, R. Saritha, K. Chandru, P. K. Mandali and S. Ramesh, *J. Org. Chem.*, 2021, **86**, 16473–16484.