


 Cite this: *RSC Adv.*, 2025, 15, 21872

Catalytic system-controlled divergent reactions of pyrazolidinones with 3-alkynyl-3-hydroxyisoindolinones to construct diversified nitrogen-containing heterocyclic scaffolds†

 Zhenwei Zhang,^{‡*abc} Xiaoxue Zhang,^{‡bcd} Hao Wang,^{bc} Geyao Xie^{abc} and Yu Zhou^{‡*abcd}

A catalytic system-controlled divergent reaction was reported to construct three distinct nitrogen-containing heterocycles from readily available starting materials *via* a precise chemical bond activation and annulation cascade. Notably, this is the first capture of pyrazolidinones and propargyl alcohols to construct tetrahydropyrimidinones *via* selective N–N bond activation and to generate previously unreported 3-acylindoles. This protocol demonstrates a broad substrate scope, moderate to good yields, and valuable transformations, laying a robust foundation for drug discovery applications.

 Received 21st March 2025
 Accepted 30th May 2025

DOI: 10.1039/d5ra01992c

rsc.li/rsc-advances

Introduction

Nitrogen-containing heterocyclic structures are highly valued patterns present in a wide range of natural products, pharmaceuticals, agrochemicals, and organic materials.^{1–3} Recently, pyrazolidinones with a free NH moiety have been employed as a traceless directing group for constructing diverse privileged nitrogen-containing heterocyclic frameworks, including N-substituted indoles,^{4–6} pyrazolo[1,2-*a*]pyrazolones,^{7–9} *N,N*-bicyclic pyrazolidinones,^{10,11} diazepines^{12–14} and CF₃-tethered indazoles.¹⁵ The propargylic alcohols have also been successfully involved in constructing diverse nitrogen-containing heterocyclic compounds.^{16,17} In 2020, Fan's group developed a pioneering Rh(III)-catalyzed redox-neutral coupling of 1-phenylpyrazolidinones with alkynyl cyclobutanols, enabling selective construction of 2-acylindoles and pyrazolo[1,2-*a*]pyrazolones through [3 + 2] and [4 + 1] annulations, respectively, by adjusting reaction conditions (Scheme 1a, left).¹⁸ In 2018, Ji's group reported three examples where Rh(III)-catalyzed [4 + 1] annulation of substituted 1-phenylpyrazolidinones with 1-

phenylbut-2-yn-1-ol was used to construct desired pyrazolo[1,2-*a*]indazolones.¹⁹ In 2020, the groups of Cui and Fan independently accomplished the Rh(III)-catalyzed [4 + 3] annulations of pyrazolidinones with propargyl alcohols or propargylic acetates to construct benzodiazepines (Scheme 1a, right).^{20,21} Despite these elegant works, they mainly focused on C–H bond activation/annulation cascades, with limited exploration of N–N bond activation for heterocyclic framework construction from these substrates.

In our continuous efforts to pursue rapid construction of diverse drug-like heterocyclic skeletons *via* transition-metal catalysed chemical bond activation/annulation cascades,^{22–27} we sought to develop more distinct and structurally complex

^aSchool of Pharmaceutical Science and Technology, Hangzhou Institute for Advanced Study, University of Chinese Academy of Sciences, Hangzhou 310024, China. E-mail: zhouyu@simm.ac.cn; zhangzhenwei@ucas.ac.cn


^bDrug Discovery & Development Center, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203, China

^cUniversity of Chinese Academy of Sciences, Beijing 100049, China

^dSchool of Chinese Materia Medica, Nanjing University of Chinese Medicine, Nanjing 210023, China

† Electronic supplementary information (ESI) available. CCDC 2410303, 2410305 and 2410625. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d5ra01992c>

‡ These authors contributed equally to this work.



Scheme 1 Previous works and this work.




nitrogen-containing heterocycles by employing the polycyclic propargyl alcohols. Interestingly, we observed an interesting phenomenon when the propargylic alcohols (**2**) with a large sterically hindered group were introduced as the coupling partner, 1-phenylpyrazolidinones (**1**) looked like a generalist that could be precisely and independently assembled into three distinct types of nitrogen-containing heterocyclic skeletons at different reaction sites of itself under different catalytic conditions (Scheme 1b). Of note, except for passing through a C(sp²)-H/N-H activation to generate pyrazolo[1,2-*a*]pyrazolones (**4**), we have captured a new N-N bond activation product tetrahydropyrimidinones (**3**) and an unusual product 3-acylindoles (**5**). These findings provide a versatile strategy for synthesizing diverse and biologically active nitrogen-containing heterocycles.

Results and discussion

We initially investigated this transformation by employing the 4,4-dimethyl-1-phenylpyrazolidin-3-one (**1a**, 0.24 mmol) and 3-hydroxy-3-(phenylethynyl)isoindolin-1-one (**2a**, 0.2 mmol) as model substrates, and the results showed that the desired N-N bond activation product **3aa** could be constructed with a 20% yield under [Cp*RhCl₂]₂ (5 mol%) as the catalyst and 1,2-dichloroethane (DCE) as the solvent at 90 °C for 4 h (Table 1, entry 1). Surprisingly, adding silver acetate and acetic acid into the catalytic system would result in a new [4 + 1] annulation

product **4aa** with 8% yield (Table 1, entry 2), while the addition of cesium pivalate and acetic acid into the catalytic system simultaneously formed **4aa** (28% yield) and a [3 + 2] annulation product **5aa** (10% yield) (Table 1, entry 3). Interestingly, the yield of **5aa** could be increased to 50% when we used dioxane as the solvent (Table 1, entry 4). Encouraged by these results, we further screened different catalysts, including Rh₂(esp)₂, CyJohnPhos AuCl, (Ph₃P)AuCl, Ph₃PAuNTf₂, CyJohnPhos AuNTf₂ and AuCl(IPr) (Table 1, entries 5–10). The results showed that the AuCl(IPr) offered the best catalytic activity (Table 1, entry 10) for the formation of **3aa**, the yield of which could be increased to 54%. Subsequent solvent screening indicated that DCE was the most conducive to this transformation (Table 1, entries 11–13). Next, lowering the reaction temperature to 70 °C could further increase the yield to 65% (Table 1, entry 14), while increasing the reaction temperature to 100 °C resulted in a slight decrease in yield (Table 1, entry 15). In brief, the optimal reaction conditions for **3aa** were as follows: **1a** (0.24 mmol), **2a** (0.2 mmol), and AuCl(IPr) (5 mol%) in DCE at 70 °C for 4 h (condition A). Likewise, the optimal reaction conditions for **4aa** and **5aa** were also separately investigated (see Tables S1 and S2†). The results displayed that when using **1a** (0.24 mmol), **2a** (0.2 mmol), [Cp*Rh(CH₃CN)₃](SbF₆)₂ (5 mol%), LiOAc (0.2 mmol), BzOH (0.1 mmol) at 100 °C for 8 h (Table 1, entry 16, condition B) and **1a** (0.2 mmol), **2a** (0.24 mmol), [Cp*RhCl₂]₂ (5 mol%), CsOPiv (0.2 mmol) and AcOH (0.2 mmol)

Table 1 Optimization of reaction conditions^a



| Entry | Catalytic system | Solvent | Yield (%) | | |
|-----------------|---|--------------------|-----------|-----|-------|
| | | | 3aa | 4aa | 5aa |
| 1 | [Cp*RhCl ₂] ₂ | DCE | 20 | — | — |
| 2 ^b | [Cp*RhCl ₂] ₂ /AgOAc/AcOH | DCE | — | 8 | — |
| 3 ^b | [Cp*RhCl ₂] ₂ /CsOPiv/AcOH | DCE | — | 28 | 10 |
| 4 ^b | [Cp*RhCl ₂] ₂ /CsOPiv/AcOH | Dioxane | — | — | 50 |
| 5 | Rh ₂ (esp) ₂ | DCE | 44 | — | — |
| 6 | CyJohnPhos AuCl | DCE | 31 | — | — |
| 7 | CyJohnPhos AuNTf ₂ | DCE | 44 | — | — |
| 8 | (Ph ₃ P)AuCl | DCE | 44 | — | — |
| 9 | Ph ₃ PAuNTf ₂ | DCE | 36 | — | — |
| 10 | AuCl(IPr) | DCE | 54 | — | — |
| 11 | AuCl(IPr) | MeCN | 15 | — | — |
| 12 | AuCl(IPr) | CHCl ₃ | 41 | — | — |
| 13 | AuCl(IPr) | HFIP | Trace | — | — |
| 14 ^c | AuCl(IPr) | DCE | 65 | — | — |
| 15 ^d | AuCl(IPr) | DCE | 50 | — | — |
| 16 ^e | [Cp*Rh(CH ₃ CN) ₃](SbF ₆) ₂ /LiOAc/BzOH | CH ₃ CN | — | 60 | Trace |
| 17 ^f | [Cp*RhCl ₂] ₂ /CsOPiv/AcOH | Diglyme | — | — | 80 |


^a Reaction conditions: **1a** (0.24 mmol), **2a** (0.2 mmol), catalyst (5.0 mol%) in solvent (4 mL) at 90 °C for 4 h, NMR yield (1,3,5-trimethoxybenzene as the internal standard). ^b **1a** (0.2 mmol), **2a** (0.24 mmol), inorganic salts (0.4 mmol), AcOH (0.2 mmol) at 80 °C for 8 h, isolated yield. ^c 70 °C. ^d 100 °C. ^e **1a** (0.24 mmol), **2a** (0.2 mmol), [Cp*Rh(CH₃CN)₃](SbF₆)₂ (5 mol%), LiOAc (0.2 mmol), BzOH (0.1 mmol) at 100 °C for 8 h, isolated yield. ^f **1a** (0.2 mmol), **2a** (0.24 mmol), [Cp*RhCl₂]₂ (5 mol%), CsOPiv (0.2 mmol), AcOH (0.2 mmol) at 80 °C for 2 h, isolated yield.




in diglyme at 80 °C for 2 h (Table 1, entry 17, condition C) as their respective catalytic systems, the products **4aa** and **5aa** were effectively prepared with 60% and 80%, respectively. Their structures were confirmed by ^1H and ^{13}C NMR, MS and X-ray spectra.

With the optimized reaction conditions in hand, we firstly conducted a comprehensive investigation into the scope of pyrazolidinones **1** and coupling partners **2** under the optimal reaction condition to build diversified tetrahydropyrimidinones **3**. As shown in Scheme 2, whether introducing halogen atoms (F, Cl and Br) or electron-donating groups (CH_3 and OCH_3) into 4-position of benzene ring of substrate **1**, the reaction efficiencies were maintained at a moderate yield (Scheme 2, **3ba–3fa**, 36–64%). However, introducing an F atom into 3-position of phenyl ring resulted in a decrease in yield (**3ia**, 25%). Subsequently, we investigated the effect of various substituents on the coupling partners **2**. Notably, substrates **2** bearing halogen atoms (F, Cl and Br) or electron-donating groups (CH_3 , OCH_3 and $(\text{CH}_2)_4\text{CH}_3$) at the 4-position of the phenyl ring afforded corresponding products in moderate to good yields (**3ab–3ag**, 49–72%). Introducing CH_3 and Cl groups into the 3-position of the phenyl ring also gave satisfactory results (**3ai–3ak**, 69–72%). Replacing the phenyl ring of **2** with the thiophene ring also provided a good yield (**3al**, 66%).

During the reaction condition screening, we observed the pyrazolo[1,2-*a*]pyrazolone derivative **4aa** could be selectively



Scheme 2 The investigation of substrate scope for the synthesis of **3**. General reaction conditions (condition A): **1** (0.24 mmol), **2** (0.2 mmol), $\text{AuCl}(\text{IPr})$ (5 mol%) in DCE (4.0 mL) at 70 °C under air for 4 h, isolated yields.




Scheme 3 The investigation of substrate scope for the synthesis of **4**. General reaction conditions (condition B): **1** (0.24 mmol), **2** (0.2 mmol), $[\text{Cp}^*\text{Rh}(\text{CH}_3\text{CN})_3](\text{SbF}_6)_2$ (5 mol%), LiOAc (0.2 mmol), BzOH (0.1 mmol) in MeCN (4.0 mL) at 100 °C under air for 8 h, isolated yields.

constructed *via* a $\text{C}(\text{sp}^2)\text{-H}$ bond activation/[4 + 1] annulation cascade. As shown in Scheme 3, the reaction worked well with electron-withdrawing and electron-donating groups at the 4-position of the phenyl ring of substrates **1**, affording **4ba–4ha** with 47–71% yields (Scheme 3). Translocating Cl or Br to the 3-position of the phenyl ring could also successfully generate corresponding products with 45–68% yields (**4ia–4ka**). Next, various substituted groups at the 4-position of the phenyl ring of substrate **2** were explored, and the results indicated that all substrates could react smoothly with **1a** to provide the corresponding products **4ab–4ag** in moderate to good yields (40–60%). Likewise, introducing CH_3 or Cl into the *meta* or *ortho* position of the benzene ring was also satisfactory (**4ai–4ak**). However, replacing benzene ring of **2** with thiophene ring gave a relatively low yield (**4al**, 21%).

Another discovery was that we successfully obtained previously unreported 3-acylindoles *via* a $\text{C}(\text{sp}^2)\text{-H}$ bond activation and [3 + 2] annulation cascade from pyrazolidinone **1** and its coupling partner **2**. As shown in Scheme 4, whether introducing electron-donating or electron-withdrawing groups into pyrazolidinones **1** or 3-alkynyl-3-hydroxyisoindolinones **2** could






Scheme 4 The investigation of substrate scope for the synthesis of **5**. General reaction conditions (condition C): **1** (0.2 mmol), **2** (0.24 mmol), $[\text{Cp}^*\text{RhCl}_2]_2$ (5 mol%), CsOPiv (0.2 mmol), AcOH (0.2 mmol) in diglyme (4.0 mL) at 80 °C under air for 2 h, isolated yields.

generate desired products in good yields (**5ba–5fa**, 81–91%; **5ab–5ak**, 62–72%). In particular, in this transformation, replacing the benzene ring of **2** with the thiophene ring could provide a good yield (Scheme 4, **5al**, 75%). Additionally, we also attempted to introduce the group into R^3 position of substrate **2**, the corresponding product **5am** was obtained with good yield.

Given the potential application of these intriguing heterocycles, the gram-scale preparations for the products **3aa**, **4aa**, and **5aa** were performed with 61%, 47%, and 67% isolated yields, respectively (Scheme 5a). Then, the synthetic transformations of compounds **4aa** and **5aa** were also explored. For example, the C–C double bond of **4aa** could be selectively reduced using hydrogen gas and Pd/C, generating the desired product **6** with 80% yield (Scheme 5b). Interestingly, we successfully fulfilled a simultaneous transformation of two amide groups in **5aa**, in which compound **5aa** could be converted into nitrile product **7** with 84% yield in the presence of cyanuric chloride (Scheme 5b). Additionally, treatment of **5aa** with triethylsilane and trifluoroacetic acid afforded a ring-closed product **8** (Scheme 5b).

Having established the substrate scope and utility of the product, we then conducted a series of experiments to explore the preliminary mechanism (Scheme 6). Firstly, 95% of *ortho* H could be deuterated by treating **1a** with D_2O under standard condition B, 25% of the *ortho* H could be deuterated by treating **1a** with D_2O under standard condition C, suggesting the reversibility of the C(aryl)–H bond cleavage during the




Scheme 5 Gram-scale preparations and conversions.


formation of products **4** and **5** (Scheme 6a). Next, the kinetic isotope effect (KIE) values of 1.70 for **4aa** and 1.17 for **5aa** indicated that the C–H bond cleavage was likely not the turnover-limiting step (Scheme 6b). Moreover, three competitive experiments within three reaction conditions by introducing different substrates with electron-withdrawing and electron-donating groups were carried out. Treating a 1:1 mixture of fluorine/methoxy substituted pyrazolidinone (**1b/1f**) with **2a** under conditions A and C, respectively, showed that there is no apparent difference in the reaction rate between two substrates **1** with electron-withdrawing and electron-donating groups (Scheme 6c). These results indicated that these different groups are well tolerated in these transformations. However, we observed a high ratio (8.0) between **4ga** and **4fa** when treating a 1:1 mixture of trifluoromethyl/methoxy substituted pyrazolidinone (**1g/1f**) with **2a** under condition B (Scheme 6c), suggesting that the electron-withdrawing substrates reacted faster in this transformation.

Based on the above observations and literature precedents,^{18,21,24,28–32} three plausible catalytic mechanisms are proposed for N–N bond activations and C–H functionalizations. As shown in Scheme 7a, the propargylic alcohol **2a** initially forms the intermediate **A** by coordinating with the Au(I) catalyst. Upon the entry of the pyrazolidinone **1a** into the system, it





Scheme 6 Preliminary mechanism studies.



Scheme 7 Plausible mechanisms.

undergoes alkyne hydroamination with Au(I) to yield complex **B**, which further forms the intermediate **C** after the intramolecular electrophilic addition. Finally, elimination of the Au(I) catalyst and the hydroxy group from **C** gives the desired product **3aa**. The speculated mechanisms of the desired products **4aa** and **5aa** are shown in Scheme 7b. The NH directing group of **1a** assists the activated Rh catalyst with the cleavage of the *ortho* C(sp²)-H bond, enabling the formation of a five-membered rhodacycle **D**. Next, the complexation of Rh(III) with the triple bond of **2a** gives intermediate **E**. The following regioselective insertion of the alkynyl unit of **2a** into the C-Rh bond of **E** generates intermediate **F**, which is most likely driven by additional coordination of the OH group with Rh(III) to overcome the unfavorable steric interactions between the isoindolin-1-one group and the Cp ligand. Further, the cleavage of the C-Rh bond and elimination of water under the assistance of benzoic acid offers the allenyl intermediate **G**. Finally, **G** undergoes nucleophilic addition and followed by protodemetalation to form target product **4aa**. Specially, using [Cp*RhCl₂]₂ as the catalyst in diglyme results in the opposite regioselective insertion of the alkyne unit of **2a** into the C-Rh bond of **E** to form intermediate **H**, probably because the solvent prevents the additional coordination of the OH group of propargylic alcohol with Rh(III) complex. Then protonation of intermediate **H** leads to the generation of intermediate **I**, which then undergoes an oxidative addition with N-N bond to form intermediate **J**. Finally, intermediate **J** undergoes a reductive elimination and a following *N*-demethylation/protonation to produce the desired product **5aa**.

Conclusions

In summary, we have successfully fulfilled a catalytic system-controlled divergent reaction strategy to construct three types of nitrogen-containing heterocyclic skeletons through chemical bond activation/annulation routes of pyrazolidinones with 3-alkynyl-3-hydroxyisoindolinones. More importantly, this is the first capture of pyrazolidinones and propargyl alcohols to construct tetrahydropyrimidinones and unusual 3-acylindoles. These strategies exhibit a broad range of substrates, moderate to good yields, and valuable transformations, offering structural and biological potentials for further drug discovery.

Data availability

The data supporting this article have been included as part of the ESI,[†] which includes experimental details, characterization data, NMR spectra, and HPLC spectra, along with single-crystal X-ray data for compounds **3aa** (CCDC 2410303), **4aa** (CCDC 2410625), and **5aa** (CCDC 2410305).

Conflicts of interest

The authors declare no competing interests.



Acknowledgements

We thank the National Natural Science Foundation of China (82173656, 82373709, 82151219), Shandong Laboratory Program (SYS202205), and China Postdoctoral Science Foundation (2023M730821) for financial support.

References

- 1 C. Wallwey and S.-M. Li, Ergot alkaloids: structure diversity, biosynthetic gene clusters and functional proof of biosynthetic genes, *Nat. Prod. Rep.*, 2011, **28**, 496–510.
- 2 C. Yang, F. Song, J. Chen and Y. Huang, Construction of Pyridazine Analogues via Rhodium-mediated C-H Activation, *Adv. Synth. Catal.*, 2017, **359**, 3496–3502.
- 3 W. Zi, Z. Zuo and D. Ma, Intramolecular dearomative oxidative coupling of indoles: a unified strategy for the total synthesis of indoline alkaloids, *Acc. Chem. Res.*, 2015, **48**, 702–711.
- 4 Z. Zhang, H. Jiang and Y. Huang, Ruthenium-catalyzed redox-neutral C-H activation via N-N cleavage: synthesis of N-substituted indoles, *Org. Lett.*, 2014, **16**, 5976–5979.
- 5 Z. Yang, Q. Yue, M. Yang, H. Zhang and X. Cui, Ru(II)-Catalyzed Tunable Cascade Reaction via C-H/C-C Bond Cleavage, *J. Org. Chem.*, 2020, **85**, 12960–12970.
- 6 Z. Fan, H. Lu, W. Li, K. Geng and A. Zhang, Rhodium-catalyzed redox-neutral coupling of phenidones with alkynes, *Org. Biomol. Chem.*, 2017, **15**, 5701–5708.
- 7 N. Strašek, L. Lavrenčič, A. Oštrek, D. Slapšak, U. Grošelj, M. Klemenčič, H. Brodnik Žugelj, J. Waggener, M. Novinec and J. Svete, Tetrahydro-1H,5H-pyrazolo[1,2-a]pyrazole-1-carboxylates as inhibitors of Plasmodium falciparum dihydroorotate dehydrogenase, *Bioorg. Chem.*, 2019, **89**, 102982.
- 8 L. Yang, Y. Lv, F. Wang and G. Zhong, Chiral NHC-catalyzed 1,3-dipolar [3 + 2] cycloaddition of azomethine imines with α -chloroaldehydes for the synthesis of bicyclic pyrazolidinones, *Org. Biomol. Chem.*, 2018, **16**, 4433–4438.
- 9 R. Na, C. Jing, Q. Xu, H. Jiang, X. Wu, J. Shi, J. Zhong, M. Wang, D. Benitez, E. Tkatchouk, W. A. Goddard, H. Guo and O. Kwon, Phosphine-catalyzed annulations of azomethine imines: allene-dependent [3 + 2], [3 + 3], [4 + 3], and [3 + 2 + 3] pathways, *J. Am. Chem. Soc.*, 2011, **133**, 13337–13348.
- 10 M. P. Clark, S. K. Laughlin, M. J. Laufersweiler, R. G. Bookland, T. A. Brugel, A. Golebiowski, M. P. Sabat, J. A. Townes, J. C. VanRens, J. F. Djung, M. G. Natchus, B. De, L. C. Hsieh, S. C. Xu, R. L. Walter, M. J. Meikel, S. A. Heitmeyer, K. K. Brown, K. Juergens, Y. O. Taiwo and M. J. Janusz, Development of orally bioavailable bicyclic pyrazolones as inhibitors of tumor necrosis factor- α production, *J. Med. Chem.*, 2004, **47**, 2724–2727.
- 11 W. Lewgowd and A. Stanczak, Cinnoline derivatives with biological activity, *Arch. Pharm.*, 2007, **340**, 65–80.
- 12 V. A. Assimon, Y. Tang, J. D. Vargas, G. J. Lee, Z. Y. Wu, K. Lou, B. Yao, M. K. Menon, A. Pios, K. C. Perez, A. Madriaga, P. K. Buchowiecki, M. Rolfe, L. Shawver, X. Jiao, R. Le Moigne, H. J. Zhou and D. J. Anderson, CB-6644 Is a Selective Inhibitor of the RUVBL1/2 Complex with Anticancer Activity, *ACS Chem. Biol.*, 2019, **14**, 236–244.
- 13 J. Velcicky, U. Bodendorf, P. Rigollier, R. Epple, D. R. Beisner, D. Guerini, P. Smith, B. Liu, R. Feifel, P. Wipfli, R. Aichholz, P. Couttet, I. Dix, T. Widmer, B. Wen and T. Brandl, Discovery of the First Potent, Selective, and Orally Bioavailable Signal Peptide Peptidase-Like 2a (SPPL2a) Inhibitor Displaying Pronounced Immunomodulatory Effects In Vivo, *J. Med. Chem.*, 2018, **61**, 865–880.
- 14 F. A. Romero, A. M. Taylor, T. D. Crawford, V. Tsui, A. Côté and S. Magnuson, Disrupting Acetyl-Lysine Recognition: Progress in the Development of Bromodomain Inhibitors, *J. Med. Chem.*, 2016, **59**, 1271–1298.
- 15 H. Li, M. Shen, B. Li, X. Zhang and X. Fan, Solvent-Dependent Selective Synthesis of CF(3)-Tethered Indazole Derivatives Based on Multiple Bond Activations, *Org. Lett.*, 2023, **25**, 720–725.
- 16 X.-R. Song, R. Yang and Q. Xiao, Recent Advances in the Synthesis of Heterocyclics via Cascade Cyclization of Propargylic Alcohols, *Adv. Synth. Catal.*, 2021, **363**, 852–876.
- 17 P. Sihag and M. Jeganmohan, Regioselective Synthesis of Isocoumarins via Iridium(III)-Catalyzed Oxidative Cyclization of Aromatic Acids with Propargyl Alcohols, *J. Org. Chem.*, 2019, **84**, 2699–2712.
- 18 Y. Xu, M. Shen, X. Zhang and X. Fan, Selective Synthesis of Pyrazolo[1,2-a]pyrazolones and 2-Acylindoles via Rh(III)-Catalyzed Tunable Redox-Neutral Coupling of 1-Phenylpyrazolidinones with Alkynyl Cyclobutanols, *Org. Lett.*, 2020, **22**, 4697–4702.
- 19 X. Wu and H. Ji, Rhodium-Catalyzed [4 + 1] Cyclization via C-H Activation for the Synthesis of Divergent Heterocycles Bearing a Quaternary Carbon, *J. Org. Chem.*, 2018, **83**, 4650–4656.
- 20 L. Zhang, Y. Xu, X. Zhang, X. Zhang and X. Fan, Synthesis of pyrazolone fused benzodiazepines via Rh(III)-catalyzed [4 + 3] annulation of 1-phenylpyrazolidinones with propargyl alcohols, *Org. Chem. Front.*, 2020, **7**, 2284–2290.
- 21 T. Li, Z. Yang, Z. Song, R. Chauvin and X. Cui, Rhodium(III)-Catalyzed [4+3] Annulation of N-Aryl-pyrazolidinones and Propargylic Acetates: Access to Benzo[c][1,2]diazepines, *Org. Lett.*, 2020, **22**, 4078–4082.
- 22 F. Fang, C. Zhang, C. Zhou, Y. Li, Y. Zhou and H. Liu, Rh(III)-Catalyzed C-H Activation of Benzoylacetonitriles and Tandem Cyclization with Diazo Compounds to Substituted Benzo[de]chromenes, *Org. Lett.*, 2018, **20**, 1720–1724.
- 23 X. Wu, B. Wang, S. Zhou, Y. Zhou and H. Liu, Ruthenium-Catalyzed Redox-Neutral [4 + 1] Annulation of Benzamides and Propargyl Alcohols via C-H Bond Activation, *ACS Catal.*, 2017, **7**, 2494–2499.
- 24 F. Fang, S. Hu, C. Li, Q. Wang, R. Wang, X. Han, Y. Zhou and H. Liu, Catalytic System-Controlled Divergent Reaction Strategies for the Construction of Diversified Spiropyrazolone Skeletons from Pyrazolidinones and Diazopyrazolones, *Angew. Chem., Int. Ed.*, 2021, **60**, 21327–21333.



- 25 H. Wang, Y. Li, L. Huang, H. Xu, Y. Jiao, Z. Zhou, Z. Tang, F. Fang, X. Zhang, K. Ding, W. Yi, H. Liu, X. Wu and Y. Zhou, The synthesis of spirocyclopropane skeletons enabled by Rh(III)-catalyzed enantioselective C–H activation/[4 + 2] annulation, *Chem Catal.*, 2023, **3**, 100822.
- 26 J. Gao, K. Luo, X. Wei, H. Wang, H. Liu and Y. Zhou, Rh(III)-Catalyzed Successive C–H Activations of 2-Phenyl-3H-indoles and Cyclization Cascades to Construct Highly Fused Indole Heteropolycycles, *Org. Lett.*, 2023, **25**, 3341–3346.
- 27 X. Hou, R. Wang, F. Fang, Z. Qu, J. Zhou, T. Yu, D. Wang, H. Liu and Y. Zhou, Rh(III)-Catalyzed C–H Activation/Annulation for the Construction of Quinolizinones and Indolizines, *Org. Lett.*, 2024, **26**, 4451–4456.
- 28 Y. Yang, N. Li, J. Zhao, Y. Jiang, X. Zhang and X. Fan, Selective Synthesis of 3-(α -Fluorovinyl)indoles and 3-Acylindoles via the Cascade Reactions of 1-Phenylpyrazolidinones with α,α -Difluoromethylene Alkynes, *Adv. Synth. Catal.*, 2021, **363**, 3600–3606.
- 29 Z. Yang, Z. Song, L. Jie, L. Wang and X. Cui, Iridium(III)-catalysed annulation of pyrazolidinones with propiolates: a facile route to pyrazolo[1,2-*a*] indazoles, *Chem. Commun.*, 2019, **55**, 6094–6097.
- 30 D. Campeau, D. F. León Rayo, A. Mansour, K. Muratov and F. Gagosz, Gold-Catalyzed Reactions of Specially Activated Alkynes, Allenes, and Alkenes, *Chem. Rev.*, 2021, **121**, 8756–8867.
- 31 A. S. K. Hashmi, Gold-Catalyzed Organic Reactions, *Chem. Rev.*, 2007, **107**, 3180–3211.
- 32 S. K. Yadav, B. Ramesh and M. Jeganmohan, Cobalt(III)-Catalyzed Chemo- and Regioselective [4 + 2]-Annulation of Aromatic Sulfoxonium Ylides with 1,3-Diynes, *J. Org. Chem.*, 2022, **87**, 4134–4153.

