


 Cite this: *RSC Adv.*, 2025, 15, 21872

# Catalytic system-controlled divergent reactions of pyrazolidinones with 3-alkynyl-3-hydroxyisoindolinones to construct diversified nitrogen-containing heterocyclic scaffolds†

 Zhenwei Zhang,<sup>‡\*abc</sup> Xiaoxue Zhang,<sup>‡bcd</sup> Hao Wang,<sup>bc</sup> Geyao Xie<sup>abc</sup>  
 and Yu Zhou<sup>‡\*abcd</sup>

A catalytic system-controlled divergent reaction was reported to construct three distinct nitrogen-containing heterocycles from readily available starting materials *via* a precise chemical bond activation and annulation cascade. Notably, this is the first capture of pyrazolidinones and propargyl alcohols to construct tetrahydropyrimidinones *via* selective N–N bond activation and to generate previously unreported 3-acylindoles. This protocol demonstrates a broad substrate scope, moderate to good yields, and valuable transformations, laying a robust foundation for drug discovery applications.

 Received 21st March 2025  
 Accepted 30th May 2025

DOI: 10.1039/d5ra01992c

[rsc.li/rsc-advances](https://rsc.li/rsc-advances)

## Introduction

Nitrogen-containing heterocyclic structures are highly valued patterns present in a wide range of natural products, pharmaceuticals, agrochemicals, and organic materials.<sup>1–3</sup> Recently, pyrazolidinones with a free NH moiety have been employed as a traceless directing group for constructing diverse privileged nitrogen-containing heterocyclic frameworks, including N-substituted indoles,<sup>4–6</sup> pyrazolo[1,2-*a*]pyrazolones,<sup>7–9</sup> *N,N*-bicyclic pyrazolidinones,<sup>10,11</sup> diazepines<sup>12–14</sup> and CF<sub>3</sub>-tethered indazoles.<sup>15</sup> The propargylic alcohols have also been successfully involved in constructing diverse nitrogen-containing heterocyclic compounds.<sup>16,17</sup> In 2020, Fan's group developed a pioneering Rh(III)-catalyzed redox-neutral coupling of 1-phenylpyrazolidinones with alkynyl cyclobutanols, enabling selective construction of 2-acylindoles and pyrazolo[1,2-*a*]pyrazolones through [3 + 2] and [4 + 1] annulations, respectively, by adjusting reaction conditions (Scheme 1a, left).<sup>18</sup> In 2018, Ji's group reported three examples where Rh(III)-catalyzed [4 + 1] annulation of substituted 1-phenylpyrazolidinones with 1-

phenylbut-2-yn-1-ol was used to construct desired pyrazolo[1,2-*a*]indazolones.<sup>19</sup> In 2020, the groups of Cui and Fan independently accomplished the Rh(III)-catalyzed [4 + 3] annulations of pyrazolidinones with propargyl alcohols or propargylic acetates to construct benzodiazepines (Scheme 1a, right).<sup>20,21</sup> Despite these elegant works, they mainly focused on C–H bond activation/annulation cascades, with limited exploration of N–N bond activation for heterocyclic framework construction from these substrates.

In our continuous efforts to pursue rapid construction of diverse drug-like heterocyclic skeletons *via* transition-metal catalysed chemical bond activation/annulation cascades,<sup>22–27</sup> we sought to develop more distinct and structurally complex

<sup>a</sup>School of Pharmaceutical Science and Technology, Hangzhou Institute for Advanced Study, University of Chinese Academy of Sciences, Hangzhou 310024, China. E-mail: zhouyu@simm.ac.cn; zhangzhenwei@ucas.ac.cn


<sup>b</sup>Drug Discovery & Development Center, Shanghai Institute of Materia Medica, Chinese Academy of Sciences, Shanghai 201203, China

<sup>c</sup>University of Chinese Academy of Sciences, Beijing 100049, China

<sup>d</sup>School of Chinese Materia Medica, Nanjing University of Chinese Medicine, Nanjing 210023, China

† Electronic supplementary information (ESI) available. CCDC 2410303, 2410305 and 2410625. For ESI and crystallographic data in CIF or other electronic format see DOI: <https://doi.org/10.1039/d5ra01992c>

‡ These authors contributed equally to this work.



Scheme 1 Previous works and this work.




nitrogen-containing heterocycles by employing the polycyclic propargyl alcohols. Interestingly, we observed an interesting phenomenon when the propargylic alcohols (**2**) with a large sterically hindered group were introduced as the coupling partner, 1-phenylpyrazolidinones (**1**) looked like a generalist that could be precisely and independently assembled into three distinct types of nitrogen-containing heterocyclic skeletons at different reaction sites of itself under different catalytic conditions (Scheme 1b). Of note, except for passing through a C(sp<sup>2</sup>)-H/N-H activation to generate pyrazolo[1,2-*a*]pyrazolones (**4**), we have captured a new N-N bond activation product tetrahydropyrimidinones (**3**) and an unusual product 3-acylindoles (**5**). These findings provide a versatile strategy for synthesizing diverse and biologically active nitrogen-containing heterocycles.

## Results and discussion

We initially investigated this transformation by employing the 4,4-dimethyl-1-phenylpyrazolidin-3-one (**1a**, 0.24 mmol) and 3-hydroxy-3-(phenylethynyl)isoindolin-1-one (**2a**, 0.2 mmol) as model substrates, and the results showed that the desired N-N bond activation product **3aa** could be constructed with a 20% yield under [Cp\*RhCl<sub>2</sub>]<sub>2</sub> (5 mol%) as the catalyst and 1,2-dichloroethane (DCE) as the solvent at 90 °C for 4 h (Table 1, entry 1). Surprisingly, adding silver acetate and acetic acid into the catalytic system would result in a new [4 + 1] annulation

product **4aa** with 8% yield (Table 1, entry 2), while the addition of cesium pivalate and acetic acid into the catalytic system simultaneously formed **4aa** (28% yield) and a [3 + 2] annulation product **5aa** (10% yield) (Table 1, entry 3). Interestingly, the yield of **5aa** could be increased to 50% when we used dioxane as the solvent (Table 1, entry 4). Encouraged by these results, we further screened different catalysts, including Rh<sub>2</sub>(esp)<sub>2</sub>, CyJohnPhos AuCl, (Ph<sub>3</sub>P)AuCl, Ph<sub>3</sub>PAuNTf<sub>2</sub>, CyJohnPhos AuNTf<sub>2</sub> and AuCl(IPr) (Table 1, entries 5–10). The results showed that the AuCl(IPr) offered the best catalytic activity (Table 1, entry 10) for the formation of **3aa**, the yield of which could be increased to 54%. Subsequent solvent screening indicated that DCE was the most conducive to this transformation (Table 1, entries 11–13). Next, lowering the reaction temperature to 70 °C could further increase the yield to 65% (Table 1, entry 14), while increasing the reaction temperature to 100 °C resulted in a slight decrease in yield (Table 1, entry 15). In brief, the optimal reaction conditions for **3aa** were as follows: **1a** (0.24 mmol), **2a** (0.2 mmol), and AuCl(IPr) (5 mol%) in DCE at 70 °C for 4 h (condition A). Likewise, the optimal reaction conditions for **4aa** and **5aa** were also separately investigated (see Tables S1 and S2†). The results displayed that when using **1a** (0.24 mmol), **2a** (0.2 mmol), [Cp\*Rh(CH<sub>3</sub>CN)<sub>3</sub>](SbF<sub>6</sub>)<sub>2</sub> (5 mol%), LiOAc (0.2 mmol), BzOH (0.1 mmol) at 100 °C for 8 h (Table 1, entry 16, condition B) and **1a** (0.2 mmol), **2a** (0.24 mmol), [Cp\*RhCl<sub>2</sub>]<sub>2</sub> (5 mol%), CsOPiv (0.2 mmol) and AcOH (0.2 mmol)

Table 1 Optimization of reaction conditions<sup>a</sup>



Entry	Catalytic system	Solvent	Yield (%)		
			3aa	4aa	5aa
1	[Cp*RhCl <sub>2</sub> ] <sub>2</sub>	DCE	20	—	—
2 <sup>b</sup>	[Cp*RhCl <sub>2</sub> ] <sub>2</sub> /AgOAc/AcOH	DCE	—	8	—
3 <sup>b</sup>	[Cp*RhCl <sub>2</sub> ] <sub>2</sub> /CsOPiv/AcOH	DCE	—	28	10
4 <sup>b</sup>	[Cp*RhCl <sub>2</sub> ] <sub>2</sub> /CsOPiv/AcOH	Dioxane	—	—	50
5	Rh <sub>2</sub> (esp) <sub>2</sub>	DCE	44	—	—
6	CyJohnPhos AuCl	DCE	31	—	—
7	CyJohnPhos AuNTf <sub>2</sub>	DCE	44	—	—
8	(Ph <sub>3</sub> P)AuCl	DCE	44	—	—
9	Ph <sub>3</sub> PAuNTf <sub>2</sub>	DCE	36	—	—
10	AuCl(IPr)	DCE	54	—	—
11	AuCl(IPr)	MeCN	15	—	—
12	AuCl(IPr)	CHCl <sub>3</sub>	41	—	—
13	AuCl(IPr)	HFIP	Trace	—	—
14 <sup>c</sup>	AuCl(IPr)	DCE	65	—	—
15 <sup>d</sup>	AuCl(IPr)	DCE	50	—	—
16 <sup>e</sup>	[Cp*Rh(CH <sub>3</sub> CN) <sub>3</sub> ](SbF <sub>6</sub> ) <sub>2</sub> /LiOAc/BzOH	CH <sub>3</sub> CN	—	60	Trace
17 <sup>f</sup>	[Cp*RhCl <sub>2</sub> ] <sub>2</sub> /CsOPiv/AcOH	Diglyme	—	—	80


<sup>a</sup> Reaction conditions: **1a** (0.24 mmol), **2a** (0.2 mmol), catalyst (5.0 mol%) in solvent (4 mL) at 90 °C for 4 h, NMR yield (1,3,5-trimethoxybenzene as the internal standard). <sup>b</sup> **1a** (0.2 mmol), **2a** (0.24 mmol), inorganic salts (0.4 mmol), AcOH (0.2 mmol) at 80 °C for 8 h, isolated yield. <sup>c</sup> 70 °C. <sup>d</sup> 100 °C. <sup>e</sup> **1a** (0.24 mmol), **2a** (0.2 mmol), [Cp\*Rh(CH<sub>3</sub>CN)<sub>3</sub>](SbF<sub>6</sub>)<sub>2</sub> (5 mol%), LiOAc (0.2 mmol), BzOH (0.1 mmol) at 100 °C for 8 h, isolated yield. <sup>f</sup> **1a** (0.2 mmol), **2a** (0.24 mmol), [Cp\*RhCl<sub>2</sub>]<sub>2</sub> (5 mol%), CsOPiv (0.2 mmol), AcOH (0.2 mmol) at 80 °C for 2 h, isolated yield.




in diglyme at 80 °C for 2 h (Table 1, entry 17, condition C) as their respective catalytic systems, the products **4aa** and **5aa** were effectively prepared with 60% and 80%, respectively. Their structures were confirmed by  $^1\text{H}$  and  $^{13}\text{C}$  NMR, MS and X-ray spectra.

With the optimized reaction conditions in hand, we firstly conducted a comprehensive investigation into the scope of pyrazolidinones **1** and coupling partners **2** under the optimal reaction condition to build diversified tetrahydropyrimidinones **3**. As shown in Scheme 2, whether introducing halogen atoms (F, Cl and Br) or electron-donating groups ( $\text{CH}_3$  and  $\text{OCH}_3$ ) into 4-position of benzene ring of substrate **1**, the reaction efficiencies were maintained at a moderate yield (Scheme 2, **3ba–3fa**, 36–64%). However, introducing an F atom into 3-position of phenyl ring resulted in a decrease in yield (**3ia**, 25%). Subsequently, we investigated the effect of various substituents on the coupling partners **2**. Notably, substrates **2** bearing halogen atoms (F, Cl and Br) or electron-donating groups ( $\text{CH}_3$ ,  $\text{OCH}_3$  and  $(\text{CH}_2)_4\text{CH}_3$ ) at the 4-position of the phenyl ring afforded corresponding products in moderate to good yields (**3ab–3ag**, 49–72%). Introducing  $\text{CH}_3$  and Cl groups into the 3-position of the phenyl ring also gave satisfactory results (**3ai–3ak**, 69–72%). Replacing the phenyl ring of **2** with the thiophene ring also provided a good yield (**3al**, 66%).

During the reaction condition screening, we observed the pyrazolo[1,2-*a*]pyrazolone derivative **4aa** could be selectively



Scheme 2 The investigation of substrate scope for the synthesis of **3**. General reaction conditions (condition A): **1** (0.24 mmol), **2** (0.2 mmol),  $\text{AuCl}(\text{IPr})$  (5 mol%) in DCE (4.0 mL) at 70 °C under air for 4 h, isolated yields.




Scheme 3 The investigation of substrate scope for the synthesis of **4**. General reaction conditions (condition B): **1** (0.24 mmol), **2** (0.2 mmol),  $[\text{Cp}^*\text{Rh}(\text{CH}_3\text{CN})_3](\text{SbF}_6)_2$  (5 mol%), LiOAc (0.2 mmol), BzOH (0.1 mmol) in MeCN (4.0 mL) at 100 °C under air for 8 h, isolated yields.

constructed *via* a  $\text{C}(\text{sp}^2)\text{–H}$  bond activation/[4 + 1] annulation cascade. As shown in Scheme 3, the reaction worked well with electron-withdrawing and electron-donating groups at the 4-position of the phenyl ring of substrates **1**, affording **4ba–4ha** with 47–71% yields (Scheme 3). Translocating Cl or Br to the 3-position of the phenyl ring could also successfully generate corresponding products with 45–68% yields (**4ia–4ka**). Next, various substituted groups at the 4-position of the phenyl ring of substrate **2** were explored, and the results indicated that all substrates could react smoothly with **1a** to provide the corresponding products **4ab–4ag** in moderate to good yields (40–60%). Likewise, introducing  $\text{CH}_3$  or Cl into the *meta* or *ortho* position of the benzene ring was also satisfactory (**4ai–4ak**). However, replacing benzene ring of **2** with thiophene ring gave a relatively low yield (**4al**, 21%).

Another discovery was that we successfully obtained previously unreported 3-acylindoles *via* a  $\text{C}(\text{sp}^2)\text{–H}$  bond activation and [3 + 2] annulation cascade from pyrazolidinone **1** and its coupling partner **2**. As shown in Scheme 4, whether introducing electron-donating or electron-withdrawing groups into pyrazolidinones **1** or 3-alkynyl-3-hydroxyisindolinones **2** could






**Scheme 4** The investigation of substrate scope for the synthesis of **5**. General reaction conditions (condition C): **1** (0.2 mmol), **2** (0.24 mmol), [Cp\*RhCl<sub>2</sub>]<sub>2</sub> (5 mol%), CsOPiv (0.2 mmol), AcOH (0.2 mmol) in diglyme (4.0 mL) at 80 °C under air for 2 h, isolated yields.

generate desired products in good yields (**5ba–5fa**, 81–91%; **5ab–5ak**, 62–72%). In particular, in this transformation, replacing the benzene ring of **2** with the thiophene ring could provide a good yield (Scheme 4, **5al**, 75%). Additionally, we also attempted to introduce the group into R<sup>3</sup> position of substrate **2**, the corresponding product **5am** was obtained with good yield.

Given the potential application of these intriguing heterocycles, the gram-scale preparations for the products **3aa**, **4aa**, and **5aa** were performed with 61%, 47%, and 67% isolated yields, respectively (Scheme 5a). Then, the synthetic transformations of compounds **4aa** and **5aa** were also explored. For example, the C–C double bond of **4aa** could be selectively reduced using hydrogen gas and Pd/C, generating the desired product **6** with 80% yield (Scheme 5b). Interestingly, we successfully fulfilled a simultaneous transformation of two amide groups in **5aa**, in which compound **5aa** could be converted into nitrile product **7** with 84% yield in the presence of cyanuric chloride (Scheme 5b). Additionally, treatment of **5aa** with triethylsilane and trifluoroacetic acid afforded a ring-closed product **8** (Scheme 5b).

Having established the substrate scope and utility of the product, we then conducted a series of experiments to explore the preliminary mechanism (Scheme 6). Firstly, 95% of *ortho* H could be deuterated by treating **1a** with D<sub>2</sub>O under standard condition B, 25% of the *ortho* H could be deuterated by treating **1a** with D<sub>2</sub>O under standard condition C, suggesting the reversibility of the C(aryl)–H bond cleavage during the




**Scheme 5** Gram-scale preparations and conversions.

formation of products **4** and **5** (Scheme 6a). Next, the kinetic isotope effect (KIE) values of 1.70 for **4aa** and 1.17 for **5aa** indicated that the C–H bond cleavage was likely not the turnover-limiting step (Scheme 6b). Moreover, three competitive experiments within three reaction conditions by introducing different substrates with electron-withdrawing and electron-donating groups were carried out. Treating a 1:1 mixture of fluorine/methoxy substituted pyrazolidinone (**1b/1f**) with **2a** under conditions A and C, respectively, showed that there is no apparent difference in the reaction rate between two substrates **1** with electron-withdrawing and electron-donating groups (Scheme 6c). These results indicated that these different groups are well tolerated in these transformations. However, we observed a high ratio (8.0) between **4ga** and **4fa** when treating a 1:1 mixture of trifluoromethyl/methoxy substituted pyrazolidinone (**1g/1f**) with **2a** under condition B (Scheme 6c), suggesting that the electron-withdrawing substrates reacted faster in this transformation.


Based on the above observations and literature precedents,<sup>18,21,24,28–32</sup> three plausible catalytic mechanisms are proposed for N–N bond activations and C–H functionalizations. As shown in Scheme 7a, the propargylic alcohol **2a** initially forms the intermediate **A** by coordinating with the Au(I) catalyst. Upon the entry of the pyrazolidinone **1a** into the system, it





Scheme 6 Preliminary mechanism studies.

undergoes alkyne hydroamination with Au(I) to yield complex **B**, which further forms the intermediate **C** after the intramolecular electrophilic addition. Finally, elimination of the Au(I) catalyst and the hydroxy group from **C** gives the desired product **3aa**. The speculated mechanisms of the desired products **4aa** and **5aa** are shown in Scheme 7b. The NH directing group of **1a** assists the activated Rh catalyst with the cleavage of the *ortho* C(sp<sup>2</sup>)-H bond, enabling the formation of a five-membered rhodacycle **D**. Next, the complexation of Rh(III) with the triple bond of **2a** gives intermediate **E**. The following regioselective insertion of the alkynyl unit of **2a** into the C-Rh bond of **E** generates intermediate **F**, which is most likely driven by additional coordination of the OH group with Rh(III) to overcome the unfavorable steric interactions between the isoindolin-1-one group and the Cp ligand. Further, the cleavage of the C-Rh bond and elimination of water under the assistance of benzoic acid offers the allenyl intermediate **G**. Finally, **G** undergoes nucleophilic addition and followed by protodemetalation to form target product **4aa**. Specially, using [Cp\*RhCl<sub>2</sub>]<sub>2</sub> as the catalyst in diglyme results in the opposite regioselective insertion of the alkyne unit of **2a** into the C-Rh bond of **E** to form intermediate **H**, probably because the solvent prevents the additional coordination of the OH group of propargylic alcohol with Rh(III) complex. Then protonation of intermediate **H** leads to the generation of intermediate **I**, which then undergoes an oxidative addition with N-N bond to form intermediate **J**. Finally, intermediate **J** undergoes a reductive elimination and a following *N*-demethylation/protonation to produce the desired product **5aa**.



Scheme 7 Plausible mechanisms.

## Conclusions

In summary, we have successfully fulfilled a catalytic system-controlled divergent reaction strategy to construct three types of nitrogen-containing heterocyclic skeletons through chemical bond activation/annulation routes of pyrazolidinones with 3-alkynyl-3-hydroxyisoindolinones. More importantly, this is the first capture of pyrazolidinones and propargyl alcohols to construct tetrahydropyrimidinones and unusual 3-acylindoles. These strategies exhibit a broad range of substrates, moderate to good yields, and valuable transformations, offering structural and biological potentials for further drug discovery.

## Data availability

The data supporting this article have been included as part of the ESI,<sup>†</sup> which includes experimental details, characterization data, NMR spectra, and HPLC spectra, along with single-crystal X-ray data for compounds **3aa** (CCDC 2410303), **4aa** (CCDC 2410625), and **5aa** (CCDC 2410305).

## Conflicts of interest

The authors declare no competing interests.



## Acknowledgements

We thank the National Natural Science Foundation of China (82173656, 82373709, 82151219), Shandong Laboratory Program (SYS202205), and China Postdoctoral Science Foundation (2023M730821) for financial support.

## References

- 1 C. Wallwey and S.-M. Li, Ergot alkaloids: structure diversity, biosynthetic gene clusters and functional proof of biosynthetic genes, *Nat. Prod. Rep.*, 2011, **28**, 496–510.
- 2 C. Yang, F. Song, J. Chen and Y. Huang, Construction of Pyridazine Analogues via Rhodium-mediated C-H Activation, *Adv. Synth. Catal.*, 2017, **359**, 3496–3502.
- 3 W. Zi, Z. Zuo and D. Ma, Intramolecular dearomative oxidative coupling of indoles: a unified strategy for the total synthesis of indoline alkaloids, *Acc. Chem. Res.*, 2015, **48**, 702–711.
- 4 Z. Zhang, H. Jiang and Y. Huang, Ruthenium-catalyzed redox-neutral C-H activation via N-N cleavage: synthesis of N-substituted indoles, *Org. Lett.*, 2014, **16**, 5976–5979.
- 5 Z. Yang, Q. Yue, M. Yang, H. Zhang and X. Cui, Ru(II)-Catalyzed Tunable Cascade Reaction via C-H/C-C Bond Cleavage, *J. Org. Chem.*, 2020, **85**, 12960–12970.
- 6 Z. Fan, H. Lu, W. Li, K. Geng and A. Zhang, Rhodium-catalyzed redox-neutral coupling of phenidones with alkynes, *Org. Biomol. Chem.*, 2017, **15**, 5701–5708.
- 7 N. Strašek, L. Lavrenčič, A. Oštrek, D. Slapšak, U. Grošelj, M. Klemenčič, H. Brodnik Žugelj, J. Waggener, M. Novinec and J. Svete, Tetrahydro-1H,5H-pyrazolo[1,2-a]pyrazole-1-carboxylates as inhibitors of Plasmodium falciparum dihydroorotate dehydrogenase, *Bioorg. Chem.*, 2019, **89**, 102982.
- 8 L. Yang, Y. Lv, F. Wang and G. Zhong, Chiral NHC-catalyzed 1,3-dipolar [3 + 2] cycloaddition of azomethine imines with  $\alpha$ -chloroaldehydes for the synthesis of bicyclic pyrazolidinones, *Org. Biomol. Chem.*, 2018, **16**, 4433–4438.
- 9 R. Na, C. Jing, Q. Xu, H. Jiang, X. Wu, J. Shi, J. Zhong, M. Wang, D. Benitez, E. Tkatchouk, W. A. Goddard, H. Guo and O. Kwon, Phosphine-catalyzed annulations of azomethine imines: allene-dependent [3 + 2], [3 + 3], [4 + 3], and [3 + 2 + 3] pathways, *J. Am. Chem. Soc.*, 2011, **133**, 13337–13348.
- 10 M. P. Clark, S. K. Laughlin, M. J. Laufersweiler, R. G. Bookland, T. A. Brugel, A. Golebiowski, M. P. Sabat, J. A. Townes, J. C. VanRens, J. F. Djung, M. G. Natchus, B. De, L. C. Hsieh, S. C. Xu, R. L. Walter, M. J. Meikel, S. A. Heitmeyer, K. K. Brown, K. Juergens, Y. O. Taiwo and M. J. Janusz, Development of orally bioavailable bicyclic pyrazolones as inhibitors of tumor necrosis factor- $\alpha$  production, *J. Med. Chem.*, 2004, **47**, 2724–2727.
- 11 W. Lewgowd and A. Stanczak, Cinnoline derivatives with biological activity, *Arch. Pharm.*, 2007, **340**, 65–80.
- 12 V. A. Assimon, Y. Tang, J. D. Vargas, G. J. Lee, Z. Y. Wu, K. Lou, B. Yao, M. K. Menon, A. Pios, K. C. Perez, A. Madriaga, P. K. Buchowiecki, M. Rolfe, L. Shawver, X. Jiao, R. Le Moigne, H. J. Zhou and D. J. Anderson, CB-6644 Is a Selective Inhibitor of the RUVBL1/2 Complex with Anticancer Activity, *ACS Chem. Biol.*, 2019, **14**, 236–244.
- 13 J. Velcicky, U. Bodendorf, P. Rigollier, R. Epple, D. R. Beisner, D. Guerini, P. Smith, B. Liu, R. Feifel, P. Wipfli, R. Aichholz, P. Couttet, I. Dix, T. Widmer, B. Wen and T. Brandl, Discovery of the First Potent, Selective, and Orally Bioavailable Signal Peptide Peptidase-Like 2a (SPPL2a) Inhibitor Displaying Pronounced Immunomodulatory Effects In Vivo, *J. Med. Chem.*, 2018, **61**, 865–880.
- 14 F. A. Romero, A. M. Taylor, T. D. Crawford, V. Tsui, A. Côté and S. Magnuson, Disrupting Acetyl-Lysine Recognition: Progress in the Development of Bromodomain Inhibitors, *J. Med. Chem.*, 2016, **59**, 1271–1298.
- 15 H. Li, M. Shen, B. Li, X. Zhang and X. Fan, Solvent-Dependent Selective Synthesis of CF(3)-Tethered Indazole Derivatives Based on Multiple Bond Activations, *Org. Lett.*, 2023, **25**, 720–725.
- 16 X.-R. Song, R. Yang and Q. Xiao, Recent Advances in the Synthesis of Heterocyclics via Cascade Cyclization of Propargylic Alcohols, *Adv. Synth. Catal.*, 2021, **363**, 852–876.
- 17 P. Sihag and M. Jeganmohan, Regioselective Synthesis of Isocoumarins via Iridium(III)-Catalyzed Oxidative Cyclization of Aromatic Acids with Propargyl Alcohols, *J. Org. Chem.*, 2019, **84**, 2699–2712.
- 18 Y. Xu, M. Shen, X. Zhang and X. Fan, Selective Synthesis of Pyrazolo[1,2-a]pyrazolones and 2-Acylindoles via Rh(III)-Catalyzed Tunable Redox-Neutral Coupling of 1-Phenylpyrazolidinones with Alkynyl Cyclobutanols, *Org. Lett.*, 2020, **22**, 4697–4702.
- 19 X. Wu and H. Ji, Rhodium-Catalyzed [4 + 1] Cyclization via C-H Activation for the Synthesis of Divergent Heterocycles Bearing a Quaternary Carbon, *J. Org. Chem.*, 2018, **83**, 4650–4656.
- 20 L. Zhang, Y. Xu, X. Zhang, X. Zhang and X. Fan, Synthesis of pyrazolone fused benzodiazepines via Rh(III)-catalyzed [4 + 3] annulation of 1-phenylpyrazolidinones with propargyl alcohols, *Org. Chem. Front.*, 2020, **7**, 2284–2290.
- 21 T. Li, Z. Yang, Z. Song, R. Chauvin and X. Cui, Rhodium(III)-Catalyzed [4+3] Annulation of N-Aryl-pyrazolidinones and Propargylic Acetates: Access to Benzo[c][1,2]diazepines, *Org. Lett.*, 2020, **22**, 4078–4082.
- 22 F. Fang, C. Zhang, C. Zhou, Y. Li, Y. Zhou and H. Liu, Rh(III)-Catalyzed C-H Activation of Benzoylacetonitriles and Tandem Cyclization with Diazo Compounds to Substituted Benzo[de]chromenes, *Org. Lett.*, 2018, **20**, 1720–1724.
- 23 X. Wu, B. Wang, S. Zhou, Y. Zhou and H. Liu, Ruthenium-Catalyzed Redox-Neutral [4 + 1] Annulation of Benzamides and Propargyl Alcohols via C-H Bond Activation, *ACS Catal.*, 2017, **7**, 2494–2499.
- 24 F. Fang, S. Hu, C. Li, Q. Wang, R. Wang, X. Han, Y. Zhou and H. Liu, Catalytic System-Controlled Divergent Reaction Strategies for the Construction of Diversified Spiropyrazolone Skeletons from Pyrazolidinones and Diazopyrazolones, *Angew. Chem., Int. Ed.*, 2021, **60**, 21327–21333.



- 25 H. Wang, Y. Li, L. Huang, H. Xu, Y. Jiao, Z. Zhou, Z. Tang, F. Fang, X. Zhang, K. Ding, W. Yi, H. Liu, X. Wu and Y. Zhou, The synthesis of spirocyclopropane skeletons enabled by Rh(III)-catalyzed enantioselective C–H activation/[4 + 2] annulation, *Chem Catal.*, 2023, **3**, 100822.
- 26 J. Gao, K. Luo, X. Wei, H. Wang, H. Liu and Y. Zhou, Rh(III)-Catalyzed Successive C–H Activations of 2-Phenyl-3H-indoles and Cyclization Cascades to Construct Highly Fused Indole Heteropolycycles, *Org. Lett.*, 2023, **25**, 3341–3346.
- 27 X. Hou, R. Wang, F. Fang, Z. Qu, J. Zhou, T. Yu, D. Wang, H. Liu and Y. Zhou, Rh(III)-Catalyzed C–H Activation/Annulation for the Construction of Quinolizinones and Indolizines, *Org. Lett.*, 2024, **26**, 4451–4456.
- 28 Y. Yang, N. Li, J. Zhao, Y. Jiang, X. Zhang and X. Fan, Selective Synthesis of 3-( $\alpha$ -Fluorovinyl)indoles and 3-Acylindoles via the Cascade Reactions of 1-Phenylpyrazolidinones with  $\alpha,\alpha$ -Difluoromethylene Alkynes, *Adv. Synth. Catal.*, 2021, **363**, 3600–3606.
- 29 Z. Yang, Z. Song, L. Jie, L. Wang and X. Cui, Iridium(III)-catalysed annulation of pyrazolidinones with propiolates: a facile route to pyrazolo[1,2-*a*] indazoles, *Chem. Commun.*, 2019, **55**, 6094–6097.
- 30 D. Campeau, D. F. León Rayo, A. Mansour, K. Muratov and F. Gagosz, Gold-Catalyzed Reactions of Specially Activated Alkynes, Allenes, and Alkenes, *Chem. Rev.*, 2021, **121**, 8756–8867.
- 31 A. S. K. Hashmi, Gold-Catalyzed Organic Reactions, *Chem. Rev.*, 2007, **107**, 3180–3211.
- 32 S. K. Yadav, B. Ramesh and M. Jeganmohan, Cobalt(III)-Catalyzed Chemo- and Regioselective [4 + 2]-Annulation of Aromatic Sulfoxonium Ylides with 1,3-Diynes, *J. Org. Chem.*, 2022, **87**, 4134–4153.

