RSC Advances



REVIEW

View Article Online
View Journal | View Issue



Cite this: RSC Adv., 2025, 15, 18034

Vistas in the domain of 3-acetyl-4-hydroxy-2-quinolinone derivatives (AHQ) and their applications

Nedaa N. Elnaggar, Wafaa S. Hamama ** and Eslam A. Ghaith ** ab

This review discusses the significant advances in the current status and latest synthesis techniques for *N*-substituted 3-acetyl-4-hydroxyquinolinones. They are an important class of alkaloids possessing different electrophilic and nucleophilic centers. In this review, we comprehensively summarize the synthesis of 3-acetylquinolones *via* various reactions, emphasizing recent developments and challenges. Synthetic transformations of quinolone are focused on electrophilic substitution reactions supported by their mechanistic pathways, as well as nucleophilic substitution reactions and cycloaddition reactions, which have allowed access to an extensive scope of binary and fused heterocyclic scaffolds, including pyrazoles, imidazoles, tetrazoles, pyridines, pyrimidines, triazines, azepines, and pyranones. Also, the utilization of quinolone reactions as key intermediates in the ongoing and forthcoming marketed pharmaceutical syntheses is discussed. Additionally, this review covers various potential applications, including complexation, fluorescence sensing, chemical pH sensors, agricultural, and anticorrosion.

Received 22nd April 2025 Accepted 20th May 2025

DOI: 10.1039/d5ra02813b

rsc.li/rsc-advances

Introduction and scope

Heterocyclic skeletons which contain a nitrogen atom represent a common unit of a vast amount of marketed drugs, constituting the most extensive and diverse collection of natural, synthetic organic compounds, agrochemicals, and pharmaceuticals.2-4 Furthermore, the chemistry of heterocycles aggregates an essential branch of the field of drug design and the evolution of novel biologically active compounds,5-11 including antiparasitic, 12,13 anti-HIV, 14,15 antiepileptic drugs, 16,17 analgesic, 18,19 anthelmintic, 20,21 inhibitors of platelet aggregation, antiinflammatory,22-25 antibacterial and antiviral drugs,26-29 atherosclerosis, 30,31 inhibitors for Parkinson's disease, 32 proliferative, pulmonary antifibrotic agents³³ and antitumor.³⁴⁻³⁶ Additionally, they provide one of the most fruitful sources for drug discovery and development, owing to synthesizing various scaffolds via robust synthetic approaches. 37-40 These are important in synthesizing dyes, pigments, and polymeric materials 41-43 due to their optical and fluorescence properties.44 The ongoing research in synthetic organic chemistry is dedicated to advancing these goals and addressing the demand for improved and sustainable approaches in synthesizing aromatic heterocyclic compounds.2-4 Easily accessible nitrogenous scaffolds are fundamental feedstocks; among these, quinoline and quinolones

have piqued the interest of scientists since the 19th century, as over 600 quinoline derivatives have been extracted from natural sources to date.

Quinolinones have been perceived for their fascinating structural notions and versatile applications. 45-56 Also, the quinolinones structures are pivotal building blocks, 46 as they are commonly conspicuous scaffolds in a wide range of pharmacologically active synthetic and natural compounds. 57-68 Quinolin-2-one, also referred to as benzo[b]pyridine, and 1-azanaphthalene, is important classes of natural product as N-based heterocycles which extracted from Toddalia asiatica leaves and plants, marine organisms and microorganisms which containing the quinolinone derivatives alkaloid findersine, 52,69-75 consists of a benzene ring fused with a pyridine ring, sharing two carbon atoms between them and nitrogen atom not present as ring junction atom.

Quinoline is a weak base capable of forming salts with acids. It reacts similarly to pyridine and benzene and undergoes either nucleophilic or electrophilic substitution processes. The importance of the quinoline moiety is being a vital component in many naturally occurring heterocyclic and medicinal plant families. In addition, bucharidine 1 and foliosidine 2 are quinoline alkaloids that are extracted from *Haplophyllum foliosum* and *Haplophyllum bucharicum*, respectively (Fig. 1). Also, both of them have viral-RNA polymerase inhibitory action and estrogenic action, such as compounds 3 and 4, which strongly inhibit the replication of the hepacivirus C. 82-84

Additionally, quinolines have various potential biological and pharmacological activities including antimalarial,^{84–86} antibacterial,⁸⁷ antifungal,^{88,89} anticancer,^{90–92} anti-HIV (Human

^aChemistry Department, Faculty of Science, Mansoura University, Mansoura, 35516, Egypt. E-mail: wshamama@mans.edu.eg; wshamama53@gmail.com; Tel: +20-1093427024

^bChemistry Department, Faculty of Science, New Mansoura University, New Mansoura City, Egypt

Fig. 1 Some natural alkaloids quinolinone-based molecules.

Fig. 2 Chemical structure of biologically active quinolinone scaffolds.

Immunodeficiency Virus), ⁹³ antiviral, ⁹⁴ antitumor, ^{95–97} anthelmintic, ⁹⁸ antioxidant, ⁴⁹ cardiotonic, ^{76,99–101} anticonvulsant, ^{102,103} anti-inflammatory and analgesic properties ^{76,77,104,105} (Fig. 2).

2. Chemical reactivity

2.1. 3-Acetyl-*N*-substituted/(un)-4-hydroxy-quinolin-2(1*H*) one derivatives (AHQ)

AHQ is the basic ring structure of numerous alkaloids and is a highly versatile scaffold with an extensive range of applications. Progress in synthetic technologies is leading to the production of various functionalized quinolinone derivatives as biosteric analogues to linomide (4-hydroxyquinolinone) derivatives. 82,106-112

2.2. Structural feature of AHQ

2.2.1. Tautomeric structures. Whereas, the β , β' -tricarbonyl (TC) group in AHQ compound provides an analogue of 3-acyl pyrrolidine-2,4-diones (tetramic acid), and has suitable sites for interaction with different nucleophilic reagents, with its

Fig. 3 Tautomeric form of AHQ 12a and its geometrical structures.

Table 1 Spectral data of AHQ 12a-c

Spectroscopic technique	OH O 5 4a 4 4 9 10 8 8 N O 12a	OH O N O CH ₃ 12b	OH O NO Ph 12c
¹ H NMR (DMSO- <i>d</i> ₆)	2.72 (s, 3H, COCH ₃), 7.23 (t, 1H, H-6), 7.30 (d, 1H, H-8, $J = 8.3$ Hz), 7.65 (t, 1H, H-7), 7.99 (dd, 1H, H-5, $J = 8.0$, $J = 1.2$ Hz), 11.53 (s, 1H, NH) and 17.04 (s, 1H, OH) ¹¹⁶	2.79 (s, 3H, COCH ₃), 3.52 (s, 3H, N-CH ₃), 7.30 (t, 1H, H-6), 7.50 (d, 1H, H-8, $J = 8.1$ Hz), 7.78 (t, 1H, H-7), 7.96 (dd, 1H, H-5, $J = 8.4$ Hz, $J = 1.5$ Hz) and 17.04 (s, 1H, OH) ¹¹⁶	2.77 (3H, s, 3-COCH ₃), 7.12–7.41 (9H _{Arom}), 8.23 (s, OH) ¹¹⁷ Our work 2.79 (s, 3H, COCH ₃ , 6.58 (d, 1H, $J = 9$ Hz), 7.23 (m, 1H), 7.27 (d, 2H, $J = 8$ Hz), 7.45 (t, 1H, $J = 8$ Hz), 7.53 (t, 1H, $J = 7$ Hz), 7.61 (t, 2H, $J = 8$ Hz), 8.24 (d, 1H, $J = 8$ Hz), 17.16 (s, 1H, OH)
13 C NMR (DMSO- d_6)	205.7 (C-9), 174.7 (C-4), 161.1 (C-2), 140.5 (C-8a), 134.8 (C-7), 124.7 (C-5), 122.0 (C-6), 115.5 (C-8), 113.3 (C-4a), 105.7 (C-3) and 30.5 (C-10) ¹¹⁶	206.7 (C-9), 173.3 (C-4), 160.6 (C-2), 141.6 (C-8a), 135.7 (C-7), 125.4 (C-5), 122.3 (C-6), 114.4 (C-8), 115.3 (C-4a), 105.7 (C-3), 31.3 (C-10) and 28.9 (N-CH ₃) ¹¹⁶	206.9 (C-9), 175.0 (C-4), 161.9 (C-2), 142 (1C), 137.4 (1C), 134.4 (1C), 130.3 (2C), 129.0 (2C), 125.8 (1C), 122.3 (1C), 117.7 (1C), 116.0 (1C), 115.2 (1C), 106.0 (1C), 31.4 (1C)
IR (KBr, $\nu_{\text{max/cm}}^{-1}$)	3360 (OH), 3160 (NH), 1661 (C=O, acetyl), 1622 (C=O, amide), 1606 (C=C) ¹¹⁶	3250 (OH), 1658 (C=O, acetyl), 1623 (C=O, amide 1598 (C=C) ¹¹⁷	3072 (CH _{Arom}), 2924 (CH _{Aliph}), 1654(C=O, acetyl, enol form), 1618 (C=O, amide). ^{114,118}




Fig. 4 X-ray structural analysis 12a.120

regioselectivity of TC for AHQ [(A)-(D)] given below in a framework of molecular orbital perturbation.¹¹³

Detsi and coworkers¹¹⁴ predicting the four tautomeric structures A-D. of AHQ [R=H] **12a** (Fig. 3). The tautomeric equilibrium showed the presence of tautomeric forms that could be in one of four possible structures, A-D (Fig. 3). Whereas, semiempirical quantum calculations showed that the tautomer D was the most stable favorable one according to the total energy values.^{114,115} Whereas, the constitutions of AHQ

derivatives were fully elucidated via different spectroscopic analyses as shown in Table 1. $^{114,116-118}$

2.2.2. X-ray crystallographic. The constitution of AHQ 12a was supported by X-ray analysis, as the non-hydrogen atoms in compound 12a are located in a single plane with an accuracy of 0.02 Å, this phenomena indicated the formation of an intramolecular hydrogen bond $O(3)-H(3O)\cdots O(2)$ (H···O 1.41 Å, O-H···O 157°). Also, this phenomenon displayed the reestimation of the intramolecular electronic charge as shown




Fig. 5 Common synthetic methods of quinolines.

by the bond lengths O(2)–C(10) 1.255(2) and C(7)–C(8) bonds 1.393(2) (Fig. 4). 119,120

3. Aspects of the AHQ synthetic approach

Quinolines can be synthesized via different synthetic strategies involving [3 + 3], [4 + 2] and cyclization methodologies (I-III).¹²¹

(I) [3 + 3] Annulation: This strategy includes established procedures, for instance, the Skraup, Conrad-Limpach, Doebner-von Miller, Combes, and Gould-Jacobs syntheses. However, the drawback of these techniques is that they are restricted in regioselectivity for the synthesis of multisubstituted quinolines.¹²¹⁻¹²³

(II) [4 + 2] Annulation: There are two techniques to build pyridine nuclei annulated to the benzene ring as shown in designed forms (II-a and II-b), as the first one has minimal regioselectivity with a narrow substrate range. ^{121,124,125} While the other method is more prevalent and employs easily available substrates to produce products with high regioselectivity as Ptzinger and Friedlander reactions. ^{121,122,126,127}

(III) Cyclization: There are four distinct ways to generate the pyridine ring of quinolines (III-a to III-d) using cyclization. However, they have not been achieved due to the complicated reaction methodology and the limited availability of the starting substrates (Fig. 5). ^{128,129}

Synthetic approaches to construct AHQ derivatives have been achieved *via* many routes: aniline, alkaline hydrolysis of pyranoquinoline, anthranilic acid derivatives, heterocyclization of 3,1-benzoxazine-4-one and furo[3,2-*c*]quinoline.

3.1. Synthesis from amines/pyrano[3,2-c]quinolindione

Quinoline derivatives 12a-f were prepared by cyclocondensation of substituted amines and diethyl malonate in a molar ratio of 1: 2 for 3-6 h. Till the calculated amount of ethanol was collected using Dean Stark by a short Vigreux column as distillation system afforded the lactone 11, followed by alkaline hydrolysis [NaOH (2 N)], of lactone 11 and subsequent decarboxylation of the intermediate β -oxocarboxylate then acidified with HCl (2 N) to yield the products 12a-f in high yields (86-96%). In the case of 12b, compound 11 was synthesized following the methods of Kappe and Stadlbauer, which involved reacting N-methylaniline with diethyl malonate in diphenyl ether. However, Kappe's method utilized an excess of diethyl malonate, which served a dual purpose as both a reagent and a solvent (Scheme 1).115,130-135 Also, the ring opening of 11b-f by NaOH and subsequent spontaneous decarboxylation afforded 4-hydroxyquinolinone derivatives 12b-f (Scheme 1).112,133,135-137

3.2. Synthesis from 4-hydroxyquinolin-2-ones

Cyclocondensation reaction of secondary amine and diethyl malonate afforded 4-hydroxyquinolin-2-ones **15a–d.** After that, acetylation reaction of *N*-unsubstituted AHQ **15a–d** with acetyl chloride using acetic acid or pyridine in the presence of polyphosphoric acid (PPA) or AcOH with phosphorus oxychloride (POCl₃) yielded acetylquinoline derivatives **12a–d** (Scheme 1). ^{135–140}

3.3. Synthesis from methyl 2-aminobenzoate/o-aminobenzaldehydes/o-aminoarylketones

Synthesis of acetylquinoline **12a** was accomplished *via* acylating methyl anthranilate **13** using ethyl acetoacetate or methyl acetoacetate and triethanolamine by continuous removal of

Scheme 1 Synthesis of AHQ 12a-f.

RSC Advances

Scheme 2 Synthesis of 12a from anthranilic acid.

Scheme 3 Suggested mechanism for the synthesis of benzoxazinone derivatives 16

Scheme 4 Synthesis of 12b through acylation of 20 followed by cyclization of compound 22

methanol or ethanol using Dean–Stark apparatus, resulting in the formation of 2-methoxycarbonyl anilide **13**, followed by Dieckmann intramolecular cyclization of anilide derivative **14** (Scheme 1).^{120,141–143}

3.4. Synthesis from 2-methyl-3,1-benzoxazin-4-one

The synthesis of **12a** was performed through a two-step. First, 2-methyl-3,1-benzoxazin-4-one **16** was synthesized by [4 + 2] annulation of anthranilic acid or by using the iminium cation from a mixture of cyanuric chloride and dimethylformamide as a cyclizing agent, and was achieved under mild conditions.¹⁴⁴ Next, C-acylated with ethyl acetoacetate by 3,1-benzoxazin-4-one **16** to form the ester **17**. Finally, cyclization of ester **17** in a basic medium [aqueous Na₂CO₃/NaOH] at room temperature furnished the target compound **12a**. This reaction pathway involves the nucleophilic addition of the active methylene of ethyl acetoacetate (EAA) to the carbonyl group of the benzoxazine ring **16**, followed by intramolecular cyclization (Scheme 2).^{108,116,145-147}

N-Acylated anthranilic acid intermediate **18** is created from anthranilic acid and acid chloride in the presence of a catalytic

amount of triethylamine (TEA) as the HCl scavenger *via* N-acylation reaction, affording *N*-acylated anthranilic acid. After that, intramolecular nucleophilic attack at intermediate **18** yielded benzoxazin-4-one ring **16**. The cyclization reaction can be achieved by converting the carboxylic group of acetylated derivative **18** into a dynamic ester through either microwave irradiation or traditional heating (Scheme 3).¹⁴⁴

3.5. Synthesis from 1,3-dioxinone with methyl 2-(methylamino)benzoate

The synthesis of **12b** was accomplished *via* an acylation reaction of secondary amine **20** with trimethyl-1,3-dioxinone **21**, followed by a cyclization reaction of the acylation product **22** under basic conditions (Scheme 4).¹⁰⁸

3.6. Synthesis from angular furo[3,2-c]quinolinone derivatives

In 2021, the group of Elgogary described the photooxygenation reaction of furo [3,2-c] quinolin-4(5H)-one derivatives 23 in

Scheme 5 Photooxygenation reaction of 23.

Scheme 6 Synthesis of 12a from acetoacetanilide.

Scheme 7 Synthesis of dehydroxylated-3-acetylquinoline (AQ) 28.

CHCl₃ led to the formation of the photocleaved product **24** *via* the intermediate **24A**. Next, acid hydrolysis of quinolinone derivative **24** furnished **12b** (Scheme 5).¹³⁹

3.7. Synthesis from ketene dithioacetal

Compound **12a** was synthesized through *in situ* direct addition of carbon disulfide (CS_2) to acetoacetanilide **25** in the presence of K_2CO_3 and tetrabutylammonium bromide (TBAB) as a green ionic liquid. The alkylation process was achieved using dimethyl sulfate, which yielded the ketene dithioacetal **26**. In the same context, thermal cyclization of ketene **26** afforded the quinolinone **27** in fair yield. Finally, basic hydrolysis of the quinolone **27** furnished 3-acetyl-4-hydroxyquinolinone **12a** (Scheme 6).¹⁴⁰

3.8. Synthesis of dehydroxylated-3-acetylquinoline (AQ)

Also, synthesis of dehydroxylated-3-acetylquinoline (AQ) **28** via the reaction of 2-aminobenzaldehydes with α,β -unsaturated carbonyl compounds as methyl vinyl ketone in high catalytic

Scheme 8 Synthesis of 2-acetylguinoline 31.

efficiency (56–86% yield), through sequence process as aza-Michael reaction gave **28 A** followed by Aldol reaction afforded **28 B** then aromatized in presence of α -amylase catalyzed (Scheme 7).¹⁴⁸

3.9. Synthesis of 2-acetylquinoline derivatives

The reaction of *o*-aminoarylketones **29** with butanedione as a symmetrical 1,2-diketone **30** according to Friedländer synthesis. Similarly, *o*-aminoarylketones condensed with pentanedione (unsymmetrical 1,2-diketone) under acidic conditions led to the formation of regioselective 2-acetylquinoline derivatives **31** (Scheme 8). The suggested reaction mechanism (Scheme 9) is analyzed using density functional theory calculations at the B3LYP/6-311G (d,p) level. It was determined that all relative energy barriers and activation energies for the reaction steps are minimized at these theoretical levels.¹⁴⁹

A plausible mechanism for forming product **31** is shown in Scheme 9. Initially, o-aminoaryl ketone (**29**) condenses with butan-2,3-dione (**30**) in the presence of TFA with conc. H_2SO_4 to give an imine intermediate **31A** as geometrical isomer **E** by eliminating H_2O . Then, tautomerization of the **31A** into a more activated form, the enamine intermediate **31B**. Followed by nucleophilic attack of β -position of enamino group on the carbonyl carbon and this gives the benzylic carbocation (which stabilise through eight resonance hybrid structures **31(G-H)** and subsequently loses a proton to form final product subsequently loses a proton νia aromatization to form final product **31**.

3.10. Synthesis of quinolinones

Zhang *et al.*¹⁵⁰ reported an efficient methodology for preparing of 3,4-dihydroquinolinone 34 and quinolinone derivatives 35 through intramolecular cyclization of *N*-aryl cinnamides 33. Compound 33 was synthesized through Knoevenagel condensation of β -oxo-amides 32 with various aryl aldehydes, which were then subjected to intramolecular cyclization. The

31M

Scheme 9 Proposed mechanism for synthesis of 2-acetylquinoline 31.

Scheme 10 Triflic anhydride-mediated synthesis of quinolinone.

optimized conditions for cyclization, including solvent, the reaction temperature, and various catalysts such as Tf_2O and DDQ, were investigated, whereas the optimum temperature is

80 °C, and *N*,*N*-dimethyl trifluoroacetamide (DTA) as solvent afforded the most satisfactory results. Under the optimised conditions, the reaction of a wide range of substrates such as α -acetyl-*N*-aryl cinnamides and α -benzoyl *N*-aryl secondary cinnamides proceeded efficiently to afford the corresponding quinoline-2(1*H*)-ones 34 and 35 (Scheme 10).¹⁵⁰

31L

4. Reactivity of AHQ

According to the current literature survey, the reactivity of AHQ **12a-f** is part of scientific research with many reagents, as illustrated below. AHQ is a quinolin-2-one derivative with a unique chemical structure consisting of a β , β' -tricarbonyl (TC) group and thus a quinolinone ring with a carbonyl (acetyl), hydroxyl (enolate), and carbonyl (lactamic) group attached to it.

Scheme 11 Formation of angular N-methylfuro[3,2-c]quinolinones 36

Scheme 12 Formation of linear furo[2,3-b]quinolinones 38.

4.1. Electrophilic substitution reactions

An electrophilic substitution reaction is a chemical reaction in which an electrophile replaces a functional group attached to a compound, typically displacing a hydrogen atom. These reactions usually follow a three-step mechanism that includes forming an electrophile, forming a carbocation (an intermediate), and removing a proton from this intermediate. AHQ derivatives can efficiently undergo electrophilic substitution reactions such as alkylation, halogenation, Friedel–Crafts, and formylation.¹⁵¹

4.1.1. Alkylation

4.1.1.1. N_1 -Alkylation. Compound 12a can potentially undergo methylation at either the nitrogen atoms at positions 1 or the oxygen atom at C_4 . However, the methylation exclusively occurs at the nitrogen atom rather than the oxygen atom. ^{115,116,143,145} Following that, angular N-methylfuro[3,2-c]

quinolinone derivatives 36 were synthesized from 12b through the Rap–Stöermer reaction, with α -chlorocarbonyl substrates in the presence of K_2CO_3 in DMF (Scheme 11).

Whereas. the reaction of 3-acetyl-6-chloro-4phenylquinolinone 37 with different α-halocarbonyl compounds afforded linear tetrasubstituted furo[2,3-b]quinolines 38 (Scheme 12). Thus, angular fluoroquinolinones 36 have been synthesized via the reaction of 12b derivatives with α chlorocarbonyl compounds, as applied to the Rap-Stöermer reaction through both conventional and/or microwave methods. Thus, the microwave irradiation provided higher yields (88-93%) compared to the conventional method (64-72%) (Scheme 12).143,152,153

4.1.1.2. O-Alkylation and its transformation. Compounds 12b,c underwent thionation using phosphorus pentasulfide, aluminum oxide as a catalyst/CH₃CN. The thionated compound 39 by dimethyl sulfoxide (DMSO) underwent o-methylation of the hydroxyl group (enol), yielding 40 as a sole product, which prevents keto-enol tautomerism at the 3rd and 4th positions, which could have interfered with the Aldol condensation reaction. Whereby, compound 40 was further reacted with various ketones through Aldol condensation in piperidine as a basic medium, to produce 1-(4-methoxy-2-thioxoquinolin-3-yl) ethenone 41 (Scheme 13).¹⁵⁴

Scheme 13 Synthetic route of 41 from 12b,c.

Scheme 14 Transformation of the hydroxyl group of 12b.

RSC Advances Review

Scheme 15 Mechanism of the Staudinger reaction.

Scheme 16 Synthesis of phenylethylaminoquinolinone 48

Scheme 17 Synthesis of isoxazoloquinolinone.

Tosylation of the sodium salt of **12b** furnished a very reactive compound 3-acetyl-4-tosyloxyquinolone **42**, which was used as the key compound to synthesize *o*-acetylazido-quinolinone **43**

via transformation of the tosyloxy group to the azido group by using NaN₃/N-methylpyrrolidone as solvent. Followed by ring closure through elimination of nitrogen gas by thermal heating in bromobenzene at 156 °C, 3-acetylazido-quinolinone afforded angular isoxazolo[4,3-*c*]quinolones 44. The azide 43 undergoes the Staudinger reaction with triphenylphosphane, yielding the phosphazene 45 *via* nitrogen loss. The hydrolysis of 45 in AcOH (80%) produced the 4-aminoquinolone 46. This amine can also be produced by reacting tosyloxyquinolone 42 with ammonia. Similarly, benzylamine combines with 42 to yield substituted quinoline 46 (Scheme 14).¹⁵⁵⁻¹⁵⁷

Mechanism of Staudinger reaction *via* very mild reduction of the azide **43** with triphenylphosphane yields the phosphazene **43B** by loss of nitrogenous gas. The hydrolysis of **43C/43D** in AcOH (80%) produced the 4-aminoquinolone **46** (Scheme 15).¹⁵⁵

Whereas, methylation of compound **12b,c** by dimethyl sulphate, a potent methylating agent, in acetone, which replaced a hydrogen atom in the hydroxyl group. This process

Scheme 18 Claisen-Schmidt reaction of compound 47.

Scheme 19 Cyclization reaction of 54 to yield 4-thiazolidinone derivatives 55.

Scheme 20 Synthesis of oxadiazolylquinolinone derivatives 59.

yielded a single product 3-acetyl-4-methoxyquinolinone **47**. Following this interpretation, the product reacted with a primary amine in EtOH, yielding phenylethylaminoquinolin-2-one **48** (Scheme 16). 158

Treatment of 3-acetyl-4-methoxyquinolinone 47 was observed by the Chimichi group 158 with 1,2-bisnucleophiles, such as hydroxylamine, to synthesize fused isoxazolo[4,5-c]-and/or isoxazolo[4,3-c]quinolin-4(5H)ones, which gave three different products 49–51 depending on the conditions (Scheme 17). As a result, the utilization of either hydroxylamine hydrochloride or its free base leads to the formation of two distinct products. The obtained product was through a reaction of 47 with NH₂OH·HCl is structurally assigned to the regioisomeric

Scheme 21 Synthesis of Mannich base 4-hydroxy-3-(3-morpholino-propanoyl)quinolinone 60.

isoxazolo[4,3-c] or isoxazolo[4,3-c]quinolin-4(5H)one in the following ways:

- (a) The presence of a three-bond connection between C-3a and the methyl group at position 3 of compounds **49** and **50** led to the exclusion of the oxazole structure **51**.
- (b) The chemical shift of the C-3 atom being analyzed, which resembles that of a 3-substituted isoxazole more than a 5-substituted one, was used as a preliminary criterion to distinguish between the two regioisomeric isoxazole skeletons. In contrast, compound 47 exclusively and preferentially generates the regioisomeric isoxazolo[4,5-*c*]- or isoxazolo[4,3-*c*]quinolin-4(5*H*)on e (compounds 49 and 50, respectively).¹⁵⁸

Whereby, synthesis of α,β -unsaturated ketones **52** was achieved through Claisen–Schmidt reaction of compound **47** with various aromatic aldehydes in glacial AcOH. Michael addition reaction of o-phenylenediamine to α,β -unsaturated ketone **52** afforded binary diazepin-quinolinones **53** (Scheme 18), whereas derivatives of **53** have shown gratifying sedative, anxiolytic, and muscle relaxant activities. ¹⁵⁹

Methylated compound 47 was reacted with substituted phenylhydrazines, which resulted in the formation of hydrazone 54, which were then cyclized with thioglycolic acid in methanol using zinc chloride (ZnCl₂) as catalyst, affording 4-thiazolidinone derivatives 55 (Scheme 19).¹⁶⁰

The o-alkylation of 12b with ethyl bromoacetate in anhydrous acetone and K_2CO_3 led to the formation of ethyl oxoquinolinoxyacetate 56. Hydrazinolysis of 56 with hydrazine hydrate in methanol at room temperature (RT) yielded hydrazide compounds 57 in 80%, after which heating hydrazide compound 57 with different aromatic aldehydes in refluxing EtOH yielded derivatives 58. Finally, compounds 58 were refluxed with Ac_2O to obtain the corresponding substituted 1,3,4-oxadiazolyl)methoxy)-1-methylquinolinone 59 (Scheme 20). 161

4.1.1.3. Aminoalkylation. AHQ **12a-f** were subjected to Mannich reaction with formaldehyde and morpholine salt in boiling MeOH for 6 h, furnishing Mannich base 4-hydroxy-3-(3-morpholinopropanoyl) quinolinone **60** (Scheme 21).¹⁶²

The Mannich reaction mechanism starts with the reaction between formaldehyde and amine, leading to the formation of the iminium ion. The enol form of the organic compound is obtained *via* tautomerization of the ketone form. The reactive iminium ion is attacked by this enol form, which ultimately

Scheme 22 The proposed Mannich reaction mechanism.

Scheme 23 Chlorination of acetylquinolinone 12d and pyranoquinolindione 11b

produces the necessary β -amino-carbonyl molecule or Mannich base (Scheme 22).

4.1.2. Halogenation and its synthetic value for constructing various heterocyclic systems

4.1.2.1. Synthesis of 3-dichloroacetyl derivative and its utility. Among organic synthesis transformations, chlorination holds one of the most fundamental reactions that can be achieved directly by using molecular chlorine or chlorinating agents.

Thus, the electrophilic chlorination reaction of **12b,d** with sulfuryl chloride (SO_2Cl_2) in 1,4-dioxane yielding 3-dichloroacetyl derivative **61**, also by heating pyranoquinolindione **11b** with SO_2Cl_2 (Scheme 23). ^{163,164}

Treatment of compound **61** with some 1,3-binucleophilic reagents, namely, guanidine hydrochloride, cyanoguanidine, and thiourea, was carried out under reflux in DMF, to afford the corresponding 3-imidazolylquinolinones **62**. While the reaction

Scheme 24 Treatment of 3-dichloroacetyl 61 with nucleophilic nitrogenous reagents.

NHNH₂NH₂ NH₂NH₂ CH(OEt)₃ fusion СН3 ĊН₃ ĊH₃ 70 (60%) 71 (63%) 69 X | (%) H₃CS NH 73 SCH₃ H₃CS[′] CN. 72 DMF 73 (61%)

Scheme 25 Hydrazinolysis of the 1,2,4-triazine derivative 69.

Scheme 26 Synthesis of pyrimidine, diazepine, and pyrazole heterocycles from 68

of **61** with NaN $_3$ in DMF occurred at ambient temperature, the intermediate diazido compound **63A** yielded the tetrazole **63** under loss of N $_2$ gas. Condensation of compound **61** with 1,4-N,N-binucleophiles such as o-phenylenediamine and 1,6-diaminopyridine derivative **64** under reflux in DMF, afforded the quinoxalinylquinolinone **65** and the pyridotriazine derivative **66**, respectively. ¹⁶³


Reaction of **61** with 4-aminotriazine derivative **67** was carried out in boiling pyridine. Interestingly, the only product that was obtained revealed the absence of chlorine; analysis results indicate that a nucleophilic replacement took place with the leaving of a chlorine atom, which is followed by elimination of hydrogen chloride, leading to the α -iminone **68** (Scheme 24).¹⁶⁴

Reaction of compound **61** with aminoguanidine and thiosemicarbazide as 1,4-binucleophiles, under reflux in DMF, yielded the corresponding 1,2,4-triazine derivatives **69**. Hydrazinolysis of **69** in DMF produced the hydrazinotriazine **70**, Thermal cyclocondensation reaction of **70** with CH(OEt)₃ under fusion condition was carried out to get the triazolotriazine derivative **71** whereas, reaction with [*bis*(methylthio)methylene] malononitrile **72**, in DMF under reflux, gave the compound **73** (Scheme 25).¹⁶³

 α -Iminone **68**, as an interesting starting material, was subjected to interaction with some binucleophilic reagents to synthesize new quinolinone systems fused with pyrimidine,

benzodiazepine, and pyrazole heterocycles. Therefore, treatment of α -iminone **68** with guanidine, o-phenylenediamine, and hydrazine hydrate, in DMF under reflux, afforded the pyrimidoquinolinone **74** and benzodiazepinoquinolinone **75**, and pyrazoloquinolinone **76**, respectively (Scheme **26**). ¹⁶³

4.1.2.2. Synthesis of 3-(bromoacetyl)-4-hydroxyquinolinone and its benefits. The bromination of AHQs **12b-d** can be performed using bromine or *N*-bromosuccinimide (NBS) in the presence of benzoyl peroxide. 3-(Bromoacetyl)-4-hydroxyquinolinone derivatives 77 were synthesized by the free radical bromination mechanism of acetyl **12b-d** using NBS in dry carbon tetrachloride (CCl₄) with a catalytic amount of benzoyl peroxide (3 mol%) via homolytic fission, ¹⁶⁵ also reaction of **12b-d** with bromine in the presence of AcOH via electrophilic bromination gave the same product 77 (Scheme 27). ^{115,134,166,167}



Scheme 27 Bromination of *N*-substituted-3-acetyl-4-hydrox-yquinolinones **12b-d**.

RSC Advances Review

Scheme 28 Reactions of bromo-derivatives 77 with variable reagents.

Scheme 29 Vilsmeier-Haack reaction of skeletons 12b-d.

The desired thiocyanate **78** was obtained from the reaction of 77 with potassium thiocyanate. Meanwhile, phenacyl halides were used for the alkylation of 2-aminopyridine, followed by cyclization to imidazo[1,2-a]pyridines. Thus, when 77 was heated with 2-aminopyridine in ethanol, 4-hydroxy-3-imidazoyl methylquinolinone **79** was formed. The Moreover, condensation of bromo-derivatives 77 with o-vaniline gave 4-hydroxy-3-(7-methoxybenzofuran-2-carbonyl)-1-methylquinolin-2(1H)-one **80**. Whereas, compound 77 is a versatile intermediate in synthesizing compounds **83** and **84**. So that, the synthesis of **83** were achieved through condensation reaction of 77 with various thiadiazole **81** and the observed compound benzo[d]thiazol-2-yl-phenylglycylquinolinone **84** were synthesized in good yield (60–

81%) through the reaction of disubstituted-2-benzo[d]thiazole 82 with 77 in the presence of AcOH (Scheme 28). ^{166,168}

4.1.3. Formylation of AHQ and its uses. The Vilsmeier–Haack reaction was known since 1927,¹⁶⁹ which uses DMF, an acid chloride and an aqueous work-up to change an aromatic ring with a lot of electrons into an aryl aldehyde. The "Vilsmeier reagent" is an iminium salt that is created when DMF reacts with acid chloride at the start of the mechanism (Scheme 30). Moreover, double formylation of the methyl group present in acetylquinolone **12b–d** with subsequent *in situ* cyclization of intermediates **85A** and **85B** yielded pyrano[3,2-*c*]quinoline-3-carboxylaldehyde derivatives **85** *via* Vilsmeier–Haack reaction of acetylquinolinones **12b–d** (Scheme 29).^{137,170}

Plausible mechanism of Vilsmeier reaction

Scheme 31 Synthesis of pyrazolylquinolinone skeletons

Ibrahim et al. 170 independently reported the treatment of 85 with various nucleophilic reagents affording pyrano[3,2-c] quinolines via ring-opening followed by ring closure (RORC).

Also, reaction of carboxaldehyde 85 with hydrazine hydrate in refluxing **EtOH** yielded 4-hydroxypyrazol-4ylcarbonylquinolin-2-one 86, via the non-isolable hydrazone intermediate, which underwent intramolecular nucleophilic attack of NH₂ at the C-2 position with concomitant γ -pyrone ring opening in situ. In the same way, the condensation reaction of 85 with phenyl hydrazine in EtOH containing a catalytic amount of TEA afforded phenylpyrazole derivative 87. Whereby, the condensation of carboxaldehyde 85 with hydrazinylquinoline quinoline 88 and 3-hydrazinyl-1,2,4-triazine-1,2,4-triazine 89 under the same reaction conditions was achieved yielding quinolinylpyrazolylquinolinone 90 and triazinylpyrazolylquinolinone 91, respectively (Scheme 31).170

Furthermore, aldehyde 85 was subjected to react with some 1,3-N,N-binucleophiles such as thiourea, guanidine hydrochloride and cyanoguanidine in ethanolic KOH yielded the corresponding pyrimidine derivatives 92, 93 and 94, respectively. Whereby, refluxing of 85 with 5-aminotetrazole in EtOH afforded 3-[(2-azidopyrimidin-5-yl)carbonyl]4-hydroxyquinolin-2-one 95 through non-isolable tetrazolo[1,5-a]pyrimidine intermediate 95A (Scheme 32).

Meanwhile, condensation of aldehyde 85 with 3-aminotriazole in EtOH yielded triazolo[4,3-a]pyrimidine 96 containing the quinolinylcarbonyl moiety. Ultimately, condensation of 85 with 2-aminobenzimidazole in EtOH containing one crystal of p-toluenesulfonic acid afforded pyrimido[1,2-a]pyrimidine derivative 97. Also, reaction of 85 with 1H-benzimidazol-2ylacetonitrile and N-benzyl-2-cyanoacetamide in EtOH containing catalytic drops of TEA yielded 98 and 99, respectively.

Scheme 32 Synthesis of six-membered heterocyclic scaffolds.

Scheme 33 Synthesis of different heterocyclic systems containing quinolone scaffolds.

Next, condensation reaction **85** with *o*-phenylenediamine, *o*-aminophenol, and *o*-aminothiophenol in AcOH yielded the corresponding heteroannulated pyrano[3,2-*c*]quinoline

derivatives **101**. Whereby, reaction of **85** with 6-aminouracil and 6-amino-1,3-dimethyluracil in EtOH afforded pyrido[2,3-*d*] pyrimidines **100**. Furthermore, the chemical reactivity of **85** was

Scheme 34 Synthesis of tricyclic carbonitrile derivative 103.

Scheme 35 Synthesis of 4-hydroxypyrano[3,2-c]quinoline-3-carbonitrile derivative 104

investigated towards different 1,4-binucleophiles. By the way, the condensation reaction of **85** with ethylenediamine in EtOH yielded 1,4-diazepinylcarbonylquinolin-2-one **102**. The reaction was accomplished through the synthesis of the corresponding Schiff base intermediate, followed by an intramolecular nucleophilic addition at the C-2 position associated with γ -pyrone ring opening to afford **102** as the final product (Scheme 33).¹⁷⁰

In refluxing pyridine, the reaction of tricyclic aldehyde 85 with NH₂OH·HCl yielded the corresponding carbonitrile

derivative **103**, through the formation of non-isolable oxime **103A**, after *in situ* dehydration step (Scheme 34).¹³⁷

Reaction of aldehyde **85** with NH₂OH·HCl, in AcOH, did not yield either oxime **103A** or carbonitrile **103** but gave unexpected product showed that dioxopyrano[3,2-c]quinoline-3-carbonitrile **104** through two molecules of hydroxylamine reacted stepwisely with **85** yielded **103** which is regarded as a highly reactive cyclic push-pull system that quickly produces the adduct **104B**, which exists in equilibrium due to ring-chain tautomerism. The dehydration of oxime tautomer readily afforded the

Scheme 36 Formation of 1-amino-5-ethylisoxazolopyrano[3,2-c]quinolindione 105.

Scheme 37 Complexation of compounds 12b,d.

corresponding intermediate **104C**. Intramolecular 6-exo-dig cyclization type of intermediate **104C** gave 2-iminopyranoquinoline **104D** following that hydrolysis of **104D** yielded the final product **104** (Scheme 35).¹³⁷

Reaction of aldehyde 85 with $NH_2OH \cdot HCl$, in ethanolic KOH (1%) as a basic catalyst, instead of an acidic medium, gave 105 in 46% yield. The reaction gave the same intermediate 104D, in a basic catalyzed route, which underwent the addition of another molecule of $NH_2OH \cdot HCl$ led to the formation of 2-

Scheme 38 Condensation reactions on complex 106.

	OH N OH CONTROL OH OH C	2 ^{ry} amir CH ₂ O,H	ne,MeOl	→	√ _N	- ,x				
R	R ₁	(%)			uted aniline					
Н	4-methylpyridin-2-amine	72 pyridine,DMF, reflux 17-24 h								
н	5-methylpyridin-2-amine	75		renux	17-24 11					
н	6-methylpyridin-2-amine	74		R.	_X					
н	4-methyl-3-nitropyridin-2-amine	74 70 OH N R1 X								
н	Pr	52		\wedge \downarrow \downarrow \rangle	<td></td>					
н	i-Pr	63 N								
н	Bu	17 NO								
н	i-Bu	78								
н	sec-Bu	56		Ř						
Н	(CH ₂) ₂ OH	70	_	112						
H	CH ₂ Cy	72	R	R ₁	Х	(%)				
Н	(CH ₂) ₄ CH ₃	11	CH ₃	C ₆ H ₅ -NH-	0	81				
CH₃	C ₆ H ₅ -NH-	85	CH ₃	4-CI-C ₆ H ₄ -NH-	0	93				
CH₃ CH₃	4-CI-C ₆ H ₄ -NH-	89	CH ₃	C ₆ H ₅ -NH-	N-methyl	84				
CH ₃	4-Br-C ₆ H ₄ -NH- Bu	78	CH ₃	4-CI-C ₆ H ₄ -NH-	N-methyl	88				
CH ₃	Бu CH₂Ph	86 51	CH ₃	C ₆ H ₅ -NH-	CH ₂	92				
CH ₃	Ph	51 58	C ₆ H ₅	C ₆ H ₅ -NH-	0	90				
C ₆ H ₅	C ₆ H ₅ -NH-	82	C ₆ H ₅	4-CI-C ₆ H ₄ -NH-	0	88				
C ₆ H ₅	4-CI-C ₆ H₄-NH-	79	C ₆ H ₅	4-NO ₂ -C ₆ H ₄ -NH-	N-methyl	86				
C ₆ H ₅	4-Br-C ₆ H₄-NH-	84	C ₆ H ₅	4-CI-C ₆ H ₄ -NH-	N-methyl	79				
CH ₂ Ph	CH ₂ Ph (64%)	64	C ₆ H ₅	C ₆ H ₅ -NH-	CH ₂	81				
CH ₂ Ph	Ph	61								

Scheme 39 Synthesis of imine 111, Mannich base (QVIR) 60 and its imine analogue 112.

Scheme 40 Synthesis of pyrimido[5,4-c]quinolinones 113a-c.

imino-5-oxopyrano[3,2-c]quinoline-3-carboximidamide **105A**. Whereas, the cyclization of imidamide through a condensation reaction yielded isoxazole derivative **105B**. After that, hydrolysis of imine **105B** was achieved, leading to the final product **105**. Additionally, compound **105** was accurately prepared from the reaction of **104** with NH₂OH·HCl in refluxing EtOH (Scheme 36).¹³⁷

4.1.4. Synthesis of 1,3,2-dioxaborinino[5,4-c]quinolinone *via* metallation and its utility. The metallation reaction of acetylquinolinone 12b,d with boron trifluoride etherate BF₃·(OC₂-H₅)₂ in benzene gave its boron difluoride chelate complex 106 (Scheme 37).¹⁷¹

Reaction of complex **106** with DMF gave enamine **107**, which can serve as a valuable synthon for transformations of the two-quinolinone fragment. Moreover, complex **106** was treated with different carbonyl compounds to yield the corresponding condensation products at the methyl group. For example, a reaction of two moles of complex **106** with CH(OEt)₃ with a catalytic amount of TEA afforded symmetric polymethine dye **108**. In acetonitrile, complex **108** absorbs at 581 nm, displaying noticeable fluorescence. Lastly, boron-containing styryl dyes **109** were produced by the condensation reaction of several aromatic and heterocyclic aldehydes with complex **106**.

Compounds **109** were hydrolyzed to produce novel 4-hydroxy-2-quinolinone derivatives **110**, which exhibit fluorescence in both solid and solution states (Scheme 38).¹⁷¹


4.2. Nucleophilic substitution reactions

The substitution of a nucleophile at a tetrahedral or sp³ carbon is known as aliphatic nucleophilic substitution. Substitutions involving aliphatic nucleophiles are not glamorous or essential in the field of chemistry. The way carbonyl additions and carboxyloid replacements seem to occur in biochemistry, they do not occur in all significant processes. Rather, they are commonplace, minor reactions with significant, minor effects everywhere.

4.2.1. Reactions with nitrogen bases

4.2.1.1. With amines. Condensation reaction of *N*-substituted (un)acetylquinolinone **12a–c,e** with different amines under various conditions afforded enaminones **111.**^{143,155,162,172,173} While, addition of a solution of substituted aniline in dimethylformamide (DMF) and pyridine to **60** and refluxed for 17–24 h to give **112** (Scheme 39). ^{162,173}

The reaction of quinolinone **12a** with nitrogen bases derivatives such as guanidine nitrate, urea and thiourea in boiling ethanol using AcONa as a catalyst through nucleophilic



Scheme 41 Reactions of guinolinones 12a-c with variable hydrazines.

RSC Advances Review

Scheme 42 Condensation and cyclization of 114 and 118 to synthesize compounds 119 and 120

Scheme 43 Synthesis of pyrazolo[4,3-c]quinoline 121

addition of the amino group with the regioselectively at the C=O of acetyl function of **12a** accompanied by intramolecular cyclocondensation afforded regioisomer pyrimido[5,4-*c*]quinolinones **113a-c** (Scheme 40).¹⁷⁴

4.2.1.2. With hydrazines. Condensation reaction of hydrazines and arylhydrazines with AHQ derivatives **12a–c** in boiling EtOH or DMF yielded corresponding hydrazones **114**. ¹¹⁵, ¹¹⁸, ¹³⁴, ¹⁷² The reaction conditions varied depending on the aryl substituents, with phenylhydrazine reacting in dimethylformamide

EtOH 123 (%) NHNH₂ CH₃ C₂H₅ 90 **AcONa** ĊН₃ 12b,d 122 (%) CH₃ C₂H₅ 55 57 AcOH AcONa k 124

Scheme 44 Synthesis of compounds 123 and 124 through condensation and cyclization.

(DMF) at ambient temperature, while nitro- and chlorophenyl hydrazines required refluxing in BuOH (Scheme 41). 135,155

The formation of compound 115 can be accomplished through various methods. This reaction involves two straightforward and efficient approaches. The first method involves a one-step condensation of acetylquinoline 12a with hydroxyquinolone-3-carboxylic acids hydrazides 116, resulting in high yields. The second method consists of two stages: initially, treatment of acetylquinoline 12a to quinolinone, followed by acylation using ethyl hydroxyquinoline-3-carboxylates 117. This method also proves to be effective in the production of the hydrazones 115 described in this study; both methods were found to be equally valuable (Scheme 41). 115,120,134

Similarly, compounds **119** and **120** were not achieved immediately by the condensation of **12a-c** with the following hydrazines as hydrazine hydrate, semicarbazide, or thiosemicarbazide but gave acyclized hydrazone **114** and **118**, respectively. These products were cyclized with Conc. H₂SO₄ to yield angular fused pyrazoloquinolinone **119** and angular triazepinoquinolinone **120** (Scheme 42). ^{118,174}

Angular pyrazolo[4,3-*c*]quinoline derivatives **121** were synthesized under *via* enolizable 3-acetylquinolinone **12a,b** with phenylhydrazine derivatives, utilizing InCl₃ as an effective Lewis acid in providing greater yields of products **121** under microwave irradiation (Scheme 43).¹⁷⁵

Also, bis quinolinone **123** and **124** were obtained by reaction of **12b,d** with hydrazinyl derivative **122** in different conditions,

Scheme 45 Synthesis of phenylhydrazonoquinolinylmethylbenzenesulfonate 126

Scheme 46 Synthesis of functionalized quinolone derivatives 128–131.

in hot ethanol or glacial acetic acid and sodium acetate (AcONa) (Scheme 44).¹⁷⁶

Initially, the reaction of hydrazone derivatives **114** with CH₃ONa in diethyl ether converted the hydroxy group at position 4 to the sodium enolate group, affording **125**. The reaction of sodium salts with toluenesulfonyl chloride in dry CH₃CN yielded targeted toluenesulfonyloxy quinolones **126**. Conversely, the attempts to obtain the toluenesulfonyloxy compounds **126** (R=H) were not avail However, the separation of sodium salts **125** (R=H) in good yields may be attributed to the high reactivity, in both cases, after reaction with TsCl yielded non-isolable mixture of products (Scheme 45). ¹³⁵

For the synthesis of pyrazole-4-carbaldehydes 127, the hydrazones 114 were exposed to Vilsmeier–Haack reagent (DMF-POCl₃).^{177,178} While traditional condensation of pyrazole-4-carbaldehydes 127 with hydrazine hydrate or thiosemicarbazide furnished hydrazone derivatives 128 in

acceptable yields. Another system in this area was created in two steps. Condensing acetylquinolinone **12b** with aldehyde **127** (R = Ph) furnished the α , β -unsaturated ketone, which was then treated with NH₂NH₂, affording polynuclear heterocyclic skeleton **129** (Scheme 46).

Unexpectedly, the condensation of 127 (R = Ph) with NH₂-OH·HCl in AcOH did not produce the predicted oxime or the likely carbonitrile 130A but yielded phenylpyrazolopyrano[3,2-c] quinolindione 130, while the oxime transformed to carbonitrile via a dehydration process. The carbonitrile 130A was then cycloadditionally converted to an iminopyranoquinolinone, hydrolyzed to yield the isolated tetracyclic system 130. Another option is that the hydrolysis of the intermediate 130A into the equivalent carboxylic acid causes the pyrone ring to form via intramolecular condensation. Anyway, both routes are rational and feasible.¹⁷⁸ Condensation of equimolar amounts of carbaldehyde 127 (R = Ph) with the 3-acetylcoumarin in the presence

Scheme 47 Reaction of hydrazone 114 with sulfur-containing phosphorus reagents.

Scheme 48 Treatment of hydrazone 114 with P(OCH₂CH₂Cl)₃.

of piperidine as catalyst afforded the corresponding chalcone 131 (Scheme 46). 177

Additionally, the synthesis of fused phosphorus heterocycles with quinolinone moieties was achieved through an efficient protocol *via* the reaction of hydrazone **114** with some

phosphorus reagents. As the treatment of hydrazone **114** with Lawesson's reagent (LR) [which forms two reactive dithiophosphine ylides (R-PS₂)], phosphorus pentasulfide (P_2S_5) and diethyldithiophosphoric acid in refluxing dioxane afforded the

Scheme 49 Annulation of hydrazone 114 with diethyl phosphite and phosphohalogenating agents.

Scheme 50 Condensation and thermal Beckmann rearrangement of oxime product

corresponding fused tricyclic oxadiazaphosphepino[6,7-c]quinolinones **132–134** (Scheme 47).¹⁷⁹

Synthesis of cyclic phosphonic ester 135 was accomplished through treatment of hydrazone 114 with tris(2-chloroethyl) phosphite P(OCH₂CH₂Cl)₃. The credible mechanism for synthesis of compound 135 starting with Michael addition of the phosphite to the azomethine bond to produce the nonisolatable intermediate 135A, which undergoes cyclization through nucleophilic attack of NH functionality at position 4 of the quinoline ring to remove a water molecule yielding the pyrazolyl non-isolatable intermediate 135B which underwent cyclization through elimination of HCl, followed by acidic hydrolysis yielding 135 as final product (Scheme 48).¹⁷⁹

Furthermore, Pudovik reaction of hydrazone **114** with diethyl phosphite in the presence of BF $_3$ etherate at 80–90 °C furnished oxadiazophosphepino[6,7-c]quinolinone **136A,B**. The plausible mechanism involves a cyclization of hydrazone **114** through the nucleophilic attack of OH and NH groups at the phosphorus atom, followed by stripping off two molecules of EtOH. Additionally, heterocyclization of hydrazone **114** with phenylphosphonic dichloride ($C_6H_5Cl_2PO$) in dioxane afforded the corresponding oxadiazaphosphepinoquinolinone **137**(Scheme 49).¹⁷⁹

Finally, reaction of hydrazone 114 with equimolar amounts of with diethyl phosphite, $PhPOCl_2$, phosphorus tribromide (PBr_3) or $POCl_3$ in dry dioxane in molar ratio (1:1) containing

OH O
PhNCS/H₂NCH₂CO₂H
AcOH
N
O
R
12b,d

O
N
O
CICH₂COOH
Or
CH₃COONa
Ac₂O
R
144
R=CH₃, C₂H₅

O
PhNCS/H₂NCH₂CO₂H
AcOH
AcOH
N
O
N
O
R
143A

H₂O
O
H
N
N
O
H
N
O
R
143A

143 (70-75%)

Scheme 51 Formation of thiazolidine derivatives 144.

catalytic amount of TEA afforded **138**, The proposed mechanism for the synthesis of fused tricyclic **138** may be attributed to halogenation of OH group process through the at C₄. A phosphorus halide of the quinolinone ring, cascade by cyclization of intermediate **138A** through dehydrohalogenation elimination of HX using TEA yielding **138** (Scheme 49).¹⁷⁹

4.2.1.3. With hydroxylamine. Reaction of acetylquinolinone 12a with NH₂OH·HCl was accomplished in an ethanolic solution in the presence of AcONa furnished the corresponding oxazole 139. Whereas, repeating the reaction in the presence of NaOH instead of AcONa furnished corresponding oxime 140 (Scheme 50).^{115,118,134,145,172,174}

3-Oximatoacyl-4-hydroxyquinolones **140** were synthesized by reacting AHQ **12a,b,e** with excess hydroxylammonium chloride and hydrogenearbonate, yielding hydroxyiminoquinolones **140** ($R_1 = H$). To obtain the oximethers **140** ($R_1 = methyl$), quinolinone **12a,b** were reacted with methyloxyammonium chloride and hydrogenearbonate. Both types of compounds **140** structurally resemble biologically active compounds like alloxydim or sethoxy. When oximes are thermolyzed according to Beckmann rearrangement and converted into isomeric oxazolo[5,4-c] quinolinones **141** in high-boiling solvents like 1,2-dichlorobenzene and diphenyl ether. Whereby, the hydrolysis process of oxazole derivatives **141** with dil. HCl led to formation of 3-acylamino-4-hydroxyquinolones **142** (Scheme 50).¹⁷²

4.2.2. With phenylisothiocyante. Multicomponent reaction (MCR) of **12b,d**, phenylisothiocyanate and glycine furnished *N*-pyrano[3,2-*c*]quinolinedion phenylthiourea **143** in acceptable yields *via* condensation of the carbonyl group of **12b,d** with the active methylene group in glycine to yield the intermediates **143A** which readily cyclized through stripping off water molecule to afford **143**. Whereas, the condensation of **143** with chloroacetic acid and Ac₂O afforded thiazolidine derivatives **144** (Scheme 51).¹⁵¹

4.3. Condensation reactions

4.3.1. With aldehydes (synthesis of α,β -unsaturated ketones). A series of α,β -unsaturated ketones of quinolinone analogues **145** were synthesized through Claisen–Schmidt condensation of **12a–d** with aldehydes under reflux or microwave irradiation. ^{115,131,146,180–184} The mechanism of the reaction started by reacting of formyl group with L-proline to form an

OH O R_5 O OH O R_1 R_2 OH O R_5 R_4 R_5 O OH O R_5 R_4 R_5 O OH O R_5 R_4 R_5 O OH O R_5 R_5 R_4 R_5 O OH O R_5 R_5 R_4 R_5 R_5 R_4 12a-d													
R	R ₁	R ₂	R_3	R ₄	R ₅	(%)	R	R ₁	R ₂	R ₃	R ₄	R ₅	(%)
H (() H () H () H () H ()	H H H H H H H H H H H H H H H H H H H	H OCH ₃ OCH ₃ H C(CH ₃) ₃ H H H (CH=CH) ₂ OCH ₃ H OCH ₃ H	H OH OCH ₃ (OCH ₃) ₂ OH NO ₂ N(CH ₃) ₂ (CH=CH) ₂ CI H OH CI H OCH ₃ N(C ₂ H ₅) ₂ CO ₂ CH ₃	H H H C(CH ₃) ₃ H H (CH=CH) ₂ H H H H H H H	H H H H H OCH ₃ H H H	89 92 72 78 89 86 88 58 90 78 68 90 91 72	H CH3 CH3 CH3 CH3 CH3 CH3 CH3 CH3 CH3 CH	H	Br H H H H H CH ₃ CI H OCH ₃ OCH ₃ H H	H H CI H COH ₃ OH CH ₃ H COCH ₃ COCH COCH COCH COCH COCH COCH COCH COC			60 74 63 85 78 70 87 78 85 71 48 72 65 75

i) C_2H_5 ONa,MW/320 W/1 min or 12h stirring; ii) piperidine, EtOH; iii) BuOH, drops piperidine and reflux iv) neutral alumina and ultrasonic waves; v) L-Proline/ DMSO/ 30-40 min/ RT;vi) 5% aq.NaOH, 24 h /RT vii) 5% Alc.NaOH. 12 h/ RT

Scheme 52 Synthesis of α,β -unsaturated ketones 145

Scheme 53 Proposed mechanism for the synthesis of α, β -unsaturated ketone 145 in the presence of L-proline.

Scheme 54 Reaction of quinolinone 12a,e with cinnamaldehyde.

iminium carboxylate ion **146**, after that this ion eliminating a proton from active methylene of **12a** affording the carboxylic acid containing iminium ion **148** and quinoline anion **147** as ion pairs that combine to form the adduct **149** which loses proline to afford α,β -unsaturated ketone **145** (Schemes 52 and 53). ¹⁸³

Similarly, reaction of quinolone **12a,e** with cinnamaldehyde in the presence of piperidine in refluxing EtOH afforded α,β -unsaturated ketone **150**. After that, treatment of **150** with

hydrazine hydrate led to the formation of styrylpyrazolylquinolinone **151** (Scheme 54). 146,185

The predominant transformations for generating pyrazole rings typically involve the use of α,β -unsaturated carbonyl group derivatives as parent skeletons. Chalcones **145** were treated with appropriate hydrazines in AcOH, furnished functionalized pyrazoline analogues **152** (Scheme 55). ^{108,146,184,186}

Whereby, the hydrazinoquinolinone 122 underwent condensation followed by cycloaddition reaction with α,β -

OH O R ₂ NO R ₄ NH ₂ NHR ₁ .H ₂ O R ₄ NH ₂ NHR ₁ .H ₂ O R R 145											
R	R ₁	R ₂	R ₃	R ₄		R	R_1	R ₂	R ₃	R ₄	
H H H H H CH ₃ CH	4-CN-C ₆ H ₄ 4-(OCH ₃)C ₆ H ₄ 4-(CN)C ₆ H ₄ 4-(CN)C ₆ H ₄ 4-(benzyloxy)C ₆ H ₄ 4-(benzyloxy)C ₆ H ₄ 4-(benzyloxy)C ₆ H ₄ C ₆ H ₅ COCH ₃ 4-(CN)C ₆ H ₄ 3,4-(CH ₃) ₂ C ₆ H ₃ 4-(CN)C ₆ H ₄ 3,4-(CH ₃) ₂ C ₆ H ₃ 4-(CH ₃)C ₆ H ₄ 4-(CI)C ₆ H ₄ 4-(CI)C ₆ H ₄ 4-(CI)C ₆ H ₄ 4-(COCH ₃)C ₆ H ₄ 4-(OCH ₃)C ₆ H ₄ 4-(OCH ₃)C ₆ H ₄		OCH ₃ OCH ₃	OCH ₃ OCH ₃ OH OH OCH ₃ OH OCH ₃ H OCH ₃ OCH ₃ OCH ₃ OCH ₃ OCH OH OH OH OCH OCH OCH OCH OCH OCH OC		CH ₃ CH ₃ CH ₃ CH ₃ CH ₃ CH ₃ Ph Ph Ph	H H H H H H H H H H	HCHFHCHHCHFHC	H H H H H H H H H H H H H H H H H H H	H H CI H F CI OCH₃ OH H CI H CI	

Scheme 55 Synthesis of pyrazolylquinolinones through the reaction of 145 with hydrazines.

Scheme 56 Synthesis of binary pyrazolylquinolinone derivatives.

Scheme 57 Synthesis of fused and binary isoxazolo and isoxazolyl quinolinone 154 and 155

unsaturated ketone derivatives **145**, affording binary quinolinylpyrazoloquinolinone derivatives **153** up to yield 59% (Scheme 56). ¹⁷⁶

Whereas, the reaction of 145 with NH₂OH·HCl in glacial AcOH yielded a mixture of two products 154 and 155. Conversely, repeating the same reaction in other solvents such

i) alcoholic KOH and reflux; ii) basic alumina and microwave irradiation.

Scheme 58 Synthesis of pyrimidinylquinolinone 156

Scheme 59 Synthesis of 1,5-benzothiazepin ylquinolinones 157 anchored on quinolone

Scheme 60 Synthesis of 4-hydroxy-oxoguinolin-4-arylpyridine-3-carbonitriles 158

as benzene, MeOH, and EtOH was unsuccessful. Since the iso-xazolo[4,5-*c*]quinolin-4(5*H*)-ones **155** formation could not be controlled to get only isoxazolines **154**, increasing the reaction duration to 19–20 h led to isoxazoles **155** as major products, and the minor products could not be isolated (Scheme 57).¹⁸⁰

Pyrimidinylquinolinone quinolinyl pyrimidine derivatives **156** have been synthesized through both traditional methods and microwave irradiation. These compounds were obtained *via* condensing α,β -unsaturated ketone derivatives **145** with urea or thiourea in basic media (Scheme 58).¹⁸²

Furthermore, substituted 1,5-benzothiazepinylquinolinones 157 were synthesized by cyclocondensation reaction followed by Michael addition. Whereas, a mixture of α,β -unsaturated ketone 145, o-aminothiophenol in absolute EtOH then added glacial AcOH (5 mol%). The obtained reaction mixture was refluxed for 5–7 h, furnishing the target product 1,5-

Scheme 61 Reaction of guinolinone 12b with 3-formylchromone 159.

benzothiazepinylquinolinones 157 in yields 62–82% (by the optimized condition) (Scheme 59).¹³³

MCRs of α,β -unsaturated ketone **145**, malononitrile, and ammonium acetate in DMF at 75 °C afforded functionalized pyridine binary with quinoline systems **158** (Scheme 60).¹⁸⁷

The desired precursor chromenacryloylquinolinone **160** was obtained smoothly *via* a one-pot Aldol condensation dehydration reaction of **12b** with 3-formylchromone **159** (Scheme 61). ¹⁸⁸

In refluxing EtOH, the treatment of compound **160** with an equimolar amount of hydrazine hydrate led to the opening of the pyrone ring and the enone side chain unaffected, affording a new product assigned as 4-hydroxy-3-[3-(2-hydroxyphenyl) pyrazolyl]acryloylg quinolineone **161**. Whereas, using AcOH as the solvent oriented the reaction away from the γ -pyrone nucleus, yielding 3-[1-acetyl-5-(4-oxo-4*H*-chromen-3-yl)-4,5dihydropyrazolyl]-4-hydroxyquinolinone **162**.

While by, compound **160** was reacted with of hydrazine hydrate (1:2) in refluxing DMF affording 4-hydroxy-3-[30-(2-hydroxyphenyl)bipyrazolyl]quinolinone **163** in 73% yield, through double nucleophilic attack at both the α,β -unsaturated ketone side chain and chromone ring, where by, the same product was prepared by reacting the pyrazolylacryloylquinolinone **161** with an excess amount of hydrazine hydrate in boiling DMF in 60% yield. Interestingly, the reaction of compound **160** with hydrazine (1:2) in AcOH did

Scheme 62 Synthetic pathway to generate pyrazolo and bipyrazoloquinolone.

Scheme 63 Reaction of 160 with NH₂OH·HCl in different conditions.

not give either compound **162** or compound **163** but yielded the N-acetyl derivative of compound **163** and formulated as binary bipyrazoles **164**. Annotative synthesis of **164** was preceded by reaction of **161** with NH_2NH_2 in AcOH or treatment of compound **162** with NH_2NH_2 in boiling DMF (Scheme 62). ¹⁸⁸

Reaction of equimolar amounts of **160** with NH₂OH·HCl in a mixed solvent of ethanol–DMF led to the formation of 4-hydroxy[5-(2-hydroxyphenyl)isoxazolyl]acryloylgquinolinone **165**. Additionally, the reaction of both compounds **165** with NH₂OH·HCl in boiling DMF or pyridine afforded 4-hydroxy-3-[5-(2-hydroxypheny1)biisoxazoly1]quinolinone **166**, which is obtained directly from **160**, when using an excess of NH₂-OH·HCl (1:2) in refluxing pyridine. Additionally, polycyclic skeleton **167** was obtained upon reacting isoxazolylacryloy1quinolinone **165** with NH₂NH₂ in boiling DMF.

Moreover, boiling compound **167** in AcOH led to acetylation, yielding acetyl-5-[5-(2-hydroxypheny1)isoxazol-4-y1]pyrazol-3-y1-4-hydoxyquinolinone **168**, a compound that was also obtained during heterocyclization of the acryloy1 derivative **165** by means of hydrazine hydrate in AcOH. In the same line, the reaction of *N*-acetylpyrazoline **162** with NH₂OH·HCl furnished the same compound **168** (Scheme 63).

Whereas, the reaction of **160** with an equimolar amount of NH₂OH·HCl in AcOH proceeds in a different manner, and afforded 4-hydroxy-1-methy1-3-[5-(4-oxochromeny1)isoxazol-3-yl)quinolinone] **169**. After that, the reaction of chromeny1isoxazoliny1quinolinone **169** with NH₂OH·HCl in boiling DMF afforded 4-hydroxy-3-[5-(2-hydroxypheny1)biisoxazol-y1] quinolinone **166**. Also, chromeny1isoxazoliny1quinolinone **169** was treated with NH₂NH₂ in boiling DMF yielded 4-hydroxy

Scheme 64 Synthetic pathway to generate pyrimidine, pyrazole and oxazole scaffolds

Scheme 65 Reaction of 160 with cyano thioacetamide in different reaction conditions.

[3-(2-hydroxy phenyl) -pyrazol-4-yl]isoxazol-3-y1-quinolinone **170** (Scheme 63). 188

Equimolar ratio of **160** and thiourea was refluxed in the presence of sodium ethoxide, affording thioxopyrimidinquinolin-2-one **171** under various reaction conditions. In refluxing EtOH containing a catalytic amount of HCl or in refluxing AcOH, the same reaction resulted in the formation of chromenthioxopyrimidinquinolinone **172**. The reaction of compound **172** with both hydrazine hydrate and $NH_2OH \cdot HCl$ proceeded smoothly using ethanolic potassium hydroxide to afford pyrazolyl **173** and its isoxazolyl analogue **174**, respectively (Scheme 64).¹⁸⁸

2-Thioxopyridine-3-carbonitriles have great interest due to their usage as intermediates for the synthesis of various medicines. Thus, synthesis of 4-hydroxy-1-methyl-2-oxoquinolin thioxo-3-carbonitrile 175 was achieved by reaction of an equimolar amount of cyano thioacetamide with compound 160, using sodium ethoxide as a catalyst. Whereas, in a relatively moderate basic medium using piperidinium acetate, the reaction approaching another possible product accessible by

a Michael route led to the formation of 6-(4-hydroxy-2-oxo-1,2-dihydroquinolinyl)-4-(4-oxo-4*H*-chromen-3-yl)-2-thioxopyridine-3-carbonitrile **176.** The heterocyclization reaction of both compounds **175** and **176** with NH₂OH·HCl, in boiling pyridine and/or DMF, yielded two important triheterocyclic systems: 5-[3-(4-hydroxy-2-oxo-quinolin-3-yl)-4,5-dihydro-isoxzolyl]-6-(2-hydroxyphenyl)-2-thioxopyridine-3-carbonitrile **177** and 6-(4-hydroxyl-2-oxoquinolin-3-yl)-4-[5-(2-hydroxyphenyl)-isoxazolyl]-2-thioxopyridine-3-carbonitnile **178** (Scheme 65).¹⁸⁸

2-Thiobarbituric acid is recognized as a reactive cyclic methylene compound of the 1,3-dione type, which is utilized to synthesize pyranopyrimidines when cyclized with enone systems or 1,3-dicarbonyl compounds, and more recently with 3-substituted chromones. When compound 160 was treated with 2-thiobarbituric acid in an equimolar ratio in the presence of sodium ethoxide, it yielded (4-hydroxy-2-oxoquinolin-3-yl)-3-oxopropenyl]thioxopyrano[2,3-*d*]pyrimidinone 179.

While by, the same reaction occurred in the presence of piperidinium acetate, a polycyclic molecule, thioxopyrano[2,3-d] pyrimidinone derivative **180** was obtained. Furthermore, both

Scheme 66 Reaction of 160 with thiobarbituric acid in different reaction conditions

Scheme 67 Reaction series of enaminone 183 with different heterocyclic reagents.

179 and 180 were reacted with NH₂OH·HCl in DMF affording the [3-(4-hydroxy-2-oxoquinolin-3-yl)isoxazol-5-yl]-7-(2-hydroxyphenyl)-2-thioxopyrano[2,3-d]-pyrimidinone 181 and (4-hydroxy-2-oxo-quinolin-3-yl)-5-[5-(2-hydroxyphenyl)isoxazol-4-yl]-2-thioxopyrano[2,3-d]pyrimidinone 182, respectively (Scheme 66). 188

4.3.2. With DMF/DMA. *N,N*-Dimethylformamide dimethyl acetal (DMFDMA), also known as 1,1-dimethoxy-*N,N*-dimethylmethanamine, The chemical structure of DMF/DMA has two significant sites: a partially positive carbon that functions as an electrophile in condensation processes and a partially

negative nitrogen that acts as a nucleophile. Owing to its structure, DMF/DMA reacts with a wide range of organic groups and can be a useful reagent as a C1 synthon, particularly in the production of heterocycles. Also, it is used to formylate aromatic compounds. The process involves initial conversion of DMF to a chloroiminium ion, [(CH₃)₂N=CH(Cl)]⁺, known as a Vilsmeier reagent, which attacks arenes. Thermal condensation of **12b,d** with DMF/DMA, afforded the corresponding dimethylaminoacryloylquinolinone derivatives as (enaminone compound) **183**. ¹⁹⁰ Reaction of enaminone **183** with 4-aminoantipyrine, furnished 3-(3-((1,5-dimethyl-3-oxo-2-phenyl-2,3-dihydro-1*H*-

Plausible mechanism for synthesis of compond 187 and 189

Scheme 69 Coupling of enaminone 183 with 4-nitrobenzenediazonium chloride followed by cyclization.

pyrazol-4-yl)amino)acryloyl)-4-hydroxyquinolinone 184. Whereas, the reaction of the enaminone 183 with morpholine and piperidine as cycloaliphatic secondary amines smoothly proceeded, leading to the corresponding 4-hydroxy-1-alkyl-3-(3morpholino/piperidinoacryloyl)quinolinone 185.

While by, treating enaminone with hippuric acid (N-benzoylglycine) or aceturic acid (N-acetylglycine) 186 in Ac₂O resulted in the production of 187. It is suggested that this process begins with the cyclization of 186 into oxazolone, which then adds to the activated double bond system of enaminone 183, followed by further spontaneous rearrangement yielding 187.177,191

Furthermore, compound 189 was synthesized through condensing the active methylene group in diethyl acetonedicarboxylate 188 with the carbonyl group in compound 183, via water elimination and the formation of an intermediate 189B. This intermediate is then cyclized in the presence of AcONH₄, resulting in the stripping of a water molecule yielding angular tetracyclic compound 188 (Scheme 67).190

A probable mechanism for the synthesis of 187 started with the cyclization of hippuric acid into oxazolone 186A, which then adds to the activated double bond system of the enaminone 183, yielding 187A, followed by further rearrangement of this

intermediate to give 187. Whereas, compound 189 is thought to be formed via condensation of the active methylene in 188 with the carbonyl function in 183 with water elimination, forming the intermediate 189A that was cyclized in the presence of ammonium acetate via elimination of a molecule of EtOH and water to afford the final product 189 (Scheme 68).

Biological activities of pyridazinone ring systems represent one of the most active classes of heterocycles. Consequently, enaminone 183 was employed as a promising building block for synthesizing pyridazinone 191. Compound 183 coupled with 4nitrobenzenediazonium chloride, leading to the formation of the arylazo intermediate 190A, which undergoes hydrolysis during the reaction to produce α -hydrazono- β -oxopropanal **190**. Cyclization of the hydrazone derivative 190 with hippuric acid 186 in Ac₂O produced the 4-benzoylamino-6-[(quinolin-3-yl) carbonyl]-pyridazinone 191 in 66% yield. Whereas, the reaction of compound 183 with malononitrile in absolute ethanol in the presence of piperidine yielded carbonitrile derivative 192 in 55% yield (Scheme 69).178

Additionally, treatment of the enaminone 183 with thiourea in ethanolic potassium hydroxide solution did not give the expected pyrimidinylquinolinone 193. Similarly, reaction of enaminone 183 with cyanoguanidine as amidine reagent gave

Reaction series of enaminone 183 analogues synthesis.

Scheme 71 Reaction of enaminone 183 with binucleophilic reagents.

another pyrimido[5,4-c]quinolinone derivative 194. Moreover, when enaminone 183 was heated in AcOH, it went through intramolecular cyclocondensation, resulting in the formation of 6-methylpyrano[3,2-c]quinolindione **195**. Finally, when enaminone 183 was reacted with compounds containing active methylene like malononitrile and ethyl cyanoacetate in the presence of KOH, it yielded the pyranoquinolinone derivatives 196 (Scheme 70).178

Moreover, treatment of the enaminone 183 with thiosemicarbazide and thiocarbodihydrazide yielded 197. The cyclocondensation of 7-chloro-4-hydrazinoquinoline 88 with enaminone 183 in acetic acid resulted in the formation of 4hydroxy-1-methyl-3-[1-(7-chloroquinolin-4-yl)-1H-pyrazol-3-yl] quinolinone 198. The proposed structure for compound 198 was further validated through an alternative synthesis starting from acetylquinolinone 12b. Specifically, treating compound

Scheme 72 Reaction of enaminone 183 with different hydrazine derivatives.

205 (68%)

Scheme 73 Synthesis of fused heterocyclic scaffolds 202, 203, and 205

205A

Scheme 74 Synthesis and cyclization reactions of compound 206.

12b with compound **88** yielded hydrazone **199**. Subsequently, the *in situ* thermal condensation of hydrazone **199** with DMF/DMA smoothly produced compound **198** (Scheme 71).¹⁷⁸

Reaction of enaminone **183** with different hydrazines and hydrazides, such as hydrazine hydrate, phenylhydrazine, and 4-amino-5-hydrazineyl-4H-1,2,4-triazole-3-thiol afforded binary pyrazole **200** in good yield. Whereas, treatment of **183** with heterocyclic hydrazine as 2-hydrazineyl-1H-benzo[d]imidazole or 2-imino-2H-chromene-3-carbohydrazide yielded chemoselective fused pyrazole (E)-3-(2-(dimethylamino)-5-phenyl-4H-pyrazolo[4,3-c]quinolin-4-one (**201**) (Scheme 72).

Reaction of enaminone **183** with triethylenetetramine (TETA) as highly active linear polyamines in CH_2Cl_2 led to the formation of benzo[h][1,6]naphthyridine-4,5-dione derivative **202** instead of the anticipated transamination products. Furthermore, refluxing of **183** with malononitrile led to the formation of tetrahydrobenzo-[h][1,6]naphthyridine-3-carbonitrile **203** through 1,2-addition mechanism followed by the nucleophilic attack of OH group to the cyano function. After that, the Dimroth rearrangement was achieved instead of an alternative path

involving the replacement of the active methylene with the dimethylamino group (Scheme 73). Refluxing of enaminone **183** with pyrazolone **204** in AcOH gave fused triazaindenoanthracenol **205** (Scheme 73). The mechanistic pathway for the formation of **205** was recognized through a condensation reaction afforded 1,5-dicarbonyl intermediate **205A**, followed by cyclization reactions of intermediate **205A** through double nucleophilic attacks of the both two OH groups to the endoketonic carbonyl group of quinolinone and β -position of enaminone led to affording the fused polycyclic system **205** in 68% yield (Scheme 73).¹⁹²

4.3.3. With esters. Compound **206** was synthesized through the condensation of quinolinone derivative **12b** with methyl/ethyl propionate in the presence of Na metal. This reaction occurred through the nucleophilic attack of the active methylene group on the carbonyl group of the ester reagent. Cyclization of compound **206** with sulfuric acid gave 2-ethyl-6-methylpyrano[3,2-*c*]quinolindione **207** (Scheme 74).¹⁵¹ The bifunctional ammonia derivatives such as semicarbazide, thiosemicarbazide or aminoguanidine reacted with **206** in

Scheme 75 Claisen condensation of compound 12b,d with different esters

Scheme 76 Synthesis of pyrazolonyl moiety anchored on quinolone 213 and 214

Scheme 77 Synthesis of furo[3,2-c]quinolone 215.

refluxing ethanol to furnish a binary triazepinone, triazepinthione and iminotriazepine **208**, respectively. Furthermore, reaction of compound **206** with hydrazinoquinolinone **122** afforded the intended dihydromethylquinolinyl)-3-pyrazolyl)-4-hydroxyquinolinones **209** (Scheme 74).¹⁷⁶

Additionally, Claisen condensation reaction of **12b,d** with diethyl carbonate led to **1,**3-diketone; 3-ethoxycarbonylacetyl-4-hydroxyquinolin-2-one **210** in the presence of sodium metal. Cyclization of compound **210** with sulfuric acid afforded 4-hydroxypyrano[3,2-*c*]quinolindione **11b,d**. In a similar process, Claisen condensation of compound **12d** with diethyl oxalate in the presence of Na metal produced the diketo-ester; (ethylhydroxy-1,2-dihydroquinolinyl)dioxobutyrate **211**. Cyclization of **211**, in dry pyridine, or AcOH/AcONa (freshly fused) yielded ethyl 6-ethyldihydropyrano[3,2-*c*]quinoline-2-carboxylate **212** (Scheme **75**). ¹⁵¹

The synthesis of binary pyrazolylquinolinone **213** was achieved by reacting the β -ketoester **210** with hydrazine hydrate to

give the target compound **213** in 48% yield.^{193a} While, direct treatment of pyranoquinolindione **11** with hydrazine hydrate gave the desired pyrazolylquinoline **213** in 92% yield. The possibility of forming the isomeric quinolinylpyrazolin-3-yl) quinolinone **214** was ruled out by independently synthesizing this derivative through the cyclization of hydrazinoquinolinone **122** with 3-ethoxycarbonylacetyl-4-hydroxy-1-methylquinolinone **210** in 74% yield (Scheme 76).^{176b}

Whereas, the reaction of acetylquinolinone **12b** with ethyl (triphenylphosphoranylidene)acetate in refluxing xylene afforded **215A** followed by bromination to give compound **215B** that underwent hydrolysis gave furo[3,2-*c*]quinolinone **215** (Scheme 77). 193*b*

The proposed reaction mechanism for the molecular rearrangement of **215B** to acids **215** begins with the opening of the pyrane ring in a basic medium, resulting in the formation of dianion **A**. Tautomer **B** is considered more likely than **A** due to the greater distance between the two negatively charged

RSC Advances Review

Scheme 78 Plausible mechanism for the synthesis of 215

Scheme 79 Reaction of guinolone 12b with ethyl (triphenylphosphoranylidene)chloroacetate.

functional groups. Following this, the elimination of a halogen atom from the sp² hybridized carbon atom occurs alongside the migration of the substituent in a trans position relative to the departing halogen. This migrating group exhibits partial carbanionic character due to its potential tautomerism. This rearrangement, which can be classified as a Wagner-Meerwein type, leads to the formation of intermediate C, which subsequently yields compound D and ultimately product 215 after acidification (Scheme 78).

Treatment of 12b with ethyl (triphenylphosphoranylidene) chloroacetate in refluxing xylene afforded a complex mixture of compounds 216 and 217. Compound 216 was isolated by repeated column chromatography and identified as an ethyl

ester. The second reaction product 217 contained an unsaturated lactone moiety (Scheme 79).193,194

4.3.4. With activated olefins as synthetic routes to pyrano skeletons and their benefits (vlidene nitriles). A Michael reaction refers to a type of organic chemistry reaction involving the addition of a nucleophile (Michael donor) to an activated olefin or alkyne (Michael acceptor), such as an acrylate. This reaction is known for its rapid occurrence, tolerance of various functional groups, and its utility in synthesizing novel step-growth compounds with tailored macromolecular architectures. Derivatives 219 are similarly synthesized via Michael type addition of the anion ion of the acetyl group in 12c,e to the activated double bond in ylidenenitriles 218 in (1:1) or (1:2)molar ratios in ethanol/piperidine to yield the acyclic intermediate 219A, which dehydrogenated and undergo cyclization into the adducts 219B which afford compounds 219 under basic conditions by adding one molecule of CH₂(CN)₂, which exists in equilibrium with two moles (Scheme 80).191

On the other hand, the nucleophilic addition of 3-acetylquinolinones 12b,d to 220 in EtOH, using piperidine as a catalyst, produced intermediate 221A. This intermediate underwent hydrolysis, losing the acetyl group, to form quinolone intermediate 221. Subsequently, cyclization of this

OH O CN CN EtOH, piperidine reflux
$$\frac{R_1}{R}$$
 CN $\frac{EtOH, piperidine}{reflux}$ $\frac{R_1}{R}$ CN $\frac{R_1}{R}$ CN

Scheme 80 Michael addition of compounds 12c.e.

Scheme 81 Catalyzed nucleophilic addition of 3-acetylquinolinones 12b,d

Scheme 82 Synthetic pathway to access tetracyclic systems 222 and 223.

intermediate yielded 2-amino-5-oxopyrano[3,2-c]quinoline derivatives **221** (Scheme 81).^{115,134,195}

2-Amino-4-(2-chlorophenyl)oxopyrano[3,2-c]quinolinecarbonitrile **221**, as a typical enaminonitrile derivative, reacted with HCOOH to afford tetracyclic 5-methylpyrimidopyrano[3,2-c]quinolindione **222**. Compound **221** also reacted with Ac₂O to yield 5,10-dimethylpyrimidopyrano[3,2-c]quinolindione **223** (Scheme 82).¹⁹⁵

6-Methyldioxodihydrospiro[indoline-3,4'-pyrano[3,2-*c*]-quinoline]-3'-carboxylate **225** was synthesized through the reaction of compound **12b** with **224** in EtOH, pyridine (Scheme 83).¹¹⁵

4.4. Miscellaneous reactions

4.4.1. Deacetylated quinolinone (HQ) and its utility. Hydrolysis reaction of acetylquinolinone **12b–f** with sulfuric acid, accompanied by deacetylation produced 4-hydroxyquinoline derivatives (HQ) **15** (Scheme 84). 115,130,132,141,145,155

Refluxing of 4-hydroxyphenylquinolinone 15 with POCl₃ for 2 h yielded the corresponding 4-chloroquinolinone derivative 226, reaction of 226 with three equivalent piperazine in DMF in the presence of TEA was heated at 80 °C for 6 h yielded 4-piperazinylquinolinone 227, heating 227 with formaldehyde and the appropriate amine in EtOH at reflux then leave overnight afforded the target *N*-Mannich bases 228. Alkylation of the piperazinyl nitrogen of compounds 227 with 2-bromo-*N*-phenylacetamide derivatives 229 was achieved by stirring overnight in DMF using K₂CO₃ as a base to yield the target compounds 230 (Scheme 85).¹⁹⁴

Refluxing of 15 with propargyl bromide in acetone in the presence of anhydrous K₂CO₃ for 14–16 h afforded a mixture of 231, 232, and 233, which were separated by HPLC.¹⁹⁶ Basecatalysed Claisen rearrangement of propynyloxyquinolone 231 in hexamethylphosphoric triamide (HMPT) in the presence of NaHCO₃ (2 equiv), refluxed for 10–18 h afforded angular furo [3,2-c]quinolinone 234 according to the shown mechanism (Schemes 86 and 87). Whereas, 231 products followed by

Scheme 84 Deacetylation of 3-acetyl-4-hydroxyquinolone derivatives 12b-f.

Scheme 83 Synthetic approach of spiro compound 225.

Scheme 85 Chloronation of 15 followed by synthesis of N-Mannich base

Scheme 86 O-alkylation followed by Suzuki coupling reaction.

Scheme 87 Mechanistic pathway for the synthesis of 234.

Sonogashira coupling yielded 4-((3-arylprop2-yn-1-yl)oxy) quinolin-2-onederivatives 235 in good yield. Then, electrophilic cyclization between 235 and iodine to obtain 236, the reaction was initially performed in various solvents (DMF, CH₃CN, CH₃OH, THF, DCM, and CH₃NO₂) at 25 $^{\circ}$ C, using sodium bicarbonate as a base (Scheme 86).¹⁹⁷

Benzoylation reaction of **15a-c** with benzoyl chloride and a catalytic amount of TEA afforded 4-benzoyloxy-2-quinolinones **237**. ¹⁹⁸ Compound **237** underwent Fries rearrangement with

AlCl₃, yielding 3-benzoyl-4-hydroxy-2-quinolinones **238** in excellent yields. Reaction of β -hydroxyketone compound **238** with phenylhydrazines in dimethylformamide (DMF) at room temperature and also at 0 °C yielded mixtures of compounds that can be separated, which contained cyclized pyrazolo[4,3-c] quinolinones **239**. To obtain pyrazolo[4,3-c]quinolinones **239** directly from 3-benzoyl-4-hydroxy-2-quinolinones **240** pure and in good yield (87–99%). A suspension of 3-benzoyl compounds **238** was treated in AcOH with a few drops of conc. H₂SO₄ to

NHNH₂ ОН PhCOCI/ TEA AICI₃/ neat 15a,c toluene/ reflux N R R 239 R₁ 238 237 (%) R (%) H CH₃ CH₃ C₆H₅ C₆H₅ 3-CI H 4-NO₂ H 4-NO₂ 3-CI 52 72 85 68 95 98 90 99

Scheme 88 Synthesis of pyrazolo[4,3-c]quinolinones 239 through o-benzoylation.

Scheme 89 Synthesis of chlorohydrazonoquinolinones 242.

enhance the reaction rate and obtain good to excellent yields. Heating for 2 h yielded pyrazolo[4,3-c]quinolones 239 (Scheme 88). 135

The synthetic pathway for compound **242** begins from 3-phenylaminomethylene quinolindiones **240**, which were synthesized from 4-hydroxy 2-quinolones, aniline and triethyl

				QΗ	I										
	OH		ОН) OH	4							Ó	O, R	2
	OHC R		N R 243	00	N-R	I	R ₁						N-N N	N	R_3
15 —	0 244 CHO R ₁		Br i		OH		R ₃	246	N ₃	R ₁ \		OH	0	ОН	R ₁
	ii i) TEA/EtOH/reflux; ii) DMF/Na ₂ CO ₃ /Stiri		R 245	00	N R		J	Stir	F/ Cul r. 24h			R 247		N R	
		R	R ₁	(%)		R	R ₁	R ₂	R ₃	(%)	R	R ₁	R ₂	R ₃	(%)
		H H CH₃	H ОСН₃ Н	85 90 78		H H H	Н Н Н	Н Н Н	H CH ₃ OCH ₃	75 80 82	H H CH₃	OCH ₃ OCH ₃ H	H CH ₃ H	OCH₃ H H	83 69 72
				-		H H	H OCH₃	CH ₃ H	H H	70 77	CH ₃ CH ₃	H	H	CH ₃ OCH ₃	77 71
						ΗΙ	OCH ₂	Н	CH₃	82	CH ₃	Н	CH ₃	Н	66

Scheme 90 Synthesis of triazolyl-bis(4-hydroxy-quinolin-2(1H)-ones) 247.

1mol of 15 ОН СНО 249(%) R ĊH₃ н 91 248 3-CI 84 reflux 4-CI 83 2-OH 72 ĊН₃ 76 OH 4-OH 15a,b 3-NO₂ 85 1mol of 15 or 85 4-NO 2mol of 15 4-OCH 75 3,4-(OCH₃)₂ ĊН₃ ĊH₃ 249

Scheme 91 Reaction of 4-hydroxyquinoline derivatives 15 with various aromatic aldehydes.

Scheme 92 Mechanistic pathway for the synthesis of 249

Scheme 93 Synthesis of 250 and 252 via Knoevenagel-Michael reaction.

orthoformate (CH(OEt)₃), after that converted to 4-chloro-3-formylquinolinones **241**.^{135,199} The reaction of carboxaldehydes **241** with a specific amount of phenylhydrazines in a solution of DMF at room temperature or in refluxing BuOH afforded 4-chloro-3-arylhydrazinonoquinolinones **242** in good to excellent yields (Scheme 89).¹³⁵

The synthesis of a new series of triazolylquinolinones, specifically 3,3'-(oxoquinolin-4-yl)-1,2,3-triazol-4-yl)methoxy) phenyl)methylene)bis(4-hydroxyquinolinones) 247, was achieved by reacting 4-azido-2-quinolinones 246 with 3,3'-(prop-2-yn-1-yloxy)phenyl)methylene)bis(4-hydroxyquinolinones) 245 through a Cu-catalyzed [3 + 2] cycloaddition (Huisgen-Meldal-

Sharpless reaction). Initially, the synthesis of terminal alkynes 245 was performed via interaction between 4-hydroxyquinoline-2-ones 15 and p-hydroxybenzaldehyde in different molar ratios under reflux to afford 3,3'-(hydroxy-phenyl)methylene)bis(4-hydroxyquinolin-2-ones) 243. Phenol compound 243 reacts with propargyl bromide in DMF to produce alkynes 245. On the other hand, 4-hydroxy-2-quinolinones 15, interacted with aldehyde 244 in different molar ratios (2:1) to produce the desired terminal alkynes 245 (Scheme 90).

Reaction of compound 15 with several aromatic and heterocyclic aldehydes, which contain both electronwithdrawing and electron-donating groups, was investigated Review RSC Advances

Scheme 94 Suggested mechanism for the synthesis of 250 and 252.

Scheme 95 Photoinduced reaction of compound 15b with TEA producing 254a and the Plausible mechanism.

under catalyst-free conditions using water as the eco-medium to produce compound **249**, through Knoevenagel condensation followed by a Michael-type addition with compound **15**, leading to the smooth formation of a wide range of substituted bisquinolinones. In these cases, aldehydes with electron-withdrawing groups produced higher product yields, ranging from 85% to 90%. The study revealed that compound **249** was the primary product, rather than compound **248**, when using different moles of compound **15** (Scheme 91).²⁰¹⁻²⁰⁴

A believable mechanism for the formation of compound 15 has been suggested. Initially, one equivalent of HQ 15b undergoes a straightforward Knoevenagel condensation with aromatic and heterocyclic aldehydes to produce carbonyl intermediate 249A. This intermediate serves as a strong Michael acceptor and reacts with another equivalent of 15b through Michael addition, resulting in the keto-enol intermediate 249B,

followed by isomerization to yield the final product (Scheme 92).²⁰¹

Furthermore, treatment of **15b** with the aromatic and heterocyclic aldehydes in different molar ratios (2:1) led to the formation of **250**. Whereas, the three-component Knoevenagel–Michael reaction of **15b**, aldehyde, and aminocyclohex-2-enone **251** produced skeleton **252**. But, when the reactants were added sequentially, only a small amount of the desired product **252** was obtained, with compound **250** being the major product (Scheme 93).²⁰³

A plausible mechanism for the formation of compounds 250 and 252 started with simple Knoevenagel condensation of 15 with the aromatic or heterocyclic aldehydes (known to occur under solvent and catalyst-free conditions) to generate an adduct 253, which acts as a strong Michael acceptor. After that, another molecule of HQ 15 (two-component reaction) or 3-

RSC Advances Review

Scheme 96 Oxidation of guinolone derivatives 12b,d.

Proposed mechanism for the synthesis of 255 and 256

aminocyclohex-2-enone (three-component reaction) attacks the electron-deficient β position of 253 in a Michael addition fashion to afford 250 and 252 (Scheme 94).203

Pyrano[3,2-c]quinolinone derivatives represent a significant category of naturally occurring alkaloids. As their synthetic counterparts are currently the focuses of research due to their diverse biological activities with potential medicinal applications. Additionally, these compounds are frequently employed as synthetic precursors in producing dimeric quinoline and polycyclic heterocycles.205 While by, 2-hydroxydimethylpyrano [3,2-c]quinolin-5-one (254a) was synthesized by a photoinduced reaction of 15b in the presence of TEA, which generated DEA and acetaldehyde derivatives in situ, whereas, photoinduced electron-transfer reactions with TEA as an monodentate electron donating through a redox reactions were achieved according sketeched mechanism in Scheme 93.206 Reaction of 4hydroxyquinolinone 15 with aliphatic aldehyde gave the corresponding quinolone methide, base-catalyzed condensation of a quinolinone 15 with an aldehyde yields the corresponding 4hydroxy-3-(1-hydroxyethyl)quinolin-2-one 254A which underwent dehydration in basic medium to furnish the highly electrophilic quinone methide intermediate 254B. After that, Michael's addition reaction at the exocyclic methylene carbon of quinone methide 254B by the generated enamine (from DEA and the aldehyde) and the carbanion 15A derived from the deprotonation of 15. The Michael-type addition of the enamine

proceeds in a 1,4-fashion and results in an intramolecular cyclization to yield 2-(diethylamino)pyrano[3,2-c]quinolin-5-one 254C which hydrolyzed to afford 254 (Scheme 95).206

4.4.2. Oxidation. Oxidation of quinolone derivatives 12b,d under the classic Riley conditions afforded a mixture of two products: α -keto acid 255 and its dehydrated dimer derivatives 256. The major product is soluble in aq. Na₂CO₃; besides, it can be isolated after neutralization to yield the α -keto acid 255, whereby the minor product is insoluble in aq. Na₂CO₃ solution, but soluble in boiling NaOH and crystallizable from AcOH and identified as spiropolyketonic scaffold 256 (Scheme 96).207

A probable mechanism for the synthesis of the α -keto acids 255 can be simply explained by the fast oxidation of quinolone derivatives **12b,d** to α -keto aldehyde intermediates **255A**, which cyclize into the hemiacetal intermediates 255B, which can afford the corresponding 5-alkylfuro[3,2-c]quinolinetriones 255C. These furoquinolinetriones 255C and 255D are easily hydrolyzed, furnishing the formation of 255 as a major product and 256 as a minor product (Scheme 97).

Specifically, the reactivity of 255 with various nucleophilic reagents was investigated, as the reaction of 255 with p-toluidine in refluxing EtOH with a few drops of piperidine produced (4-hydroxy-2-oxoquinolin-3-yl)-p-tolyliminoacetic Similarly, reaction of 255 with p-phenylenediamine in DMF bis[(hydroxy-1,2-dihydroquinolinyl)methylidene]-1,4yielded phenylenediamine 258. While condensing α -ketoacid 255 with

Review **RSC Advances**

Scheme 98 Reaction of α -keto acids **255** with different reagents

Reduction of the acetyl group of 12b to form compound 263, in addition to the plausible mechanism of the Clemmensen reduction.

thiocarbohydrazide in different molar ratios in boiling EtOH containing TEA afforded bis[2-(4-hydroxy-2-oxoquinolin-3-yl)-2oxoacetic acid]thiocarbohydrazone 259. Additionally, the reaction of α -ketoacid 255 with various carbon nucleophiles was explored through its reaction with active methylene such as dichloroacetic acid, chloroacetonitrile, and diethyl malonate, which led to the synthesis of quinoline-3-carboxylic acids 260 and fused tricyclic scaffolds (3-chloro-2-hydroxymethylbenzo) carboxylic acid **261** and 2,5-dioxopyrano[3,2-c]quinoline **262**, respectively. Whereas, the hydrolysis process of quinoline-3carboxylic acids 260 by NaOH was continued for a day, yielding 4-hydroxy-2-oxoquinoline 15 (Scheme 98).208

4.4.3. Reduction of AHQ. Whereas, the reduction is one of the most effective chemical transformation processes in organic

chemistry209 as the development in this field has been tremendous, progressing from the use of stoichiometric reagents to get novel organic systems, whereas, Clemmensen-type reduction of acetylquinolinone 12b in the presence of zinc dust (particle size <45 µm) afforded 3-ethyl-4-hydroxyquinolinone 263 in acceptable yield 75% (Scheme 99).210 The Clemmensen reduction is a chemical reaction used to convert a carbonyl group (C=O) into a methylene group directly. Firstly, the carbonyl group is protonated by acidic hydrogen to be more electrophilic and easily attacked. Then, nucleophilic Zn attacks the protonated carbocation, leading to the formation of a tetrahedral carbinol containing the carbon-zinc bond, which is protonated by HCl through a carbenoid mechanism. After that, the reduction process happens on the surface of the zinc metal, leading to the

он он substituted NaBH₄ phenyhydrazine MeOH k 264 265 12b,c (%) (%) CH₃ 3-CI CH₃ 2-CH₃ 3-CI 2-CH₃ 95 CH₃ 84 CH₃ 3-CH₃ 82 3-CH₃ CH₃ 91 4-CN 90 76 93 4-CN 4-CH₃ C₆H₅ CH₃ 4-CH₃ 80 CH₃ 4-Br 2,4-CI 77 CH₃ 4-Br 78 80 2,4-CI C₆H₅ 2,4-CH₃ CH₃ 87 90 80 75 CH₃ 2,4-CH₃ н 3-NO₂ 3-NO₂ CH₃ 2,4-NO₂ 92 78 2,4-NO₂ 82

Scheme 100 Reduction of compounds 12b,c using NaBH₄

formation of the corresponding alkane through undergoing two sequential 2e⁻ reduction steps involving a dehydration step and anionic intermediates (Scheme 99).²¹¹

Reduction of derivatives 12b,c via using sodium borohydrate (NaBH₄) as hydrogen source and electron donor afforded 4-hydroxy-3-(hydroxyethyl)quinolinones 264. Whereby, a chemoselective reduction of 264 was achieved by the treatment of 264 with substituted phenylhydrazines, yielding hydrazono-3-(1-hydroxyethyl)quinolin-2-one derivatives 265 (71–92%) (Scheme 100).

5. Applications

5.1. Medicinal perspective of quinolinones

Quinolinone derivatives are significant heterocyclic systems with multiple medicinal applications. As these compounds possess various pharmacological properties, anti-inflammatory, including analgesic effects, including anti-inflammatory, including antialler-genic, including antipolar antipolar effects, including analgesic effects, including effects, including effects, including effects, in

OH N CN OH N O O	H ₃ CO N N N N N N N N N N N N N N N N N N N	H ₃ C COOH Levofoxacin Antibacterial
O O O O O O O O O O O O O O O O O O O	OH O H S F N N N N N N N N N N N N N N N N N N	Moxifloxacin Antibiotic ²⁴⁸⁻²⁵⁰
F COOH N N N Ciprofloxacin Antibiotic 51	H ₃ C N N O COOH HN O Gatifloxacin Antibiotic ¹⁰⁶	OH O NH O H Anticancer activity ¹⁶⁸
NC NO CH ₃ Anticancer ²⁵²	OH O H N S CI Ph N CI Anticancer and antimicrobial 168	OH O N H H CH ₃ Linomide Anticancer ²¹⁶

Fig. 6 Biologically active compounds containing quinolinone scaffolds.

Review RSC Advances

Fig. 7 Design an approach to increase anti-tuberculosis activity.




Fig. 8 The history of quinolinones and modifications to improve their pharmacokinetics

antioxidant, ^{107,192,216,242} anti-tubercular activity ^{106,243–246} and other potential biological applications ^{247–252} as shown in Fig. 6.

Globally, tuberculosis (TB) is a highly lethal infectious disease caused by the bacterium Mycobacterium tuberculosis, which is responsible for approximately three million deaths. The recent report from the World Health Organization (WHO) identifies TB as the second leading cause of death from infectious diseases worldwide. About one-third of the global population is at risk due to latent infections with Mycobacterium tuberculosis, according to the WHO. 106,253,254 On the other hand, quinolines are a crucial active pharmaceutical ingredient that plays a vital role in discovering new drug candidates. Many quinoline-based compounds are currently in clinical and preclinical development for tuberculosis treatment. Additionally, several quinoline-derived medications, such as moxifloxacin, gatifloxacin, and TMC207, are utilized to treat tuberculosis.²⁵⁵⁻²⁵⁷ Interestingly the incorporation of pyrazole derivatives into the quinolinone scaffold enhances its biological efficacy, as it alters modes of action, improves selectivity profiles, and reduces unwanted side effects. Those compounds may be further changed to have better pharmacokinetics and oral bioavailability (Fig. 7).106,258

The accidental discovery of nalidixic acid during chloroquine synthesis and its role in the development of numerous quinolinone analogues, including flumequine, rosoxacin, ofloxacin, ciprofloxacin, moxifloxacin, levofloxacin, trovafloxacin, and marbofloxacin. All of the aforementioned pharmaceutical candidates demonstrated significant antibiotic activities.259-261 Over the years, the history of quinolones is characterized by numerous iterations, innovations, and expansions, as evidenced by the many potent drugs available today. In 1962, nalidixic acid (1st generation) was discovered and then approved in 1967 for treating uncomplicated urinary tract infections (UTIs), but resistance quickly developed among various species.262 Due to adverse bioeffects as low serum concentrations and high minimum inhibitory concentrations, nalidixic acid was largely abandoned till the emergence of fluoroquinolones in the 1970s and 1980s. 263,264 Modifying quinolones fluoroquinolone scaffolds improved pharmacokinetics and expanded their antimicrobial spectrum.263,264 Interestingly, certain second-generation marketed antibiotics, such as ofloxacin, ciprofloxacin, and norfloxacin, are still in use today (Fig. 8).250

RSC Advances Review

Fig. 9 Quinolinone scaffolds with biological efficacy.

Fleroxacin signifies the start of the "3rd generation" of quinolinones, which demonstrated meaningfully enhanced antimicrobial activity due to various modifications in its chemical structure, as shown to be photocarcinogenic and photomutagenic. Finally, Moxifloxacin, the "4th generation" of quinolones, is characterized by robust antimicrobial activity, including high efficacy against pathogens due to improved anaerobic coverage. 550,263,266,267

Also synthesized *N*-(3-acetyl-2-oxoquinolin-1(2*H*)-yl)benzamide derivatives **266** were tested for their antitubercular and antimicrobial activities. The majority of the tested derivatives exhibited encouraging antitubercular efficacy in comparison to the standard drugs (isoniazid and streptomycin). The inclusion of electron-donating groups such as methyl, amino, hydroxy, and dimethylamino has enhanced its antitubercular activity. Notably, most of the investigated derivatives of **266** displayed considerable antibacterial efficacy against both Gram-positive and Gram-negative microorganisms. They also showed marked antifungal efficacy against *C. albicans* and *A. niger* (Fig. 9).²⁶⁸

Krishnakumar *et al.*²⁶⁹ reported the evaluation of *in vitro* antibacterial properties for ethyl-2-oxoquinoline-3-carboxylate **267**, showing moderate activity against the *Vibrio cholerae* and *Bacillus subtilis*. Likewise, 1-methyl-3-(3-oxo-3-phenylprop-1-enyl) quinoline-2-(1*H*)-ones **268** were examined *in vitro* as antimicrobial agents exhibiting significant antibacterial activity against *S. Typhi,S. aureus*, *P. aeruginosa*, and *B. subtilis* (Fig. 10).²⁷⁰

Whereas, the next generation of anticancer, antimicrobial, and anti-HIV-1 drugs is expected to be characterized by quinolinone-pyrimidine-based. Anticancer evaluation of pyrimidoquinolinone 269 and pyrimidotetrazinoquinoline 270 was examined *in vitro* against the human liver cancer cell line

Fig. 11 Quinolone-pyrimidine based molecular hybrids as potential anticancer agents.

Fig. 12 Quinolone-pyrimidine based molecular hybrids as potential antibacterial agents.

Fig. 13 Pyrimidinylquinolinones based molecular hybrids as potential anticancer agents.

(HEPG2). The anticancer activity results indicated that these compounds showed inhibitory activity against the tested cell line with IC₅₀ values of 38.30 μ M for 269 skeleton and 39.8 μ M for 270 (Fig. 11).²⁷¹

Compound anilinouracil-fluoroquinolone **271** demonstrated Gram-positive antibacterial potency at least 15 times that of the corresponding [3-(4-hydroxybutyl)-6-(3-ethyl-4-methylanilino)

R	R ₁	R ₂	(%)	R ₂ O	0
H H H Br H	CI H CI CI Br CI	C ₆ H ₅ CH ₃ 2-CI-C ₆ H ₄ 2-F-C ₆ H ₄ H 4-F-C ₆ H ₄	80 85		R CH ₃ 268 I, 3-NO ₂ , 4-NO ₂ , 4-NH ₂ , 4-OCH ₃ , I-CH ₃ , 4-CI, 2-OH, 4-C ₂ H ₅

Fig. 10 Quinolinone skeletons with antibacterial activity.

Agrochemical compounds with quinolinone scaffolds.

Quinolinone scaffolds with agrochemical efficacy.

uracil], effectively inhibited pol IIIC and topoisomerase/gyrase, inhibited gyrase and B. subtilis topoisomerase IV with IC₅₀ of 31 and 43.6 µM, respectively, and, as anticipated, had a selective effect on bacterial DNA. Additionally, this compound showed the ability to attack both of its possible targets in the bacterium by being active against a wide range of Gram-positive pathogens that were resistant to antibiotics, as well as a number of Gramnegative organisms. It was also active against Gram-positive organisms that were resistant to fluoroquinolones and anilinouracils. Moreover, it lacked toxicity in vitro and was bactericidal for Gram-positive bacteria. With a unique dual mechanism of action and strong activity against Gram-positive bacteria that are both susceptible to and resistant to antibiotics, this class of anilinouracil-fluoroquinolone hybrids offers a promising new pharmacophore (Fig. 12).272

Furthermore, the synthesized derivatives of 3-(2-amino-6arylpyrimidin-4-yl)-4-hydroxy-1-methylquinolin-2(1H)-ones 272 had shown the best activity in the series, with anticancer efficacy against HepG2 cell lines (IC₅₀ values of 1.32 and 1.33 μM, respectively) (Fig. 13).131

Agricultural applications

In recent years, the discovery of agrochemicals based on the quinoline scaffold structure has led to great progress. Some new quinoline pesticides that have been commercialized or are under development not only inject new vitality into the market but also can replace unfriendly old pesticides.⁵¹ For example, ethyl-4-hydroxy-1-(4-methoxyphenyl)quinolinone-3-carboxylate 273, has been investigated for the first time as a sensitization

chromophore for Tb(III) to improve selectivity and sensitivity for organophosphorus pesticide detection.273

The insecticidal activity of pyrazologuinolone compounds was investigated, both in vitro and in vivo, against the cotton leafworm, S. littoralis, and cotton aphids. The most effective compounds were (E)-4-(2-(dimethylamino)phenyl)-2,5-dioxo-6phenyl-1,2,5,6-tetrahydrobenzo[h][1,6] naphthyridin-3carbonitrile 203 and 3-(1-(4-amino-5-mercapto-4H-1,2,4-trizol-3-yl)-1*H*-pyrazol-3-yl)-4-hydroxy-1-phenylquinoline-2(1*H*)-one **200**, with LC_{50s} of 119.79 and 164.63 mg L⁻¹ against S. littoralis. 192 Additionally, 1,4-dihydro-quinolinecarboxylate 275 is isolated from Beauveria sp. for the first time and has insecticidal activity against Bemisia tabaci with remarkable toxicity in contact and feeding assays, as its LC₅₀ values were 10.59 μg mL^{-1} (contact) and 5.66 µg mL^{-1} for feeding. Whereas, no adverse effect on plant height/growth or phytotoxicity was detected on pepper, tomato, cotton, and cucumber during the treatment.274 Also, Liu et al.275 reported that the utilization of Schinifoline 276, the fruit pericarp of Zanthoxylum schinifolium possessed remarkable feeding toxicity against Sitophilus zeamais and Tribolium castaneum through reducing their food consumption and growth rate, leading to weakness of their reproductive ability and low resistance (Fig. 14).275

Additionally, a series of (E)-3-acyl-quinoline-2,4-(1H,3H)dione imine derivatives 277 displayed remarkable herbicidal activity against monocot and dicot species by affecting PSII electron transport inhibitors and inhibiting the electron transport chain, leading to the prevention of the production of NADPH and ATP. As the majority of the compounds showed good to excellent herbicidal activities against a number of dicot or monocot species, with an inhibition percentage of over 50%

RSC Advances Review

Fig. 16 Quinolinone scaffolds with photosynthesis-inhibiting activity.

Fig. 17 Quinolinone scaffolds with insecticidal activity

at a dosage of 94 g per ha or even lower. Notably, the compounds featuring short straight-chain alkyl groups display the highest activity, comparable with the longer alkyl chains; insecticidal activity diminishes with increasing chain length (Fig. 15).²⁷⁶

A series of substituted 4-hydroxy-1*H*-quinolin-2-one derivatives **278–280** were analyzed using RP-HPLC to determine lipophilicity and photosynthesis-inhibiting activity (the inhibition of photosynthesis in *Spinacia oleracea* L. As the synthesized quinolinones displayed high to moderate inhibitory effects on

the photosynthesis activity process. In addition, *in vitro* antifungal screening of these scaffolds was evaluated against various fungal strains, displaying moderate efficacy (Fig. 16).²⁷⁷

Quinolactacide 281 is structurally analogous to quinolactacins derived from Penicillium species, which are recognized for exhibiting various biological activities, particularly insecticidal properties. This compound exhibited 88% mortality against the green peach aphid (Myzus persicae) at a concentration of 250 ppm. The insecticidal and miticidal efficacy of quinolactacide was assessed against five distinct insects and one mite at a concentration of 500 ppm. The insects utilized for the experiments included the green peach aphid, silverleaf whitefly (Bemisia argentifolii), diamondback moth (Plutella xylostella), common cutworm (Spodoptera litura), western flower thrips (Frankliniella occidentalis), and two-spotted spider mite (Tetranychus urticae). The synthetic quinolactacide exhibited no activity against the other insects, but it demonstrated 100% and 42% mortality against the green peach aphid and diamondback moth, respectively (Fig. 17).278

5.3. Complex formation

Quinolinones bind metal ions to create complexes where they can function as bidentate, unidentate, or bridging ligands. Quinolinone molecules have a basic side nucleus that becomes

Fig. 18 Coordination modes of quinolinone complexes with high biological activity.

Fig. 19 Bioactive complexes with quinolinone scaffolds.

protonated and shows up as cations in the ionic complexes under highly acidic conditions. Quinolone solubility, pharmacokinetics, and bioavailability are all significantly impacted by interaction with metal ions, which also plays a role in the bactericidal agents' mode of action. Many metal complexes have shown various biological activities, including antifungal, anticancer, antiparasitic, and antimicrobial (Fig. 18).²⁷⁹

A series of Cu(II) **282** and Co(II) **283** complexes were designed and synthesized by Khalaf *et al.*²⁸⁰ as 4-hydroxy-2*H*-pyrano[3,2-c] quinolin-2,5(6*H*)-dione acts as monobasic didentate ligand and form tetrahedral and octahedral complexes at molar ratio of 1: 2. Interestingly, the synthesized complexes displayed greater efficacy as antibacterial, antifungal and antioxidant candidates in comparison with the free pyrano[3,2-c]quinolin-2,5(6*H*)-dione as ligand (Fig. 19). ²⁸⁰

The potential antibacterial and antifungal properties of the quinoline ligands (SL_1 – SL_4) and their $Cu(\pi)$ and $Zn(\pi)$ complexes (Fig. 20) were evaluated *in vitro*. The ligands (SL_1 – SL_4) demonstrated moderate antibacterial activity against Gram-positive bacteria but exhibited no effectiveness against Gram-negative bacteria or fungal strains. Among them, the SL_3 ligand displayed the highest activity, achieving a zone of inhibition of 24 mm against *S. aureus*. However, the antibacterial activity of these ligands was significantly lower than that of the standard drugs amoxicillin and fluconazole. These metal




Fig. 20 $\,$ Hydroxyquinoline functionalized Schiff base copper and zinc complexes ${\rm SL}_{1-4}.$

Fig. 21 The chemical structures of lanthanide complexes.

complexes generally enhanced antibacterial activity against both Gram-positive *S. aureus* and *E. faecalis* strains. On comparing the metal complexes, it was found that $[Cu(SL_1)_2]$ was the most harmful substance to Gram-positive bacteria and had moderate antifungal properties.²⁸¹

Fouad *et al.*²⁸² studied lanthanide nano-complexes (Gd³⁺, Er³⁺, Pr³⁺, Y³⁺, Dy³⁺, and La³⁺) containing quinolinone. The obtained results reveal that the quinolinone **284** acts as a bidentate *via* OO donor sites, creating octahedral complexes. The complexes are nanoscale, having crystalline or amorphous structures. The synthesized nano-complexes have acceptable anticancer efficacy against the hepatocellular carcinoma cell line. Whereby, the maximum anticancer activity of Pr³⁺ nanocomplex (Fig. 21).²⁸²

A series of quinolinone-derived Schiff bases 286 were synthesized via the condensation of 7-amino-4-methylquinolinone 285 with different aromatic aldehydes. Then, complexation of Schiff ligands with cupric salts involving $Cu(\pi)$ acetate or $Cu(\pi)$ perchlorate hexahydrate was achieved, affording $Cu(\pi)$ complexes of the corresponding ligands 287 (Scheme 101). These complexes showed significant antifungal effects, suppressing C. albicans growth by 50% at a dosage of 4 μ M.²⁸³

5.4. Corrosion inhibitors

Hussein *et al.*¹⁸⁵ reported the utilization of 4-hydroxyquinolinone derivatives as antioxidant agents for various lubricating oils through using the ASTM D-942 and ASTM D-664 tests. The compositions that they synthesized are: dichlorophenyl-pyrazol-4-hydroxy-1-methylquinolinone **288**, 4-hydroxy-8-methyl-2-oxoquinolin-1-phenyl-2-

hydroxyhydropyrimidin-4,6-dione **289**, and 1-butyl-4-hydroxy-3-1*H*-pyrazol-3-yl)quinolinone **290** (Fig. 22). The obtained results indicate that the hydroxyquinolinone scaffolds reduce both the total acid number and the oxygen pressure drop in lubricating oils. On the other hand, it has been discovered that the antioxidant activity is most efficient when quinolinones have both butyl and hydroxyl groups. These quinoline scaffolds (**288–290**) inhibit radical processes during oil oxidation, as well as their continuation and production. A comparison of the antioxidant activity of compounds **288**, **289**, and **290** reveals that the efficacy and antioxidant benefits of the third compound. ^{184,284}

Quinolinone chalcone (PPQ) 291 showed effective corrosion inhibition for high carbon steel (HCS) in 1.0 M HCl, and the

Scheme 101 Complexation of ligand 286 with Cu(II) salts.

Fig. 22 Quinolinone scaffolds with antioxidant effect in lubricating oils.

Fig. 23 Quinolinone scaffolds as corrosion inhibitors

inhibition efficiency improved as the inhibitor concentration was raised. The compound's efficiency is attributed to the presence of heteroatoms and phenyl rings with delocalized π electrons that can act as adsorption centers. These factors support both physical and chemical interactions and

adsorption of inhibitor scaffolds with the metal surface according to the Langmuir adsorption isotherm (Fig. 23).^{285,286}

The dihydroquinoline-3-carboxylate derivatives **292** and **293** (**NODC**, **AODC**, **CODC**, and **MODC**) display potential corrosion inhibition for carbon steel in 1 M HCl, due to the presence of a substituent attached to the benzene portion of the 4-quinolinone (Fig. 24).²⁸⁷

Imino quinolinone IQ **294** acts as a corrosion inhibitor on low carbon steel (LCS) in 0.5 M HCl solution with inhibition efficiency 93.2% at the optimal concentration of IQ (30×10^{-3} Mm), adhered to the Langmuir adsorption model, signifying a monolayer adsorption mechanism. The characterization of IQ as a mixed-type inhibitor was evidenced due to its capacity to impede both cathodic and anodic processes, predominantly functioning as an anodic inhibitor (Fig. 25).²⁸⁸

Fig. 24 Quinolinones scaffolds as potential corrosion inhibitors.





Fig. 25 Imino quinolinone IQ 294.

Review RSC Advances

Fig. 26 Quinolinone/hydroxyquinoline scaffolds as chemosensors.

Scheme 102 Hydroxyquinoline scaffolds as fluorescence sensors



Scheme 103 Complexation mechanism of sensor 299 and Fe³⁺ ions.

5.5. Sensors

5.5.1. Chemosensors. Quinolinone derivative-based chemosensors designed to detect various metal ions were discussed. There is considerable interest in improving these chemosensors due to their ease of synthesis, high sensitivity, and stability. Nonetheless, there remains substantial potential for enhancement in their *in vivo* applications, particularly regarding water solubility, selectivity, and fluorescent bioimaging capabilities. Consequently, the development of receptors tailored for different ions is essential. For instance, skeleton **295** employed an alternative bonding model to identify Zn²⁺ and Cd²⁺. Also, **296** act as a Cr³⁺ chemosensor. Additionally, two-photon laser sources can be used to excite an extended conjugated quinoline system (Fig. 26).²⁸⁹

5.5.2. Fluorescence sensors. Fluorescence-based techniques have become essential tools in metalworking because of their sensitivity and selectivity benefits.^{290–292} A lot of work has gone into creating fluorescent Fe³⁺ indicators that are highly sensitive and selective. The majority of documented fluorescent

sensors for Fe³⁺ are based on a fluorescence quenching mechanism due to the paramagnetic nature of Fe³⁺,^{293,294} and the majority of these have interference issues brought on by transition metal cations like Cu²⁺, Co²⁺, or Hg²⁺. Thus, looking for highly selective chemosensors for Fe³⁺ based on the fluorescence enhancement is crucial. While using the ether derivative **297** as a starting material, the synthesis of Schiff base **299** was utilized. Whereby, quinoline **299** is a chemosensor that made up of the quinoline fluorophore **298** and *p*-anisidine binding site led to formation of bearing polarized C–C, C–N and C–O bonds conjugated to the quinoline moiety, it displays a strong fluorescence increase of Fe³⁺ ions, demonstrating excellent sensitivity and good selectivity for Fe³⁺ over a broad spectrum of other ecologically and physiologically significant metal ions (Schemes 102 and 103).²⁹⁵

Sensor **299** is an excellent sensitive chemosensor for the Fe³⁺ ion as it exhibits no fluorescence in MeOH, but when Fe³⁺ is added, it affords a highly effective fluorescence sensor **300**, and there is a noticeable increase in fluorescence of about 44-fold

RSC Advances Review

(Scheme 103). As a result, the fluorescence response is explained by Fe³⁺ blockage of electron transfer to the quinoline group, which encouraged sensor 299 to produce high fluorescence. A color shift from colorless to brown that is readily apparent to the unaided eye resulted from the identification of Fe³⁺. Consequently, the Fe³⁺ ion can be readily distinguished in visible light from all other metal ions. Additionally, under light from a 365 nm UV lamp, Fe³⁺ shows an increase in the intensity of fluorescence, vivid green, while other metal ions show no change under the same circumstances.295

Conclusion

ever-increasing importance of 3-acetyl-4hydroxyquinolinone and its N-substituted derivatives as structurally decisive scaffolds in bioactive natural products and pharmaceutical drugs, so in this review enormous efforts have been made to summarize the synthesis, structural features and chemical reactivity of 3-acetyl-4-hydroxy-2-quinolinone entities and its N-substituted derivatives as structurally decisive scaffolds in bioactive natural products and pharmaceutical drugs, accredited by reaction mechanisms. Also, we highlight the most important breakthroughs of 3-acetyl-4-hydroxy-2-quinolinone derivatives as an auspicious class displaying a wide range of potential pharmacological activities and their developments in the various clinical stages. Generally, this review supplies an overview of the chemistry for 3-acetyl-4-hydroxy-2-quinolinone derivatives and documents more than two hundred references over the last decade of research, covering mainly the period from 2020 to the beginning of 2025.

Abbreviations

AHQ 3-Acetyl-4-hydroxyquinolinone

BQB 1,4-Bis-(quinolin-6-ylimino methyl)benzene

 $BF_3 \cdot (OC_2H_5)_2$ Boron trifluoride etherate

Carbon disulfide CS_2 CCl_4 Carbon tetrachloride DMF Dimethylformamide **DMA** Dimethylamine **DMSO** Dimethyl sulfoxide EAA Ethyl acetoacetate HCS High carbon steel

NH₂OH·HCl Hydroxylamine hydrochloride **HMPT** Hexamethylphosphoric triamide

HQ 4-Hydroxyquinoline LCS Low-carbon steel LR Lawesson's reagent Multicomponent reaction MCR

DMF/DMA N,N-Dimethylformamide dimethyl acetal

DTA N,N-Dimethyl trifluoroacetamide

NBS N-Bromosuccinimide POCl₃ Phosphorous oxychloride Phosphorus pentasulfide P_2S_5 PBr₃ Phosphorus tribromide PhPOCl₂ Phosphonic dichloride

PPA Polyphosphoric acid

RORC Ring opening and ring closure

RT Room temperature **AcONa** Sodium acetate Sulfuryl chloride SO_2Cl_2 NaBH₄ Sodium borohydrate CH(OEt)₃ Triethyl orthoformate

Triethylamine TEA

P(OCH₂CH₂Cl)₃ Tris(2-chloroethyl)phosphite

TB **Tuberculosis** TC β,β -Tricarbonyl

TBAB Tetrabutylammonium bromide TsCl Toluenesulfonyl chloride WHO World Health Organization

Zinc chloride ZnCl₂

Data availability

No primary research results, software or code have been included, and no new data were generated or analyzed as part of this review.

Conflicts of interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this review.

Acknowledgements

The authors are thankful to Mansoura University, Egypt, for their support under project ID: MU-SCI-23-64.

References

- 1 A. Gulraiz, M. Sohail, M. Bilal, N. Rasool, M. Usman Qamar, C. Ciurea, L. G. Marceanu and C. Misarca, Molecules, 2024, 29, 2232.
- 2 X. Zhang, M. Liu, W. Qiu and W. Zhang, Molecules, 2024, 29,
- 3 W. S. Hamama, M. E. Ibrahim, A. A. Gooda and H. H. Zoorob, RSC Adv., 2018, 8, 8484.
- 4 P. Borah, S. Hazarika, A. Chettri, D. Sharma, S. Deka, Venugopala, P. Shinu, N. A. Al-Shar'i, S. K. Bardaweel and P. K. Deb, In Viral, Parasitic, Bacterial, and Fungal Infections, 2023, pp. 781-804.
- 5 (a) D. A. Sabbah, H. H. Samarat, E. Al-Shalabi, S. K. Bardaweel, R. Hajjo, K. Sweidan, R. Abu Khalaf, A. M. Al-Zuheiri and G. Abushaikha, ChemistrySelect, 2022, 7, e202200662; (b) R. S. Keri, S. Budagumpi and V. Adimule, ACS Omega, 2024, 9, 42630-42667.
- 6 M. Rana, R. Ranjan, N. S. Ghosh, D. Kumar and R. Singh, Curr. Cancer Ther. Rev., 2024, 20, 372-385.
- 7 (a) M. Ramanathan and Z. Moussa, Org. Chem. Front., 2025, 12, 256-327; (b) S. M. Alsafy and N. Abd Alrazzak, Eng. Proc., 2024, 59, 178.

Review **RSC Advances**

- 8 W. S. Hamama, A. E. Hassanien, M. G. El-Fedawy and H. H. Zoorob, J. Heterocycl. Chem., 2016, 53, 945-952.
- 9 N. Bhusare and M. Kumar, Oncol. Res., 2024, 32, 849-875.
- 10 K. Kajal, R. Shakya, M. Rashid, V. Nigam, B. D. Kurmi, G. D. Gupta and P. Patel, Sustain. Chem. Pharm., 2024, 37, 101374.
- 11 R. B. Bakr, I. H. El Azab and N. A. A. Elkanzi, Chem. Biodivers., 2024, 21, e202400200.
- 12 A. Darque, A. Dumètre, S. Hutter, G. Casano, M. Robin, C. Pannecouque and N. Azas, Bioorg. Med. Chem. Lett., 2009, 19, 5962-5964.
- 13 D. Dize, M. B. T. Tali, C. A. N. Ngansop, R. Keumoe, E. A. M. Kemgne, L. R. T. Yamthe, P. V. T. Fokou, B. P. Kamdem, K. Hata and F. F. Boyom, Future Pharmacol., 2024, 4, 188-198.
- 14 J. Sun and J. J. Kessl, Viruses, 2024, 16, 200.
- 15 D. Shi, S. Xu, D. Ding, K. Tang, Y. Zhou, X. Jiang, S. Wang, X. Liu and P. Zhan, Expert Opin. Drug Discovery, 2024, 19, 139-146.
- 16 W. S. Hamama, A. E. Hassanien, M. G. El-Fedawy and H. H. Zoorob, J. Heterocycl. Chem., 2015, 52, 492-496.
- 17 B. A. Mısır, Y. Derin, S. Ökten, A. Aydın, Ü. M. Koçyiğit, H. Şahin and A. Tutar, J. Biochem. Mol. Toxicol., 2024, 38, e23706.
- 18 M. M. Hammouda, M. M. Rashed, W. S. Abo El-Yazeed and K. M. Elattar, ChemistrySelect, 2024, 9, e202401384.
- 19 S. Siddique, K. R. Ahmad, S. K. Nawaz, A. R. Raza, S. N. Ahmad, R. Ali, I. Inayat, S. Suleman, M. A. Kanwal and M. Usman, Sci. Rep., 2023, 13, 8639.
- 20 P. Kołodziej, M. Wujec, M. Doligalska, A. Makuch-Kocka, Khylyuk, J. Bogucki, M. Demkowska-Kutrzepa, M. Roczeń-Karczmarz, M. Studzinska, K. Tomczuk, Kocki, P. Reszka-Kocka, S. Granica, R. Typek, A. L. Dawidowicz, J. Kocki and A. Bogucka-Kocka, J. Adv. Res., 2024, 60, 57-73.
- 21 R. Das, D. K. Mehta, S. Gupta, S. Mujwar, V. Sharma, A. Goyal, S. Patel and A. Patel, Lett. Org. Chem., 2023, 20, 1182-1191.
- 22 Y. M. Abdel Aziz, M. S. Nafie, P. A. Hanna, S. Ramadan, A. Barakat and M. Elewa, *Pharmaceuticals*, 2024, 17, 710.
- 23 M. C. Costas-Lago, P. Besada, R. Mosquera, E. Cano and C. Teran', Bioorg. Chem., 2024, 150, 107615.
- 24 T. M. Ramsis, M. A. Ebrahim and E. A. Fayed, Med. Chem. Res., 2023, 32, 2269-2278.
- 25 S. Siddique, K. Hussain, N. Shehzadi, M. Arshad, M. N. Ashad, S. Iftikhar, F. Saghir, A. Shaukat, M. Sarfraz and N. Ahmed, Org. Biomol. Chem., 2024, 22, 3708-3724.
- 26 H. E. Hashem, S. Ahmad, A. Kumer and Y. El Bakri, Sci. Rep., 2024, 14, 1152.
- 27 Y. U. Cebeci, O. O. Batur and H. Boulebd, J. Mol. Struct., 2024, 1299, 137115.
- 28 S. K. Verma, S. Rangappa, R. Verma, F. Xue, S. Verma, K. S. S. Kumar and K. S. Rangappa, Bioorg. Chem., 2024, 145, 107241.
- 29 K. B. Gangurde, R. A. More, V. A. Adole and D. S. Ghotekar, J. Mol. Struct., 2024, 1299, 136760.

- 30 Y. Jia, Y. Zhao, M. Niu, C. Zhao, X. Li and H. Chen, Anim. Models Exp. Med., 2024, 7, 419-432.
- 31 M. El Mesky, H. Zgueni, Y. Rhazi, O. El-Guourrami, O. Abchir, M. Jabha, A. Nakkabi, S. Chtita, S. Achamlale, M. Chalkha, D. Chebabe and El. Mabrouk, J. Mol. Struct., 2024, 1313, 138705.
- 32 (a) Swati, A. Raza, B. Singh and Dr. Pankaj Wadhwa, ChemistrySelect, 2025, 10, e202404978; (b) I. Ahmad, H. Khalid, A. Perveen, M. Shehroz, U. Nishan, F. Ur Rahman, Sheheryar, A. A. Moura, R. Ullah, E. A. Ali, M. Shah and S. C. Ojha, ACS Omega, 2024, 9, 16262-16278.
- 33 M. A. Fawzy, K. H. Ibrahim, A. A. Aly, A. H Mohamed, S. M. N. Abdel Hafez, W. Y. Abdelzaher, E. B Elkaeed, A. A Alsfouk and El. MN Abdelhafez, Future Med. Chem., 2024, 16, 2211-2230.
- 34 A. M. Mohassab, H. A. Hassan, H. A. Abou-Zied, M. Fujita, M. Otsuka, H. A. M. Gomaa, B. G. M. Youssif and M. Abdel-Aziz, J. Mol. Struct., 2024, 1297, 136953.
- 35 W. S. Hamama, M. E. Ibrahim, A. A. Gooda and H. H. Zoorob, J. Heterocycl. Chem., 2018, 55, 2623-2634.
- 36 M. Mashhadinezhad, M. Mamaghani, M. Rassa and F. Shirini, ChemistrySelect, 2019, 4, 4920-4932.
- 37 G. L. Monica, A. Bono, F. Alamia, A. Lauria and A. Martorana, Bioorg. Med. Chem., 2024, 109, 117791.
- 38 E. A. Ghaith, H. H. Zoorob, M. E. Ibrahim, M. Sawamura and W. S. Hamama, ChemistrySelect, 2020, 5, 14917-14923.
- 39 S. Nadar and T. Khan, Chem. Biol. Drug Des., 2022, 100, 818-842.
- 40 W. S. Hamama, A. E. Hassanien, M. G. El-Fedawy and H. H. Zoorob, J. Heterocycl. Chem., 2017, 54, 859-863.
- 41 X. Guo, Q. Zhou, X. Fan, Q. Zhu and M. Jin, ACS Appl. Polym. Mater., 2024, 6, 5566-5575.
- 42 M. Olesiejuk, A. Kudelko and M. Świątkowski, Dyes Pigm., 2023, 220, 111721.
- 43 W. S. Hamama, M. A. Waly, I. I. EL-Hawary and H. H. Zoorob, J. Heterocycl. Chem., 2016, 53, 953-957.
- A. Pordel, 44 S. Bozorgnia, M. Davoodnia and S. A. Beyramabadi, Opt. Mater., 2024, 148, 114762.
- 45 M. Escolano, D. Gaviña, G. Alzuet-Piña, S. Díaz-Oltra, M. Sánchez-Roselló and C. del Pozo, Chem. Rev., 2024, 124, 1122-1246.
- 46 (a) A. A. Aly, H. A. Abd El-Naby, E. Kh. Ahmed, S. A. Gedamy, M. B. Alshammari, A. Ahmad and S. Bräse, Curr. Org. Chem., 2025, 29, 181-212; (b) A. Mandal and A. T. Khan, Org. Biomol. Chem., 2024, 22, 2339-2358.
- 47 O. O. Ajani, K. T. Iyaye and O. T. Ademosun, RSC Adv., 2022, 12, 18594-18614.
- 48 S. Rajendran, K. Sivalingam, R. P. K. Jayarampillai, W. Wang and C. O. Salas, Chem. Biol. Drug Des., 2022, 100, 1042-1085.
- 49 L. F. Hernández-Ayala, E. G. Guzmán-López and A. Galano, Antioxidants, 2023, 12, 1853.
- 50 R. Kumar, A. Thakur, Sachin, D. Chandra, A. K. Dhiman, P. K. Verma and U. Sharma, Coord. Chem. Rev., 2024, 499,
- 51 Q. Cai, H. Song, Y. Zhang, Z. Zhu, J. Zhang and J. Chen, J. Agric. Food Chem., 2024, 72, 12373-12386.

RSC Advances

- 52 M. Owais, A. Kumar, S. M. Hasan, K. Singh, I. Azad, A. Hussain, Suvaiv and M. Akil, Mini-Rev. Med. Chem., 2024, 24, 1238-1251.
- 53 S. Ghosh, S. Mallick, D. Karolly and S. D. Sarkar, ACS Org. Inorg. Au, 2024, 5, 492-497.
- 54 (a) W. Al Zoubi and Y. G. Ko, J. Colloid Interface Sci., 2020, 565, 86-95; (b) Y. Senpradit, S. Wacharasindhu and M. Sukwattanasinitt, Spectrochim. Acta, Part A, 2025, 326, 125128.
- 55 H. M. Abd El-Lateef, A. G. A. Gaafar, A. S. Algahtani, A. A. Al-Mutairi, D. S. Alshaya, F. G. Elsaid, E. Fayadg and N. A. Farouk, RSC Adv., 2024, 14, 24781-24790.
- 56 P. A. Jagtap, V. R. Sawant and B. M. ChemCatChem, 2024, 16, e202400979.
- 57 S. Sharma, K. Singh and S. Singh, Curr. Org. Synth., 2023, 20, 606-629.
- 58 M. Mohasin, M. Z. Alam, Q. Ullah, A. Ahmad, P. F. Rahaman and S. A. Khan, *Polycycl*, 2024, 44, 6369–6398.
- 59 A. Dorababu, Arch. Pharm., 2021, 354, 2000232.
- 60 D. Mabire, S. Coupa, C. Adelinet, A. Poncelet, Y. Simonnet, M. Venet, R. Wouters, A. S. J. Lesage, L. V. Beijsterveldt and F. Bischoff, J. Med. Chem., 2005, 48, 2134-2153.
- 61 T. L. Viveka, G. Angajala, V. Aruna, M. Nakka and Y. Aparna, J. Mol. Struct., 2024, 1303, 137482.
- 62 F. H. Al-Ostoot, Zabiulla, S. Salah and S. A. Khanum, J. Iran. Chem. Soc., 2021, 18, 1839-1875.
- 63 J. Bergwik, J. Liu, M. Padra, R. K. V. Bhongir, L. Tanner, Y. Xiang, M. Lundblad, A. Egesten and M. Adner, Respir. Res., 2024, 25, 146.
- 64 P. Y. Wang, H. Chen, Y. Wanga and Y. K. Lyu, J. Chem. Technol. Biotechnol., 2020, 95, 2171-2179.
- 65 C. M. Al-Matarneh, A. Nicolescu, I. C. Marinas, M. D. Găboreanu, S. Shova, A. Dascălu, M. Silion and M. Pinteală, Molecules, 2024, 29, 772.
- 66 G. Shumi, T. B. Demissie, R. Eswaramoorthy, R. F. Bogale, G. Kenasa and T. Desalegn, ACS Omega, 2024, 9, 25014-25026.
- 67 J. C. Coa, A. Yepes, M. Carda, L. Conesa-Milián, Y. Upegui, S. M. Robledo, Dr. and W. Cardona-G, ChemistrySelect, 2020, 5, 2918-2924.
- 68 I. A. Bala, O. F. Al Sharif, A. M. Asiri and R. M. El-Shishtawy, Results Chem., 2024, 7, 101529.
- 69 M. H. El-Shershaby, K. M. El-Gamal, A. H. Bayoumi, K. El-Adl, M. Alswah, H. E. A. Ahmed, A. A. Al-Karmalamy and H. S. Abulkhair, New J. Chem., 2021, 45, 13986-14004.
- 70 S. Kondaparla, A. Soni, A. Manhas, K. Srivastava, S. K. Purib and S. B. Katti, RSC Adv., 2016, 6, 105676-105689.
- 71 A. Rezvanian, B. Khodadadi, S. Tafreshi and P. Shiri, Mol. Diversity, 2024, 28(1), 197-207.
- 72 A. D. Sonawane, D. R. Garud, T. Udagawa and M. Koketsu, Org. Biomol. Chem., 2018, 16, 245-255.
- 73 K. V. Belyaeva, L. P. Nikitina, L. A. Oparina, V. S. Saliy, D. N. Tomilin, A. V. Kuzmin, A. V. Afonin and B. A. Trofimov, New J. Chem., 2024, 48, 1336-1349.
- 74 X. An, N. Li, L. Zhang, Z. Xu, S. Zhang and Q. Zhang, J. Hazard. Mater., 2024, 465, 133158.

- 75 http://www.africanplants.senckenberg.de/root/index.php? page id=78&id=4191#.
- 76 R. Kaur and K. Kumar, Eur. J. Med. Chem., 2021, 215, 113220.
- 77 P. Yadav and K. Shah, Bioinorg. Chem., 2021, 109, 104639.
- 78 D. Talwar, A. Gonzalez-de-Castro, H. Y. Li and J. Xiao, Angew. Chem., 2015, 127, 5312-5316.
- 79 B. S. Matada, R. Pattanashettar and N. G. Yernale, Bioorg. Med. Chem., 2021, 32, 115973.
- 80 A. D. Arboleda, L. M. Moreno and R. Abonia, Curr. Org. Chem., 2024, 28, 595-635.
- 81 M. Kischkewitz, B. Marinic, N. Kratena, Y. Lai, H. B. Hepburn, M. Dow, K. E. Christensen and T. J. Donohoe, Angew. Chem., Int. Ed., 2022, 61, e202204682.
- 82 k. Proisl, S. Kafka and J. Kosmrli, Curr. Org. Chem., 2017, 21, 1949-1975.
- 83 S. S. Nazrullaev, T. Α. Bessonova and Kh. S. Akhmedkhodzhaeva, Chem. Nat. Compd., 2001, 37, 551-555.
- 84 (a) S. S. Anjanikar and S. S. Chandole, Orient. J. Chem., 2023, 39, 197; (b) K. Gach-Janczak, J. Piekielna-Ciesielska, J. Waskiewicz, K. Krakowiak, K. Wtorek and A. Janecka, Molecules, 2025, 30, 163.
- 85 M. A. Barmade, P. Agrawal, S. R. Rajput, P. R. Murumkar, B. Rana, D. Sahal and M. R. Yadav, RSC Med. Chem., 2024, **15**, 572-594.
- 86 A. Uddin, S. Gupta, R. Shoaib, B. Aneja, I. Irfan, K. Gupta, N. Rawat, J. Combrinck, B. Kumar, M. Aleem, P. Hasan, M. C. Joshi, Y. S. Chhonker, M. Zahid, A. Hussain, K. Pandey, M. F. Alajmi, D. J. Murry, T. J. Egan, S. Singh and M. Abid, Eur. J. Med. Chem., 2024, 264, 115969.
- 87 J. Gao, H. Hou and F. Gao, Eur. J. Med. Chem., 2023, 247, 115026.
- 88 B. Çiftci, S. Ökten, Ü. M. Koçyiğit, V. E. Atalay, M. Ataş and O. Cakmak, Eur. J. Med. Chem. Rep., 2024, 10, 100127.
- 89 S. Guin, K. M. Alden, D. J. Krysan and M. J. Meyers, ACS Med. Chem. Lett., 2024, 15, 822-827.
- 90 S. Verma, S. Lal and R. Narang, Future Med. Chem., 2024, 16, 1283-1286.
- 91 T. Arasakumar, S. Mathusalini, S. S. Shyamsivappan, A. Ata and P. S. Mohan, Bioorg. Med. Chem. Lett., 2017, 27, 1538-1546.
- 92 E. A. E. El-Helw, M. Asran, M. E. Azab, M. H. Helal, A. Y. A. Alzahrani and S. K. Ramadan, Sci. Rep., 2024, 14, 15522.
- 93 S. G. Azimi, G. Bagherzade, M. R. Saberi and Z. A. Tehranizadeh, Bioinorg. Chem. Appl., 2023, 2881582.
- 94 M. Beus, L. Persoons, D. Daelemans, D. Schols, K. Savijoki, P. Varmanen, J. Yli-Kauhaluoma, K. Pavić and B. Zorc, Mol. Diversity, 2022, 26, 2595-2612.
- 95 H. Abdelmegeed, L. M. A. Abdel Ghany, A. Youssef, A. S. El-Etrawycd and N. Ryadc, RSC Adv., 2024, 14, 22092-22112.
- 96 X. He, J. Chen, M. Kandawa-Shultz, G. Shao and Y. Wang, Dalton Trans., 2023, 52, 4728-4736.
- 97 G. Jina, Z. Lib, F. Xiaoa, X. Qia and X. Sun, Bioorg. Chem., 2020, 99, 103837.

2014, 126, 205-212.

98 N. D. Shashikumar, G. Krishnamurthy, H. S. Bhojyanaik, M. R. Lokesh and K. S. Jithendrakumara, *J. Chem. Sci.*,

- 99 R. P. Pakhariya, A. Bhatnagar and G. Pemawat, *RSC Adv.*, 2025, **15**, 3646–3663.
- 100 J. C. d. Santos, J. E. F. Alves, R. D. S. de Azevedo, M. L. de Lima, M. R. de Oliveira Silva, J. G. da Silva, J. M. da Silva, A. C. de Carvalho Correia, M. do Carmo Alves de Lima, J. F. de Oliveira, R. O. de Moura and S. M. V. de Almeida, Int. J. Biol. Macromol., 2024, 254, 127651.
- 101 A. Saral, R. Shahidha, M. Thirunavukkarasu and S. Muthu, *Chem. Phys. Impact.*, 2023, **6**, 100193.
- 102 S. B. Wang, X. Q. Deng, Y. Zheng, H. J. Zhang and Z. S. Quan, *Arch. Pharmacal Res.*, 2013, 36, 32–40.
- 103 K. Shabana, Salahuddin, A. Mazumder, H. Singh, R. Kumar, S. Tyagi, V. Datt, A. S. Sharma, M. S. Yar, M. J. Ahsan and R. K. Yadav, *ChemistrySelect*, 2023, 8, e202300209.
- 104 J. Song, Y. Zhu, W. Zu, C. Duan, J. Xu, F. Jiang, X. Wang, S. Li, C. Liu, Q. Gaoa, H. Li, Y. Zhang, W. Tang, T. Lu and Y. Chen, *Bioorg. Med. Chem.*, 2021, 29, 115856.
- 105 S. K. Suthar, V. Jaiswal, S. Lohan, S. Bansal, A. Chaudhary, A. Tiwari, A. T. Alex and A. Joesph, *Eur. J. Med. Chem.*, 2013, 63, 589–602.
- 106 S. H. Pattanashetty, K. M. Hosamani and D. A. Barretto, *Chem. Data Collect.*, 2018, **15**, 184–196.
- 107 A. Krishna, V. Vijayakumar and S. Sarveswari, ChemistrySelect, 2020, 5, 7967–7972.
- 108 I. Kostopoulou, A. Diassakou, E. Kavetsou, E. Kritsi, P. Zoumpoulakis, E. Pontiki, D. Hadjipavlou-Litina and A. Detsi, *Mol. Diversity*, 2021, 25, 723–740.
- 109 M. Azad, M. A. Munawar and M. Athar, *J. Appl. Sci.*, 2007, 7, 1620–1625.
- 110 M. Rowley, P. D. Leeson, G. I. Stevenson, A. M. Moseley, I. Stansfield, I. Sanderson, L. Robinson, R. Baker, J. A. Kemp, G. R. Marshall, A. C. Foster, S. Grimwood, M. D. Tricklebank and K. L. Saywell, *J. Med. Chem.*, 1993, 36, 3386–3396.
- 111 T. S. Tunna, I. S. M. Zaidul, Q. U. Ahmed, K. Ghafoor, F. Y. Al-Juhaimi, M. S. Uddin, M. Hasan and S. Ferdous, S. Afr. J. Bot., 2015, 99, 144–152.
- 112 M. M. Hassan and H. M. Hassanin, *J. Heterocycl. Chem.*, 2017, 54, 3321–3330.
- 113 V. M. Krokhalev, V. I. Saloutin, A. D. Romas, B. A. Ershov and K. I. Pashkevich, *Div. Chem. Sci.*, 1990, **39**, 316–322.
- 114 A. Detsi, V. Bardakos, J. Markopoulos and O. Igglessi-Markopoulou, *J. Chem. Soc., Perkin Trans.* 1, 1996, 1, 2909–2913.
- 115 M. M. Abdou, Z. Seferoğlu, M. Fathy, T. Akitsu, M. Koketsu, R. Kellow and E. Amigues, *Res. Chem. Intermed.*, 2019, 45, 919–934.
- 116 G. Athanasellis, E. Gavrielatos, O. Igglessi-Markopoulou and J. Markopoulos, *J. Heterocycl. Chem.*, 2003, **40**, 645–648.
- 117 S. M. Abu-elwafa, E. E. Mohamed, R. M. Issa and M. Gaber, *Indian J. Chem.*, 1985, **24A**, 407–411.
- 118 P. G. Dessai, S. P. Dessai, R. Dabholkar, P. Pednekar, S. Naik, S. Mamledesai, M. Gopal, P. Pavadai,

- B. K. Kumar, S. Murugesan, S. Chandavarkar, P. Theivendren and K. Selvaraj, *Mol. Diversity*, 2023, 27, 1567–1586.
- 119 H. B. Bürgi and J. D. Dunitz, *Structure Correlation*, John Wiley & Sons, 2008.
- 120 I. V. Ukrainets, A. A. Tkach and L. Y. Yang, *Chem. Heterocycl. Compd.*, 2009, **45**, 169–175.
- 121 V. V. Kouznetsov, L. Y. V. Méndez and C. M. M. Gómez, *Curr. Org. Chem.*, 2005, **9**, 141–161.
- 122 L. M. Nainwal, S. Tasneem, W. Akhtar, G. Verma, M. F. Khan, S. Parvez, M. Shaquiquzzaman, M. Akhter and M. M. Alam, *Eur. J. Med. Chem.*, 2019, **164**, 121–170.
- 123 G. A. Ramann and B. J. Cowen, Molecules, 2016, 21, 986.
- 124 S. M. Prajapati, K. D. Patel, R. H. Vekariya, S. N. Panchal and H. D. Patel, *RSC Adv.*, 2014, 4, 24463–24476.
- 125 J. Barluenga, F. Rodriguez and F. J. FaÇanas, *Chem.-Asian J.*, 2009, **4**, 1036–1048.
- 126 A. Weyesa and E. Mulugeta, *RSC Adv.*, 2020, **10**, 20784–20793.
- 127 J. B. Bharate, R. A. Vishwakarma and S. B. Bharate, *RSC Adv.*, 2015, **5**, 42020–42053.
- 128 C. Teja and F. R. N. Khan, *Chem.-Asian J.*, 2020, **15**, 4153-4167.
- 129 E. Vessally, L. Edjlali, A. Hosseinian, A. Bekhradnia and M. D. Esrafil, *RSC Adv.*, 2016, **6**, 49730–49746.
- 130 T. Kappe, R. Aigner, P. Hohengassner and W. Stadlbauer, *J. Prakt. Chem.*, 1994, 336, 596–601.
- 131 D. N. Toan, N. D. Thanh, M. X. Truong and D. T. Van, *Arabian J. Chem.*, 2020, **13**, 7860-7874.
- 132 T. Razzaq and C. O. Kappe, *Tetrahedron Lett.*, 2007, 48, 2513–2517.
- 133 D. N. Toan, N. D. Thanh, M. X. Truong, D. N. Bang, M. T. Ngaa and N. T. T. Huong, New J. Chem., 2020, 44, 20715–20725.
- 134 D. Abdel-Kadera, M. Abass and S. Shawkat, *Russ. J. Org. Chem.*, 2024, **60**, 294–328.
- 135 W. Stadlbauer and G. Hojas, *J. Heterocyclic Chem.*, 2004, **41**, 681.
- 136 R. E. Bowman, A. Campbell and E. M. Tanner, *J. Chem. Soc.*, 1959, **81**, 444–447.
- 137 M. A. Ibrahim, H. M. Hassanin, M. Abass and S. Badran, *Arkivoc*, 2013, 4, 424–431.
- 138 K. Tomita, Yakugaku Zasshi, 1951, 71, 1100-1112.
- 139 S. Elgogary, H. Abd Elghafar and M. Mashaly, *J. Chin. Chem. Soc.*, 2021, **68**, 1082–1089.
- 140 M. M. Hassan, E. S. Othman and M. Abass, *Res. Chem. Intermed.*, 2013, **39**, 1209–1225.
- 141 K. Alla and S. Sarveswari, Iran J. Sci. Technol. Trans. Sci., 2019, 43, 465–475.
- 142 S. Sarveswari, V. Vijayakumar, R. Siva and R. Priya, *Appl. Biochem. Biotechnol.*, 2015, **175**, 43–64.
- 143 J. Brawley, E. Etter, D. Heredia, A. Intasiri, K. Nennecker, J. Smith, B. M. Welcome, R. K. Brizendine, T. W. Gould, T. W. Bell and C. Cremo, J. Med. Chem., 2020, 63, 11131– 11148.
- 144 M. Shariat, M. W. Samsudin and Z. Zakaria, *Chem. Cent. J.*, 2013, 7, 58.

RSC Advances

- 145 C. Mitsos, J. Petrou, O. Igglessi-Markopoulou and J. Markopoulos, J. Heterocycl. Chem., 1999, 36, 881-887.
- 146 M. Roussaki, B. Hall, S. C. Lima, A. C. da Silva, S. Wilkinson and A. Detsi, Bioorg. Med. Chem. Lett., 2013, 23, 6436-6441.
- 147 B. Baruah, K. Dasu, B. Vaitilingam, A. Vanguri, S. R. Casturi and K. R. Yeleswarapu, Bioorg. Med. Chem. Lett., 2004, 14, 445-448.
- 148 S. Dutt and V. Tyagi, Tetrahedron Lett., 2021, 87, 153527.
- 149 R. Satheeshkumar, R. Shankar, W. Kaminsky and K. J. R. Prasad, ChemistrySelect, 2016, 21, 6823-6829.
- 150 Q. Zhang, J. Yuan, M. Yu, R. Zhang, Y. Liang, P. Huang and D. Dong, Synthesis, 2017, 49, 4996-5002.
- 151 H. M. Hassanin, M. A. Ibrahim, Y. A. Gabr and Y. A. Alnamer, J. Heterocycl. Chem., 2012, 49, 1269-1289.
- 152 D. K. Kumar, R. Rajkumar and S. P. Rajendran, Chem. Heterocycl. Compd., 2016, 52, 322-325.
- 153 M. Abass, A. R. A. Alzandi, M. M. Hassan and N. Mohamed, Polycyclic Aromat. Compd., 2021, 41, 2120-2209.
- 154 S. D. Naik, S. K. Chandavarkar, S. S. Tawade, S. G. Shingade, M. B. Palkard and S. N. M. Desai, Indian J. Chem., 2022, 61, 544-550.
- 155 P. Roschger and W. Stadlbaue, Liebigs Ann. Chem., 1990, 821-823.
- 156 W. Steinschifter, W. Fiala and W. Stadlbauer, J. Heterocycl. Chem., 1994, 31, 1647-1652.
- 157 G. Hojas, W. Fiala and W. Stadlbauer, J. Heterocycl. Chem., 2000, 37, 1559-1569.
- 158 S. Chimichi, M. Boccalini and A. Matteucci, Tetrahedron, 2007, 63, 11656-11660.
- 159 R. R. Raiturkara, S. N. M. Desaia, V. M. Graciasa, A. D. S. Sangavkara, C. Fernandesa, B. S. Biradarb and S. K. Chandavarkar, Indian J. Chem., 2022, 61, 849-857.
- 160 M. G. Reis, S. M. Desai, S. Naik, J. Fernandes and P. Tari, Indian J. Chem., 2016, 55B, 1254-1258.
- 161 G. A. Salman, Al-Mustansiriyah J. Sci., 2018, 28, 73-79.
- 162 S. N. M. Desai, R. N. S. Priolkar, H. A. N. Karmali, P. D. Amambe and B. S. Birdar, Int. J. Pharm. Pharm. Sci., 2017, 9, 240-244.
- 163 H. M. Hassanin and S. M. El-edfawy, Heterocycles, 2012, 85, 2421-2436.
- 164 K. Faber and T. Kappe, J. Heterocycl. Chem., 1984, 21, 1881-1883.
- 165 S. Goswami, K. Ghosh, R. Mukherjee, A. K. Adak and A. K. Mahapatra, J. Heterocycl. Chem., 2001, 38, 173-177.
- 166 B. P. Nandeshwarappa, S. K. Manjunatha, D. K. Ramesh, M. Suchitra and S. O. Sadashiv, J. Pharm. Sci., 2020, 10, 33-38.
- 167 M. M. Girges, M. A. Hanna, H. M. Hassan and E. B. Moawad, Collect. Czechoslov. Chem. Commun., 1988, **53**, 3179-3183.
- 168 G. Bolakatti, M. Palkar, M. Katagi, G. Hampannavar, R. V. Karpoormathd, S. Ninganagouda and A. Badiger, J. Mol. Struct., 2021, 1227, 129413.
- 169 A. Vilsmeier and A. Haack, Ber. Dtsch. Chem. Ges., 1927, 60,
- 170 M. A. Ibrahim, H. M. Hassanin and Y. A. Alnamer, Synth. Commun., 2014, 44, 3470-3482.

- 171 A. V. Manaev, I. N. Okhrimenko, K. A. Lyssenko and V. F. Traven, Russ. Chem. Bull., 2008, 57, 1734-1739.
- 172 T. kappe, R. Aigner, M. Jobstl, P. Hohengassner and W. Stadlbauer, Heterocycl. Commun., 1995, 1, 341-352.
- 173 M. Islamuddin, O. Afzal, W. H. Khan, M. Hisamuddin, A. S. A. Altamimi, I. Husain, K. Kato, M. A. Alamri and S. Parveen, ACS Omega, 2021, 6, 9791-9803.
- 174 M. Sankaran, C. Kumarasamy, U. Chokkalingam and P. S. Mohan, Bioorg. Med. Chem. Lett., 2010, 20, 7147-7151.
- 175 T. Arasakumar, S. Mathusalini, K. Lakshmi, P. S. Mohan, A. Ata and C. H. Lin, Synth. Commun., 2016, 46, 232-241.
- 176 (a) S. S. Ibrahim, Z. M. El-Gendy, H. A. Allimony and E. S. Othman, Chem. Pap., 1999, 53, 53-64; (b) M. Abass, Synth. Commun., 2000, 30, 2735-2757.
- 177 M. M. Hassan, A. R. A. Alzandi and M. M. Hassan, Arabian J. Chem., 2020, 13, 6184-6190.
- 178 M. Abdel-Megid, M. Abass and M. Hassan, J. Heterocycl. Chem., 2007, 44, 315-322.
- 179 M. M. Hassan, S. M. Abdel-Kariem and T. E. Ali, Phosphorus, Sulfur Silicon Relat. Elem., 2017, 192, 866-873.
- 180 S. Sarveswari and V. Vijayakumar, Arabian J. Chem., 2016, 9, S835-S839.
- 181 G. V. Kalechits, V. K. Ol'khovik, I. I. Kalosha, E. D. Skakovskii, A. A. Pap, A. A. Zenyuk and Y. V. Matveenko, Russ. J. Gen. Chem., 2001, 71, 1257.
- 182 M. A. Munawar, M. Azad, H. L. Siddiquia and F. Nasim, J. Chin. Chem. Soc., 2008, 55, 394-400.
- 183 R. S. Bhupathi, B. R. Devi and P. K. Dubey, *Indian J. Chem.*, 2012, 51, 855.
- 184 H. Maruthesh, M. S. Katagi and B. P. Nandeshwarappa, Russ. J. Org. Chem., 2023, 49, 1059-1067.
- 185 M. F. Hussein, M. A. Ismail and R. A. El-Adly, Int. J. Org. Chem., 2016, 6, 207-219.
- 186 V. F. Traven, A. V. Manaev, I. V. Voevodina and I. N. Okhrimenko, Russ. Chem. Bull., 2008, 57, 1508-1515.
- 187 A. Krishna and S. Sarveswari, ChemistrySelect, 2019, 4, 9987-9992.
- 188 M. Abass, M. Abdel-Megid and M. Hassan, Synth. Commun., 2007, 37, 329-352.
- 189 M. Abass and A. Hassan, Chem. Pap., 2003, 57, 267-277.
- 190 M. Abass and B. B. Mostafa, Bioorg. Med. Chem., 2005, 13, 6133-6144.
- 191 F. M. A. A. El-Taweel, J. Heterocycl. Chem., 2005, 42, 943-946.
- 192 N. N. Elnaggar, W. S. Hamama, M. Abd El Salam and E. A. Ghaith, RSC Adv., 2025, 15(8), 6050-6067.
- 193 (a) M. Abass and E. S. Othman, Synth. Commun., 2001, 31, 3361-3376; (b) A. Klásek, K. Kořistek, P. Sedmera and P. Halada, Heterocycles, 2003, 60, 799-815.
- 194 A. Hassan, M. Badr, D. Abdelhamid, H. A. Hassan, M. A. S. Abourehab and G. A. Abuo-Rahma, Bioinorg. Chem., 2022, 120, 105631.
- 195 A. A. Elagamey, F. M. A. El-Taweel and M. H. M. Khalil, Sci. J. Damietta Fac. Sci., 2012, 1, 33-41.
- 196 V. S. Rao and M. Darbarwar, Synthesis, 1989, 139–141.
- 197 Z. Chen and Z. Wang, Tetrahedron, 2016, 72, 4288-4293.

198 T. Kappe and B. Schnell, J. Heterocycl. Chem., 1996, 33, 663-

- 199 W. Fiala and W. Stadlbauer, J. Prakt. Chem., 1993, 335, 128-134.
- 200 E. M. El-Sheref, M. A. I. Elbastawesy, A. B. Brown, A. M. Shawky, H. A. M. Gomaa, S. Bräse and B. G. M. Youssif, Molecules, 2021, 26, 6798.
- 201 B. Madhu, C. V. R. Reddy and P. K. Dubey, Synth. Commun., 2017, 47, 421-427.
- 202 B. Madhu, B. R. Sekar, C. V. R. Reddy and P. K. Dubey, Res. Chem. Intermed., 2017, 43, 6993-7012.
- 203 S. I. Bhat and D. R. Trivedi, RSC Adv., 2014, 4, 11300-11304.
- 204 R. Bhupathi, B. Madhu, B. R. Devi, C. V. R. Reddy and P. K. Dubey, J. Heterocyclic Chem., 2016, 53, 1911-1916.
- 205 S. A. Barr, C. F. Neville, M. F. Grundon, D. R. Boyd, J. F. Malone and T. A. Evans, J. Chem. Soc., Perkin Trans. 1, 1995, 1, 445.
- 206 J. H. Ye, K. Q. Ling, Y. Zhang, N. Li and J. H. Xu, J. Chem. Soc., Perkin Trans. 1, 1999, 14, 2017-2024.
- 207 M. Abass, H. A. Allimony and H. Hassan, Phosphorus, Sulfur Silicon Relat. Elem., 2013, 188, 1799-1810.
- 208 J. M. Morsy, H. M. Hassanin, M. M. Ismail and M. M. A. Abd-Alrazk, J. Chem. Res., 2016, 40, 239-246.
- 209 T. Ali, H. Wang, W. Iqbal, T. Bashir, R. Shah and Y. Hu, Adv. Sci., 2023, 10, 2205077.
- 210 T. Kappe, R. Aigner, P. Roschger, B. Schnell and W. Stadlbauer, Tetrahedron, 1995, 51, 12923-12928.
- 211 Q. R. Luca and A. Q. Fenwick, J. Photochem. Photobiol., B, 2015, 152, 26-42.
- 212 V. Fonsecaa, S. Chandavarkar, Dabholkara, P. G. Dessaia, M. Deshpandec and S. N. M. Desai, Indian J. Chem., 2021, 60B, 267-272.
- 213 N. Azzman, S. Anwar, W. A. S. Mohamed and N. Ahemad, Med. Chem., 2024, 24, 1134-1157.
- 214 A. Pervaiza, M. M. Athara, M. A. Khana, M. Pervaizb, M. Sagirc and M. Y. Naz, Sci. Int., 2014, 26, e202300209.
- 215 Y. Al-Majedy, A. A. Kadhum, H. Ibraheem, A. Al-Amiery, A. A. Moneim and A. B. Mohamad, Sys. Rev. Pharm., 2018, 9, 49-54.
- 216 B. A. Subhasrao, Master's thesis, Rajiv Gandhi University of Health Sciences, India, 2010.
- 217 K. C. Prousis, A. Tzani, N. Avlonitis, T. Calogeropoulou and A. Detsi, J. Heterocycl. Chem., 2013, 50, 1313-1321.
- 218 I. V. Ukrainets, N. L. Bereznyakova and E. V. Mospanova, Chem. Heterocycle. Cmpd., 2007, 43, 856-862.
- 219 X. Collin, J. M. Robert, M. Duflos, G. Wielgosz, G. Le Baut, C. Robin-Dubigeon, N. Grimaud, F. Lang and J. Y. Petit, J. Pharm. Pharmacol., 2001, 53, 417-423.
- 220 A. Zaman, I. Ahmad, M. Pervaiz, S. Ahmad, S. Kiran, M. A. Khan, T. Gulzar and T. Kamal, J. Mol. Struct., 2019, 1180, 227-236.
- 221 V. Vijayakumar, Int. J. Chemtech Res., 2016, 9, 629-634.
- 222 Y. A. M. M. Elshaier, A. A. Aly, M. Abd El-Aziz, H. M. Fathy, A. B. Brown and M. Ramadan, Mol. Diversity, 2021, 26, 2341-2370.

- 223 I. V. Ukrainets, M. Y. Golik, L. V. Sidorenko, V. I. Korniyenko, L. A. Grinevich, G. Sim and O. V. Kryvanych, Sci. Pharm., 2018, 86, 31.
- 224 V. Yadav, J. Reang, V. Sharma, J. Majeed, P. C. Sharma, K. Sharma, N. Giri, A. Kumar and R. K. Tonk, Chem. Biol. Drug Des., 2022, 100, 389-418.
- 225 N. Priya, A. Gupta, K. Chand, P. Singh, A. Kathuria, H. G. Raj, V. S. Parmar and S. K. Sharma, Bioorg. Med. Chem., 2010, 18, 4085-4094.
- 226 T. Shiro, T. Fukaya and M. Tobe, Eur. J. Med. Chem., 2015, **97**, 397–408.
- 227 S. H. Kwak, S. Shin, J. H. Lee, J. K. Shim, M. Kim, S. D. Lee, A. Lee, J. Bae, J. H. Park, A. Abdelrahman, C. E. Müller, S. K. Cho, S. G. Kang, M. A. Bae, J. Y. Yang, H. Ko, W. A. Goddard III and Y. C. Kim, Eur. J. Med. Chem., 2018, 151, 462-481.
- 228 L. Senerovic, D. Opsenica, I. Moric, I. Aleksic, M. Spasić and B. Vasiljevic, Adv. Exp. Med Bio., 2020, 14, 37-69.
- 229 A. A. Aly, E. M. El-Sheref, A. E. Mourad, M. E. M. Bakheet and S. Bräsem, Mol. Diversity, 2020, 24, 477-524.
- 230 K. Arya and M. Agarwal, Bioorg. Med. Chem. Lett., 2007, 17, 86-93.
- 231 M. S. Katagi, G. S. Bolakatti, A. M. Badiger, D. Satyanarayana, S. N. Mamledesai and M. L. Sujatha, Drug Res., 2013, 63, 53-59.
- 232 V. Sharma, R. Das, D. K. Mehta, D. Sharma, S. Aman and M. U. Khan, Mol. Diversity, 2024, 1-27.
- 233 Z. S. Quan, J. M. Wang, J. R. Rho, K. C. Kwak, H. C. Kang, C. S. Jun and K. Y. Chai, Bull. Korean Chem. Soc., 2005, 26, 1757-1760.
- 234 A. Pradhanand and S. K. Vishwakarma, Chem. Int., 2020, 6, 224-231.
- 235 S. Smita, G. Anand, S. Ranjit and V. Vikrant, Int. J. Pharm. Sci. Res., 2011, 3, 164-171.
- 236 M. S. Katagi, J. Fernandes, D. Satyanarayana, G. Bolakatti and S. N. Mamledesai, J. Pharm. Sci., 2014, 4, 57-61.
- Katagi, J. Fernandes, S. Mamledesai, 237 S. M. D. Satyanarayana, P. Dabadi and G. Bolakatti, J. Pharm. Res., 2015, 14, 51-56.
- 238 H. Bi, Q. Ouyang, Z. Wei and Z. Zheng, Bioinorg. Chem., 2020, 100, 103902.
- 239 M. S. Katagi, S. Mamledesai, G. Bolakatti, J. Fernandes, S. ML and P. Tari, Chem. Data Collect., 2020, 30, 100560.
- 240 D. N. Toan, N. D. Thanh, M. X. Truong and D. T. Van, Med.Chem., 2022, 18, 36-50.
- 241 Q. Li, K. W. Woods, W. Wang, N. H. Lin, A. Claiborne, W. Gu, J. Cohen, V. S. Stoll, C. Hutchins, D. Frost, S. H. Rosenberg and H. L. Sham, Bioorg. Med. Chem. Lett., 2005, 15, 2033-2039.
- 242 A. Detsi, D. Bouloumbasi, K. C. Prousis, M. Koufaki, G. Athanasellis, G. Melagraki, A. Afantitis, O. Igglessi-Markopoulou, C. Kontogiorgis and D. J. Hadjipavlou-Litina, J. Med. Chem., 2007, 50, 2450-2458.
- 243 A. S. Ginsburg, J. H. Grosset and W. R. Bisha, Lancet Infect. Dis., 2003, 3, 432-442.

244 A. Aubry, N. Veziris, E. Cambau, C. Truffot-Pernot, V. Jarlier and L. M. Fisher, *Antimicrob. Agents Chemother.*, 2006, **50**, 04–112.

RSC Advances

- 245 G. C. Muscia, M. Bollini, A. M. Bruno and S. E. Asís, *J. Chil. Chem. Soc.*, 2006, **51**, 859–863.
- 246 A. F. B. Cheng, W. W. Yew, E. W. C. Chan, M. L. Chin, M. M. M. Hui and R. C. Y. Chan, *Antimicrob. Agents Chemother.*, 2004, 48, 596.
- 247 I. Dine, E. Mulugeta, Y. Melaku and M. Belete, *RSC Adv.*, 2023, **13**, 8657–8682.
- 248 H. Yin, Y. Wu, X. Gu, Z. Feng, M. Wang, D. Feng, M. Wang, Z. Chenga and S. Wang, *RSC Adv.*, 2022, **12**, 21066–21078.
- 249 M. A. Elanany, E. E. A. Osman, E. M. Gedawy and S. M. Abou-Seri, Sci. Rep., 2023, 13, 4144.
- 250 F. Thurner and F. A. Alatraktchi, *Chemosensors*, 2023, 11, 493.
- 251 C. Ji, I. Sharma, D. Pratihar, L. L. Hudson, D. Maura, T. Guney, L. G. Rahme, E. C. Pesci, J. P. Coleman and D. S. Tan, *ACS Chem. Biol.*, 2016, 11, 3061–3067.
- 252 K. Alla, V. Vijayakumar and S. Sarveswari, *Polycycl*, 2023, 43, 2844–2865.
- 253 S. Khasnobis, V. E. Escuyer and D. Chatterjee, *Expert. Opin. Ther. Targets*, 2002, **6**, 21–40.
- 254 M. J. Ahsan, M. Y. Ansari, S. Yasmin, S. S. Jadav, P. Kumar, S. K. Garg, A. Aseri and H. Khalilullah, *Infect. Disord. Drug Targets*, 2015, **15**, 32–41.
- 255 N. Minovski, M. Vracko and T. Šolmajer, *Mol. Diversity*, 2011, **15**, 417–426.
- 256 A. Lilienkampf, J. Mao, B. Wan, Y. Wang, S. G. Franzblau and A. P. Kozikowski, *J. Med. Chem.*, 2009, **52**, 2109–2118.
- 257 B. Villemagne, C. Crauste, M. Flipo, A. R. Baulard, B. Déprez and N. Willand, *Eur. J. Med. Chem.*, 2012, **51**, 1–16.
- 258 D. S. Reddy, K. M. Hosamani and H. C. Devarajegowda, *Eur. J. Med. Chem.*, 2015, **101**, 705–715.
- 259 M. K. Sharma, M. K. Kumawat, A. Diwan, S. Sardana, N. Yadav and B. Kumar, *Herb. Med. J.*, 2024, **9**.
- 260 V. T. Andriole, Infect. Dis. Clin. Pract., 1994, 3, S211-S218.
- 261 P. S. Dube, L. J. Legoabe and R. M. Beteck, *Mol. Diversity*, 2023, 27, 1501–1526.
- 262 P. C. Appelbaum and P. A. Hunter, *Int. J. Antimicrob. Agents*, 2000, **16**, 5–15.
- 263 T. D. M. Pham, Z. M. Ziora and M. A. T. Blaskovich, *Med. Chem. Commun.*, 2019, **10**, 1719–1739.
- 264 A. M. Emmerson and A. M. Jones, *J. Antimicrob. Chemother.*, 2003, **51**, 13–20.
- 265 R. Stahlmann, Toxicol. Lett., 2002, 127, 269-277.
- 266 F. M. E. Wagenlehner and K. G. Naber, *Curr. Infect. Dis. Rep.*, 2005, 7, 9–16.
- 267 Y. Pommier, E. Leo, H. L. Zhang and C. Marchand, *Chem. Biol.*, 2010, 17, 421–433.
- 268 A. Kumar, A., J. Fernandes and P. Kumar, *Orient. J. Chem.*, 2014, 4, 1993–1997.
- 269 V. Krishnakumar, F. R. N. Khan, B. K. Mandal, S. Mitta, R. Dhasamandha and V. N. Govindan, Res. Chem. Intermed., 2012, 38, 1819–1826.
- 270 C. P. Kumar, M. S. Katagi and B. P. Nandeshwarappa, *Chem. Data Collect.*, 2022, 42, 100955.

- 271 S. A. Al-Issa, Saudi Pharm. J., 2013, 21(3), 305-316.
- 272 M. M. Butler, W. A. LaMarr, K. A. Foster, M. H. Barnes,
 D. J. Skow, P. T. Lyden, L. M. Kustigian, C. Zhi,
 N. C. Brown, G. E. Wright and T. L. Bowlin, *Antimicrob. Agents Chemother.*, 2007, 51(1), 119–127.
- 273 H. A. Azaba, I. A. Ibrahim, N. Hassan, A. M. Abbas and H. M. Darwish, *J. Lumin.*, 2017, **192**, 376–384.
- 274 R. An, M. Ahmed, H. Li, Y. Wang, A. Zhang, Y. Bi and Z. Yu, *Sci. Rep.*, 2021, **11**(1), 12020.
- 275 Z. L. Liu, S. S. Chu and G. H. Jiang, *J. Agric. Food Chem.*, 2009, 57(21), 10130–10133.
- 276 Y. X. Liu, H. P. Zhao, Z. W. Wang, Y. H. Li, H. B. Song, H. Riches, D. Beattie, Y. C. Gu and Q. M. Wang, *Mol. Diversity*, 2013, 17, 701–710.
- 277 J. Jampilek, R. Musiol, M. Pesko, K. Kralova, M. Vejsova, J. Carroll, A. Coffey, J. Finster, D. Tabak, H. Niedbala, V. Kozik, J. Polanski, J. Csollei and J. Dohnal, *Molecules*, 2009, 14(3), 1145–1159.
- 278 M. Abe, T. Imai, N. Ishii and M. Usui, *Biosci., Biotechnol., Biochem.*, 2006, **70**(1), 303–306.
- 279 V. Uivarosi, Molecules, 2013, 18(9), 11153-11197.
- 280 M. M. Khalaf, M. Gouda, K. Shalabi, S. Shaaban and H. M. Abd El-Lateef, *ACS Omega*, 2024, **9**, 6466–6481.
- 281 H. A. Althobiti and S. A. Zabin, *Open Chem.*, 2020, **18**, 591–607.
- 282 R. Fouad and H. F. El-Shafiy, *J. Mol. Struct.*, 2019, **1190**, 68–76.
- 283 B. S. Creaven, B. Duff, D. A. Egan, K. Kavanagh, G. Rosair, V. R. Thangella and M. Walsh, *Inorg. Chim. Acta*, 2010, 363(14), 4048–4058.
- 284 K. Habibi, M. Mamaghani and M. Nikpassand, *Bulg. Chem. Commun.*, 2018, **50**, 76–82.
- 285 M. Azad, J. Appl. Sci., 2007, 7, 2485-2489.
- 286 G. Ayyannan, K. Karthikeyan, S. S. Vivekananthan, M. Gopiraman and A. Rathinavelu, *Ionics*, 2013, 19, 919– 932.
- 287 C. M. Fernandes, A. R. Costa, M. C. Leite, V. Martins, H. S. Lee, F. da CS Boechat, M. C. B. V. de Souza, P. N. Batalha, H. Lgaz and E. A. Ponzio, *J. Mol. Liq.*, 2023, 375, 121299.
- 288 M. M. Abdou and M. N. EL-Haddad, *J. Mol. Struct.*, 2025, 1324, 140889.
- 289 X. M. Meng, S. X. Wang and M. Z. Zhu, Quinoline-Based Fluorescence Sensors, *Molecular Photochemistry Various Aspects*, 2012, ch. 2, DOI: 10.5772/31771.
- 290 B. Bag and A. Pal, Org. Biomol. Chem., 2011, 9, 4467-4480.
- 291 I. Grabchev, J. M. Chovelon and C. Petkov, *Spectrochim. Acta, Part A*, 2008, **69**, 100–104.
- 292 L. Zhang, J. Fan and X. Peng, *Spectrochim. Acta, Part A*, 2009, 73, 398–402.
- 293 U. Diwan, A. Kumar, V. Kumar and K. K. Upadhyay, *Dalton Trans.*, 2013, 42, 13889–13896.
- 294 Y. Yang, K. Yu, L. Yang, J. Liu, K. Li and S. Luo, *Sensors*, 2014, **15**, 49–58.
- 295 B. Li, J. Tian, D. Zhang and F. Tian, *Lumin.*, 2017, 32, 1567–1573.