RSC Advances



PAPER

View Article Online



Cite this: RSC Adv., 2025, 15, 34334

Design and synthesis of Indol-PHOX: a new class of modular phosphine-oxazoline ligands for palladium-catalyzed enantioselective decarboxylative allylation

Anzhela Karapetyan, a Laura Borrel, b Virginie Ratovelomanana-Vidal, Phannarath Phansavath, (1) *a Tahar Ayad (1) b and David Virieux (1) *b

A novel class of chiral and modular P,N-ligands featuring an indolizine core (Indol-PHOX) has been developed, successfully enabling the highly efficient and enantioselective palladium-catalyzed decarboxylative allylation of various α -fluoro- β -ketoesters to access valuable α -allyl- α -fluoro ketones with yields up to 99% and 96% ee. This process exhibits a broad substrate scope, demonstrating good tolerance for both electron-withdrawing and electron-donating groups, as well as substitution patterns of the substrates. The reaction is scalable to the gram level, and subsequent post-functionalization reactions proceed successfully without erosion of enantiopurity.

Received 4th July 2025 Accepted 2nd September 2025

DOI: 10.1039/d5ra04775g

rsc.li/rsc-advances

Introduction

The development of numerous single-enantiomer drugs, along with a broad spectrum of other biologically active substances agrochemicals, pheromones, fragrances-has been driven by significant advancements in asymmetric catalysis conducted over the past decades.1 Central to this progress is the design and synthesis of cost-effective ligands that are readily prepared and can be swiftly tailored for a specific chemical transformation. Among the plethora of reported ligands,2 chiral P,N-ligands, in particular, have gained particular prominence due to their dual functionality, combining the benefits of both phosphorus and nitrogen coordinating atoms.3 One significant example is the chiral phosphine-oxazoline (PHOX) ligands, which were introduced independently by Pfaltz, ⁴ Helmchen, ⁵ and Williams ⁶ in 1993. These ligands coordinate to transition metals through both electronically hard nitrogen and soft phosphorus atoms. They have demonstrated remarkable efficacy as chiral inducers across a wide range of transition metal-catalyzed reactions. Moreover, their modular design allows for easy fine-tuning of geometric, steric and electronic properties by modifying various components of the ligand scaffold, such as the phosphine group, oxazoline ring, or backbone moiety. With their easy accessibility and the virtually limitless selection of readily available precursors, it is not surprising that, since the initial

Nevertheless, despite these significant advancements, it is important to recognize that no universal ligand exists, and that the scaffold plays a determining role in profoundly influencing catalytic performance. As a result, the quest for novel ligands with unique backbones to enhance the selectivity, activity, and




Fig. 1 Representative examples of phosphine-oxazoline (PHOX) ligands (R, R^1 , R^2 = alkyl, Ar = aryl).

studies on phosphinooxazolines, a remarkable array of new linkers—only a few major classes of which are shown in Fig. 1 -connecting the oxazoline ring to the phosphine backbone has been developed and successfully employed in various asymmetric catalytic transformations.7

^aPSL University, Chimie ParisTech, CNRS UMR 8060, Institute of Chemistry for Life and Health Sciences, CSB2D Team, 75005 Paris, France. E-mail: phannarath. phansavath@chimieparistech.psl.eu

 $[^]bICGM,\,Univ.\,Montpellier,\,ENSCM,\,CNRS\,Montpellier,\,France.\,\,E-mail:\,david.virieux @$ enscm.fr

2-benzylpyridine (2.0 eq.) Acetone, reflux, 72 h NaOH (2.0 eq.) FtOH 130 °C 50 W 30 min quant. 1) H₂N ОН DIPEA (1.0 eq.), PyBOP (1. eq.) (1.0 eq.) DCM, RT, 2 h 2) MsCl (1.2 eq.), NEt₃ (6.0 eq.), DCM, 0 °C to RT, 16 h 4a : R = Ph, 86% 4b : R = tBu, 60% LDA (1.1 eq.), CIPAr₂ (1.1eq.) THF, -78 °C to RT, 2 h 5a : Ar = Ph. R = Ph. 82% Ar = furyl, R = Ph, 82% Ar = C_6H_5 , R = tBu, 80% : Ar = 4-(CF₃)C₆H₄, R = tBu, **64%** : Ar = 3,5-(CH₃)₂C₆H₃, R = tBu, **66%**

Ar = furyl, R = tBu, 70%

Scheme 1 Synthesis of Indol-PHOX ligands.

reactivity of transition metal catalysts still remains a formidable challenge. To address this, we have recently initiated a new program dedicated to the design and synthesis of ligands featuring heteroaryl scaffolds, which are underexplored in the realm of chiral ligand design.8 As part of this effort, we have chosen to focus on the indolizine motif for several compelling reasons, as we believe it holds great promise as a framework for the synthesis of P,N-ligands. Indolizines are electron-rich heteroaromatic compounds that belong to the family of Nfused 5/6-membered heterocycles. They can be easily synthesized on a gram-scale using highly efficient and cost-effective

methods, allowing for the introduction of tailored substituents at specific positions. Additionally, indolizines display high and controllable reactivity, enabling straightforward modulation for fine-tuning of steric and electronic properties. In this paper, we present the concise and modular synthesis of a new series of Indol-PHOX ligands, which, to the best of our knowledge, represent the first examples of chiral P,N-chelating indolizine ligands.

Results and discussion

The streamlined synthesis pathway for the preparation of ligands 5 is outlined in Scheme 1. The reaction of commercially available and cost-effective ethyl bromopyruvate 1 with 2-benzylpyridine in refluxing acetone for 72 h, followed by alkaline hydrolysis of the resulting product 2 under microwave irradiation, afforded 3-phenyl-indolizine-2-carboxylic acid 3 on a multi-gram scale.

Condensation of 3 with enantiomerically pure (S)-t-leucinol (S)-2-amino-2-phenylethan-1-ol in the presence benzotriazol-1-yloxytripyrrolidinophosphonium hexafluorophosphate (PyBOP) and diisopropyethylamine in dichloromethane, yielded the corresponding amide. Without further purification, these amides were converted to the oxazolines 4a and 4b via treatment with triethyl amine and methanesulfonyl chloride in dichloromethane, in 86% and 60% isolated yields, respectively. Finally, target ligands 5a-5f were obtained in yields ranging from 64% to 82% through a standard ortho-lithiation process using lithium isopropylamide at -78 °C, followed by reaction with readily accessible chlorodiarylphosphines bearing electron-rich and electron-poor substituents or chlorodifurylphosphine.

Table 1 Optimization of the reactions conditions for the enantioselective decarboxylative allylation a

Entry	$[Pd]^b/L (mol\%)$	T (°C)	t (h)	Conv. (%)	7:8 ratio ^c	$\operatorname{Yield}^{d}\left(\% ight)$	ee ^e (%)
1	[Pd]/5a (2.5/6.25)	25	48	45	91:9	25	14
2	[Pd]/ 5b (2.5/6.25)	25	48	45	91:9	25	14
3	[Pd]/ 5c (2.5/6.25)	25	48	33	85:15	20	76
4	[Pd]/ 5c (5.0/12.5)	25	15	100	93:7	85	80
5	[Pd]/ 5d (5.0/12.5)	30	30	36	78:22	25	87
6	[Pd]/5e (5.0/12.5)	30	30	59	88:12	52	76
7	[Pd]/5e (5.0/12.5)	40	24	100	83:17	65	80
8^f	[Pd]/ 5e (5.0/12.5)	40	3	100	100:0	95	88
9 ^f	[Pd]/ 5f (5.0/12.5)	40	3	100	100:0	99	94
10 ^f	[Pd]/5f(2.5/6.25)	40	3	100	100:0	99	94

^a Conditions: 6a (0.11 mmol), [Pd₂(dba)₃] (2.8 or 5.5 μmol, 2.5 or 5.0 mol%), ligand 5 (7.0 or 14.0 μmol, 6.25 or 12.5 mol%), THF or MTBE (0.05 M), 25–40 °C, 3–48 h. [Pd] refers to [Pd₂(dba)₃]. Continuous Determined by HNMR spectroscopy of the crude product. Isolated yield. Enantiomeric excess for 7a was determined by HPLC analysis. The reaction was run in MTBE (0.05 M).

Scheme 2 Substrate scope of the enantioselective decarboxylative allylation. a a Conditions: 6 (0.13 mmol), $[Pd_2(dba)_3]$ (3.3 μ mol, 2.5 mol%, 3.0 mg), ligand 5f (8.3 μ mol, 6.25 mol%, 4.0 mg), MTBE (0.05 M, 2.6 mL), 40 °C, 3 h. Isolated yields. Enantiomeric excesses were determined by HPLC analysis. b Reaction time of 20 h. c X-ray crystallography of 7n. Displacement ellipsoids are shown at the 30% probability level.

We chose to evaluate these new series of Indol-PHOX ligands in the palladium-catalyzed enantioselective decarboxylative allylation⁹⁻¹⁷ of α -fluoro- β -ketoesters aiming to access valuable α -allyl- α -fluoro ketones. Compound **6a** served as a model substrate for the optimization of the reaction parameters for the enantioselective decarboxylative allylation (Table 1). The reaction was first carried out at 25 °C in THF in the presence of 2.5 mol% of [Pd₂(dba)₃] and 6.25 mol% of ligand **5a** for 48 h

(Table 1, entry 1). Under these conditions, a conversion of 45% was achieved and the desired product 7a was isolated in 25% yield and 14% ee alongside the by-product 8a resulting from the protonation of the enolate generated during the reaction (with a 7a/8a ratio of 91:9). The reaction carried out under the same conditions with ligand 5b produced identical results in terms of yield and enantioselectivity (Table 1, entry 2). However, a significant increase in enantioinduction was observed with ligand 5c bearing a tert-butyl substituent on the oxazoline ring instead of the phenyl substituent present in ligands 5a and 5b. Indeed, although a lower conversion of 33% with a 7a/8a ratio of 85:15 was achieved, the enantiomeric excess of 7a could be improved to 76% (Table 1, entry 3). Increasing the catalyst loading to 5 mol% for [Pd₂(dba)₃] and 12.5 mol% for 5c, allowed to reach a full conversion within 15 h, with a 7a/8a ratio of 93:7 (85% yield and 80% ee for 7a) (Table 1, entry 4). When the reaction was run with ligands 5d and 5e, incomplete conversions of 36% and 59%, respectively, were observed even at a higher temperature of 30 °C (Table 1, entries 5 and 6, 25% and 52% yield respectively for 7a). However, when the transformation was operated at 40 °C using ligand 5e, a complete conversion was achieved and 7a was isolated in 65% yield with 80% ee and a 7a/8a ratio of 83:17 (Table 1, entry 7). Interestingly, the formation of by-product 8a could be prevented by using MTBE as a solvent. In that case the rearrangement product 7a was obtained in 95% yield and 88% ee using ligand 5e within a shorter reaction time of 3 h (Table 1, entry 8), whereas ligand 5f allowed both a higher yield of 99% and a better enantioselectivity of 94% ee (Table 1, entry 9).

Notably, we found that the catalyst loading could be decreased to 2.5 mol% for [Pd₂(dba)₃] and 6.25 mol% for 5f with no detrimental effect (Table 1, entry 10). However when the catalyst loading was further lowered to 1.0 mol% of [Pd₂(dba)₃] and 2.5 mol% of the ligand, a 7a/8a ratio of 95:5 was observed with an isolated yield of 90% for 7a even though the enantioselectivity of 94% ee remained unaffected. On the other hand, a short solvent screening confirmed MTBE as the more suitable solvent for the reaction since 7a was obtained in 85% yield with 88% ee (7a/8a = 100:0) in toluene, and in 69% yield and 90% ee in diethyl ether (7a/8a = 95:5). The use of protic solvents significantly lowered the reaction rate, with MeOH affording only 15% conversion (7a/8a = 40:60) and iPrOH giving 18% conversion (7a/8a = 50:50) at 40 °C after 3 h. Thus, the optimized reaction conditions were set as follows: 2.5 mol% of [Pd₂(dba)₃], 6.25 mol% of ligand **5f** in MTBE (0.05 M) at 40 °C for 3 h. It should be noted that when the commercially available (S)-i-Pr-PHOX ligand was used instead of ligand 5f, the reaction led to a 7a/8a ratio of 90: 10 and the rearrangement product was isolated in 87% yield and 84% ee. When [Pd(allyl)Cl]2 or Pd(OAc)₂ were employed as alternative palladium sources, the reaction proceeded with only 13% conversion in the case of [Pd(allyl)Cl]₂ and 3% conversion with Pd(OAc)₂.

With the optimized reaction conditions in hand, the scope of the enantioselective decarboxylative allylation using the newly developed ligand $\mathbf{5f}$ was evaluated using the previously synthesized family of substituted β -ketoesters $\mathbf{6a-6t}$ (Scheme 2). Allyl 1-oxo-1,2,3,4-tetrahydronaphthalene-2-carboxylate

Scale-up experiment: crystallographic analysis, and by analogy, we conjectured that the remainder of the rearrangement products followed the same trend.

Scheme 3 Scale-up experiment and post-functionalization reactions

derivatives bearing electron-donating groups such as a methoxy or a methyl substituent on the phenyl ring led to high yields (94-97%) of the rearrangement products 7b-7d that were formed with excellent enantioinductions of 94-96% ee. In the same manner, the parent substrates having halogen atoms such as bromine, chlorine or fluorine at various positions of the phenyl ring were obtained in 90-95% yield and 93-94% ee (7e-7i). The more sterically hindered derivative 7j having a naphthyl ring instead of the benzene ring was isolated in a lower yield of 52% whereas the enantioselectivity remained similar (91% ee). On the other hand, the enantioselective decarboxylative allylation carried out with the tetralone substrate bearing an isoprenyl substituent failed to give any conversion into 7k whereas the reaction of a 7-membered carbocycle led to (R)-6-allyl-6fluoro-6,7,8,9-tetrahydro-5*H*-benzo[7]annulen-5-one 71 in 97% yield and 92% ee. Indanone derivatives were also suitable substrates for this transformation giving products 7m-7s in yields ranging from 66% to 95% and high enantiomeric excesses of 84-94%. As previously observed for the tetralone series, the enantioselectivity of the reaction with indanone derivatives was not dependent on either the electronic nature or the positions of the substituents present on the benzene ring. Indeed, electron-withdrawing substituents such as chlorine or trifluoromethyl (7n-7o) led to the same levels of enantioinduction than electron-donating groups such as methoxy or methyl (7p-7r). On the other hand, the sterically hindered derivative 7s bearing a naphthyl ring was formed in 79% yield with an excellent enantiomeric excess of 94%. The reaction was also extended to a heterocyclic substrate, affording the fluorinated thiochromanone 7t which was obtained in 48% yield and 92% ee. The absolute configuration of compounds 7a, 7l and 7m were assigned by comparison of the optical rotation values with those reported.¹⁸ Additionally, the absolute configuration of compound 7n was unambiguously assigned as (R) by X-ray

The efficiency of the new ligand 5f in the enantioselective decarboxylative allylation was further supported by a scale-up experiment performed with substrate 6a on a gram scale. Under the optimized reaction conditions, the desired product 7a was obtained in 92% yield and 94% ee (Scheme 3). Furthermore, the rearrangement products 7 can serve as useful intermediates for further post-functionalization reactions. Thus, aryl bromide 7h was engaged in a Suzuki-Miyaura coupling reaction with 4-methoxyphenylboronic acid using $Pd(OAc)_2$ as a catalyst, cataCXium A as a ligand, and K_2CO_3 as a base, affording the corresponding binaphthyl 9 in 93% yield while maintaining the 94% ee. On the other hand, a crossmetathesis reaction between (R)-2-allyl-2-fluoro-3,4-dihydronaphthalen-1(2H)-one 7a and styrene in the presence of 5.0 mol% of the Grubbs catalyst 2nd generation afforded the corresponding alkene (E)-10 in 70% yield and 94% ee.

Conclusions

In summary, we have developed a new, efficient and modular series of chiral P,N ligands, termed Indol-PHOX ligands, representing the first examples of chiral oxazolidinyl indolizinebased ligands. We also demonstrated their successful application in palladium-catalyzed enantioselective decarboxylative allylation. The reactions proceed smoothly under mild reaction conditions and accommodate a wide substrate scope providing valuable α-allyl-α-fluoro ketones that bear an α-quaternary stereocenter. Such compounds were isolated in high yields (up to 99%) and with excellent enantioselectivities (up to 96% ee). The scalability of the process was confirmed through a gramscale experiment, and the synthetic utility of the resulting chiral α-allyl-α-fluoro ketone products was further demonstrated through successful post-functionalization reactions (Suzuki-Miyaura coupling and cross-metathesis) without erosion of enantiopurity.

Conflicts of interest

There are no conflicts of interest to declare.

Data availability

CCDC 2434440 contains the supplementary crystallographic data for this paper.¹⁹

The data supporting this article have been included as part of the SI. Supplementary information: Experimental procedures, NMR spectra, spectral data and X-ray data. See DOI: https://doi.org/10.1039/d5ra04775g.

Acknowledgements

Financial support from the ANR (BOOSTS project, ANR-21-CE07-0034) is gratefully acknowledged. We thank G. Gontard

for the X-ray analysis (Sorbonne Université, Paris) and Dr Céline Fosse for HRMS analysis (Chimie ParisTech-PSL).

References

- 1 (*a*) N. Senkuttuvan, B. Komarasamy, R. Krishnamoorthy, S. Sarkar, S. Dhanasekaran and P. Anaikutti, The Significance of Chirality in Contemporary Drug Discovery— a Mini Review, *RSC Adv.*, 2024, **14**, 33429; (*b*) J. Clayden, N. Greeves and S. G. Warren, *Organic Chemistry*, Oxford University Press, Oxford, 2nd edn, 2012, pp. 1102–1133.
- 2 R. Connon, B. Roche, B. V. Rokade and P. J. Guiry, Further Developments and Applications of Oxazoline-Containing Ligands in Asymmetric Catalysis, *Chem. Rev.*, 2021, **121**, 6373.
- 3 M. P. Carroll and P. J. Guiry, P,N Ligands in Asymmetric Catalysis, *Chem. Soc. Rev.*, 2014, 43, 819.
- 4 P. Von Matt and A. Pfaltz, Chiral Phosphinoaryldihydrooxazoles as Ligands in Asymmetric Catalysis: Pd-Catalyzed Allylic Substitution, *Angew. Chem., Int. Ed.*, 1993, 32, 566.
- 5 J. Sprinz and G. Helmchen, Phosphinoaryl- and Phosphinoalkyloxazolines as New Chiral Ligands for Enantioselective Catalysis: Very High Enantioselectivity in Palladium Catalyzed Allylic Substitutions, *Tetrahedron Lett.*, 1993, 34, 1769.
- 6 G. J. Dawson, C. G. Frost, J. M. J. Williams and S. J. Coote, Asymmetric Palladium Catalysed Allylic Substitution Using Phosphorus Containing Oxazoline Ligands, *Tetrahedron Lett.*, 1993, 34, 3149.
- 7 (a) Y. Imai, W. Zhang, T. Kida, Y. Nakatsuji and I. Ikeda, Diphenylphosphinooxazoline Ligands with Binaphthyl Backbone for Pd-Catalyzed Allylic Alkylation, Tetrahedron Lett., 1998, 39, 4343; (b) W. Zhang, F. Xie, H. Yoshinaga, T. Kida, Y. Nakatsuji and I. Ikeda, A Novel Axially Chiral Phosphine-Oxazoline Ligand with an Axis-Unfixed Biphenyl Backbone: Preparation, Complexation, and Application in an Asymmetric Catalytic Reaction, Synlett, 2006, 8, 1185; (c) H. Danjo, M. Higuchi, M. Yada and T. Imamoto, P-Stereogenic P/N Hybrid Ligands: a Remarkable Switch in Enantioselectivity in Palladium-Catalyzed Asymmetric Allylation, Tetrahedron Lett., 2004, 45, 603; (d) S.-F. Zhu, J.-B. Xie, Y.-Z. Zhang, S. Li and Zhou, Well-Defined Chiral Spiro Iridium/ Q.-L. Phosphine-Oxazoline Cationic Complexes for Highly Enantioselective Hydrogenation of Imines at Ambient Pressure, J. Am. Chem. Soc., 2006, 128, 12886; (e) Y. Zhang, Z. Han, F. Li, K. Ding and A. Zhang, Highly Enantioselective Hydrogenation of α-Aryl-β-Substituted Acrylic Acids Catalyzed by Ir-SpinPHOX, Chem. Commun., 2010, 46, 156; (f) S. P. Smidt, F. Menges and A. Pfaltz, SimplePHOX, a Readily Available Chiral Ligand System for Iridium-Catalyzed Asymmetric Hydrogenation, Org. Lett., 2004, 6, 2023; (g) X.-L. Hou, D. X. Dong and K. Yuan, Synthesis of New Chiral Benzylically Substituted P,N-Ligands and Their Applications in the Asymmetric Heck Reaction, Tetrahedron: Asymmetry, 2004, 15, 2189; (h)

- D. J. Richards, T. Damalidis, D. E. Hibbs and M. B. Hursthouse, Synthesis of 2-[2-(Diphenylphosphino) ferrocenyl]oxazoline Ligands, *Synlett*, 1995, 74; (*i*) Y. Nishibayashi and S. Uemura, Asymmetric Synthesis and Highly Diastereoselective ortho-Lithiation of Oxazolinylferrocenes, *Synlett*, 1995, 79; (*j*) Y. Wang, A. Hämäläinen, J. Tois and R. Franzén, Preparation of Indole-Phosphine Oxazoline (IndPHOX) Ligands and Their Application in Allylic Alkylation, *Tetrahedron: Asymmetry*, 2010, 21, 2376.
- 8 (a) C. Laborde, M.-M. Wei, A. van der Lee, E. Deydier, J.-C. Daran, J.-N. Volle, R. Poli, J.-L. Pirat, E. Manoury and D. Virieux, Double [3+2]-Dimerisation Cascade Synthesis of Bis(triazolyl)bisphosphanes, a New Scaffold for Bidentate Bisphosphane, Dalton Trans., 2015, 44, 12539; (b) N. Sevrain, J.-N. Volle, J.-L. Pirat, T. Ayad and D. Virieux, 1,1'-Dibenzyl-Bis-(Triazolyl)diphenylphosphine a New Efficient Organocatalyst for Silicon Tetrachloride-Mediated Enantioselective Abramov-Type Phosphonylation of Aldehydes with Trialkyl Phosphites, RSC Adv., 2017, 7, 52101; (c) N. Sevrain, J.-N. Volle, J.-L. Pirat, T. Ayad and D. Virieux, Chiral Bisdiphenylphosphine Dioxides Bearing an Original Bis(triazolyl) Backbone as Promising Lewis Base for Asymmetric Organocatalysis, Eur. J. Org Chem., 2018, 2267; (d) A. Gernet, V. Ratovelomamana-Vidal, J.-L. Pirat, D. Virieux and T. Ayad, Efficient Synthesis of 2-Amino-1-Arylethanols through a Lewis Base-Catalyzed SiCl₄-Mediated Asymmetric Passerini-Type Reaction, Eur. J. Org Chem., 2020, 6497; (e) A. Gernet, N. Sevrain, J.-N. Volle, T. Ayad, J.-L. Pirat and D. Virieux, Diversity-Oriented Synthesis Toward Aryl- and Phosphoryl-Functionalized Imidazo[1,2-a]Pyridines, J. Org. Chem., 2020, 85, 14730.
- 9 G. Laidlaw and V. Franckevičius, Palladium-Catalyzed Decarboxylative Asymmetric Allylic Alkylation of Thietane 1,1-Dioxides, Org. Lett., 2022, 24, 400.
- 10 J. James, M. Jackson and P. J. Guiry, Palladium-Catalyzed Decarboxylative Asymmetric Allylic Alkylation: Development, Mechanistic Understanding and Recent Advances, Adv. Synth. Catal., 2019, 361, 3016.
- 11 R. Kuwano, N. Ishida and M. Murakami, Asymmetric Carroll Rearrangement of Allyl α-Acetamido-β-Ketocarboxylates Catalysed by a Chiral Palladium Complex, *Chem. Commun.*, 2005, **31**, 3951.
- 12 J. D. Weaver, A. Recio, A. J. Grenning and J. A. Tunge, Transition Metal-Catalyzed Decarboxylative Allylation and Benzylation Reactions, *Chem. Rev.*, 2011, **111**, 1846.
- 13 D. C. Behenna and B. M. Stoltz, The Enantioselective Tsuji Allylation, *J. Am. Chem. Soc.*, 2004, **126**, 15044.
- 14 B. M. Trost and J. Xu, Regio- and Enantioselective Pd-Catalyzed Allylic Alkylation of Ketones through Allyl Enol Carbonates, J. Am. Chem. Soc., 2005, 127, 2846.
- 15 J. T. Mohr, D. C. Behenna, A. M. Harned and B. M. Stoltz, Deracemization of Quaternary Stereocenters by Pd-Catalyzed Enantioconvergent Decarboxylative Allylation of Racemic β-Ketoesters, *Angew. Chem., Int. Ed.*, 2005, **44**, 6924.

- 16 M. Nakamura, A. Hajra, K. Endo and E. Nakamura, Synthesis of Chiral α-Fluoroketones through Catalytic Enantioselective Decarboxylation, *Angew. Chem., Int. Ed.*, 2005, 44, 7248.
- 17 E. C. Burger, B. R. Barron and J. A. Tunge, Catalytic Asymmetric Synthesis of Cyclic α -Allylated α -Fluoroketones, *Synlett*, 2006, 17, 2824.
- 18 É. Bélanger, K. Cantin, O. Messe, M. Tremblay and J. F. Paquin, Enantioselective Pd-Catalyzed Allylation
- Reaction of Fluorinated Silyl Enol Ethers, *J. Am. Chem. Soc.*, 2007, **129**, 1034.
- 19 A. Karapetyan, L. Borrel, V. Ratovelomanana-Vidal, P. Phansavath, T. Ayad and D. Virieux, CCDC 2434440: Experimental Crystal Structure Determination, 2025, DOI: 10.5517/ccdc.csd.cc2mq7by.