


 Cite this: *RSC Adv.*, 2025, 15, 4331

DOI: 10.1039/d5ra90012c

[rsc.li/rsc-advances](https://rsc.li/rsc-advances)

## Correction: Synthesis of an exfoliated kaolinite–poly(urea–formaldehyde) nanocomposite

 Hervé Barye Tatang,<sup>\*a</sup> Jacques Richard Mache,<sup>b</sup> Cyrill Joël Ngally Sabouang,<sup>c</sup> Angelina Razafitianamaharavo,<sup>d</sup> Renaud Gley,<sup>d</sup> Sakeo Kong<sup>a</sup> and Jean Aimé Mbey<sup>\*a</sup>

 Correction for 'Synthesis of an exfoliated kaolinite–poly(urea–formaldehyde) nanocomposite' by Hervé Barye Tatang *et al.*, *RSC Adv.*, 2025, 15, 3026–3039, <https://doi.org/10.1039/D4RA08707K>.

The authors regret that an incorrect version of eqn (2) and (3) were included in the original article. The correct version of eqn (2) and (3) are presented here.

Eqn (2):

$$\%KUF = \frac{\sum \frac{I}{I_0}(\text{UF bands})}{\frac{I}{I_0}(3621)} \times \frac{100}{7.42}$$

Eqn (3):

$$P_0 = \frac{\left(\frac{I}{I_0}\right) 3620 \text{ cm}^{-1}}{\left(\frac{I}{I_0}\right) 3690 \text{ cm}^{-1}}$$

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

<sup>a</sup>Laboratory of Applied Inorganic Chemistry, Department of Inorganic Chemistry, University of Yaoundé I, P. O. Box 812, Yaoundé, Cameroon. E-mail: mbey25@yahoo.fr; jean-aime.mbey@facsciences-uy1.cm; barye.tatang@facsciences-uy1.cm

<sup>b</sup>School of Geology and Mining Engineering, University of Ngaoundere, P.O. Box 115, Meiganga, Cameroon

<sup>c</sup>Departments of Chemistry, Higher Teacher Training College, University of Bamenda, P. O. Box 39, Bamili, Cameroon

<sup>d</sup>Université de Lorraine, CNRS, LIEC, F-54000 Nancy, France

