

CORRECTION

[View Article Online](#)
[View Journal](#) | [View Issue](#)

Cite this: *RSC Adv.*, 2025, **15**, 27524

DOI: 10.1039/d5ra90091c

rsc.li/rsc-advances

Correction: Glycyrrhetic acid-modified redox-sensitive polymeric mixed micelles for tumor-specific intracellular delivery of cantharidin

Yu Hu,^a Tian Lan,^b Ji Li,^c Lingjun Li^{*a} and Jizheng Song^{*a}

Correction for 'Glycyrrhetic acid-modified redox-sensitive polymeric mixed micelles for tumor-specific intracellular delivery of cantharidin' by Yu Hu *et al.*, *RSC Adv.*, 2024, **14**, 28753–28767, <https://doi.org/10.1039/D4RA03171G>.

Authors regret that an incorrect version of Fig. 6A was included in the originally published article. The correct version of Fig. 6 is shown herein.

^aSchool of Pharmacy, Shandong University of Traditional Chinese Medicine (TCM), 250355 Jinan, Shandong, China. E-mail: sdzyylilingjun@163.com; songjizheng345@163.com

^bInnovative Institute of Chinese Medicine, Shandong University of TCM, 250355 Jinan, Shandong, China

^cAffiliated Hospital of Shandong University of TCM, 250011 Jinan, Shandong, China





Fig. 6 (A) Size distribution and morphology of F127-SS-PDLA/CTD (a) and GA/F127-SS-PDLA/CTD (b) micelles detected with DLS and TEM. The scale bar for TEM represents 100 nm. (B) Size distributions of F127-SS-PDLA/CTD (a) and GA/F127-SS-PDLA/CTD (b) micelles under reductive environment (10 mM GSH) for 24 h. (C) Changes in particle size of CTD-loaded micelles incubated with 10% FBS at 37 °C for 24 h. (D) Changes in particle size and PDI of F127-SS-PDLA/CTD (a) and GA/F127-SS-PDLA/CTD (b) micelles stored at 4 °C for 30 days. (E) Cumulative release profiles of free CTD and CTD in CTD-loaded micelles in normal saline with different GSH concentrations (0 and 10 mM). Data were presented as mean \pm SD; $n = 3$.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.