# Chemical Science



### **EDGE ARTICLE**

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2025, 16, 4014

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 19th October 2024 Accepted 15th January 2025

DOI: 10.1039/d4sc07116f

rsc li/chemical-science

## CO<sub>2</sub> hydrosilylation catalyzed by an N-heterocyclic carbene (NHC)-stabilized stannyliumylidene†

Dechuang Niu, Arseni Kostenko, John A. Kelly, Debotra Sarkar, Huihui Xu and Shigeyoshi Inoue \*

The di-N-heterocyclic carbene (NHCs) stabilized stannyliumylidene, [MesTerSn(IMe<sub>4</sub>)<sub>2</sub>][BArF], (MesTer = 2,6-Mes<sub>2</sub>C<sub>6</sub>H<sub>3</sub>, Mes = 2,4,6-Mes<sub>3</sub>-C<sub>6</sub>H<sub>2</sub>, IMe<sub>4</sub> = 1,3,4,5-tetramethylimidazol-2-ylidene, BArF = (3,5-(CF<sub>3</sub>)<sub>2</sub>-C<sub>6</sub>H<sub>5</sub>)<sub>4</sub>B), was isolated from the reaction of (MesTer)SnCl with two equivalents of IMe<sub>4</sub>, followed by one equivalent of Na[BArF]. This stannyliumylidene acts as a precatalyst for the homogeneous hydrosilylation of CO<sub>2</sub>. Experimental mechanistic studies and quantum chemical calculations have been conducted to elucidate the catalytically active species and the mechanism for the transformation, revealing the stannyliumylidene [MesTerSn(CO<sub>2</sub>IMe<sub>4</sub>)<sub>2</sub>][BArF], which is formed in the presence of CO<sub>2</sub>, as the catalytically active species.

#### Introduction

A major goal in contemporary chemical research is the utilization of carbon dioxide in the creation of fine chemicals, due to its high production/abundance and its status as a greenhouse gas. Recent advances in main group chemistry have seen the successful isolation of highly reactive compounds with unprecedented oxidation states and bonding modes, which in turn have demonstrated the remarkable capability of activating  $CO_2$ .<sup>1-5</sup> For example, silylenes can react with  $CO_2$  yielding the corresponding silicon<sup>IV</sup> carbonates  $[C(=O)(O-)_2]^{2-}$ .<sup>6-7</sup> Germylene and stannylene species tend to afford  $E^{II}$  (E=Ge and Sn) carboxylate compounds, owing to the weak nucleophilicity of their lone pair of electrons.<sup>8,9</sup>


While main group-mediated CO<sub>2</sub> activation is becoming more commonplace, the catalytic transformation of carbon dioxide is still rare, especially when it comes to hydrosilylation reactions. Main-group catalysis has attracted tremendous attention recently, due to the higher abundance and lower toxicity of such elements. Currently, the most potent catalysts for the hydrosilylation of CO<sub>2</sub> using main group compounds are: Lewis bases (LBs), Lewis acids (LAs), and frustrated Lewis pairs (FLPs).¹ The use of N-heterocyclic carbenes (NHCs) as catalysts for the hydrosilylation of CO<sub>2</sub> was first reported by Zhang, Ying and coworkers (Fig. 1, A).¹¹0 Later, experimental and theoretical studies showed that the NHC-CO<sub>2</sub> adduct acts as the active catalytic species in the hydrosilylation of CO<sub>2</sub>, wherein

TUM School of Natural Sciences, Department of Chemistry, Institute of Silicon Chemistry and Catalysis Research Center, Technische Universität München, Lichtenbergstraße 4, 85748 Garching bei München, Germany. E-mail: s.inoue@tum.de † Electronic supplementary information (ESI) available. CCDC 2255434, 2255353 and 2255355. For ESI and crystallographic data in CIF or other electronic format see DOI: https://doi.org/10.1039/d4sc07116f

the Si-H bond is activated *via* coordination of the oxygen atom of the NHC-CO<sub>2</sub> adduct (Fig. 1, B).<sup>11,12</sup>

Main-group Lewis acids have been shown to be capable of the catalytic hydrosilylation of  $CO_2$ , examples include the silylium cation,  $C^{13}$  and two-coordinate group 13 cations D, E, and F

#### (1) NHC catalyzed CO<sub>2</sub> hydrosilylation



#### (2) LA system for CO<sub>2</sub> hydrosilylation

Fig. 1 (1) Proposed mechanism for NHC-catalyzed hydrosilylation of CO<sub>2</sub>; (2) LA systems effective in CO<sub>2</sub> hydrosilylation.

**Edge Article Chemical Science** 

#### (1) FLP catalyzed CO<sub>2</sub> hydrosilylation

Fig. 2 (1) FLP systems; (2) heavier tetryliumylidene cations and K catalyzed CO2

(Fig. 1(2)). 14-16 Okuda and coworkers found BPh3 could catalyze the hydrosilylation of CO<sub>2</sub> to form silyl formate, but only in highly polar, aprotic solvents, such as acetonitrile. 17 They posit the mechanism involves weak/dynamic coordination of BPh3 to CO2, as well as to the Si-H moiety, generating two partially polarized transient species.

Another pertinent contribution to main-group based catalysis was reported in 2009 by the Piers group, in which amine/ borane FLP (Fig. 2, G) was effective in the hydrosilylation of CO2.18 Since then, a multitude of various FLP catalytic systems have been reported, such as  $\{[Tism^{Pribenz}]M\}$   $[HB(C_6F_5)_3][M =$ Zn, Mg; Tism<sup>Pribenz</sup> = tri[(1-isopropylbenzimidazol-2-yl)-dimeand [(Dipp nacnac)Ga(Ad)][HB(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>](thylsilyl]methyl)19,20 <sup>Dipp</sup>nacnac  $= [\{N(Dipp)CMe\}_2CH]^-,$ Dipp diisopropylphenyl, Ad = 1-adamantyl).21 A noteworthy example from the Kato group is an N,P-heterocyclic germylene/B(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub> Lewis pair that can be used as a catalyst for the selective CO<sub>2</sub> hydrosilylation to H<sub>2</sub>C(OSiEt<sub>3</sub>)<sub>2</sub>, through Si-H bond activation to give the highly reactive cationic (amino)(phosphino) germylene with  $HB(C_6F_5)_3$  as a counter anion, followed by activation of CO<sub>2</sub> (Fig. 2, H).<sup>22</sup>

An emerging class of compounds that is of high interest to main group chemists, from the point of view of their catalytic capabilities, are the tetryliumylidenes. Tetryliumylidenes (RE:<sup>+</sup>), are mono-substituted E<sup>II</sup> (E = Si, Ge, Sn, Pb) atoms with a lone pair of electrons and two vacant orbitals, that exhibit high reactivity due to their unsaturated and amphiphilic nature. 4,23,24 There has been a growing number of reports of these compounds ever since the first was reported by Jutzi and coworkers, 23,25 whereby they used pentamethyl-cyclopentadienyl to stabilize the heavier tetryliumylidenes (Fig. 2, I, Cp\*E:<sup>+</sup>, E =  $Si,^{26} Ge,^{27} Sn,^{27} Pb;^{28} Cp^* = Me_5C_5$ ). These were shown to be able to activate small molecules (e.g., CO2, H2, N2O, alkenes, ketones, etc.).24 Silyliumylidene ions stabilized by Lewis bases (e.g., Nheterocyclic carbene (NHC) (J),29 PMe3) could reduce CO2, resulting in the silaacylium ions,  $[RSi(O)(IMe_4)_2]Cl$   $(R = {}^{Mes}Ter$ or Tipp;  $^{\text{Mes}}$ Ter = 2,6-Mes<sub>2</sub>-C<sub>6</sub>H<sub>3</sub>, Mes = 2,4,6-Me<sub>3</sub>-C<sub>6</sub>H<sub>2</sub>, Tipp =

2,4,6-i $Pr_3$ - $C_6H_2$ ,  $IMe_4 = 1,3,4,5$ -tetramethylimidazol-2-ylidene),30,31 via the initial coordination of the C atom to the Si center. We reported on the CO<sub>2</sub> insertion into the Si-C<sup>NHC</sup> bond of the dicationic silicon<sup>IV</sup> compound [MesTer(IMe<sub>4</sub>)<sub>2</sub>SiH][OTf]<sub>2</sub>  $(OTf = CF_3SO_3^-).^{32}$ 

Recently, our group has isolated the germyliumylidene ion (K, Fig. 2), which was shown to react with N2O to produce a germa-acylium ion  $[(^{Mes}Ter)Ge(O)(IMe_4)_2]X$  (X = Cl or BArF, BArF =  $(3.5-(CF_3)_2C_6H_5)_4B$ ).<sup>33</sup> This germa-acylium ion could then be utilized in the catalytic functionalization of CO<sub>2</sub>, the catalytically active germylene, Ge(OSiHPh<sub>2</sub>)(IMe<sub>4</sub>). It is also worth noting that K catalyzes the reduction of CO<sub>2</sub> with silane via a three-component transition state (Fig. 2, L).34 The mechanism is analogous to the cooperative silane/CO<sub>2</sub> mechanism for the hydrosilylation of CO<sub>2</sub> catalyzed by NHC (Fig. 1, B).12

The heavier tetryliumylidene analogues, i.e. stannyliumylidene cations, are also an emerging class of low valent main group species that that may have the potential to be used in catalysis. However, examples of the utilization of stannyliumylidene in the activation of small molecules are rare. Herein, we present the first substantiated example of stannyliumylidene acting as a catalyst for CO2 hydrosilylation. We report the synthesis of the NHC-stabilized stannyliumylidene [2]+ - the tin analogue of J and K, which reacts with CO2 to form the catalytically active stannyliumylidene [4]+. We present the mechanism of conversion of [2]+ to the catalytically active stannyliumylidene species, isolating and fully describing the reactive intermediates of this process, as well as the catalytically active species. Additionally, we examine the catalytic performance of the stannyliumylidene species in the catalytic hydrosilylation CO2 using various silanes and conditions. Based on these studies and quantum chemical calculations we propose a mechanism for the catalytic reaction.

#### Results and discussion

#### Synthesis of stannyliumylidene

NHC-stabilized stannyliumylidene [MesTerSn(IMe<sub>4</sub>)<sub>2</sub>][BArF] ([2] [BArF]) was obtained by the reaction of chlorostannylene, Mes-TerSnCl (1) with two equivalents of IMe<sub>4</sub> in toluene, followed by adding one equivalent of Na[BArF] in a one-pot procedure (Scheme 1). It was isolated as a colorless crystalline solid in

Scheme 1 Synthesis of NHC-stabilized stannyliumylidene ion [2] [BArF].

Chemical Science Edge Article

a 72% yield. [2][BArF] has poor solubility in nonpolar organic solvents, *e.g.*, benzene and toluene, but it is soluble in fluorobenzene and THF. In acetonitrile, substitution occurs between the IMe<sub>4</sub> and solvent molecules leading to a mixture of coordination compounds. This is in stark contrast to silyliumylidene (Fig. 2, **J**) and germyliumylidene ions (Fig. 2, **K**), where ligand substitution does not occur.<sup>29,33</sup> In the <sup>119</sup>Sn{<sup>1</sup>H} NMR spectrum, one sharp signal was observed at  $\delta = -235.72$  ppm in THF-d<sub>8</sub>, similar to the cationic stannylene, [<sup>Tipp</sup>TerSn(IMe<sub>2</sub>)<sub>2</sub>] [HB(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>](<sup>Tipp</sup>Ter = 2,6-Tipp<sub>2</sub>-C<sub>6</sub>H<sub>3</sub>) ( $\delta = -234$  ppm), reported by the Wesemann group.<sup>35</sup> The <sup>13</sup>C{<sup>1</sup>H} NMR spectrum displays a resonance at  $\delta = 170.18$  ppm for the carbon of IMe<sub>4</sub>, which is downshifted compared to that of the germyliumylidene (**K**) ( $\delta = 164$  ppm) and the silyliumylidene (**J**) ( $\delta = 160$  ppm).<sup>29,33</sup>

X-ray quality crystals of [2][BArF] were obtained from a toluene/fluorobenzene ( $C_6H_5F$ ) solution stored at -35 °C. Single crystal X-ray diffraction (SC-XRD) analysis revealed that the Sn central is tricoordinate with two IMe4 moieties and one Mes Ter-substituent (Fig. 3). The sum of the bond angles around the Sn1 atom is 300.4°. The Sn1-B1 distance is 8.587 Å, indicating no interaction between the cation and anion. Additionally, we found the shortest distance between the Sn atom and a fluorine atom at 4.032 Å. This also indicates there is no Sn-F interaction as the sum of the van der Waals radii of these atoms amounts to 3.56 Å (2.16 Å for Sn and 1.40 Å for F). The Sn– $C^{NHC}$ bond distances are Sn1-C25 = 2.339(4) Å and Sn1-C32 = 2.308(4) Å. They are similar to that found in IPr<sub>Me</sub>SnCl<sub>2</sub> (2.290(5)  $\mathring{A}$ ) (IPr<sub>Me</sub> = 1,3-diisopropyl-4,5-dimethylimidazol-2-ylidene),<sup>36</sup>  $IPr_{Me}Sn[C(Cl)PMes^*]_2$  (Mes\* = 2,4,6-tri-tert-butylphenyl) (2.316(2) Å)<sup>37</sup> and slightly longer than (Dipp Ter)Sn(IMe<sub>4</sub>)  $CH_2P(CH_3)_2 = P(^{Mes}Ter) (2.274(3) \text{ Å}) (^{Dipp}Ter = 2,6-Dipp_2-C_6H_3,$  $Dipp = 2,6-iPr_2-C_6H_3$ ).<sup>38</sup> As expected, they are longer than the observed distances in the NHC-stabilized silyliumylidene ion (J) (1.948(19) Å, 1.966(19) Å) and NHC-stabilized germyliumylidene ion (K) (2.063(3) Å, 2.093(3) Å).

To gain further insight into the electronic structure of [2]<sup>†</sup>, as well as to analyze the thermodynamics and kinetics in the processes involving this species, quantum chemical




Fig. 3 The molecular structure of [2]<sup>+</sup>. Ellipsoids are set at the 30% probability level; hydrogen atoms and [BArF]<sup>-</sup> anion are omitted for clarity. Selected bond lengths [Å] and bond angles [°]: Sn1–C1 2.244(4), Sn1–C32 2.308(4), Sn1–C25 2.339(4), C1–Sn1–C25 107.7(1), C1–Sn1–C32 96.7(1), C25–Sn1–C32 96.0(1).




Fig. 4 NBOs of  $[2]^+$  (iso = 0.04) at the PBE0/def2-TZVP// $r^2$ SCAN-3c level of theory. The mesityl substituents and the methyl substituents are shown as wireframes for clarity.

calculations were carried out. For details regarding the computational methods see the ESI.† Natural Bond Orbital (NBO) analysis of the optimized structure of [2]<sup>+</sup> (Fig. 4) shows that the stannyliumylidene cation exhibits the expected bonding situation around the tin center with a  $\sigma$ -type lone pair of electrons at Sn (NBO 55), and a polarized Sn-CAr bond (NBO 57) with Wiberg Bond Index (WBI) of 0.67. The bonding between the Sn1 and C25 (of IMe<sub>4</sub>) where the WBI = 0.58, indicates a very polarized bonding interaction with electron density of 18.25% on Sn and 81.75% on C (NBO 55). This type of polarization, as well as the sp<sup>1.45</sup> hybridization of the carbon, which interacts with the almost pure p orbital of Sn (sp<sup>19.72</sup>), can be interpreted as carbene coordination to the empty p-orbital of the Sn center. The situation is almost identical to that of the second carbene (Sn1-C32, NBO 56), in terms of bond polarization, hybridization of the interacting orbitals and the WBI (0.58) (Fig. 5). The Sn center of [2]<sup>+</sup> is positively charged by 0.79 el. The aryl moiety of [2] withdraws substantial electron density being negatively charged with -0.48 el., while the Sn center is compensated by the donation from the NHC moieties with the sum of NPA charges of +0.34 and +0.35 el. In addition to the  $[2]^+$  representation as a bis(NHC)-stabilized stannyliumylidene - with dative bonds between the NHC ligands and the Sn atom - the NBO depiction may also hint at the stannyl anion character of the Sn center. In this case, the ionic representation of [2]<sup>+</sup>, where the negative charge is located on the Sn atom to which the two positively charged NHC moieties are bound covalently, should be considered. A detailed discussion of such stannyl-anion-type species in given in the ESI (Fig. S55 and S56†).

The frontier orbitals of  $[2]^+$  are presented in Fig. 6. According to the NBO analysis of the canonical MOs, the HOMO has 52%

**Edge Article Chemical Science** 

Mes 
$$\Sigma q = 0.34$$

Mes  $0.67$ 

Sn:  $q = 0.79$ 

Mes  $0.67$ 

Sn:  $0.58$ 

Mes  $0.58$ 
 $0.58$ 
 $0.58$ 
 $0.58$ 
 $0.58$ 

NPA charges and WBIs in [2]+

non-bonding and 43% bonding character and consists mostly of the non-bonding lone pair of electrons at Sn (51%). Additional important contributions arise from the Sn-Ar and Sn-IMe<sub>4</sub> bonding interactions (Sn-C1 (16%) and Sn-C25 (7%)). The LUMO is 73% anti-bonding and 19% non-bonding with the most significant contributions from  $\pi^*(N3-C25)$  25%,  $\pi^*(N1-C25)$  C32) 21%,  $\sigma^*(Sn1-C1)$  11%,  $\sigma^*(Sn1-C32)$  7%.

Calculations at the PW6B95-D4/def2-QZVPP(CPCM)/r2SCAN-3c level of theory show that the anion-cation interaction in [2] [BArF] is weak, and only 10.4 and 0.0 kcal mol<sup>-1</sup> are required to fully separate the ions in benzene and difluorobenzene, respectively (Scheme 2). The charge transfer in [2][BArF] between  $[2]^+$  cation ( $\Sigma q_{NPA} = + 0.98$  el.) and the  $[BArF]^-$  anion  $(\Sigma q_{\rm NPA} = -0.98 \text{ el.})$  is negligible. The Sn-IMe<sub>4</sub> bonding is also quite labile, with a dissociation free energy for [2][BArF] to form [2a][BArF] and free NHC of 12.0 and 9.3 kcal mol<sup>-1</sup> in benzene and difluorobenzene, respectively (Scheme 2). This bond lability may imply that [2][BArF] may be capable of reacting as a two coordinated stannyliumylidene, upon dissociation of an IMe4 ligand in solution.

#### Hydrosilylation of CO<sub>2</sub> by stannyliumylidene

Inspired by the successful hydrosilylation of CO<sub>2</sub> catalyzed by germyliumylidene (K),34 we attempted to investigate the catalytic potential of [2][BArF] in the same reaction.

We found the optimal conditions for the catalytic reaction of CO<sub>2</sub> and diphenylsilane at room temperature in the presence of 5 mol% [2][BArF] in the  $C_6D_6$ :  $C_6H_5F$  (4:1) solvent mixture. According to the <sup>1</sup>H NMR spectrum, the consumption of the




Fig. 6 Frontier molecular orbitals of [2]<sup>+</sup> (iso = 0.04) at the PBEO/ def2-TZVP//r2SCAN-3c level of theory. The mesityl substituents and the methyl substituents are shown as wireframes for clarity.

Scheme 2 Calculated free energies for anion cation interaction and for NHC dissociation in [2][BArF].

majority (94%) of the Ph<sub>2</sub>SiH<sub>2</sub> was observed within 25 hours, resulting in silvl formate, Ph<sub>2</sub>SiH(OCHO), which is then transformed into silylated methanol (Table 1, Entry 1) (Fig. S35†). The catalyst [2][BArF] exhibited poor solubility in benzene, which led to lower catalytic efficiency (Table, Entry 2). To further investigate the impact of polar solvents on the CO2 hydrosilylation catalyzed by [2][BArF], 1,2-difluorobenzene (C<sub>6</sub>H<sub>4</sub>F<sub>2</sub>) (Table 1, Entry 4) in different ratios with  $C_6D_6$  (Table 1, Entry 5) were selected as reaction solvents. The results indicated that polar solvents did not participate in the hydrosilylation reaction mechanism; their role was limited to enhancing the solubility of [2][BArF]. In comparison to fluorobenzene, the use of THF, a donor solvent, resulted in lower conversion (Table 1, Entry 3). Elevated temperature accelerated the catalytic rate, achieving 64% conversion in 2 hours (Table 1, Entry 6). However, the catalyst decomposed rapidly under these conditions (Fig. S40†).

Scheme 3 Reactions of NHC-stabilized stannyliumylidene ion [2] [BArF] with CO2.

Table 1 Screening of the conditions for the hydrosilylation ( $H_2SiPh_2$ ) of  $CO_2$ 

| Entry | Deviation from the standard conditions $^{a}$    | Conversion <sup>b</sup> (%) |  |
|-------|--|-----------------------------|--|
|       |  |                             |  |
| 1     | None   | 94                          |  |
| 2     | $C_6D_6$   | 78                          |  |
| 3     | THF-d <sub>8</sub>                               | 75                          |  |
| 4     | $C_6D_6/C_6H_4F_2$ (4:1)                         | 97                          |  |
| 5     | $C_6D_6/C_6H_4F_2$ (1:1)                         | 94                          |  |
| 6     | 50 °C  | 64 (2 hours)                |  |
| 7     | 0.5 mol%   | 4                           |  |
| 8     | 1 mol%   | 11                          |  |
| 9     | 2.5 mol%   | 74                          |  |
| 10    | $[2][Al(OC(CF_3)_3)_4]$                          | 70                          |  |
| 11    | $[3][BArF]^c$                                    | 92                          |  |
| 12    | $[4][BArF]^c$                                    | 93                          |  |
| 13    | IMe <sub>4</sub> (10 mol%)                       | 51                          |  |
| 14    | $IMe_4-CO_2$ (10 mol%), $C_6D_6/C_6H_4F_2$ (1:1) | 51                          |  |
| 15    | $\operatorname{Hg}^d$                            | 89                          |  |

 $^a$  Standard reaction conditions: H<sub>2</sub>SiPh<sub>2</sub> (80 μmol), 1,3,5-trimethoxylbenzene (8 μmol, internal standard), and 5 mol% [2][BArF] catalysis in C<sub>6</sub>D<sub>6</sub> (0.32 mL) and C<sub>6</sub>H<sub>5</sub>F (0.08 mL) under 1 bar CO<sub>2</sub> at RT for 25 hours.  $^b$  NMR yield was monitored by  $^1$ H NMR with methoxybenzene as an internal standard.  $^c$  Refer to Scheme 3 for the synthesis of [3][BArF] and [4][BArF].  $^d$  Hg to catalyst ratio 250:1.

Furthermore, to examine the effect the counter anion has on the conversion we performed this reaction under the same conditions with  $[2][Al(OC(CF_3)_3)_4]$ , and a decrease in activity was observed (Table 1, Entry 10). This decrease in catalytic activity could indicate that the very weak coordination strength between  $Al(ORf)_4$  and the Sn atom increases the electrophilicity of the Sn center, which suppresses NHC dissociation from Sn center, compared to BArF $^{-}$ . $^{39}$  Furthermore, the hydrosilylation of  $CO_2$ , catalyzed by the [2][BArF], exhibited minimal perturbation in

Table 2 Screening of the silanes for the hydrosilylation of CO<sub>2</sub>

| Entry | Silanes <sup>a</sup>             | T (°C) | Time (h) | Conversion <sup>b</sup> (%) |
|-------|----------------------------------|--------|----------|-----------------------------|
| 1     | H <sub>2</sub> SiPh <sub>2</sub> | RT     | 25       | 94                          |
| 2     | HSiMe <sub>2</sub> Ph            | RT     | 25       | 47                          |
| 3     | H <sub>3</sub> SiPh              | RT     | $12^c$   | 71                          |
| 4     | HSiEt <sub>3</sub>               | 60     | 25       | 29                          |
| 5     | HSiPh <sub>2</sub>               | 80     | d        | _                           |

 $<sup>^</sup>a$  Standard reaction conditions: H<sub>2</sub>SiPh<sub>2</sub> (80 μmol), 1,3,5-trimethoxylbenzene (8 μmol, internal standard), and 5 mol% [2][BArF] catalysis in C<sub>6</sub>D<sub>6</sub> (0.32 mL) and C<sub>6</sub>H<sub>5</sub>F (0.08 mL) under 1 bar CO<sub>2</sub> at RT for 25 hours.  $^b$  NMR yield was monitored by  $^1$ H NMR with methoxybenzene as an internal standard.  $^c$  Refer to Scheme 3 for the synthesis of [3][BArF] and [4][BArF].  $^d$  Hg to catalyst ratio 250:1.

the presence of excess mercury (Hg), strongly suggesting the nature of the catalytic cycle is homogeneous (Table 1, Entry 15).

In addition to diphenylsilane we tested the catalytic performance of [2][BArF] toward a few other silanes (Table 2). In the case of  $HSiMe_2Ph$  (Table 2, Entry 2), the corresponding silyl formate is formed with a 47% conversion within 25 hours. The catalyst deactivated after consumption of 71% of phenylsilane within 12 hours (Entry 5). The more sterically demanding silanes,  $HSiPh_3$  and  $HSiEt_3$ , were less effective in the hydrosilylation of  $CO_2$  even at elevated temperatures (Table 2, Entries 4 and 5).

[2][BArF] demonstrated superior performance in CO<sub>2</sub> hydrosilylation compared to certain Lewis acids, such as  $[Et_2Al]^+$  and  $[Et_3Si]^+$ . However, it was slightly less effective than  $[(2,6\text{-Mes}_2C_6H_3O)_2Al]^+$ ,  $[2,6\text{-Ph}_2C_6H_3AlEt]^+$ , and  $[(2,6\text{-Dipp}_2C_6H_3)GaEt]^+$  at elevated temperatures, which produced methane, toluene, and diphenylmethane as products. Here [2][BArF] was less effective than some FLP catalytic systems, such as the N,P-heterocyclic germylene/ $[E](C_6F_5)_3$  pair [E](E) and [E](E) produced bis(silyl)acetals or methane. Its performance was comparable to that of germa-acylium ions and germyliumylidene [E](E), which yielded silyl formate, bis(silyl)acetal, and silylated methanol. Here [E](E)

Although [2][BArF] exhibits moderate catalytic ability for  $CO_2$  hydrosilylation, its ability to selectively reduce  $CO_2$  to silyl formate in a one-step process is valuable for studying the mechanism of  $CO_2$  hydrosilylation.

#### Mechanistic studies

To study the mechanism of the hydrosilylation of CO<sub>2</sub> catalyzed by [2][BArF], stoichiometric reactivity studies were conducted. Exposing [2][BArF] to 1 equivalent of  $CO_2$  in toluene:  $C_6H_5F$  (3: 1) at room temperature, afforded the mono-insertion product  $[Mes TerSn(CO_2IMe_4]IMe_4][BArF]$ , [3][BArF], after 4 hours. It bears one CO<sub>2</sub> molecule inserted between the Sn center and one of the IMe4 ligands. It has increased solubility in nonpolar organic solvents such as benzene and toluene compared to [2] [BArF]. The <sup>119</sup>Sn{<sup>1</sup>H} NMR spectrum of [3][BArF] shows a resonance at  $\delta = -10.61$  ppm, which is downfield shifted compared to [2][BArF], due to the Sn being bound to the more electronegative oxygen atom. The 13C(1H) resonance of the corresponding carboxylate (IMe<sub>4</sub>-C(O)OSn) appears at  $\delta$  = 157.48 ppm (THF-d<sub>8</sub>), which is comparable to the reported adducts, *i.e.*,  $IPrCO_2$  (IPr =1,3-bis(2,6diisopropylphenyl)imidazole-2-ylidene) ( $\delta = 152.3$  ppm,  $CD_2Cl_2$ ),  $IMeCO_2$  (IMe = 1,3-dimethylimidazolium) ( $\delta =$ 158 ppm,  $D_2O$ , 40 ItBuCO<sub>2</sub> (ItBu = 1,3-di-tert-butyl-4,5dimethylimidazolin-2-ylidene) ( $\delta = 160$  ppm,  $CD_2Cl_2$ ),  $IMe_4CO_2$  ( $\delta = 155.9$  ppm,  $CD_2Cl_2$ ).<sup>41</sup>

The SC-XRD analysis of [3][BArF] revealed that the central Sn is tricoordinate with an IMe<sub>4</sub> ligand, a terphenyl substituent, and an imidazolium carboxylate (Fig. 7(a)). The Sn1–O1 bond length of 2.231(2) Å is longer than that of the reported Gibson's ( $^{\text{Dipp}}$ nacnac)SnO(iPr), (2.000(5) Å), $^{42}$  tin-carbamate complex (( $^{\text{Dipp}}$ nacnac)SnOC(O)N(iPr<sub>2</sub>)) (2.1346(16) Å) $^{43}$  and shorter than (ArNet2Ge)( $\mu$ - $k^1$ (C)- $k^2$ (O,O')-CO<sub>2</sub>)(SnArNet2) (2.301(3), 2.336(3) Å),




Fig. 7 Molecular structures of [3]<sup>+</sup> (a) and [4]<sup>+</sup> (b) in the solid state. Ellipsoids are set at the 30% probability level; hydrogen atoms and counterions are omitted for clarity. Selected bond lengths [Å] and bond angles [°]: [3]+ C1-Sn1 2.237(3), C25-Sn1 2.283(3), O1-Sn1 2.231(2), O1-C32 1.263(4), C32-O2 1.211(4), C32-C33 1.508(5), C1-Sn1-C25 96.3(1), C1-Sn1-O1 93.38(9), C25-Sn1-O1 86.39(9). [4]<sup>+</sup> C1-Sn1 2.231(2), Sn1-O1 2.168(2), O1-C39 1.270(3), O2-C39 1.223(3), C39-C25 1.450(1), Sn1-O3 2.230(2), O3-C40 1.279(2), C40-O4 1.220(3), C40-C32 1.522(5), C1-Sn1-O1 96.41(7), C1-Sn1-O3 90.89(7), O1-Sn1-O3 90.12(6)

 $(Ar^{NEt2} = [2,6-(Et_2NCH_2)_2C_6H_3]).^{44}$  The C-O bond lengths (C39-O1 = 1.263(4), C39-O2 = 1.211(4) Å) with the C-O moiety bound to the Sn center being the longer of the two, are longer than those in free  $CO_2$  (1.16 Å). This is also in the same range as other imidazolium carboxylates, 45 IPrCO<sub>2</sub> (1.221(4) and 1.225(4) Å) 46  $ItBu_{Me}$ - $CO_2$ - $B(C_6F_5)_3$  ( $ItBu_{Me} = 1,3$ -di-tert-butyl-4,5dimethylimidazolin-2-ylidene) (1.302(3) and 1.211(3) Å).46 The C-C bond length between the IMe<sub>4</sub> and CO<sub>2</sub> is 1.508(5) Å in [3] [BArF], which is close to the reported imidazolium carboxylates, 41,45,46 IPrCO2 (1.511(4) Å), IMe4CO2 (1.521 Å) and IEtMeCO2 (IEt<sub>Me</sub> = 1,3-diethyl-4,5-methyl-imidazolium) (1.535(15) Å). In addition, the carboxylic carbon atoms exhibit a trigonal-planar coordination environment, with the bond angles of 126.5(3)° for O1-C32-O2, 116.3(3)° for C33-C32-O2 and 117.0(3)° for C33-C32-O1.


The addition of excess CO<sub>2</sub> (1 bar) to a solution of [2][BArF] in Et<sub>2</sub>O and pentane at room temperature results in the doubly CO<sub>2</sub> inserted product [MesTerSn(CO<sub>2</sub>IMe<sub>4</sub>)<sub>2</sub>][BArF], [4][BArF], after 21 hours. Yellow crystals suitable for SC-XRD were obtained from an Et<sub>2</sub>O/pentane (1:2) solution stored at −35 °C under a CO<sub>2</sub> atmosphere. The <sup>119</sup>Sn{<sup>1</sup>H} NMR spectrum of [4] [BArF], shows a sharp singlet signal at  $\delta = 55.9$  ppm, which is

shifted downfield compared to [2][BArF] and [3][BArF], due to the Sn center now being bound to two electronegative O atoms. The resonance of the carboxyl carbon atom appears at  $\delta$  = 157.48 ppm in the <sup>13</sup>C{<sup>1</sup>H} NMR spectrum, which is consistent with [3][BArF]. The SC-XRD of [4][BArF] (Fig. 7(b)) shows that the bond lengths of Sn1-O1 and Sn1-O3 are 2.168(2) Å and 2.230(2) Å, respectively. The C-O bond lengths are as follows: C39-O1 (1.270(3) Å), C40–O3 (1.279(2) Å) and C39–O2 (1.223(3) Å), C40– O4 (1.220(3) Å). The lengths of C-C bonds between the IMe<sub>4</sub> carbon and the CO<sub>2</sub> group C39-C25 (1.450(1) Å) and C40-C32 (1.522(5) Å) are different compared to [3][BArF] (1.508(5) Å), but comparable to imidazolium carboxylates. 41,45,46

The isolation of large quantities of pure solid [4][BArF] is not possible as it decomposes to [3][BArF] under reduced pressure via the release of one molecule of CO<sub>2</sub>.

To study the stability of [4][BArF] in solution, <sup>1</sup>H NMRmonitoring experiments were conducted in a toluene-d<sub>8</sub>/  $C_6H_5F$  solution under a  $CO_2$  atmosphere (Fig. S29†). At 100 °C, [4][BArF] slowly converts to [3][BArF], releasing one equivalent of CO<sub>2</sub>. Leaving the sample at ambient temperature for 15 hours resulted in the reformation of [4][BArF].

Calculations at the PW6B95-D4/def2-QZVPP(CPCM)/r2SCAN-3c level of theory show that the likely scenario for the formation of [3][BArF] and [4][BArF] is a stepwise reaction, in which the consecutive dissociation of a IMe4 ligands from the Sn center of [2][BArF] occurs (Scheme 4). The IMe<sub>4</sub> ligands react with carbon dioxide to form the imidazolium carboxylates and reassociate with the Sn complex (Scheme 4). The IMe<sub>4</sub> dissociation from [2]



Scheme 4 Calculated pathway and relative Gibbs energies for the reaction of [2]<sup>+</sup> with CO<sub>2</sub> to give [3]<sup>+</sup> and [4]<sup>+</sup> (black). Gibbs energies for the reactions including the counter anion, relative to [2][BArF] are shown in blue

[BArF] is endergonic by 12.0 kcal mol<sup>-1</sup> and proceeds without a barrier yielding [2a] [BArF] and free IMe<sub>4</sub> (Scheme 4, path (a)). The IMe<sub>4</sub> reacts with a molecule of CO<sub>2</sub> to form the IMe<sub>4</sub>-CO<sub>2</sub> imidazolium carboxylate at 6.4 kcal mol<sup>-1</sup> (Scheme 4, path (b)). The barrier for the reaction of IMe4 with CO2 is only 9.3 kcal mol<sup>-1</sup>. [3][BArF] forms upon barrierless association of  $IMe_4-CO_2$  with [2a][BArF] at -4.8 kcal  $mol^{-1}$  (Scheme 4, path (c)). We also considered a mechanism in which [2][BArF] reacts with CO<sub>2</sub> concertedly, without IMe<sub>4</sub> dissociation, to give [3] [BArF]. However, this scenario requires overcoming a barrier of 29.6 kcal mol<sup>-1</sup> (Fig. S57†). The formation of [4][BArF] proceeds via similar steps from [3][BArF], i.e. dissociation of the IMe<sub>4</sub> from [3][BArF] giving [3a][BArF] + IMe<sub>4</sub> at 5.6 kcal  $\text{mol}^{-1}$ (Scheme 4, path (d)), followed by the formation of the imidazolium carboxylate at 0.1 kcal mol<sup>-1</sup> (Scheme 4, path (e)), and finally association of [3a][BArF] with IMe<sub>4</sub>-CO<sub>2</sub> to give [4][BArF] at -13.2 kcal mol<sup>-1</sup> (Scheme 4, path (f)).


Notably, comparison of the spectroscopic data indicated the formation of [3][BArF] and [4][BArF] during the [2][BArF] catalyzed hydrosilylation of  $\mathrm{CO}_2$  (Fig. S35†). Furthermore, [4][BArF] could be converted to [3][BArF] by adding  $\mathrm{Ph}_2\mathrm{SiH}_2$  under an argon atmosphere, with the simultaneous formation of silyl formate (Fig. S31†). DFT calculations for the proposed mechanism of this transformation were performed (see Fig. S58†). Conversely, [3][BArF] decomposed to unidentified compounds when treated with  $\mathrm{Ph}_2\mathrm{SiH}_2$  at room temperature. Both [3][BArF] and [4][BArF] are efficient catalysts for  $\mathrm{CO}_2$  hydrosilylation (Table 1, Entries 11 and 12), comparable to the performance of [2][BArF]. Thus, we propose [4][BArF] as the resting state of the active cycle. Furthermore, control experiments using  $\mathrm{IMe}_4$  and  $\mathrm{IMe}_4\text{-CO}_2$  as catalysts both show lower conversion (Table 1,

 $\begin{array}{lll} \textbf{Scheme 5} & \textbf{Proposed catalytic cycle of hydrosilylation of carbon} \\ \textbf{dioxide by diphenylsilane in the presence of stannyliumylidene [2]}^+. \end{array}$ 

Entries 13 and 14), suggesting that these compounds are not the catalytically active species.

Based on our reactivity studies and the quantum chemical calculations (Scheme 4, Fig. S57 and S58†) a mechanism for the catalytic hydrosilylation of CO2 using [2][BArF] is proposed (Scheme 5). Under catalytic conditions, the stannyliumylidene [2] reacts with two equivalents of carbon dioxide to form the double insertion product, [4]<sup>+</sup>, via [3]<sup>+</sup>, as described above (Scheme 5, step (a)). Compound [4]<sup>+</sup> reacts with a molecule of diphenylsilane to form intermediate INT A (Scheme 5, step (b)), which dissociates to the ((silylcarbonyl)oxy)imidazolium intermediate (INT\_B) and the stannyl hydride (INT\_C). The former can abstract a hydride from the stannyl hydride, releasing the tin complex [3a]+ and forming a zwitterionic adduct, INT\_D (Scheme 5, step (d)). This type of adduct is well known in NHC organocatalysis, as similar species can form upon the reaction of NHC with an aldehyde, which can later convert to amino enols, known as Breslow intermediates.47 In this case the reverse reaction of the zwitterionic adduct releasing the diphenylsilyl formate product, and the NHC (Scheme 5, step (e)). The NHC reacts with an additional equivalent of CO2 forming the imidazolium carboxylate (Scheme 5, step (f)), which reassociates with tin complex [3a]+ to regenerate the starting compound [4]+ (Scheme 5, step (g)).

In addition to the isolation of the two key intermediates of the catalytic cycle ([3][BArF] and [4][BArF]) and the quantum chemical calculations, we carried out additional experiments (Scheme 6), using deuterated silanes (D<sub>2</sub>SiPh<sub>2</sub>), to support the reaction mechanism proposed. Under standard conditions, the rate of the reactions with H<sub>2</sub>SiPh<sub>2</sub> and D<sub>2</sub>SiPh<sub>2</sub> were compared, giving a kinetic isotope effect (KIE) of 1.11 (Scheme 6, (a)). This value indicates a secondary isotope effect, which is consistent with the rate-determining transition state **TS4** in the proposed mechanism (Fig. S58†), in which the H/D atom is a substituent on carbon from which the NHC moiety is dissociated.<sup>48</sup>



Scheme 6 Mechanistic studies.

Furthermore, CO<sub>2</sub> hydrosilylation catalyzed by [2][BArF] was performed in the presence of 1:1 mixture of D2SiPh2 and HSiMe<sub>2</sub>Ph (Scheme 6, (b)). H/D exchange was observed at the carbon center, supporting the formation of the tin-hydride intermediate INT\_C (Scheme 5) in the proposed catalytic cycle.49 This intermediate can form either via deuterium abstraction from D2SiPh2 or via hydrogen abstraction from HSiMe<sub>2</sub>Ph. Subsequently, INT\_C, whether D/H-substituted, can transfer deuterium or hydrogen to the ((silylcarbonyl)oxy)imidazolium INT\_B, which can also from in the previous step from either D<sub>2</sub>SiPh<sub>2</sub> or HSiMe<sub>2</sub>Ph. The H/D transfer from INT\_C to INT\_B will ultimately produce Ph2SiD(OCDO), Ph2SiD(OCHO), Me<sub>2</sub>PhSi(OCDO), and Me<sub>2</sub>PhSi(OCHO).

#### Conclusions

In summary, we reported on the synthesis of a di-NHCsstabilized stannyliumylidene [2][BArF], which can act as an efficient carbon dioxide hydrosilylation catalyst under ambient conditions. Both experimental and computational studies have been performed to elucidate the reaction mechanism, revealing it to be distinct from those previously proposed for the lighter germyliumylidenes.34 In the latter case, [MesTerGe(IMe4)2]+ (K, Fig. 2) was identified as the active catalytic species, with the Si-H bond activation occurring through coordination with the germanium lone pair (L, Fig. 2), which constitutes the ratedetermining step. In comparison to K, [2][BArF] captures CO<sub>2</sub> via dissociation of NHC, followed by reassociation of the NHC-CO2 moiety to the Sn center, forming the catalytically active species, [4][BArF]. This divergent mechanism is another example of how the lighter and heavier congeners can behave differently and is something we hope to exploit in our further studies in applying tetryliumylidene cations to catalytic transformations.

## Data availability

The data supporting this article have been included as part of the ESI.†

#### Author contributions

D. N. and D. S. carried out the synthetic and reaction studies, and D. N. wrote the original draft. A. K. carried out the computational studies. J. A. K. conducted the crystallographic studies. A. K., J. A. K, H. X., and D. S. reviewed and edited the draft. S. I. managed the project.

#### Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

D. N. gratefully acknowledges financial support from the China Scholarship Council (CSC). We thank Dr Franziska Hanusch and Fiona Kiefer for SC-XRD measurements. We appreciate Ivan Antsiburov (Prof. Roland A. Fischer) and Tobias Weng for the Measurement of LIFDI-MS. The authors gratefully acknowledge the computational and data resources provided by the Leibniz Supercomputing Centre.

#### Notes and references

- 1 J. Chen, M. McGraw and E. Y. Chen, ChemSusChem, 2019, 12, 4543-4569.
- 2 S. Fujimori and S. Inoue, Eur. J. Inorg. Chem., 2020, 2020, 3131-3142.
- 3 J. A. Kelly, F. J. Kiefer, A. Kostenko and S. Inoue, Adv. Inorg. Chem., 2023, 82, 157-187.
- 4 R. Akhtar, K. Gaurav and S. Khan, Chem. Soc. Rev., 2024, 53, 6150-6243.
- 5 M. He, C. Hu, R. Wei, X. F. Wang and L. L. Liu, Chem. Soc. Rev., 2024, 53, 3896-3951.
- 6 S. Yadav, S. Saha and S. S. Sen, ChemCatChem, 2016, 8, 486-
- 7 C. Shan, S. Yao and M. Driess, Chem. Soc. Rev., 2020, 49, 6733-6754.
- 8 D. Sarkar, L. Groll, D. Munz, F. Hanusch and S. Inoue, ChemCatChem, 2022, 14, e202201048.
- 9 L. Groll, J. A. Kelly and S. Inoue, Chem.-Asian J., 2023, 19, e202300941.
- 10 S. N. Riduan, Y. Zhang and J. Y. Ying, Angew. Chem., Int. Ed., 2009, 48, 3322-3325.
- 11 F. Huang, G. Lu, L. Zhao, H. Li and Z.-X. Wang, J. Am. Chem. Soc., 2010, 132, 12388-12396.
- 12 Q. Zhou and Y. Li, J. Am. Chem. Soc., 2015, 137, 10182–10189.
- 13 A. Schafer, W. Saak, D. Haase and T. Muller, Angew. Chem., Int. Ed., 2012, 51, 2981-2984.
- 14 M. Khandelwal and R. J. Wehmschulte, Angew. Chem., Int. Ed., 2012, 51, 7323-7326.
- 15 M. Saleh, D. R. Powell and R. J. Wehmschulte, Organometallics, 2017, 36, 4810-4815.
- 16 R. J. Wehmschulte, M. Saleh and D. R. Powell, Organometallics, 2013, 32, 6812-6819.
- 17 D. Mukherjee, D. F. Sauer, A. Zanardi and J. Okuda, Chem.-Eur. J., 2016, 22, 7730-7733.
- 18 A. Berkefeld, W. E. Piers and M. Parvez, J. Am. Chem. Soc., 2010, 132, 10660-10661.
- 19 M. Rauch and G. Parkin, J. Am. Chem. Soc., 2017, 139, 18162-
- 20 M. Rauch, Z. Strater and G. Parkin, J. Am. Chem. Soc., 2019, 141, 17754-17762.
- 21 A. Caise, J. Hicks, M. Angeles Fuentes, J. M. Goicoechea and S. Aldridge, Chem.-Eur. J., 2021, 27, 2138-2148.
- 22 N. Del Rio, M. Lopez-Reyes, A. Baceiredo, N. Saffon-Merceron, D. Lutters, T. Muller and T. Kato, Angew. Chem., Int. Ed., 2017, 56, 1365-1370.
- 23 V. S. Swamy, S. Pal, S. Khan and S. S. Sen, Dalton Trans., 2015, 44, 12903-12923.
- 24 S. Stigler, S. Fujimori, A. Kostenko and S. Inoue, Chem. Sci., 2024, 15, 4275-4291.
- 25 S. L. Powley and S. Inoue, Chem. Rec., 2019, 19, 2179-2188.

- 26 P. Jutzi, A. Mix, B. Rummel, W. W. Schoeller, B. Neumann and H.-G. Stammler, *Science*, 2004, **305**, 849–851.
- 27 P. Jutzi, F. Kohl, P. Hofmann, C. Krüger and Y. H. Tsay, *Chem. Ber.*, 1980, **113**, 757–769.
- 28 P. Jutzi, R. Dickbreder and H. Nöth, Chem. Ber., 2006, 122, 865–870.
- 29 S. U. Ahmad, T. Szilvasi and S. Inoue, *Chem. Commun.*, 2014, 50, 12619–12622.
- 30 S. U. Ahmad, T. Szilvasi, E. Irran and S. Inoue, *J. Am. Chem. Soc.*, 2015, **137**, 5828–5836.
- 31 R. Nougue, S. Takahashi, A. Dajnak, E. Maerten, A. Baceiredo, N. Saffon-Merceron, V. Branchadell and T. Kato, *Chem.–Eur. J.*, 2022, **28**, e202202037.
- 32 S. Stigler, M. Park, A. Porzelt, A. Kostenko, D. Henschel and S. Inoue, *Organometallics*, 2022, **41**, 2088–2094.
- 33 D. Sarkar, C. Weetman, S. Dutta, E. Schubert, C. Jandl, D. Koley and S. Inoue, *J. Am. Chem. Soc.*, 2020, **142**, 15403– 15411.
- 34 D. Sarkar, S. Dutta, C. Weetman, E. Schubert, D. Koley and S. Inoue, *Chem.-Eur. I.*, 2021, 27, 13072–13078.
- 35 C. P. Sindlinger, F. S. Aicher and L. Wesemann, *Inorg. Chem.*, 2017, **56**, 548–560.
- 36 N. Kuhn, T. Kratz, D. Bläser and R. Boese, *Chem. Ber.*, 1995, 128, 245–250.
- 37 T. G. Kocsor, D. Matioszek, G. Nemes, A. Castel, J. Escudie, P. M. Petrar, N. Saffon and I. Haiduc, *Inorg. Chem.*, 2012, 51, 7782–7787.

- 38 T. Ochiai, D. Franz and S. Inoue, *Chem. Soc. Rev.*, 2016, **45**, 6327–6344.
- 39 I. M. Riddlestone, A. Kraft, J. Schaefer and I. Krossing, *Angew. Chem., Int. Ed.*, 2018, **57**, 13982–14024.
- 40 J. D. Holbrey, W. M. Reichert, I. Tkatchenko, E. Bouajila, O. Walter, I. Tommasi and R. D. Rogers, *Chem. Commun.*, 2003, 28–29.
- 41 E. Theuergarten, T. Bannenberg, M. D. Walter, D. Holschumacher, M. Freytag, C. G. Daniliuc, P. G. Jones and M. Tamm, *Dalton Trans.*, 2014, 43, 1651–1662.
- 42 A. P. Dove, V. C. Gibson, E. L. Marshall, A. J. P. White and D. J. Williams, *Chem. Commun.*, 2001, 283–284.
- 43 L. A. M. Harris, M. P. Coles and J. R. Fulton, *Inorg. Chim. Acta*, 2011, **369**, 97–102.
- 44 A. Caise, L. P. Griffin, C. McManus, A. Heilmann and S. Aldridge, *Angew. Chem., Int. Ed.*, 2022, **61**, e202117496.
- 45 B. R. Van Ausdall, J. L. Glass, K. M. Wiggins, A. M. Aarif and J. Louie, *J. Org. Chem.*, 2009, 74, 7935–7942.
- 46 H. A. Duong, T. N. Tekavec, A. M. Arif and J. Louie, *Chem. Commun.*, 2004, 112–113.
- 47 M. Pareek, Y. Reddi and R. B. Sunoj, *Chem. Sci.*, 2021, **12**, 7973–7992.
- 48 D. Bai, F. Wu, L. Chang, M. Wang, H. Wu and J. Chang, *Angew. Chem., Int. Ed.*, 2022, **61**, e202114918.
- 49 Z. Cheng, M. Li, X. Y. Zhang, Y. Sun, Q. L. Yu, X. H. Zhang and Z. Lu, *Angew. Chem.*, 2023, **135**, e202215029.