



Cite this: *Chem. Sci.*, 2025, 16, 6317

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 3rd December 2024
Accepted 3rd March 2025

DOI: 10.1039/d4sc07923j
rsc.li/chemical-science

Facile, general allylation of unactivated alkyl halides *via* electrochemically enabled radical-polar crossover†


Haifeng Chen and Magnus Rueping *

Electrochemically driven carbon–carbon formation is receiving considerable interest in organic synthesis. In this study, we present an electrochemically driven method for the formation of C(sp³)-C(sp³) bonds using readily available allylic carbonates, as well as primary, secondary, and tertiary alkyl bromides as electrophiles. This approach offers a highly selective route for synthesizing a broad range of allylic products with excellent functional group tolerance, all without the need for transition metal catalysts. Remarkably, this method also enables the smooth late-stage functionalization of various natural product- and drug-derived substrates, yielding the corresponding complex allylalkanes.

Introduction

The development of new methodologies for constructing C(sp³)-C(sp³) bonds remains a significant challenge in the field of C-C coupling chemistry.^{1–3} Transition-metal-catalyzed cross-coupling has emerged as a powerful approach for rapid construction and increasing molecular complexity.^{4–8} However, traditional cross-coupling reactions often require preformed organometallic reagents, which limits their application and substrate scope.⁹ To address this, reductive cross-coupling using electrophiles as coupling partners has proven to be an efficient strategy,^{10–13} significantly enhancing molecular complexity by circumventing the need for activated substrates.^{14–17} Recently, the metallaphotocatalyzed^{18–21} and metallaelectrocatalyzed^{22–28} reactions have offered alternative pathways for forming C(sp³)-C(sp³) bonds under mild conditions. Despite these advances,^{29–33} constructing C(sp³)-C(sp³) bonds remains challenging due to intrinsic obstacles (Scheme 1a). For instance, these methods often rely on expensive metal-ligand catalysts or stoichiometric metal reductants.^{34,35} Therefore, there is a strong demand for developing new, efficient, mild, metal-ligand-catalyst-free, and environmentally friendly strategies for C(sp³)-C(sp³) bond formation.

Electrochemistry has emerged as a versatile and powerful platform for sustainable synthesis, leveraging finely-tuned electron-transfer processes and the use of electrons as traceless redox reagents.^{36–42} In this context, electroreduction has shown great promise, enabling direct interactions between substrates and electrode surfaces to generate alkyl radicals




*Broad Scopes Transition-metal-ligand free Sterically hindered C(sp³)-C(sp³)
Readily available Mild conditions Complex structure functionalization*

Scheme 1 The development of electrocatalyzed directly formation of the C(sp³) radical.

KAUST Catalysis Center (KCC), King Abdullah University of Science and Technology (KAUST), Thuwal, 23955-6900, Saudi Arabia. E-mail: magnus.rueping@kaust.edu.sa
† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d4sc07923j>



Table 1 Optimization of electrochemically driven alkyl-allylation^a

Optimization of reaction

Entry	Variables	Yield ^b [%]	Entry	Variables	Yield ^b [%]
1	Standard	65 ^c	10	DMF as solvent	50
2	RVC(+)//Ni foam(-) w/TMEDA	8	11	MeCN as solvent	17
3	RVC(+)//RVC(-) w/TMEDA	12	12	3 : 1	51
4	(+)Zn//(-)Ni foam	25	13	2 : 1	40
5	(+)Fe//(-)RVC	51	14	1 : 1	36
6	(+)Zn//(-)RVC	27	15	TBACl as electrolyte	23
7	1 mA	44	16	TBAI as electrolyte	45
8	2 mA	63	17	w/o current	ND
9	6 mA	42	18	w/o TBAB	32

^a Reactions were performed with 0.2 mmol of **1a**, 0.4 mmol of **2a**, 0.2 M of TBAB, 4 mL of DMA in an undivided cell at R. T. and $i = 4$ mA, 5.2 F mol⁻¹.

^b GC-FID yield using dodecane as internal standard. ^c Isolated yield.




Fig. 1 Allylic carbonate scope. Reactions were performed with 0.2 mmol of **1**, 0.4 mmol of **2**, 0.2 M of TBAB, and 4 mL of DMA in an undivided cell at R. T. and $i = 4$ mA, 5.2 F mol⁻¹.

through the cathodic reduction of alkyl halides.^{43–47} Elegant studies have explored this concept.^{48–51} Alkyl halides play a crucial role in organic chemistry due to their diverse reactivity and ease of synthesis.⁵² As a result, significant progress has been made in the electroreduction of alkyl halides, including applications such as the Giese reaction, cross-electrophile

coupling (e-XEC), deuteration, borylation, and bifunctionalization of alkenes, among others.^{53–57}

Allylic moieties are essential substructures in organic molecules and versatile functional groups due to their convertible C–C double bonds and allylic single bonds.⁵⁸ The allylation of various molecules *via* the construction of a C–C bond has been thoroughly investigated.^{3,59} To date, transition-metal catalyzed electrophile allylation of alkyl halides is one of the most important protocols for delivering allylated-alkane compounds (Scheme 1b),^{60–65} In comparison, the electrochemical allylation of alkyl halides with allylcarbonates, which benefits from transition-metal-free and sustainable conditions, has been relatively underexplored. Thus, direct electrochemical allylation of unactivated alkyl halides presents a practical and attractive alternative for the synthesis of allylated alkane compounds.

In this regard, we present a novel and versatile electroreduction protocol for the formation of allylated alkanes (Scheme 1c). This reaction offers a straightforward and efficient route to allylated products under mild conditions. Moreover, the successful late-stage functionalization of natural products highlights the potential of our methodology. This strategy not only expands the toolkit for C(sp³)-C(sp³) bond formation but also paves the way for future developments in this area.

Results and discussion

We initiated the exploration with methyl 2-(4-(*tert*-butyl) phenyl) allyl ethyl carbonate (**1a**) and 2-bromo-2-methylpropane (**2a**) as the coupling partners. As outlined in Table 1, a combination of (+) iron//(-) Ni foam as electrodes, TBAB (tetrabutylammonium bromide) as an electrolyte, and DMA (dimethylacetamide) as solvent with a 4 mA current in an undivided cell delivered the






Fig. 2 Alkyl-Br substrate scope. Reactions were performed with 0.2 mmol of **1**, 0.4 mmol of **2**, 0.2 M of TBAB, and 4 mL of DMA in an undivided cell at R. T. and constant current $i = 4$ mA, 5.2 F mol^{-1} .^a Detected by ¹H NMR.

desired product **3a** in 65% isolated yield (entry 1). Altering the anode or cathode significantly decreased the reaction efficiency (entries 2–6). Adjusting the current also succeeded in this alkyl-allylation process, albeit with lower efficiency (entries 7–9). Conducting the reaction in alternative solvents, such as DMF or MeCN, did not improve the reaction outcomes (entries 10 & 11). Screening the substrates ratio also led to lower yields (entries 12–14). Applying other electrolytes did not improve the reaction efficiency (entries 15 & 16).

Control experiments revealed that the current and electrolyte were essential for obtaining high reaction results (entries 17 & 18). In order if the iron ions formed at the anode affect the reaction outcome,⁶⁶ we added different amounts of FeBr₂ and FeBr₃ as well as additional FeBr₂ in combination with different electrodes. However, no reaction enhancement was observed (Fig. S1 & S2[†]).

With the optimized condition in hand, we moved attention to evaluating the scope of this protocol. As shown in Fig. 1, the






Fig. 3 Modification of structurally complex molecules. Reactions were performed with 0.2 mmol of **1**, 0.4 mmol of **2**, 0.2 M of TBAB, and 4 mL of DMA in an undivided cell at R. T. and constant current $i = 4$ mA, 5.2 F mol⁻¹.

installation of electro-donating (**3b–f**), electron-neutral (**3g & h**) and electron-withdrawing (**3i–m**) groups onto the 2-phenylallyl carbonate showed high coupling efficiency and excellent functional group compatibility. Hetero-quinoline substituted substrate also afforded the corresponding product in acceptable yield (**3n**).

Next, we investigated the scope of the alkyl halides (Fig. 2). To our delight, a series of primary, secondary and tertiary alkyl bromides, in spite of different electronic properties, all could be engaged in the reaction efficiently. For the primary alkyl

bromide, alkyl bromides tethered with phenyl and methoxyl substituted phenyl, ether, ester, cyano, benzoate, acetal, trifluoromethyl, *N*-boc-piperidine and phthalimidyl groups could all be transferred to the cascade products in moderate to excellent yields (**4a–k**). The use of secondary alkyl bromides with acyclic, cyclic and heteroatoms gave good yields (**5a–5l**). Unexpectedly, isomerizations (**5d & 5i**) were detected when bromocyclohexane and 4-bromo-*N*-boc-piperidine were used. Bicyclic and spiro compounds reacted efficiently and delivered the corresponding products (**5j–l**) in good yields. In addition, the






Fig. 4 Mechanistic studies. ^aGC-FID yield using dodecane as internal standard. ^bIsolated yield. (a) Radical scavengers; (b) fragmentation cyclopropyl substrates; (c) radical cyclization experiment; (d) competition reaction; (e) reaction monitor; (f) CV measurement; (g) proposed mechanism.

linear (6a–h) and cyclic tertiary alkyl bromides (6i–j) also reacted efficiently in this system. Significantly, the pentafluoro-substituted substrate could furnish the desired product (6d) in 40% yield. Next, our focus shifted to further demonstrating the practicability and synthetic utility of this protocol, we

explored modifications on structurally complex molecules. As displayed in Fig. 3, various primary, secondary, and tertiary natural or pharmaceutical alkyl bromides derived from *l*-menthol (7a), naproxen (7b), adapalene (7c), indometacin (7d), gemfibrozil (7e), flubiprofen (7f), dehydroabietic acid (7g),



acetobromo-*a*-*D*-glucose (**7h**), epiandrosterone (**7i**), lithocholic acid (**7j**), tetrahydrolinalool (**7k**), probenecid (**7l**), γ -terpineol (**7m**) and sclareolide (**7n**) were effectively converted into the corresponding alkyl radical *via* this protocol, and subsequently delivered a variety of structurally complex allylated-alkane derivatives in moderate to good yields.

To further elucidate the possible mechanism of this electrochemically driven formation of allylated-alkanes, a range of control experiments were performed. Firstly, the use of stoichiometric amounts of radical scavengers (1, 1-diphenylethylene and TEMPO) under the standard conditions was applied. Both desired products and alkyl radical captured products could be observed (Fig. 4a). In principle, the allylic carbonate might also be reduced by the cathode to release the allyl radical. However, we did not observe the formation. Alkyl radical clock experiments were performed by using the cyclopropyl tethered (bromomethyl) cyclopropane (**2d**) and ethyl 2-bromo-2-cyclopropylacetate (**2e**). Here, we obtained the ring-opening product **14** in 55%, although the secondary alkyl substrate only furnished the desired product **15** in trace yield and by-product **15a** was formed in 45%, which indicates that the secondary alkyl radical is involved in the reaction (Fig. 4b). Thus, radical cyclization experiments were also performed. The cyclization products **16** and **17** were obtained in 67% and 31%, respectively (Fig. 4c). These observations suggest the involvement of alkyl radicals in this protocol. To further investigate, a competition reaction was performed (Fig. 4d), yielding products **3a**, **4a**, and **5a** with yields of 59%, 8%, and 17%, respectively. These results demonstrated that the alkyl reactivity followed the sequence: tertiary > secondary > primary. Additionally, reaction monitoring indicated that the allylation reaction was completed within 7 hours (Fig. 4e). To explore the electrochemical behavior of the reactants, cyclic voltammetry (CV) measurements were conducted. As shown in Fig. 4f, the reduction peaks for 3-phenylpropyl bromide, 2-bromopropane, and *tert*-butyl bromide were observed at -2.70 V, -2.66 V, and -2.52 V *versus* the saturated calomel electrode (SCE), and the onset values were -2.4 V, -2.3 V and -2.2 V, respectively. For 2-(4-(*tert*-butyl)phenyl) allyl ethyl carbonate, two onset reduction peaks were recorded at -2.74 V and -2.40 V *versus* SCE. These findings indicate that alkyl halides can be reduced to generate the corresponding alkyl radicals under the conditions of this protocol.

Mechanism

Based on the above analysis results and literature report,^{53,65} we propose a plausible mechanism for the new electrochemical C(sp³)-C(sp³) bond formation (Fig. 4g). Initially, the allylic carbonate reacts with the cathode-activated alkyl radicals, forming a stable tertiary benzyl radical. This radical is subsequently reduced at the cathode to generate a carbanion. The carbanion intermediate can then undergo β -elimination, leading to the formation of the desired products.

Conclusions

In conclusion, we have successfully developed an efficient electrochemically-driven radical polar crossover method for the

formation of allylated alkanes using allylic carbonate and unactivated alkyl bromides. This protocol operates under mild conditions, enabling the synthesis of a diverse range of synthetically valuable allylated alkanes. Given the commercial availability of the reactants and the properties of allyl and alkyl compounds, we believe that this electroreduction protocol will have broad applicability, expand the library of allylated alkanes, and inspire further exploration in the field of C(sp³)-C(sp³) coupling reactions.

Data availability

The data supporting this article have been included as part of the ESI.†

Author contributions

H. C., and M. R. conceived the idea of this work. H. C. carried out the reaction optimization and substrate scope. H. C. and M. R. co-wrote the manuscript. M. R. directed the entire research.

Conflicts of interest

The authors declare no competing financial interest.

Acknowledgements

This work was financially supported by the King Abdullah University of Science and Technology (KAUST), Saudi Arabia, Office of Sponsored Research. This publication is based upon work supported by King Abdullah University of Science and Technology (KAUST) under Award No. RFS-OPF2023-5517 and URF/1/5033.

Notes and references

- 1 E. J. C. Corey and X.-M. Cheng, *The Logic Of Chemical Synthesis*, Wiley, New York, 1995.
- 2 H. M. Huang, P. Bellotti and F. Glorius, Transition metal-catalysed allylic functionalization reactions involving radicals, *Chem. Soc. Rev.*, 2020, **49**, 6186–6197.
- 3 L. Susse and B. M. Stoltz, Enantioselective Formation of Quaternary Centers by Allylic Alkylation with First-Row Transition-Metal Catalysts, *Chem. Rev.*, 2021, **121**, 4084–4099.
- 4 R. Jana, T. P. Pathak and M. S. Sigman, Advances in transition metal (Pd, Ni, Fe)-catalyzed cross-coupling reactions using alkyl-organometallics as reaction partners, *Chem. Rev.*, 2011, **111**, 1417–1492.
- 5 L. J. Cheng and N. P. Mankad, C–C and C–X coupling reactions of unactivated alkyl electrophiles using copper catalysis, *Chem. Soc. Rev.*, 2020, **49**, 8036–8064.
- 6 M. J. Goldfogel, L. Huang and D. J. Weix, in *Nickel Catalysis In Organic Synthesis: Methods And Reactions*, ed. S. Ogoshi, Wiley-VCH, Weinheim, 2020, pp. 183–222, DOI: [10.1002/9783527813827.ch9](https://doi.org/10.1002/9783527813827.ch9).



- 7 W. Xue, X. Jia, X. Wang, X. Tao, Z. Yin and H. Gong, Nickel-catalyzed formation of quaternary carbon centers using tertiary alkyl electrophiles, *Chem. Soc. Rev.*, 2021, **50**, 4162–4184.
- 8 Y. Wei, L. Q. H. Lin, B. C. Lee and M. J. Koh, Recent Advances in First-Row Transition Metal-Catalyzed Reductive Coupling Reactions for π -Bond Functionalization and C-Glycosylation, *Acc. Chem. Res.*, 2023, **56**, 3292–3312.
- 9 W. Shi, C. Liu and A. Lei, Transition-metal catalyzed oxidative cross-coupling reactions to form C–C bonds involving organometallic reagents as nucleophiles, *Chem. Soc. Rev.*, 2011, **40**, 2761–2776.
- 10 X. Wang, Y. Dai and H. Gong, Nickel-Catalyzed Reductive Couplings, *Top. Curr. Chem.*, 2016, **374**, 43.
- 11 K. E. Poremba, S. E. Dibrell and S. E. Reisman, Nickel-Catalyzed Enantioselective Reductive Cross-Coupling Reactions, *ACS Catal.*, 2020, **10**, 8237–8246.
- 12 L. M. Chen and S. E. Reisman, Enantioselective C(sp²)-C(sp³) Bond Construction by Ni Catalysis, *Acc. Chem. Res.*, 2024, **57**, 751–762.
- 13 Y. Gong, J. Hu, C. Qiu and H. Gong, Insights into Recent Nickel-Catalyzed Reductive and Redox C–C Coupling of Electrophiles, C(sp³)-H Bonds and Alkenes, *Acc. Chem. Res.*, 2024, **57**, 1149–1162.
- 14 H. Chen, X. Jia, Y. Yu, Q. Qian and H. Gong, Nickel-Catalyzed Reductive Allylation of Tertiary Alkyl Halides with Allylic Carbonates, *Angew. Chem., Int. Ed.*, 2017, **56**, 13103–13106.
- 15 Y. Ye, H. Chen, J. L. Sessler and H. Gong, Zn-Mediated Fragmentation of Tertiary Alkyl Oxalates Enabling Formation of Alkylated and Arylated Quaternary Carbon Centers, *J. Am. Chem. Soc.*, 2019, **141**, 820–824.
- 16 C. Dorval, M. Tricoire, J.-M. Begouin, V. Gandon and C. Gosmini, Cobalt-Catalyzed C(sp²)-CN Bond Activation: Cross-Electrophile Coupling for Biaryl Formation and Mechanistic Insight, *ACS Catal.*, 2020, **10**, 12819–12827.
- 17 H. Chen, H. Yue, C. Zhu and M. Rueping, Reactivity in Nickel-Catalyzed Multi-component Sequential Reductive Cross-Coupling Reactions, *Angew. Chem., Int. Ed.*, 2022, **61**, e202204144.
- 18 C. Zhu, H. Yue, L. Chu and M. Rueping, Recent advances in photoredox and nickel dual-catalyzed cascade reactions: pushing the boundaries of complexity, *Chem. Sci.*, 2020, **11**, 4051–4064.
- 19 S. B. Beil, T. Q. Chen, N. E. Intermaggio and D. W. C. MacMillan, Carboxylic Acids as Adaptive Functional Groups in Metallaphotoredox Catalysis, *Acc. Chem. Res.*, 2022, **55**, 3481–3494.
- 20 A. Y. Chan, I. B. Perry, N. B. Bissonnette, B. F. Buksh, G. A. Edwards, L. I. Frye, O. L. Garry, M. N. Lavagnino, B. X. Li, Y. Liang, E. Mao, A. Millet, J. V. Oakley, N. L. Reed, H. A. Sakai, C. P. Seath and D. W. C. MacMillan, Metallaphotoredox: The Merger of Photoredox and Transition Metal Catalysis, *Chem. Rev.*, 2022, **122**, 1485–1542.
- 21 J. Zhang and M. Rueping, Metallaphotoredox catalysis for sp³ C–H functionalizations through single-electron transfer, *Nat. Catal.*, 2024, **7**, 963–976.
- 22 P. Gandeepan, L. H. Finger, T. H. Meyer and L. Ackermann, 3d metalla-electrocatalysis for resource economical syntheses, *Chem. Soc. Rev.*, 2020, **49**, 4254–4272.
- 23 J.-S. Zhong, Y. Yu, Z. Shi and K.-Y. Ye, An electrochemical perspective on the roles of ligands in the merger of transition-metal catalysis and electrochemistry, *Org. Chem. Front.*, 2021, **8**, 1315–1328.
- 24 A. L. Xu Cheng, T.-S. Mei, H.-C. Xu, K. Xu and C. Zeng, Recent Applications of Homogeneous Catalysis in Electrochemical Organic Synthesis, *CCS Chem.*, 2022, **4**, 1120–1152.
- 25 H. Chen, C. Zhu, H. Yue and M. Rueping, Group 14 Elements Hetero-Difunctionalizations via Nickel-Catalyzed Electroreductive Cross-Coupling, *Angew. Chem., Int. Ed.*, 2023, **62**, e202306498.
- 26 C. A. Malapit, M. B. Prater, J. R. Cabrera-Pardo, M. Li, T. D. Pham, T. P. McFadden, S. Blank and S. D. Minter, Advances on the Merger of Electrochemistry and Transition Metal Catalysis for Organic Synthesis, *Chem. Rev.*, 2022, **122**, 3180–3218.
- 27 L. Zou, X. Wang, S. Xiang, W. Zheng and Q. Lu, Paired Oxidative and Reductive Catalysis: Breaking the Potential Barrier of Electrochemical C(sp³)-H Alkenylation, *Angew. Chem., Int. Ed.*, 2023, **62**, e202301026.
- 28 N. Li, R. Sitdikov, A. P. Kale, J. Steverlynck, B. Li and M. Rueping, A review of recent advances in electrochemical and photoelectrochemical late-stage functionalization classified by anodic oxidation, cathodic reduction, and paired electrolysis, *Beilstein J. Org. Chem.*, 2024, **20**, 2500–2566.
- 29 D. Brosamlen and M. Oestreich, Regioselective Hydroalkylation of Vinyl and Allylsilanes as Well as Vinylgermanes under Ni–H Catalysis, *Org. Lett.*, 2023, **25**, 5319–5323.
- 30 A. Hossain, R. L. Anderson, C. S. Zhang, P. J. Chen and G. C. Fu, Nickel-Catalyzed Enantioconvergent and Diastereoselective Allenylation of Alkyl Electrophiles: Simultaneous Control of Central and Axial Chirality, *J. Am. Chem. Soc.*, 2024, **146**(11), 7173–7177.
- 31 R. Chen, N. E. Intermaggio, J. Xie, J. A. Rossi-Ashton, C. A. Gould, R. T. Martin, J. Alcazar and D. W. C. MacMillan, Alcohol-alcohol cross-coupling enabled by S(H)₂ radical sorting, *Science*, 2024, **383**, 1350–1357.
- 32 X. C. Gan, B. Zhang, N. Dao, C. Bi, M. Pokle, L. Kan, M. R. Collins, C. C. Tyrol, P. N. Bolduc, M. Nicastrì, Y. Kawamata, P. S. Baran and R. Shenvi, Carbon quaternization of redox active esters and olefins by decarboxylative coupling, *Science*, 2024, **384**, 113–118.
- 33 P. Li, Z. Zhu, C. Guo, G. Kou, S. Wang, P. Xie, D. Ma, T. Feng, Y. Wang and Y. Qiu, Nickel-electrocatalysed C(sp³)-C(sp³) cross-coupling of unactivated alkyl halides, *Nat. Catal.*, 2024, **7**, 412–421.
- 34 C. L. Sun and Z. J. Shi, Transition-metal-free coupling reactions, *Chem. Rev.*, 2014, **114**, 9219–9280.
- 35 J. Xie, H. Jin and A. S. K. Hashmi, The recent achievements of redox-neutral radical C–C cross-coupling enabled by visible-light, *Chem. Soc. Rev.*, 2017, **46**, 5193–5203.



- 36 A. Jutand, Contribution of electrochemistry to organometallic catalysis, *Chem. Rev.*, 2008, **108**, 2300–2347.
- 37 G. S. Sauer and S. Lin, An Electrocatalytic Approach to the Radical Difunctionalization of Alkenes, *ACS Catal.*, 2018, **8**, 5175–5187.
- 38 T. H. Meyer, L. H. Finger, P. Gandeepan and L. Ackermann, Resource Economy by Metallalectrocatalysis: Merging Electrochemistry and C–H Activation, *Trends Chem.*, 2019, **1**, 63–76.
- 39 C. Zhu, N. W. J. Ang, T. H. Meyer, Y. Qiu and L. Ackermann, Organic Electrochemistry: Molecular Syntheses with Potential, *ACS Cent. Sci.*, 2021, **7**, 415–431.
- 40 Y. Liu, P. Li, Y. Wang and Y. Qiu, Electroreductive Cross-Electrophile Coupling (eXEC) Reactions, *Angew. Chem., Int. Ed.*, 2023, **62**, e202306679.
- 41 C. Zhu, H. Chen, H. Yue and M. Rueping, Electrochemical chemo- and regioselective arylalkylation, dialkylation and hydro(deutero)alkylation of 1,3-enynes, *Nat. Synth.*, 2023, **2**, 1068–1081.
- 42 H. Chen, C. Zhai, C. Zhu and M. Rueping, Allylgermane synthesis *via* facile and general nickela-electrocatalyzed electrophile coupling, *Chem Catal.*, 2025, DOI: [10.1016/j.cheecat.2024.101257](https://doi.org/10.1016/j.cheecat.2024.101257).
- 43 J. B. Sperry and D. L. Wright, The application of cathodic reductions and anodic oxidations in the synthesis of complex molecules, *Chem. Soc. Rev.*, 2006, **35**, 605–621.
- 44 J. Yoshida, K. Kataoka, R. Horcajada and A. Nagaki, Modern strategies in electroorganic synthesis, *Chem. Rev.*, 2008, **108**, 2265–2299.
- 45 M. Yan, Y. Kawamata and P. S. Baran, Synthetic Organic Electrochemical Methods Since 2000: On the Verge of a Renaissance, *Chem. Rev.*, 2017, **117**, 13230–13319.
- 46 S. Mohle, M. Zirbes, E. Rodrigo, T. Gieshoff, A. Wiebe and S. R. Waldvogel, Modern Electrochemical Aspects for the Synthesis of Value-Added Organic Products, *Angew. Chem., Int. Ed.*, 2018, **57**, 6018–6041.
- 47 W. Zhang, W. Guan, J. I. Martinez Alvarado, L. F. T. Novaes and S. Lin, Deep Electroreductive Chemistry: Harnessing Carbon- and Silicon-Based Reactive Intermediates in Organic Synthesis, *ACS Catal.*, 2023, **13**, 8038–8048.
- 48 A. Claraz and G. Masson, Recent Advances in C(sp³)–C(sp³) and C(sp³)–C(sp²) Bond Formation through Cathodic Reactions: Reductive and Convergent Paired Electrolyses, *ACS Org. Inorg. Au*, 2022, **2**, 126–147.
- 49 B. Huang, Z. Sun and G. Sun, Recent progress in cathodic reduction-enabled organic electrosynthesis: Trends, challenges, and opportunities, *eScience*, 2022, **2**, 243–277.
- 50 T. B. Hamby, M. J. LaLama and C. S. Sevov, Controlling Ni redox states by dynamic ligand exchange for electroreductive Csp³–Csp² coupling, *Science*, 2022, **376**, 410–416.
- 51 H. Xiang, J. He, W. Qian, M. Qiu, H. Xu, W. Duan, Y. Ouyang, Y. Wang and C. Zhu, Electroreductively Induced Radicals for Organic Synthesis, *Molecules*, 2023, **28**, 857–879.
- 52 X. Sun and K. Zheng, Electrochemical halogen-atom transfer alkylation *via* alpha-aminoalkyl radical activation of alkyl iodides, *Nat. Commun.*, 2023, **14**, 6825.
- 53 W. Zhang and S. Lin, Electroreductive Carbofunctionalization of Alkenes with Alkyl Bromides *via* a Radical-Polar Crossover Mechanism, *J. Am. Chem. Soc.*, 2020, **142**, 20661–20670.
- 54 B. Wang, P. Peng, W. Ma, Z. Liu, C. Huang, Y. Cao, P. Hu, X. Qi and Q. Lu, Electrochemical Borylation of Alkyl Halides: Fast, Scalable Access to Alkyl Boronic Esters, *J. Am. Chem. Soc.*, 2021, **143**, 12985–12991.
- 55 P. Li, C. Guo, S. Wang, D. Ma, T. Feng, Y. Wang and Y. Qiu, Facile and general electrochemical deuteration of unactivated alkyl halides, *Nat. Commun.*, 2022, **13**, 3774.
- 56 W. Zhang, L. Lu, W. Zhang, Y. Wang, S. D. Ware, J. Mondragon, J. Rein, N. Strotman, D. Lehnerr, K. A. See and S. Lin, Electrochemically driven cross-electrophile coupling of alkyl halides, *Nature*, 2022, **604**, 292–297.
- 57 L. Lu, Y. Wang, W. Zhang, W. Zhang, K. A. See and S. Lin, Three-Component Cross-Electrophile Coupling: Regioselective Electrochemical Dialkylation of Alkenes, *J. Am. Chem. Soc.*, 2023, **145**, 22298–22304.
- 58 J. F. Han, P. Guo, X. G. Zhang, J. B. Liao and K. Y. Ye, Recent advances in cobalt-catalyzed allylic functionalization, *Org. Biomol. Chem.*, 2020, **18**, 7740–7750.
- 59 S. Dutta, T. Bhattacharya, D. B. Werz and D. Maiti, Transition-metal-catalyzed C–H allylation reactions, *Chem*, 2021, **7**, 555–605.
- 60 Y. Dai, F. Wu, Z. Zang, H. You and H. Gong, Ni-catalyzed reductive allylation of unactivated alkyl halides with allylic carbonates, *Chem.–Eur. J.*, 2012, **18**, 808–812.
- 61 X. Qian, A. Auffrant, A. Felouat and C. Gosmini, Cobalt-catalyzed reductive allylation of alkyl halides with allylic acetates or carbonates, *Angew. Chem., Int. Ed.*, 2011, **50**, 10402–10405.
- 62 Y. Yu, H. Chen, Q. Qian, K. Yao and H. Gong, An extension of nickel-catalyzed reductive coupling between tertiary alkyl halides with allylic carbonates, *Tetrahedron*, 2018, **74**, 5651–5658.
- 63 H. Chen, Y. Ye, W. Tong, J. Fang and H. Gong, Formation of allylated quaternary carbon centers *via* C–O/C–O bond fragmentation of oxalates and allyl carbonates, *Chem. Commun.*, 2020, **56**, 454–457.
- 64 Y. Xu, M. Zhang and M. Oestreich, Photochemical, Nickel-Catalyzed C(sp³)–C(sp³) Reductive Cross-Coupling of α -Silylated Alkyl Electrophiles and Allylic Sulfones, *ACS Catal.*, 2022, **12**, 10546–10550.
- 65 T. Constantin, M. Zanini, A. Regni, N. S. Sheikh, F. Julia and D. Leonori, Aminoalkyl radicals as halogen-atom transfer agents for activation of alkyl and aryl halides, *Science*, 2020, **367**, 1021–1026.
- 66 G. S. Kumar, A. Peshkov, A. Brzozowska, P. Nikolaienko, C. Zhu and M. Rueping, Nickel-Catalyzed Chain-Walking Cross-Electrophile Coupling of Alkyl and Aryl Halides and Olefin Hydroarylation Enabled by Electrochemical Reduction, *Angew. Chem., Int. Ed.*, 2020, **59**, 6513–6519.

