Chemical Science

EDGE ARTICLE



Cite this: Chem. Sci., 2025, 16, 11058

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 15th March 2025 Accepted 12th May 2025

DOI: 10.1039/d5sc02048d

rsc.li/chemical-science

Introduction

For many decades inorganic Sb^V complexes have been a quintessential example of a strong Lewis acid that indirectly activates nonpolar hydrocarbon bonds through protonation as a Brønsted superacid (Fig. 1a).1-3 In contrast to this Lewis acid/ superacid reactivity and mechanism, we recently disclosed that $Sb^{V}X_{5}$ (X = trifluoroacetate) can react with alkanes to generate alkyl esters without generating additional products that would be expected from a superacid/protonation reaction mechanism, and our calculations supported the idea that the Sb^V metal center can directly react with nonpolar C-H bonds to generate Sb^V-C bond intermediates through C-H activation (Fig. 1b).^{4,5} However, there was no experimental detection of the Sb^V-C bond intermediates to support the calculated C-H activation/ metalation and Sb^V-C bond reductive oxy-functionalization reaction steps. Therefore, in this work we have used both calculations and experiments to examine the reaction of inorganic Sb^V complexes with arenes that support the proposal that a Sb^v metal center can directly activate and oxidize nonpolar C-H bonds (Fig. 1c).

^bThe Herbert Wertheim UF Scripps Institute for Biomedical Innovation & Technology, University of Florida, Jupiter, Florida 33458, USA. E-mail: rperiana@ufl.edu



Anjaneyulu Koppaka, () ** Shu-Sen Chen, ** Dongdong Yang, ** Artem Marchenko, ** Sanaz Mohammadzadeh Koumleh, ** David J. Michaelis, ** Roy A. Periana*** and Daniel H. Ess **

Inorganic Sb^V complexes are one of the most well-known Lewis acids that typically indirectly activate nonpolar hydrocarbon bonds by acting as Brønsted superacids. Here we have used a combination of quantum-chemical calculations and experiments to demonstrate that inorganic Sb^V complexes can directly activate aromatic sp² C–H bonds and induce oxidative functionalization to form aryl esters. C–H activation was first demonstrated by reaction of Sb^V(TFA)₅ (TFA = trifluoroacetate) with toluene at moderate temperatures that resulted in Sb^V–C bond intermediates (TFA)₄Sb^V(*para*-tolyl) and (TFA)₃Sb^V(*para*-tolyl)₂. Calculations predicted and then experiments confirmed that at a higher temperature reductive functionalization occurs to generate oxidized aryl ester products. Theory and experiment indicate that due to a Curtin–Hammet type equilibrium the initial *para* C–H activation regioselectivity changes to mainly *ortho* and *meta* selectivity for reductive oxy-functionalization.

In addition to being a vehicle for examining Sb^V metal mediated C–H functionalization, direct and selective oxidative functionalization of unactivated aromatic sp² C–H bonds to ester (or alcohol) functional groups is itself an important and surprisingly difficult transformation.⁶ There are only a few



Fig. 1 (a) Outline of the general mechanism for alkane protonation by inorganic Sb^V complexes in strong Brønsted acid. The protonated hydrocarbon then generates a carbocation intermediate. (b) Outline of C–H activation and M–R oxy-functionalization reaction steps (R = alkyl or aryl). (c) Brief outline of experiments and calculations reported in this work demonstrating that Sb^VX₅ is capable of chemoselective and regioselective C–H activation and oxy-functionalization reaction steps. X = trifluoroacetate.



View Article Online

View Journal | View Issue

^aDepartment of Chemistry and Biochemistry, Brigham Young University, Provo, Utah 84604, USA. E-mail: Anjaneyulu.Koppaka@byu.edu; dhe@byu.edu

[†] Electronic supplementary information (ESI) available. See DOI: https://doi.org/10.1039/d5sc02048d

examples where this transformation is promoted by a homogeneous transition metal system. The most prominent examples are Pd catalyzed⁷⁻⁹ and Cu induced arene oxidative functionalization to aryl esters.¹⁰⁻¹⁷ Most germane to the work here, Gunnoe reported that Cu acetates can oxidatively functionalize benzene and toluene, and proposed a metal mediated C–H activation and reductive functionalization mechanism.¹⁸ As outlined in Fig. 1b, our goal was to determine if p-block maingroup Sb^V complexes can promote arene C–H functionalization through metal mediated C–H activation/metalation and functionalization mechanistic steps. While arene C–H activation and reductive functionalization are established reaction steps for transition metals there is much less evidence for these reaction steps with high-oxidation state p-block main-group metals.

Here we report experimental and theoretical studies that show that at moderate temperatures Sb^VX₅ in trifluoroacetic acid (TFAH) solvent induces innersphere metal mediated chemoselective and regioselective para C-H activation of toluene to give $[Sb^{V}]$ -tolyl ($[Sb^{V}] = Sb^{V}$ with four anion ligands of either TFA or aryl) intermediates (see Fig. 1c). Consistent with theoretical predictions, subsequent experiments at higher temperatures generated aryl esters with selectivity switched to orthosubstituted and meta-substituted esters rather than the expected para-substituted esters. Calculations indicate that para to ortho/meta selectivity change occurs because at a higher temperature there is a Curtin-Hammet type equilibrium between bis-tolyl and mono-tolyl Sb^V intermediates that occurs before rate limiting oxy-functionalization from the more electron deficient mono-tolyl Sb^v intermediate that is selective for ortho/meta functionalization. Independent synthesis of (TFA)₄-Sb^V(para-tolyl) showed the same selectivity in TFAH. Overall, the combination of calculations and experiments demonstrate that Sb^VX₅ is capable of both C-H activation and Sb^V-C bond oxyfunctionalization, and arenes can be directly functionalized to aryl ester products.

Results and discussion

Experimental evaluation of C-H activation and computational prediction and experimental realization of oxy-functionalization

For sixth-row p-block main-group metals, there are a few previous reports of Tl^{III} complexes reacting with arenes to generate Tl^{III}-aryl intermediates,¹⁹⁻²³ but these reactions were often proposed to occur through either electron-transfer type reaction steps or electrophilic aromatic substitution with a carbocation/Wheland type intermediate. However, our calculations showed that benzene and mono-substituted arenes likely undergo metalation with Tl^{III} metal centers through a C-H activation mechanism rather than a carbocation or electron transfer mechanism.²⁴ Added nucleophiles to the Tl^{III}-aryl intermediate were generally necessary to result in functionalization. There are also previous reports of Pb^{IV} complexes reacting with arenes. Partch reported that inorganic Pb^{IV} complexes can oxidize benzene and toluene.²⁵ For toluene it was reported that the benzylic position was oxidized potentially

suggesting a radical mechanism, although no byproducts indicative of a radical reaction were observed. Sternhell characterized Pb^{IV}–aryl intermediates from reaction of Pb^{IV} complexes with benzene derivatives.^{26,27}

While the Tl^{III} and Pb^{IV} examples outlined above indicate that oxidative functionalization of unactivated arenes is possible with high oxidation state sixth-row p-block metals there is little precedence for a similar transformation with fifthrow metals, specifically with a Sb^V metal center. Generally, as outlined in the Introduction, Sb^V complexes operate as a powerful Lewis acid to generate super Brønsted acids and the innersphere of the metal does not directly participate in bond making and breaking reaction steps with hydrocarbons. However, we have recently demonstrated that inorganic Sb^V complexes in both TFAH and sulfuric acid solvents can induce oxidative functionalization of light alkanes and based on calculations we proposed that the Sb metal center participates in C-H bond activation and functionalization reaction steps.4,5 But again, there was no experimental evidence to support the calculated C-H activation and Sb^V-C bond reductive oxyfunctionalization reaction steps.

As outlined in the Introduction, our goal was to determine if oxidative functionalization of aromatic C-H bonds can be promoted with Sb^V complexes using C-H activation and functionalization mechanistic steps. We targeted Sb^V(TFA)₅ because we previously used it to functionalize light alkane C-H bonds and it can be straightforwardly generated from reaction of [Sb^V(OMe)₅]₂ with trifluoroacetic anhydride (TFAA) and trifluoroacetic acid (TFAH).4 We examined the reaction of Sb^V(TFA)₅ with toluene because this unactivated aromatic compound provides the ability to evaluate regioselectivity/ positional C-H functionalization selectivity as well as sp² (aromatic) versus sp³ (benzylic) C-H functionalization selectivity. The aromatic versus benzylic positional selectivity provides a probe to compare non-radical versus radical type mechanisms where functionalization at the benzylic position is expected in a radical mechanism.18

Our initial experiments focused on the synthesis of Sb^V(TFA)₅ and identifying reaction conditions where we could monitor toluene C-H activation. Under an argon atmosphere 0.076 mmol of $[Sb^{V}(OMe)_{5}]_{2}$ was added to 0.2 mL of trifluoroacetic anhydride (TFAA). The reaction mixture was stirred at room temperature for 5 minutes after which 0.8 mL of TFAH was added and stirred at room temperature for another 15 min. This resulted in a solution of Sb^V(TFA)₅ indicated from the formation of 5 equivalents of methyl trifluoroacetate (MeTFA) generated from the reaction of [Sb^V(OMe)₅]₂ with TFAA. Toluene (~0.5 equivalents) was then added to the Sb(TFA)₅/TFAA/TFAH solution and heated at 60 °C for 1 h. Fig. 2, bottom spectrum, shows the ¹H NMR of the crude reaction mixture. To our surprise, this relatively mild reaction condition resulted in conversion (~50%) of toluene to the para C-H activation/ metalation product, (TFA)₄Sb^V(para-tolyl). Trace levels of Sbtolyl products from C-H activation of the ortho or meta toluene C-H bonds may be present but could not be resolved. We confirmed the signals of the mono-tolyl complex by addition of independently synthesized para-toluenestibonic acid





Fig. 2 Overlay of the ¹H NMR spectra for the crude reaction mixtures from reaction of $Sb^{V}(TFA)_{5}$ with toluene in TFAH : TFAA (4 : 1) taken after 1 hour at 60 °C (bottom spectrum) and after 17 hours at 70 °C (top spectrum). The initial product is the mono-tolyl, (TFA)₄Sb^V(*para*-tolyl), labeled as * in the bottom spectrum. The bis-tolyl product, (TFA)₃Sb^V(*para*-tolyl)₂, is labeled as \$ in the top spectrum.

 $((OH)_2OSb^V(para-tolyl))$ to TFAH solvent which quantitively generated (TFA)_4Sb^V(para-tolyl).²⁸ The ESI† shows time dependent ¹H NMR of the crude reaction mixture from room temperature up to 80 °C and from 10 minutes to 7 hours. At room temperature the same mono-tolyl complex was formed but the reaction is much slower.

Heating for longer periods or increasing the temperature to 70 °C for 17 hours (upper spectrum) led to loss of the mono-tolyl product and conversion to what we propose as the bis-tolyl intermediate, (TFA)₃Sb^V(para-tolyl)₂ (Fig. 2). It was surprising to us that the signals for $(TFA)_3Sb^{V}(para-tolyl)_2$ would be downfield relative to (TFA)₄Sb^V(para-tolyl). Therefore, we used MN15 (ref. 29)/def2-SVP30 density functional theory (DFT) gauge-independent atomic orbital31 NMR calculations in Gaussian 16 (ref. 32) to calculate the relative chemical shifts of the aryl hydrogens. These calculations confirmed the relative signal assignment of the mono-tolyl versus bis-tolyl complex and indicate that the downfield shift of the bis-tolyl structure is due to the adoption of a trans-like geometry. For both the monotolyl and bis-tolyl complexes C-H activation/metalation occurred almost exclusively at the toluene para position. This para selective C-H activation of toluene is consistent with the similar reaction between Tl^{III}(TFA)₃ and toluene where McKillop reported a 9:4:87 ratio of ortho: meta: para metalation.20

With the observation of both $(TFA)_4Sb^v(para-tolyl)$ and $(TFA)_3Sb^v(para-tolyl)_2$ and the relative time dependence of these products we considered two general mechanisms for the formation of the mono-tolyl and bis-tolyl complex. The first mechanism involves a metalation/C–H activation reaction step to generate the mono-tolyl intermediate followed by a second metalation/C–H activation reaction step to generate the bis-tolyl product. The most straightforward alternative mechanism involves two of the mono-tolyl intermediates undergoing tolyl for TFA group transfer to generate (TFA)_3Sb^v(tolyl)_2 and

 $Sb^{V}(TFA)_{5}$. To confirm this reaction sequence and potentially distinguish between double C-H activation and disproportion pathways, we independently prepared (TFA)₄Sb^V(para-tolyl), through a route using (OH)₂Sb^V(O)(*para*-tolyl), and examined its reaction and products in the presence and absence of toluene and/or toluene-d₈ in TFAH/TFAA (4:1) solvent (see ESI for results[†]). This showed that starting with (TFA)₄Sb^V(para-tolyl) generated the same bis-tolyl product as that generated from the reaction of $Sb^{V}(TFA)_{5}$ with toluene at extended times or higher temperatures. This generally confirms that the bis-tolyl complex is generated by conversion from the mono-tolyl complex. Also, this reaction indicates that the TFA for tolyl group transfer pathway is likely viable and that formation of the bis-tolyl complex does not require reaction with free toluene. However, this simple interpretation is complicated because heating $(TFA)_4Sb^{V}(p-tolyl)$ in TFAH generates low concentrations of free toluene and indicates that protonation is slower but competitive with formation of the bis-tolyl complex and therefore it cannot be ruled out that the free toluene would rapidly react with $(TFA)_4Sb^{V}(para-tolyl)$ to give the bis-tolyl complex through the double C-H activation pathway (see ESI[†]). While it is conceivable that the double C-H activation and ligand group transfer pathways could be distinguished based on rate dependence of added toluene unfortunately the poor resolution of the resonances of the mono-tolyl versus the possible bis-tolyl complexes in the ¹H NMR spectra limited this type of quantitative rate analysis (see ESI[†]).

As can be seen from Fig. 2, reaction of $Sb^{V}(TFA)_{5}$ with toluene in TFAH/TFAA at 60 °C and 70 °C generated $[Sb^{V}]$ -tolyl products but did not generate oxy-functionalized TFA ester cresol products. In retrospect this was perhaps not surprising since the calculated barriers for toluene C–H activation are relatively low (see later section describing calculations) while the reductive functionalization barriers generating tolyl trifluoroacetate cresol esters are relatively high. This comparison of experiments and calculated barriers prompted us to examine whether Sb^V(TFA)₅ could indeed be coaxed to induce oxyfunctionalization of toluene at higher temperatures. Fig. 3 shows the ¹H NMR spectrum of the crude reaction mixture of Sb^V(TFA)₅ reaction with toluene in TFAH at 170 °C for 3 hours. Consistent with the computational predictions, the ¹H NMR shows the formation of the TFA esters of cresol but with some decomposition to unidentified products. But not predicted, and relatively surprising, by comparison to independently synthesized cresol TFA esters, the major oxy-functionalized products are primarily ortho and meta cresol trifluoroacetates with only traces of the para cresol trifluoroacetate. The formation of ortho and meta cresol trifluoroacetates is surprising because we initially assumed oxy-functionalization would occur from either the para mono-tolyl or bis-tolyl complexes. To determine if the formation of ortho and meta cresol trifluoroacetates resulted from first the formation of the para cresol trifluoroacetate we examined whether this compound could be converted to ortho or meta products under reaction conditions. However, control reactions with independently synthesized para cresol trifluoroacetate showed no such conversion (see ESI[†]).

The experimental observation of (TFA)₄Sb^V(para-tolyl) and (TFA)₃Sb^V(*para*-tolyl)₂ from reaction of toluene with Sb^V(TFA)₅ at lower temperatures as well the tolyl trifluoroacetate products at higher temperatures are important for several reasons. First, in our previous examination of Sb^V(TFA)₅ reaction with methane and ethane we did not observe a [Sb^V]-alkyl intermediate (analogous to the Sb-tolyl intermediate here) presumably because the functionalization reaction step was significantly faster than the C-H activation reaction step. Rather, [Sb^V]-alkyl intermediates were only predicted based on DFT calculations of the reaction mechanism. These calculations suggested that for reactions with alkanes the C-H activation first step (other mechanisms were computationally ruled out) is rate limiting and there is a subsequent relatively fast reductive functionalization step. However, here observation of the Sb^V-tolyl bond indicates that C-H activation is a viable mechanistic step for

a Sb^V metal center,³³ which contrasts the general view of this metal and oxidation state as being primarily a Lewis acid that generates superacids and is not directly involved with C-H bond making and breaking reaction steps. Also, observation of a Sb^Varyl bond rather than a Sb^V-benzyl bond suggests a non-radical type of mechanism for C-H activation. Second, while transition metal mediated arene C-H activation is well known, the subsequent reductive functionalization to generate C-O bonds is relatively rare, with Pd and Cu being the most prominent examples. Therefore, the observation of tolvl trifluoroacetate products demonstrates a new metal and oxidation state for this type of arene oxy-functionalization reaction, and this can be compared to previous work where Tl^{III} was reported to be capable of arene C-H activation, but functionalized organic products were generally not reported without added nucleophiles. This new reaction can also be compared to work showing that Pb^{IV} can also promote oxidative functionalization of arenes,²⁵⁻²⁷ but has sometimes been proposed to occur through a radical, non-C-H activation mechanism. Third, as mentioned in the previous paragraph, in this reaction the mono-tolyl and bis-tolyl complexes show very high para selectivity, which is not surprising. However, it is surprising that subsequent functionalization generated only ortho and meta cresol trifluoroacetates.

Computational evaluation of C-H functionalization pathways

While the experiments established $(TFA)_4Sb^{v}(para-tolyl)$ and $(TFA)_3Sb^{v}(para-tolyl)_2$ as intermediates in route to oxyfunctionalization of toluene, it was not clear from experiments the exact mechanism and selectivity of forming these intermediates and their subsequent functionalization. Therefore, we decided to use DFT calculations to determine the energies and structures for reaction mechanisms leading to $(TFA)_4Sb^{v}(tolyl)$, and the metalation regioselectivity. Also, calculations were used to examine the reaction mechanisms and selectivity for reductive functionalization to determine if functionalization occurs from the mono-tolyl complex or the bis-tolyl complex.



Fig. 3 Overlay of ¹H NMR spectra of crude reaction mixture of Sb^V(TFA)₅ and toluene in TFAH : TFAA (4 : 1) taken after 3 hours at 170 °C and neat cresols dissolved in TFAH : TFAA (4 : 1). # = Unidentified products. \blacklozenge = Cresol esters.

Our previous computational evaluation⁴ of Sb^V(TFA)₅ reaction with ethane established that the likely ground state of this complex involves an equilibrium with solvent coordination to the Sb^V center to give (TFAH)Sb^V(TFA)₅. Our previous work also examined possible multinuclear ground states and in general while multi-Sb ground states are likely formed they do not change the mechanistic or reactivity model for inorganic Sb^V TFA complexes.5 Here we used MN15/def2-SVP to optimize all structures in Gaussian 16.32 Frequency calculations and intrinsic reaction coordinate (IRC) calculations were used to establish potential-energy surface connections. All intermediates have all positive vibrational frequencies while transitionstate structures have a single imaginary frequency. MN15/ def2-TZVPD single point energies were then performed on the MN15/def2-SVP structures. Final energies reported correspond to MN15/def2-TZVPD//MN15/def2-SVP. We also calculated single point energies with the M06 and PWPB95-D3(BJ) functionals. This showed that while there are a range of energies calculated structures for C-H activation and functionalization intermediates and transition states no qualitative conclusions change based on a different functional. Zero-point, thermal, and Gibbs energy corrections were applied from the MN15/def2-SVP calculations. We used a modified version of the SMD³⁴ continuum solvent method for estimation of trifluoroacetic acid (TFAH) solvent stabilization. This was used during both structure optimizations and single point energy evaluations. Additionally, we also included an explicit TFAH solvent in nearly all calculations. For each structure we used CREST/GFN2-xTB^{35,36} to perform a comprehensive search to identify Sb-ligand conformations and explicit solvent hydrogen bonding. Generally, the 15 lowest energy xTB structures were then optimized with DFT.

Fig. 4a outlines the general mechanisms examined for toluene C-H bond cleavage and formation of (TFA)₄Sb^V(paratolyl). Pathway A involves the C-H activation pathway where in one step there is proton transfer to a TFA ligand and formation of the Sb^V-C bond. A two-step electrophilic substitution pathway could be envisioned to occur by Sb-induced dearomatization to generate a Wheland type intermediate (i.e. a carbocation intermediate with a full Sb–C σ bond) followed proton transfer step. However, similar to our previous calculations with Tl^{III}(TFA)₃,²⁴ we could not locate a Wheland type intermediate on the potential energy surface because these types of structures are high in energy. Alternative to C-H activation, pathway B describes the possibility of a sequence of electron transfer (ET) and proton transfer (PT) that gives an aryl and Sb^{IV} radical pair that then combine to form the Sb^V-C bond. The one-step process that combines ET and PT is proton-coupled electron transfer (PCET) or formally hydrogen atom transfer. These oneelectron, open-shell reaction pathways are reasonable to consider because Kochi previously reported the observation of radical species when Tl^{III} was combined with arenes.^{22,23} However, our previous calculations suggested that only relatively activated arenes with more than three alkyl groups or more potent electron donor groups (e.g. OMe group) result in open-shell reaction pathways being more favorable than C-H activation.24 Lastly, pathway D involves Sb-induced hydride abstraction to generate a Sb-H anion and an arene cation.

The enthalpy/Gibbs energy change for ET from toluene to $(TFAH)Sb^{V}(TFA)_{5}$ to generate $[(TFAH)Sb^{V}(TFA)_{5}]^{-}$ and the toluene radical cation is 20.8/25.2 kcal mol⁻¹, which is a relatively low energy process for generating an ion pair and indicates that this Sb^V is a highly electrophilic species. PCET to generate (TFAH)₂Sb^{IV}(TFA)₄ and the *para* tolyl radical has a thermodynamic enthalpy/Gibbs energy change of 49.0/ 33.0 kcal mol⁻¹. While the energies of ET/PT and PCET reaction pathways are not extremely high in energy as they are in the somewhat related reaction with alkanes, they are significantly higher in energy than the C-H activation pathway (see below). As expected, due to the instability of a sp^2 carbocation, *para* hydride abstraction to form [(TFAH)Sb^V(H)(TFA)₅]⁻ and an aryl carbocation is very high in energy at ~ 60 kcal mol⁻¹. However, we did locate a one-step reaction pathway where hydride abstraction is coupled with TFA capture of the carbocation and this process requires an activation enthalpy of 46.5 kcal mol^{-1} , but this is still too high to be viable. Compared to these pathways, the C-H activation pathway was calculated to be much lower in energy.

Fig. 4b shows the C-H activation pathway (pathway A) energy landscape for reaction of (TFAH)Sb^V(TFA)₅ with toluene to give the para mono Sb^V-tolyl intermediate 2. The sequence of reaction steps involves π -coordination of toluene to give **1**, which is only slightly exergonic. There is a weak sp² C–H σ -coordination structure prior to the C–H activation transition state **1**[‡]. In this transition state there is concerted formation of the Sb^V-aryl bond and proton transfer to the non-coordinating oxygen atom of a TFA ligand (see the bottom of Fig. 4). This C-H activation barrier is surprisingly low with a ΔG^{\ddagger} value of only 9.6 kcal mol^{-1} relative to 1. However, this low barrier is consistent with some conversion observed at room temperature. Nearly all density functional methods surveyed gave a low barrier height (e.g. M06). This suggests that the barrier is likely underestimated because we used a mononuclear Sb ground state model. Most important, this C-H activation barrier is much lower in energy than ET/PT or PCET pathways. The [Sb^V]-tolyl intermediate 2 is exergonic by 10.8 kcal mol⁻¹. We also calculated the analogous C-H activation barrier at the toluene benzyl position. Consistent with the lack of experimental observation of a [Sb^V]-benzyl intermediate, this sp³ C-H bond position has a much larger barrier with a ΔG^{\ddagger} value of 33.0 kcal mol⁻¹. Further support for the C-H activation mechanism is the calculated metalation regioselectivity. Transition state 1[‡] provided correct modelling of the experimental regioselectivity with *para* selectivity. For $\mathbf{1}^{\ddagger}$ the *ortho* and *meta* versions are 3.8 and 1.0 kcal mol⁻¹ higher in Gibbs energy than the para transition state.

As previously discussed, after 2 is formed there is the possibility of either a second C-H activation reaction step or ligand exchange to form $(TFAH)(TFA)_3Sb^V(tolyl)_2$ (3). The calculated Gibbs energy for 2 $(TFAH)(TFA)_4Sb^V(tolyl)$ (2) \rightarrow $(TFAH)(TFA)_3Sb^V(tolyl)_2$ (3) + $(TFAH)Sb^V(TFA)_5$ is -3.0 kcal mol⁻¹, and this small energy change is consistent with a thermodynamic equilibrium giving both mono-tolyl and bis-tolyl complexes and favoring the latter. Despite significant searching efforts exploring the potential energy surface we



Fig. 4 (a) Outline of plausible reaction mechanisms leading to toluene oxy-functionalization promoted by (TFAH)Sb^V(TFA)₅ (energies in kcal mol⁻¹). (b) Relative Gibbs energies and enthalpies (MN15/def2-TZVPD//MN15/def2-SVP; kcal mol⁻¹) for the first and second sp² aromatic C-H activation reaction steps between (TFAH)Sb^V(TFA)₅ and toluene. Also shown is the benzylic C-H activation transition state and the intramolecular reductive functionalization transition state.

could not locate a definitive transition state for this ligand exchange process and therefore cannot easily estimate the kinetic barrier for disproportionation. In contrast, we did locate the second C-H activation transition state between 2 and toluene to give 3. The barrier for this reaction step is $12.1 \text{ kcal mol}^{-1}$. This is not a large barrier and could be accessible at room temperature and therefore unfortunately the calculations cannot definitively determine whether the mono-

tolyl to bis-tolyl conversion occurs through the double C–H activation pathway or the disproportionation pathway. Regardless of the kinetic pathway for ligand exchange, the calculated thermodynamics accurately model the mono-tolyl *versus* bistolyl thermodynamic equilibrium.

Fig. 4 shows that the functionalization transition state 3^{\ddagger} from the bis-tolyl involves a formal reductive elimination generating Sb^{III} and the aryl ester through an intramolecular

substitution type mechanism where the TFA coordinated to the Sb metal center nucleophilically displaces the Sb metal center. The barrier for 3^{\ddagger} is >45 kcal mol⁻¹, which while potentially feasible seemed too high to see significant conversion even at 170 °C. As a note, we also calculated the aryl-aryl coupling transition state from 2. This transition state is >10 kcal mol⁻¹ higher in energy than 3[‡] and explains why no C-C bond forming product is found. These large barriers from the bis-tolyl prompted us to examine the reductive functionalization transition state 2^{\ddagger} from the mono-tolyl complex. This reductive functionalization transition state requires a Gibbs barrier of only 28.0 kcal mol^{-1} (relative to 2) that is more consistent with the experimental observation of oxy-functionalization at 170 °C. This lower barrier for functionalization of the mono-tolyl intermediate versus the bis-tolvl intermediate is the result of a much larger reduction potential for the Sb^V to Sb^{III} conversion. Therefore, the lowest energy pathway for functionalization of the bis-tolyl complex is first reversion back to the mono-tolyl complex followed by transition state 2^{\ddagger} .

The potential energy surface shown in Fig. 4b with a relatively large barrier for 2[‡] provides an explanation for the *ortho/* meta functionalization selectivity that is opposite to that of the para C-H activation selectivity and that there was no para-tolyl trifluoroacetate observed. At the lower temperatures (room temperature to 70 °C) the C-H activation transition state 1^{\ddagger} controls the selectivity, and as described above the calculated barrier for para C-H activation is lower than ortho and meta C-H activation. In contrast, at high temperatures the conversion of 2 back to **1** and toluene through **1**[‡] will occur at a much faster rate than 2 proceeding through 2[‡]. This suggests a Curtin-Hammett type selectivity model where at higher temperatures there is rapid equilibration of the mono-tolyl and bis-tolyl complexes (and potentially even a tris-tolyl complex) with $Sb^{V}(TFA)_{5}$ and toluene followed by slow functionalization. Consistent with the selectivity model, calculation of the ortho 2^{\ddagger} , meta 2^{\ddagger} , and para 2^{\ddagger} shows ortho and para transition states of nearly equal energy and *para* 2^{\ddagger} is about 2 kcal mol⁻¹ higher in energy, which is just high enough in energy not to be competitive.

Conclusions

The reaction between $Sb^{V}(TFA)_{5}$ and toluene showed that $[Sb^{V}]$ is capable of C-H activation/metalation and leads to NMR detected (TFA)₄Sb^V(para-tolyl) and (TFA)₃Sb^V(para-tolyl)₂ intermediates. This importantly confirms our previous hypothesis of a C-H activation mechanism between Sb^V(TFA)₅ and light alkanes where we could not observe this type of intermediate. This toluene reaction was kinetically selective for para sp² C-H activation over ortho and meta sp² C-H activation as well as benzylic sp³ C-H activation. Experiments and theory indicate that (TFA)₄Sb^V(tolyl) is formed through a metal mediated C-H activation transition state where the Sb^V-C bond is formed simultaneous to proton transfer to a TFA ligand. Subsequently (TFA)₄Sb^V(para-tolyl) converts to (TFA)₃Sb^V(para-tolyl)₂ through a ligand exchange reaction step, although we cannot rule out sequential double C-H activation reaction steps. However, because the barriers for reductive functionalization are

relatively high there is a Curtin-Hammett type scenario where there is rapid equilibrium between the mono-tolyl and bis-tolyl complexes with Sb^V(TFA)₅ and toluene. Because of this equilibrium and the functionalization transition states the regiosechanges to *ortho/meta* lectivity selectivity for oxvfunctionalization. Overall, the combination of calculations and experiments more firmly establish a new mode of reactivity for [Sb^V] beyond the normal Lewis acid induced super Brønsted acidity. Specifically, [Sb^V] is capable of both C–H activation and [Sb^V]-C bond oxy-functionalization reaction steps. Knowledge of these reaction steps will hopefully propel new reaction discovery and catalysis based on complexes with a Sb^V metal center and dovetails with efforts by Gabbai and others demonstrating that complexes with a Sb^V metal center can operate in non-innocent reactivity modes.37,38

Data availability

The data supporting this article have been included as part of the ESI.†

Author contributions

D. Ess designed the experimental and computational studies, analyzed data, and wrote the manuscript. A. Koppaka designed experimental studies, executed experimental studies, interpreted data and wrote the manuscript. S. Chen, D. Yang, and S. Koumleh carried out calculations. A. Marchenko assisted in experimental studies. D. Michaelis and R. Periana analyzed and interpreted data.

Conflicts of interest

There are no conflicts to declare.

Acknowledgements

We thank Brigham Young University and the Office of Research computing, especially the Fulton Supercomputing Lab. This work was fully supported by the U.S. Department of Energy, Office of Science, Basic Energy Sciences, Catalysis Science Program, under award # DE-SC0018329.

Notes and references

- 1 G. A. Olah, Electrophilic methane conversion, *Acc. Chem. Res.*, 1987, **20**, 422–428.
- 2 G. A. Olah and R. H. Schlosberg, Chemistry in super acids. I. Hydrogen exchange and polycondensation of methane and alkanes in FSO_3H - SbF_5 ("magic acid") solution. Protonation of alkanes and the intermediacy of CH_5^+ and related hydrocarbon ions. The high chemical reactivity of "paraffins" in ionic solution reactions, *J. Am. Chem. Soc.*, 1968, **90**, 2726–2727.
- 3 G. A. Olah, G. K. S. Prakash, Á. Molnár and J. Sommer, *Superacid Chemistry*, John Wiley & Sons, Inc., Hoboken, New Jersey, 2nd edn, 2009, pp. 511–517.

- 4 A. Koppaka, S. H. Park, B. G. Hashiguchi, N. J. Gunsalus, C. R. King, M. M. Konnick, D. H. Ess and R. A. Periana, Selective C-H Functionalization of Methane and Ethane by a Molecular Sb(V) Complex, *Angew. Chem., Int. Ed.*, 2019, **58**, 2241–2245.
- 5 S.-S. Chen, A. Koppaka, R. A. Periana and D. H. Ess, Theory and Experiment Demonstrate that Sb(V)-Promoted Methane C-H Activation and Functionalization Outcompetes Superacid Protonolysis in Sulfuric Acid, *J. Am. Chem. Soc.*, 2021, **143**, 18242–18250.
- 6 R. J. Schmidt, Industrial catalytic processes—phenol production, *Appl. Catal., A*, 2005, **280**, 89–103.
- 7 S. R. Neufeldt and M. S. Sanford, Controlling Site Selectivity in Palladium-Catalyzed C-H Bond Functionalization, *Acc. Chem. Res.*, 2012, **45**, 936–946.
- 8 T. Yoneyama and R. H. Crabtree, Pd(II) catalyzed acetoxylation of arenes with iodosyl acetate, *J. Mol. Catal. A: Chem.*, 1996, **108**, 35–40.
- 9 Z. Li, Z. Wang, N. Chekshin, S. Qian, J. X. Qiao, P. T. Cheng, K.-S. Yeung, W. R. Ewing and J.-Q. Yu, A Tautomeric Ligand Enables Directed C–H Hydroxylation with Molecular Oxygen, *Science*, 2021, **372**, 1452–1457.
- 10 W. Wang, F. Luo, S. Zhang and J. Cheng, Copper(II)-Catalyzed Ortho-Acyloxylation of the 2-Arylpyridines sp2 C-H Bonds with Anhydrides, Using O₂ as Terminal Oxidant, *J. Org. Chem.*, 2010, 75, 2415–2418.
- 11 J. Gallardo-Donaire and R. Martin, Cu-Catalyzed Mild C(sp2)–H Functionalization Assisted by Carboxylic Acids en Route to Hydroxylated Arenes, *J. Am. Chem. Soc.*, 2013, **135**, 9350–9353.
- 12 A. M. Suess, M. Z. Ertem, C. J. Cramer and S. S. Stahl, Divergence between Organometallic and Single-Electron-Transfer Mechanisms in Copper(II)-Mediated Aerobic C-H Oxidation, *J. Am. Chem. Soc.*, 2013, **135**, 9797–9804.
- 13 S.-Z. Sun, M. Shang, H.-L. Wang, H.-X. Lin, H.-X. Dai and J.-Q. Yu, Cu(II)-Mediated C(sp2)-H Hydroxylation, *J. Org. Chem.*, 2015, **80**, 8843–8848.
- 14 B. K. Singh and R. Jana, Ligand-Enabled, Copper-Promoted Regio- and Chemoselective Hydroxylation of Arenes, Aryl Halides, and Aryl Methyl Ethers, *J. Org. Chem.*, 2016, **81**, 831–841.
- 15 X. Chen, X.-S. Hao, C. E. Goodhue and J.-Q. Yu, Cu(II)-Catalyzed Functionalizations of Aryl C-H Bonds Using O₂ as an Oxidant, *J. Am. Chem. Soc.*, 2006, **128**, 6790–6791.
- 16 X. Li, Y.-H. Liu, W.-J. Gu, B. Li, F.-J. Chen and B.-F. Shi, Copper-Mediated Hydroxylation of Arenes and Heteroarenes Directed by a Removable Bidentate Auxiliary, *Org. Lett.*, 2014, **16**, 3904–3907.
- 17 M. Shang, Q. Shao, S.-Z. Sun, Y.-Q. Chen, H. Xu, H.-X. Dai and J.-Q. Yu, Identification of monodentate oxazoline as a ligand for copper-promoted ortho-C–H hydroxylation and amination, *Chem. Sci.*, 2017, **8**, 1469–1473.
- 18 F. Kong, S. Chen, J. Chen, C. Liu, W. Zhu, D. A. Dickie, D. H. Ess, S. Zhang and T. B. Gunnoe, Cu(II) Carboxylate Arene C-H Functionalization: Tuning for Non-Radical Pathways, *Sci. Adv.*, 2022, 8, eadd1594.

- 19 A. McKillop, J. S. Fowler, M. J. Zelesko, J. D. Hunt, E. C. Taylor and G. McGillivray, Thallium in organic synthesis. IX. Facile thallation of aromatic compounds with thallium(III) trifluoroacetate, *Tetrahedron Lett.*, 1969, **10**, 2423–2426.
- 20 A. McKillop, J. D. Hunt, M. J. Zelesko, J. S. Fowler, E. C. Taylor, G. McGillivray and F. Kienzle, Thallium in organic synthesis. XXII. Electrophilic aromatic thallation using thallium(III) trifluoroacetate. Simple synthesis of aromatic iodides, *J. Am. Chem. Soc.*, 1971, **93**, 4841–4844.
- 21 E. C. Taylor, F. Kienzle, R. L. Robey, A. McKillop and J. D. Hunt, Thallium in organic synthesis. XXIII. Electrophilic aromatic thallation. Kinetics and applications to orientation control in the synthesis of aromatic iodides, *J. Am. Chem. Soc.*, 1971, **93**, 4845–4850.
- 22 I. H. Elson and J. K. Kochi, Thallium(III) in one-electron oxidation of arenes by electron spin resonance, *J. Am. Chem. Soc.*, 1973, **95**, 5060–5062.
- 23 W. Lau and J. K. Kochi, Kinetics and mechanism of aromatic thallation. Identification and proof of competing electrophilic and electron-transfer pathways, *J. Am. Chem. Soc.*, 1984, **106**, 7100–7112.
- 24 C. R. King, S. J. Gustafson, B. R. Black, S. K. Butler, M. K. Konnick, R. A. Periana, B. M. Hashiguchi and D. H. Ess, Arene C-H Functionalization by p-Block Metal Tl(III) Occurs at the Borderline of C-H Activation and Electron Transfer, *Organometallics*, 2017, 36, 109–113.
- 25 R. E. J. Partch, Comparative Lead(IV) Chemistry. Reactions of Lead Tetra(trifluoroacetate). I, J. Am. Chem. Soc., 1967, 89, 3662–3663.
- 26 J. R. Kalman, J. T. Pinhey and S. Sternhell, Reactions of Lead Tetrakistrifluoroacetate with Aromatic and Metalloaromatic Compounds. A New Route to Phenols, *Tetrahedron Lett.*, 1972, 18, 5369–5372.
- 27 J. R. Campbell, J. R. Kalman, J. T. Pinhey and S. Sternhell, Trifluoroacetoxylation of Benzene Derivatives with Lead Tetrakistrifluoroacetate (LiTFA) in Trifluoroacetic Acid (TFA), *Tetrahedron Lett.*, 1972, **18**, 1763–1766.
- 28 G. O. Doak and H. G. Steinman, The Preparation of Stibonic Acids by the Scheller Reaction, *J. Am. Chem. Soc.*, 1946, **68**, 1987–1989.
- 29 H. S. Yu, X. He, S. L. Li and D. G. Truhlar, MN15: A Kohn-Sham Global-Hybrid Exchange-Correlation Density Functional with Broad Accuracy for Multi-Reference and Single-Reference Systems and Noncovalent Interactions, *Chem. Sci.*, 2016, 7, 5032–5051.
- 30 F. Weigend and R. Ahlrichs, Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: Design and assessment of accuracy, *Phys. Chem. Chem. Phys.*, 2005, 7, 3297–3305.
- 31 R. Ditchfield, Self-consistent perturbation theory of diamagnetism.
 1. Gauge-invariant LCAO method for N.M.R. chemical shifts, *Mol. Phys.*, 1974, 27, 789–807.
- 32 M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria,
 M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone,
 G. A. Petersson, H. Nakatsuji, X. Li, M. Caricato,
 A. V. Marenich, J. Bloino, B. G. Janesko; , R. Gomperts,

B. Mennucci, H. P. Hratchian, J. V. Ortiz, A. F. Izmaylov, J. L. Sonnenberg, D. Williams-Young, F. Ding, F. Lipparini, F. Egidi, J. Goings, B. Peng, A. Petrone, T. Henderson, D. Ranasinghe, V. G. Zakrzewski, J. Gao, N. Rega, G. Zheng, W. Liang, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, 0. Kitao. H. Nakai, Т. Vreven, K. Throssell. J. A. Montgomery Jr, J. E. Peralta, F. Ogliaro, M. J. Bearpark, J. J. Heyd, E. N. Brothers, K. N. Kudin, V. N. Staroverov, T. A. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. P. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, J. M. Millam, M. Klene, C. Adamo, R. Cammi, J. W. Ochterski, R. L. Martin, K. Morokuma, O. Farkas, J. B. Foresman and D. J. Fox, Gaussian 16, Revision B.01, Gaussian, Inc., Wallingford CT, 2016.

33 C. R. King, A. Holdaway, G. Durrant, J. Wheeler, L. Suaava, M. M. Konnick, R. A. Periana and D. H. Ess, Supermetal: SbF₅-Mediated Methane Oxidation Occurs by C-H Activation and Isobutane Oxidation Occurs by Hydride Transfer, *Dalton Trans.*, 2019, **48**, 17029–17036.

- 34 A. V. Marenich, C. J. Cramer and D. G. Truhlar, Universal Solvation Model Based on Solute Electron Density and on a Continuum Model of the Solvent Defined by the Bulk Dielectric Constant and Atomic Surface Tensions, *J. Phys. Chem. B*, 2009, **113**, 6378–6396.
- 35 P. Pracht, F. Bohle and S. Grimme, Automated exploration of the low-energy chemical space with fast quantum chemical methods, *Phys. Chem. Chem. Phys.*, 2020, **22**, 7169–7192.
- 36 S. Spicher and S. Grimme, Robust Atomistic Modeling of Materials, Organometallic, and Biochemical Systems, *Angew. Chem., Int. Ed.*, 2020, **59**, 15665–15673.
- 37 J. S. Jones and F. P. Gabbai, Coordination- and Redox-Noninnocent Behavior of Ambiphilic Ligands Containing Antimony, *Acc. Chem. Res.*, 2016, **49**, 857–867.
- 38 J. M. Lipschultz, G. Li and A. T. Radosevich, Main Group Redox Catalysis of Organopnictogens: Vertical Periodic Trends and Emerging Opportunities in Group 15, *J. Am. Chem. Soc.*, 2021, 143, 1699–1721.