Chemical Science



EDGE ARTICLE

View Article Online
View Journal | View Issue



Cite this: Chem. Sci., 2025, 16, 20564

d All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 25th July 2025 Accepted 27th September 2025

DOI: 10.1039/d5sc05582b

rsc.li/chemical-science

Construction of α -quinonyl- α -hydroxy/amino acid esters through the redox chain reaction


Although quinones, α -hydroxy acids and α -amino acids are all widely used organic compounds in numerous fields, compounds based on the combination of these motifs were underexplored. In this study, the direct hydroxyalkylation of quinones with glyoxylates was developed through a redox chain reaction using SnCl₂ as the reductive initiator and catalyst, and AlCl₃ as the co-catalyst. Various α -quinonyl- α -hydroxyacetyl esters were obtained under mild conditions in up to 81% yield within a few hours. Substrates with different ester groups were accessed through *in situ* oxidation of corresponding tartrates to glyoxylates by PIDA. Moreover, aminoalkylation of quinones was achieved by the multicomponent reaction of quinones, glyoxylates and amides, affording a series of α -quinonyl- α -aminoacetyl esters in up to 80% yield. The products were easily converted to unnatural amino acids, aryl azo compounds and peptides as demonstrations of their wide potential applications.

Introduction

α-Hydroxy¹⁻⁴ and α-amino⁵⁻⁹ acids are widespread motifs in biological and pharmaceutical molecules (Scheme 1a). Artificially synthesized unnatural amino acids potentially provide distinctive functions beyond existing natural amino acids for biology and chemistry. 10,11 Despite significant research progress in this field, reports on quinone-substituted amino acid derivatives are limited. 12-14 Based on their redox properties, quinones could serve as mediators for cellular electron transfer processes. 15-20 Quinones are also a very versatile platform for further functionalization and conjugation with other molecules. Introducing quinones into the side chain of amino acids may provide new functional unnatural amino acids and related materials. In recent years, structural modifications of quinones have significantly advanced. 17,21-24 Among all the existing protocols, our group made the original contributions of a unique and efficient strategy for direct quinone functionalization with various sp³ electrophiles, the redox chain reaction (Scheme 1b).25-28 With this new strategy, quinones can be functionalized in one step, bypassing the traditional reduction-functionalization-oxidation sequence. Herein, we report that many densely functionalized α-quinonyl-α-hydroxy and amino acetyl esters (QHAE and QAAE) are constructed by the redox chain reaction of quinones with glyoxylates^{29,30} and their imines as uncharted sp² electrophiles, using SnCl₂ as both the reductive initiator and

Results and discussion

To investigate the conditions, 2,6-dimethylquinone 1a and ethyl glyoxylate 2 were chosen as the model substrates. The initial investigation used Hantzsch ester (HE) as the reductive initiator, and different Lewis acids was conducted at 30 °C in 1,2-dichloroethane (DCE) (Table 1, Entries 1–6). The combination



Scheme 1 α -Hydroxy and amino acid motifs.

the main Lewis acid catalyst, along with ${
m AlCl_3}$ as a Lewis acid cocatalyst (Scheme 1c). 31

[&]quot;School of Physical Science and Technology, ShanghaiTech University, Shanghai 201210, China. E-mail: lizhi@shanghaitech.edu.cn

^bState Key Laboratory of Advanced Medical Materials and Devices, ShanghaiTech University, Shanghai 201210, China

Table 1 Optimization of the reaction conditions^a

Entry	x	RI/(mol%)	T/°C	LA/(mol%)	Yield (%)
1	1.5	HE (10)	30	Hf(OTf) ₄ (5)	63
2	1.5	HE (10)	30	$Yb(OTf)_3(5)$	0
3	1.5	HE (10)	30	$Sc(OTf)_3$ (5)	0
4	1.5	HE (10)	30	$HfCl_4(5)$	0
5	1.5	HE (10)	30	$ZrCl_4(5)$	0
6	1.5	HE (10)	30	$AlCl_3(5)$	2
7	1.5	$SnCl_2(5)$	30	_	78
8	1.5	CuCl (5)	30	_	0
9	1.5	FeCl ₂ (5)	30	_	0
10	1.5	$SnCl_2$ (10)	30	_	73
11	1.5	$SnCl_2(5)$	30	$SnCl_4(5)$	73
12	1.5	$SnCl_2(5)$	30	FeCl ₃ (5)	69
13	1.5	$SnCl_2(5)$	30	$AlBr_3(5)$	80
14	1.5	SnCl ₂ (5)	30	AlCl ₃ (5)	97
15	1.0	$SnCl_2(5)$	30	$AlCl_3(5)$	75
16	2.0	$SnCl_2(5)$	30	$AlCl_3(5)$	98
17	1.5	$SnCl_2(5)$	0	$AlCl_3(5)$	8
18	1.5	$SnCl_2(5)$	60	$AlCl_3(5)$	85

^a Conditions: 2 (0.1 mmol, 1.0 equiv.) was added to a mixture of **1a** (1.5 equiv.), initiator and Lewis acid in 0.5 mL DCE that was pre-stirred for 10 min, stirred for 3 h, and then quenched by using MnO₂. Yields were determined by ¹H NMR using 1,1,2,2-tetrachloroethane as the internal standard.

of HE and Hf(OTf)₄ could not provide higher than 63% QHAE product **3a**, likely because of the basicity of the HE oxidation product. Reductive Lewis acids were then considered to replace both HE and Hf(OTf)₄ (Table 1, Entries 7–10). Using 5 mol% SnCl₂, the reaction afforded 78% yield of **3a**, which was not improved when doubling the amount of SnCl₂. Attempts were made using other Lewis acids to improve the reaction efficiency (Table 1, Entries 11–14), and up to 97% NMR yield was achieved in the presence of 5 mol% of AlCl₃. Reducing the amount of quinone **1a** decreased the yield, while further increasing **1a** only marginally improved the yield (Table 1, Entries 15 and 16). When the temperature was lowered to 0 °C, the efficiency significantly decreased, whereas increasing the temperature to 60 °C maintained an 85% yield (Table 1, Entries 17 and 18).

With the optimized conditions (Table 1, Entry 14) in hand, the substrate scope was investigated. Broad ranges of quinones provided various QHEAs in up to 81% isolated yields (Scheme 2). In addition to alkyl and aryl substituted (3c-g) quinones, the reaction exhibited good tolerance for groups with various electronic effects. The quinones substituted with both electron-donating groups (3h-j) and electron-withdrawing groups (3k, l) afforded reasonable yields in this system. Naphthoquinone (3m) could also serve as the reactant with a high yield. For mono-substituted benzoquinones with multiple possible reactive positions, only moderate regioselectivity was achieved (3n-q), which slightly preferred the para

Scheme 2 Product scope for QHAEs. ^aReactions were performed on a 0.20 mmol scale in 1.0 mL DCE. Isolated yields were reported.

position of the quinone substituent. Unfortunately, other carbonyl-containing substrates including pyruvates were unable to exhibit any reactivity (Scheme S1).

Next, glyoxylates with different ester groups were investigated as reactants. These compounds were typically synthesized through oxidation of the corresponding tartrates.^{32,33} However, due to the relatively high reactivity of glyoxylates, an alternative approach was implemented in which the glyoxylates were generated by *in situ* oxidation of the corresponding tartrate

Table 2 Screening of the oxidants^a

Entry	Oxidant	Yield (%)
1	0.5 equiv. H ₅ IO ₆	0
2	$0.5 \text{ equiv. NaIO}_4$	0
3	2.0 equiv. quinone 1a	0
4	0.5 equiv. PIDA	29
5	1.0 equiv. PIDA	97
6	2.0 equiv. PIDA	0
7^b	0.50 equiv. PIDA	0

^a Unless otherwise noted, 0.1 mmol **4a** (1.0 equiv.) was mixed with the oxidant in 0.5 mL DCE for 1 h, then added to a solution of 0.3 mmol **1a** (1.5 equiv.), 0.01 mmol SnCl₂ (5 mol%) and AlCl₃ (5 mol%) in 0.5 mL DCE, stirred for 3 h, and then quenched by using MnO₂. Yields were determined by ¹H NMR using 1,1,2,2-tetrachloroethane as the internal standard. PIDA = phenyliodine diacetate. ^b PIDA, **1a**, **4a**, SnCl₂ and AlCl₃ were all mixed in 1.0 mL DCE.

esters, enabling a "one-pot" reaction. Diisopropyl tartrate 4a was used as the model reactant to study the conditions (Table 2). Inorganic oxidants such as periodic acid (H₅IO₆) and sodium periodate (NaIO₄) failed to afford the desired products (Table 2, Entries 1 and 2). Increasing the amount of quinone 1a, as it could also serve as the oxidant, did not work either (Table 2, Entry 3). On the other hand, phenyliodine diacetate (PIDA) had been disclosed as a viable oxidant for the cleavage of tartrates.34,35 Stoichiometry study (Table 2, Entries 4-6) found that mixing 4a with 1.0 equiv. of PIDA for 1 h, which was then allowed to react with quinone 1a and the catalyst in the same flask, was sufficient to afford the product 5a in the highest 97% yield. The intermediate glyoxylate was confirmed by the crude NMR spectrum. A smaller amount of PIDA could not convert all the tartrate 4a, while a larger amount of PIDA inhibited the reaction likely by consumption of the reductive initiator in the following step. Likely for the same reason, if tartrate and PIDA were mixed with quinone altogether instead of stepwise mixing, no product was observed (Table 2, Entry 7). Note that the stoichiometry of the reaction between tartrate and PIDA is 1:1, and 2 equiv. of glyoxylate would be generated afterwards. In order to avoid ambiguity regarding stoichiometry, the amount of PIDA was calculated based on the tartrates as 1 equiv., while in the subsequent redox chain reaction, reagents including the quinone and catalysts were calculated taking the in situ generated glyoxylates as 1 equiv.

Investigation of the scope for esters was then possible using different tartrates as the starting materials (Scheme 3). The tartrates with primary or secondary alkyl esters (5a-e) furnished reasonable yields. Unfortunately, *tert*-butyl ester (5f) only provided a poor result, probably due to its higher reactivity for hydrolysis in the presence of Lewis acids. ³⁶ Benzyl ester (5g) and ether bonds (5h) were both tolerated in this system. Chiral (–)-menthol ester (5i) afforded the product in good yield but low diastereoselectivity.

The successful synthesis of QHAEs prompted us to investigate QAAEs from the corresponding imines. Since the isolation and purification of many imines were challenging, a three-

Scheme 3 Product scope for tartrates. ^aReactions were performed on a 0.2 mmol scale following standard conditions and procedures. Isolated yields were reported.

Scheme 4 Product scope for QAAEs. a To a mixture of 1 (2.0 equiv.), 6 (3.0 equiv.), SnCl₂ (0.1 equiv.), AlCl₃ (0.1 equiv.) and TFA (3.0 equiv.) in 1.0 mL DCE that was pre-stirred for 10 min, 2 (0.2 mmol, 1.0 equiv.) was added and stirred at 30 °C for 12 h, and then quenched by using MnO₂. b 4 (0.1 mmol, 1.0 equiv.) was mixed with 1.0 equiv. PlDA in 0.5 mL DCE for 1 h, then added to a solution of 1a (1.5 equiv.), 6 (3.0 equiv.), SnCl₂ (0.1 equiv.), AlCl₃ (0.1 equiv.) and TFA (3.0 equiv.) in 0.5 mL DCE, stirred for 12 h, and then quenched by using MnO₂. Isolated yields were reported.

component one-pot Mannich-type method³⁷ was established for *in situ* generation of imines in the presence of trifluoroacetic acid (TFA). Various QAAEs were successfully synthesized with this strategy (Scheme 4, see Table S1 also for detailed optimization of conditions). In general, the stoichiometries of 1 and 6 were increased to 2.0 equiv. and 3.0 equiv. respectively, so as to promote the conversion of 2 at a reasonable cost. The structure of the product was confirmed by the single crystal X-ray diffraction (scXRD) analysis of 7aa (CCDC 2427751). The product derived from *p*-toluenesulfonamide (6d) led to a higher yield than that from *tert*-butyl carbamate (6a). Notably, the quinones 1h and 1m would still favour the QHAE products instead of QAAEs. Monosubstituted quinones usually afforded only one regioisomer (7od and 7pd) except 7n. Aryl or alkyl amines^{38,39} showed no reactivity in this system, nor did Ellman's

Scheme 5 Synthetic applications of QHAEs & QAAEs

chiral sulfinamide. 40 Moreover, this method could be combined with the oxidation of tartrates to obtain products containing different esters (8). Similar to the previous strategy, the tartrates were first oxidized by PIDA for 1 h to form the glyoxylates *in situ*, subsequently reacting with amides, quinones and catalysts to afford the QAAEs with different ester groups.

DCM_rt_overnight

To demonstrate the practical potential of the methods, the reaction was scaled up to 50 mmol between **1a**, **2** and **6** under standard conditions, affording **11**.4 g product **7aa** in 68% yield, meanwhile recovering **4**.6 g excessive quinone **1a** (Scheme 5a). The product **7aa** could be further functionalized by the addition of thiol (**9**). The quinone was reduced with sodium dithionite to prevent possible side reactions during its hydrolysis to amino acid (**10**) (Scheme 5b). Both QAAE (**12**) and QHAE (**13**) were able to react with hydrazinium chlorides to afford aryl azo products, ⁴¹ and the derivative **14** was coupled with *N*-protected proline to form a dipeptide **15** (Scheme 5c). Note that this reaction preferred double acylation at both N and O in the same molecule even in the presence of excessive **14**.

Finally, the mechanism of the redox chain reaction was investigated by control experiments. The reaction was effectively initiated by using a catalytic amount of hydroquinone (HQ) 1a' along with non-reducing Lewis acid SnCl₄, while the efficiency was dramatically decreased without 1a' (Scheme 6a), demonstrating that the HQ intermediate was essential in the redox chain. It was also observed that AlCl₃ would act as a promoter for the SnCl₂ reduction of quinones (Scheme 6b). A plausible mechanism was then proposed taking hydroxyalkylation as an example (Scheme 6c). Quinone 1 would first be activated by AlCl₃, making it more labile towards reduction by SnCl₂. The hydroquinolate complex 18 would immediately react

a. Initiating the reaction by catalytic amount of HQ instead of the reducing agent SnCl₂

5 mol % SnCl₄
5 mol % AlCl₃
DCE, 30 °C, 3 h

1a'
1a
2
94 % yield
No HQ 1a'
35 % yield

b. Promoting the reduction of quinones by AlCl₃
DCE, 30 °C, 3 min

10 % yield
No AlCl₃
DCE, 30 °C, 3 min

10 % yield
No AlCl₃
2 % yield

c. Plausible mechanism

chain initiation

chain propagation

AlCl₃

R1

AlCl₃

AlCl₃

AlCl₂

R1

Scheme 6 Control experiments and the proposed plausible mechanism.

with the glyoxylates 2 through a Friedel–Crafts type reaction. Then the substituted hydroquinolate complex 19 would be oxidized to the product 3 by the remaining quinone 1, concomitantly generating the reduced hydroquinolate complex 18 for the next cycle of the redox chain.

Conclusions

In summary, we have developed a step-economical strategy to synthesize α -quinonyl- α -hydroxy/amino acid esters through the redox chain reaction, catalyzed by SnCl₂/AlCl₃. This method provided many potential precursors for functional unnatural α -hydroxy/amino acids.

Author contributions

The manuscript was written through the contributions of all authors. All authors have approved the final version of the manuscript.

Conflicts of interest

There are no conflicts to declare.

Data availability

CCDC 2427751 (7aa) contains the supplementary crystallographic data for this paper.⁴²

The data supporting this article have been included as part of the supplementary information (SI): materials and methods, experimental procedures, and additional characterization of synthesized compounds, experimental details and supplemental tables. See DOI: https://doi.org/10.1039/d5sc05582b.

Acknowledgements

Financial support for this work was generously provided by the ShanghaiTech University start-up funding. We thank the Analytical Instrumentation Center (grant SPST-AIC10112914), SPST, ShanghaiTech University, for providing analytical facilities and services. We thank group member Mr Huanchao Gu for his generous assistance in scXRD characterization.

References

- 1 L. Bowes, The Science of Hydroxy Acids: Mechanisms of Action, Types and Cosmetic Applications, *J. Aesthet. Nurs.*, 2013, 2(2), 77–81.
- 2 H.-M. Ng, S.-T. Bee, L. Tin Sin, C. T. Ratnam and A. R. Rahmat, Hydroxyapatite for Poly(α-Hydroxy Esters) Biocomposites Applications, *Polym. Rev.*, 2019, **59**(2), 187–239.
- 3 C. Renata, D. Katarzyna, Ś. Renata, L. Włodzimierz and H. Ś. Artur, Antimicrobial Properties of Mandelic Acid, Gallic Acid and their Derivatives, *Mini-Rev. Med. Chem.*, 2021, 21(17), 2544–2550.
- 4 C. Egli, M. Min, N. Afzal and R. K. Sivamani, The Hydroxy Acids: Where Have We Been and What's New?, *Dermatol. Rev.*, 2023, 4(6), 260–267.
- 5 G. Wu, Amino Acids: Metabolism, Functions, and Nutrition, *Amino Acids*, 2009, 37(1), 1–17.
- 6 C. T. Walsh, R. V. O'Brien and C. Khosla, Nonproteinogenic Amino Acid Building Blocks for Nonribosomal Peptide and Hybrid Polyketide Scaffolds, *Angew. Chem., Int. Ed.*, 2013, 52(28), 7098–7124.
- 7 K. Lang and J. W. Chin, Cellular Incorporation of Unnatural Amino Acids and Bioorthogonal Labeling of Proteins, *Chem. Rev.*, 2014, 114(9), 4764–4806.
- 8 C. D. Spicer and B. G. Davis, Selective Chemical Protein Modification, *Nat. Commun.*, 2014, 5(1), 4740.
- 9 J. C. Watkins and G. L. Collingridge, Phenylglycine derivatives as antagonists of metabotropic glutamate receptors, *Trends Pharmacol. Sci.*, 1994, **15**(9), 333–342.
- 10 M. A. T. Blaskovich, Unusual Amino Acids in Medicinal Chemistry, *J. Med. Chem.*, 2016, **59**(24), 10807–10836.
- 11 T. Narancic, S. A. Almahboub and K. E. O'Connor, Unnatural Amino Acids: Production and Biotechnological Potential, *World J. Microbiol. Biotechnol.*, 2019, 35(4), 67.
- 12 H. H. Ong, C. R. Creveling and J. W. Daly, Synthesis of 2,4,5-Trihydroxyphenylalanine (6-Hydroxydopa). A Centrally Active Norepinephrine-depleting Agent, *J. Med. Chem.*, 1969, 12(3), 458–461.
- 13 F. G. H. Lee, D. E. Dickson and A. A. Manian, Modified Syntheses of 2,4,5-Trihydroxyphenylalanine, 2,4,5-Trihydroxyphenethylamine, and Analogs, *J. Med. Chem.*, 1971, 14(3), 266–268.
- 14 G. A. Swan, Studies Related to the Chemistry of Melanins. Part XIV. The Alleged Formation of a *p*-Quinonoid

- Aminochrome by Oxidation of 2,4,5-Trihydroxyphenethylamine, *J. Chem. Soc.*, *Perkin Trans.* 1, 1976, (3), 339–341.
- 15 G. Lenaz, R. Fato, G. Formiggini and M. L. Genova, The role of Coenzyme Q in mitochondrial electron transport, *Mitochondrion*, 2007, 7, S8–S33.
- 16 E. A. Hillard, F. C. de Abreu, D. C. M. Ferreira, G. Jaouen, M. O. F. Goulart and C. Amatore, Electrochemical parameters and techniques in drug development, with an emphasis on quinones and related compounds, *Chem. Commun.*, 2008, (23), 2612–2628.
- 17 O. P. S. Patel, R. M. Beteck and L. J. Legoabe, Antimalarial application of quinones: A recent update, *Eur. J. Med. Chem.*, 2021, **210**, 113084.
- 18 M. V. C. Nguyen, B. Lardy, F. Rousset, F. Hazane-Puch, L. Zhang, C. Trocmé, L. Serrander, K.-H. Krause and F. Morel, Quinone compounds regulate the level of ROS production by the NADPH oxidase Nox4, *Biochem. Pharmacol.*, 2013, 85(11), 1644–1654.
- 19 D. Birth, W. C. Kao and C. Hunte, Structural analysis of atovaquone-inhibited cytochrome bc_1 complex reveals the molecular basis of antimalarial drug action, *Nat. Commun.*, 2014, 5, 4029.
- 20 M. Kawamukai, Biosynthesis, bioproduction and novel roles of ubiquinone, *J. Biosci. Bioeng.*, 2002, **94**(6), 511–517.
- 21 S. E. Walker, J. A. Jordan-Hore, D. G. Johnson, S. A. Macgregor and A.-L. Lee, Palladium-Catalyzed Direct C-H Functionalization of Benzoquinone, *Angew. Chem., Int. Ed.*, 2014, 53(50), 13876–13879.
- 22 D. R. Sutherland, M. Veguillas, C. L. Oates and A.-L. Lee, Metal-, Photocatalyst-, and Light-Free, Late-Stage C-H Alkylation of Heteroarenes and 1,4-Quinones Using Carboxylic Acids, *Org. Lett.*, 2018, **20**(21), 6863–6867.
- 23 D. R. Sutherland, N. Sharma, G. M. Rosair, I. D. W. Samuel, A.-L. Lee and E. Zysman-Colman, Synthesis and optoelectronic properties of benzoquinone-based donoracceptor compounds, *Beilstein J. Org. Chem.*, 2019, 15, 2914–2921.
- 24 R. Kumar Jha and S. Kumar, Direct Functionalization of *para*-Quinones: A Historical Review and New Perspectives, *Eur. J. Org Chem.*, 2024, 27(27), e202400535.
- 25 X.-L. Xu and Z. Li, Catalytic Electrophilic Alkylation of *p*-Quinones through a Redox Chain Reaction, *Angew. Chem., Int. Ed.*, 2017, **56**(28), 8196–8200.
- 26 X.-L. Xu and Z. Li, Deciphering the Redox Chain Mechanism in the Catalytic Alkylation of Quinones, *Synlett*, 2018, **29**(14), 1807–1813.
- 27 X.-L. Xu and Z. Li, Catalytic Redox Chain Ring Opening of Lactones with Quinones To Synthesize Quinone-Containing Carboxylic Acids, *Org. Lett.*, 2019, 21(13), 5078– 5081.
- 28 Z. Zhang, H. Gu, D.-X. Cao and Z. Li, Rapid Access to Isoprenoid Quinones through X@RONa-Catalyzed Redox Chain Reaction, *J. Am. Chem. Soc.*, 2024, **146**(42), 29064–29071.
- 29 W. Zhang and P. G. Wang, Ytterbium(III) Trifluoromethanesulfonate Catalyzed Electrophilic

- Aromatic Substitution with Glyoxalate and Lipase-Mediated Product Resolution: A Convenient Route to Optically Active Aromatic α-Hydroxy Esters, *J. Org. Chem.*, 2000, 65(15), 4732–4735.
- 30 J. Kwiatkowski, J. Majer, P. Kwiatkowski and J. Jurczak, Simple and Efficient Synthesis of Racemic Substituted Mandelic Acid Esters from Nonactivated Arenes and Ethyl Glyoxylate, *Synthesis*, 2008, 2008(20), 3237–3244.
- 31 A. Lücht, L. J. Patalag, A. U. Augustin, P. G. Jones and D. B. Werz, Reactions of Donor–Acceptor Cyclopropanes with Naphthoquinones: Redox and Lewis Acid Catalysis Working in Concert, *Angew. Chem., Int. Ed.*, 2017, **56**(35), 10587–10591.
- 32 J. Våbenø, M. Brisander, T. Lejon and K. Luthman, Diastereoselective Reduction of a Chiral *N*-Boc-Protected δ-Amino-α,β-unsaturated γ-Keto Ester Phe-Gly Dipeptidomimetic, *J. Org. Chem.*, 2002, **67**(26), 9186–9191.
- 33 X. Zhang, M. Wang, R. Ding, Y.-H. Xu and T.-P. Loh, Highly Enantioselective and Anti-Diastereoselective Catalytic Intermolecular Glyoxylate–Ene Reactions: Effect of the Geometrical Isomers of Alkenes, *Org. Lett.*, 2015, 17(11), 2736–2739.
- 34 H. Kataoka, T. Ohe, K. Takahashi, S. Nakamura and T. Mashino, Novel Fullerene Derivatives as Dual Inhibitors of Hepatitis C Virus NS5B Polymerase and NS3/4A Protease, *Bioorg. Med. Chem. Lett.*, 2016, **26**(19), 4565–4567.
- 35 S. Ni, A. F. Garrido-Castro, R. R. Merchant, J. N. de Gruyter, D. C. Schmitt, J. J. Mousseau, G. M. Gallego, S. Yang, M. R. Collins, J. X. Qiao, K.-S. Yeung, D. R. Langley, M. A. Poss, P. M. Scola, T. Qin and P. S. Baran, A General

- Amino Acid Synthesis Enabled by Innate Radical Cross-Coupling, *Angew. Chem., Int. Ed.*, 2018, 57(44), 14560–14565.
- 36 E. Marcantoni, M. Massaccesi, E. Torregiani, G. Bartoli, M. Bosco and L. Sambri, Selective Deprotection of *N*-Boc-Protected *tert*-Butyl Ester Amino Acids by the CeCl₃·7H₂O-NaI System in Acetonitrile, *J. Org. Chem.*, 2001, **66**(12), 4430–4432.
- 37 A. E. Schneider, T. Beisel, A. Shemet and G. Manolikakes, Bi(OTf)₃-catalyzed Three-component Synthesis of α -Amino Acid Derivatives, *Org. Biomol. Chem.*, 2014, 12(15), 2356–2359.
- 38 F. Zanardi, A. Sartori, C. Curti, L. Battistini, G. Rassu, G. Nicastro and G. Casiraghi, Diastereoselective Synthesis of 4,5'-Bis-proline Compounds *via* Reductive Dimerization of N-Acyloxyiminium Ions, *J. Org. Chem.*, 2007, 72(5), 1814–1817.
- 39 J. Xie, Y. Huang, H. Song, Y. Liu and Q. Wang, Copper-Catalyzed Aerobic Oxidative [2 + 3] Cyclization/Aromatization Cascade Reaction: Atom-Economical Access to Tetrasubstituted 4,5-Biscarbonyl Imidazoles, *Org. Lett.*, 2017, **19**(22), 6056–6059.
- 40 W. Huang, J.-L. Ye, W. Zheng, H.-Q. Dong and B.-G. Wei, Radical Migration–Addition of *N-tert*-Butanesulfinyl Imines with Organozinc Reagents, *J. Org. Chem.*, 2013, **78**(22), 11229–11237.
- 41 Y.-M. Ye, H.-W. Chen, H. Gu, B. Qiao and Z. Li, A Flash Conversion to Aromatic Azo Compounds Expedited by Hydrazine–Trifluoroacetate Hydrogen Bonding, *Org. Lett.*, 2025, 27(17), 4450–4456.
- 42 CCDC 2427751: Experimental Crystal Structure Determination, 2025, DOI: 10.5517/ccdc.csd.cc2mh8kz.