

CORRECTION

[View Article Online](#)
[View Journal](#) | [View Issue](#)Cite this: *Chem. Sci.*, 2025, 16, 6522

DOI: 10.1039/d5sc90072g

rsc.li/chemical-science

Correction: Low-energy electron distributions from the photoionization of liquid water: a sensitive test of electron mean free paths

Titouan Gadeyne,^{ab} Pengju Zhang,^{*a} Axel Schild^a and Hans Jakob Wörner^{*a}

Correction for 'Low-energy electron distributions from the photoionization of liquid water: a sensitive test of electron mean free paths' by Titouan Gadeyne *et al.*, *Chem. Sci.*, 2022, 13, 1675–1692, <https://doi.org/10.1039/D1SC06741A>.

The authors regret that an incorrect version of Fig. 4 was included in the original article. In the original article, the horizontal scale along the bottom plot of Fig. 4 spans a range from 0 to 50 eV, when it should read 0 to 1 eV.

The authors note that this error is only of graphical nature and does not affect the results and conclusions of the paper.

The correct version of Fig. 4 is presented here.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.




Fig. 4 Singly-differential cross sections for inelastic scattering. (Top) SDCS for electronically inelastic events, for incident kinetic energies $E = 10, 20, 30, 50$ and 100 eV. The black vertical line marks the excitation threshold for liquid water. (Bottom) SDCS for vibrationally inelastic events, for $E = 1, 2$ and 5 eV.

^aLaboratory for Physical Chemistry, ETH Zürich, Vladimir-Prelog-Weg 2, 8093 Zürich, Switzerland. E-mail: pengju.zhang@phys.chem.ethz.ch; hwoerner@ethz.ch

^bDépartement de Chimie, École Normale Supérieure, PSL University, 75005 Paris, France