Journal of Materials Chemistry B



CORRECTION

View Article Online



Cite this: J. Mater. Chem. B, 2025, **13**. 1118

Correction: Hedgehog-inspired immunomagnetic beads for high-efficient capture and release of exosomes


Jia Cheng, Nanhang Zhu, Yujia Zhang, Yue Yu, Ke Kang, Qiangying Yi* and Yao Wu*

DOI: 10.1039/d4tb90208d

rsc.li/materials-b

Correction for 'Hedgehog-inspired immunomagnetic beads for high-efficient capture and release of exosomes' by Jia Cheng et al., J. Mater. Chem. B, 2022, 10, 4059-4069, https://doi.org/10.1039/ D2TB00226D.

The authors regret that due to file compilation errors, Fig. 4D and F, and Fig. 6J-L were incorrect in the manuscript. The corrected versions of Fig. 4 and 6 are provided below.



 $Fig. 4 \quad \text{CLSM images of 500 } \mu g \text{ of (A) IMHPs } (\text{TiO}_2 @ \text{Fe}_3 \text{O}_4 - \text{anti-CD63 } \text{HPs)}, \text{ (B) } \text{TiO}_2 @ \text{Fe}_3 \text{O}_4 - \text{biotin } \text{HPs, (C) } \text{SiO}_2 @ \text{Fe}_3 \text{O}_4 - \text{anti-CD63 } \text{MPs } \text{and } \text{SiO}_2 @ \text{Fe}_3 \text{O}_4 - \text{anti-CD63 } \text{MPs}$ (D) $SiO_2@Fe_3O_4$ -biotin MPs after incubation with 2 μ g of FITC-labeled goat anti-rabbit IgG in 500 μ L of PBS at room temperature for 2 h. 200 μ L of TCEP (20 mM)-induced disulfide cleavage occurred at 37 °C for 30 min to fade away the corresponding fluorescence signals of the FITC-labeled goat anti-rabbit IgG on (E) IMHPs and (F) $SiO_2@Fe_3O_4$ -anti-CD63 MPs (excitation 488 nm, emission 520 nm).

Additionally, the figure captions of Fig. 3(E) and (F), and Fig. 8 were incorrect. The corrected versions should read: Fig. 3 (E) TiO₂@Fe₃O₄-biotin HPs and (F) SiO₂@Fe₃O₄-biotin MPs (excitation 488 nm, emission 520 nm). Fig. 8 Capture efficiency of exosomes (5.1 × 10⁸ particles per mL, 200 μL) incubated with varying amounts of (A) IMHPs, (B) TiO₂@Fe₃O₄-biotin HPs and (C) SiO₂@Fe₃O₄-anti-CD63 MPs at room temperature for 1 h.

National Engineering Research Center for Biomaterials, Sichuan University, Chengdu 610064, P. R. China. E-mail: qyi@scu.edu.cn, wuyao@scu.edu.cn




Fig. 6 Verification of the feasibility of exosome enrichment. After incubating the material and exosomes (pre-stained with PKH26) for 1 h, the CLSM $images\ of\ (A-C)\ TiO_2@Fe_3O_4-anti-CD63\ HPs\ +\ exosome,\ (D-F)\ SiO_2@Fe_3O_4-anti-CD63\ MPs\ +\ exosome,\ (G-I)\ SiO_2@Fe_3O_4-biotin\ MPs\ +exosome,\ (D-F)\ SiO_2@Fe_3O_4-biotin\ MPs\ +exosome$ and (J-L) $TiO_2@Fe_3O_4$ -biotin HPs + exosome were recorded (excitation 551 nm, emission 567 nm).

An independent expert has viewed the original and new images and has concluded that they are consistent with the discussions and conclusions presented.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.