



Cite this: DOI: 10.1039/d6bm90010k

## Correction: Tumor-triggered personalized microRNA cocktail therapy for hepatocellular carcinoma

Shiyi Shao,<sup>a</sup> Qida Hu,<sup>a</sup> Wangteng Wu,<sup>a,b</sup> Meng Wang,<sup>a,b</sup> Junming Huang,<sup>a</sup>  
Xinyu Zhao,<sup>a</sup> Guping Tang<sup>b</sup> and Tingbo Liang<sup>\*a,c,d</sup>


DOI: 10.1039/d6bm90010k

rsc.li/biomaterials-science

Correction for 'Tumor-triggered personalized microRNA cocktail therapy for hepatocellular carcinoma' by Shiyi Shao et al., *Biomater. Sci.*, 2020, **8**, 6579–6591, <https://doi.org/10.1039/D0BM00794C>.

The authors regret an error in Fig. 2a and in Fig. S3c.

In Fig. 2a the incorrect image was used for Huh7 Lipo3000/miRFAM+ and the corrected figure is shown here.



**Fig. 2** PCACP mediated cellular uptake and transfection both *in vitro* and *in vivo*. (A) Confocal images of naked miR199<sup>FAM+</sup>, PCACP/miR199<sup>FAM+</sup> pre-incubated at pH 6.5 or pH 7.4, PCAP/miR199<sup>FAM+</sup>, and Lipo3000/miR199<sup>FAM+</sup> after 4 h of incubation in HEK 293 and Huh 7 cells. miRNA was labeled with FAM (green). Cytomembrane was stained with Alexa Fluor® 647 conjugates of wheat germ agglutinin (red), while the nucleus was stained with DAPI (blue). The bar represents 20 μm.

<sup>a</sup>Department of Hepatobiliary and Pancreatic Surgery, First Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China.  
E-mail: liangtingbo@zju.edu.cn


<sup>b</sup>Institute of Chemistry Biology and Pharmaceutical Chemistry, Zhejiang University, Hangzhou 310028, China

<sup>c</sup>Zhejiang Provincial Innovation Center for the Study of Pancreatic Disease, Hangzhou 310003, China

<sup>d</sup>Zhejiang Provincial Key Laboratory of Pancreatic Disease, Hangzhou 310003, China



In Fig. S3c the incorrect image was used for PCAP/VecGFP+ pH 6.5 and the corrected figure is shown here.



**Fig. S3c** Overlapped images of bright field and fluorescent photo of HEK 293 and Huh 7 cells after transfection with naked plasmids, and plasmids loaded PCACP (pH 6.5 and pH 7.4), PCAP and Lipo3000 for 48 h. The plasmid vectors encoded GFP fluorescence (VecGFP+). The bar represents 20  $\mu\text{m}$ . Representative images in 3 replicates.

An independent expert has viewed the corrected figures and confirmed that they are consistent with the discussions and conclusions presented.

The Royal Society of Chemistry apologises for these errors and any consequent inconvenience to authors and readers.

