



Cite this: DOI: 10.1039/d6cc00157b

 Received 8th January 2026,
 Accepted 16th April 2026

DOI: 10.1039/d6cc00157b

rsc.li/chemcomm


Cu(OTf)₂-catalyzed access to 2,4,5-trisubstituted imidazoles from acyclic Reissert compounds

 Swetha Sathyendran,^a Vikraman Ganesh Moorthi,^a Sharmila Nokku,^a Aron Manick Joel,^a Suryanarayanan Chandrasekaran,^a Wei-Yu Lin^b and Gopal Chandru Senadi^{id}*^{ab}

A Cu(OTf)₂-catalyzed cascade self-coupling of *N*-acyl- α -aminonitriles (acyclic Reissert compounds) has been developed to afford 2,4,5-trisubstituted imidazoles in moderate to good yields. Single-crystal X-ray diffraction confirmed the structure of a representative 1*H*-2,4-disubstituted-5-acylimidazole, while NMR studies revealed N1/N3 tautomerism in selected cases. The synthetic utility of this protocol was further demonstrated by gram-scale synthesis and carbonyl reduction.

Imidazole, a privileged core structure, finds diverse applications in medicinal chemistry,¹ natural products,² and drug discovery.³ Substitution on the imidazole ring further extends its utility to functional materials,^{4a,b} polymers,^{4c,d} and coordination chemistry,^{4e,f} spurring interest in 2,4,5-trisubstituted imidazoles.⁵ Traditional syntheses rely on the Radziszewski multicomponent cyclocondensation of aldehydes, ammonia, and 1,2-diketones,^{4g,h} with numerous refinements enabling substituted analogues.⁵ However, methods for 5(4)-acylimidazoles bearing ketones or esters remain underexplored.^{6,7} For instance, Muthusubramanian and coworkers (2014) reported an Er(OTf)₃-catalyzed multicomponent reaction of α -azido chalcones, aryl aldehydes, and anilines (Scheme 1a).^{7a} Li and coworkers (2015) developed an FeCl₃/I₂-catalyzed aerobic oxidative coupling of amidines and chalcones for regioselective acylimidazoles (Scheme 1b).^{7e} Guchhait and coworkers (2017) devised a tandem aza-Michael addition/S_N2 redox-neutral process from α -bromo enones (Scheme 1c).^{7e} With such limited precedents, substantial opportunities remain for new approaches to C-acylimidazoles.

More than 160 years after their discovery,⁸ α -aminonitriles remain powerful synthetic linchpins, undergoing transformations⁹ *via* deprotonation or decyanation to iminium ions, as demonstrated in our amide and 2,4-diarylquinoline syntheses.^{10a-c} These findings prompted us to explore *N*-acyl- α -aminonitriles,^{10d} an open-chain



Scheme 1 Prior art on acylimidazoles and *N*-acyl- α -aminonitriles, and this work.

class of Reissert analogues. The 1,4- and 1,2-addition reactions of deprotonated Reissert compounds and their open-chain analogues have been widely exploited for the construction of diverse molecular frameworks.¹¹ The synergistic electrophilicity of the nitrile group and nucleophilicity of the amine functionality has enabled versatile access to *N*-heterocycles from these bifunctional scaffolds. In this context, cyclization reactions incorporating the nitrile moiety,^{12a-c} as well as the use of electron-deficient arenes in place of classical electrophiles,^{12d} further expanded the synthetic potential of *N*-acyl- α -aminonitriles.

In a seminal contribution, McEwen and co-workers developed a condensation of the conjugate base of *N*-acyl- α -aminonitriles with vinyltriphenylphosphonium bromide for the synthesis of substituted pyrroles (Scheme 1d).^{11ef} Later, Zhong *et al.* reported the synthesis of

^a Green and Sustainable Synthesis Laboratory, Department of Chemistry, College of Engineering and Technology, SRM Institute of Science and Technology, SRM Nagar, Kattankulathur - 603 203, Chengalpattu District, Tamil Nadu, India. E-mail: chandrug@srmist.edu.in


^b Department of Medicinal and Applied Chemistry, Kaohsiung Medical University, Kaohsiung, Taiwan



2,4-disubstituted 5-halo-1*H*-imidazoles from *N*-acyl- α -aminonitriles, which were subsequently converted into 2,4,5-trisubstituted imidazoles *via* Pd-catalyzed cross-coupling (Scheme 1e).^{12e} Nearly two decades later, we describe a simple transformation of the same *N*-acyl- α -aminonitriles into C-acylimidazoles in a single step using catalytic Cu(OTf)₂ (Scheme 1f). Our initial hypothesis targeted the synthesis of 5-amino oxazole *via* a Lewis acid-promoted 5-*exo*-dig cyclization; however, the reaction selectively delivered imidazoles through a self-coupling cascade. Given the limited availability of efficient methods for synthesizing 5(4)-acyl-substituted imidazoles, this conversion provides a useful route and highlights a different reactivity pathway of *N*-acyl- α -aminonitriles.

The optimization began with *N*-(cyano(phenyl)methyl)benzamide **1a** as the model substrate. Screening of common metal halides in 1,4-dioxane at 90 °C (Table 1, entries 1–5) delivered the desired product **2a** in modest yields (17–63%). Subsequent evaluation of various metal triflates (Table 1, entries 6–10) identified Cu(OTf)₂ as the most effective catalyst, providing **2a** in 73% yield at 20 mol% loading (Table 1, entry 7). With copper triflate selected, solvent screening was carried out next. Replacing 1,4-dioxane with ethanol, isopropanol (IPA), ethylene glycol, ethyl acetate, or acetonitrile (Table 1, entries 11–15) failed to improve the reaction efficiency. Catalyst loading studies showed that reducing Cu(OTf)₂ to 15 mol% lowered the yield (Table 1, entry 16), while increasing it to 25 mol% offered no further enhancement (Table 1, entry 17). Temperature variation (Table 1, entries 18 and 19) confirmed that 90 °C was optimal for the formation of **2a**.

Control experiments under N₂ and O₂ were performed to probe the reaction pathway (SI, Table S4). Under N₂, the reaction gave only



Scheme 2 Control experiments to study the reaction pathway.

29% yield of **2a** without water, which increased to 56% and 68% upon addition of 5 and 10 equiv. of H₂O, respectively, indicating the role of water (Scheme 2a). Under O₂, the reaction afforded **2a** in 58% yield with 30% of decyanative product **2a'** in the absence of water, while similar yields of **2a** (57%) and reduced **2a'** (24%) were observed in the presence of water, indicating competing oxygen-mediated pathways (Scheme 2b). Next, the reaction was performed¹³ in H₂¹⁸O using dry 1,4-dioxane, and LC-MS analysis showed a mixture of isotopic and non-isotopic mass peaks (SI, Fig. S10). The non-isotopic mass suggests that oxygen may also arise from water generated *in situ* during the course of the reaction and from residual moisture (Scheme 2c). Further, cross-over experiments with **1a/1b** (electron-donating) and **1a/1f** (halogen-substituted) under standard conditions afforded only the corresponding products **2a/2b** and **2a/2f**, respectively (Scheme 2d and e). No heterocoupled products were detected, indicating exclusive homocoupling (SI, Fig. S8 and S9). In the presence of TEMPO, the yield of **2a** decreased to 33%, along with formation of hydrolysed product **2''** in 41% yield likely due to the Cu/TEMPO system. In contrast, BHT afforded **2a** in 70% yield (*cf.* 73% standard). These results suggest that the reaction does not proceed *via* a radical pathway (Scheme 2f).

With the optimized conditions and control studies, the substrate scope of *N*-acyl- α -aminonitriles was examined (Table 2). A range of *para*-substituted phenyl derivatives, including *p*-Me (**1b**), *p*-Et (**1c**), *p*-OMe (**1d**), *p*-F (**1e**), *p*-Cl (**1f**), *p*-Br (**1g**), *p*-COOMe (**1h**), *p*-CF₃ (**1i**), and *p*-CN (**1j**), were well tolerated, affording the corresponding products **2b–2j** in 60–78% yield. *Meta*- and *ortho*-substituted substrates such as *m*-Me (**1k**), *m*-OMe (**1l**), and *o*-Me (**1m**) also reacted smoothly, affording **2k–2n** in 65–66% yield. Whereas *o*-OMe (**2n**) was observed only in trace amounts and could not be isolated. Polysubstituted


Table 1 Reaction optimization studies^{abc}

S. no	Catalyst (x mol%)	Solvent	Temp. (°C)/time (h)	Yield (%)
1	FeCl ₃ (20)	1,4-dioxane	90/12	63
2	CoCl ₂ ·6H ₂ O (20)	1,4-dioxane	90/24	17
3	CuCl ₂ (20)	1,4-dioxane	90/24	44
4	CuBr ₂ (20)	1,4-dioxane	90/16	56
5	ZnCl ₂ (20)	1,4-dioxane	90/24	31
6	Fe(OTf) ₂ (20)	1,4-dioxane	90/12	59
7	Cu(OTf)₂ (20)	1,4-dioxane	90/08	73
8	Sm(OTf) ₃ (20)	1,4-dioxane	90/24	48
9	AgOTf (20)	1,4-dioxane	90/24	54
10	Yb(OTf) ₃ (20)	1,4-dioxane	90/24	51
11	Cu(OTf) ₂ (20)	Ethanol	80/24	42
12	Cu(OTf) ₂ (20)	IPA	80/24	38
13	Cu(OTf) ₂ (20)	Ethylene glycol	100/12	61
14	Cu(OTf) ₂ (20)	Ethyl acetate	80/24	34
15	Cu(OTf) ₂ (20)	Acetonitrile	80/24	19
16	Cu(OTf) ₂ (15)	1,4-dioxane	90/24	64
17	Cu(OTf) ₂ (25)	1,4-dioxane	90/8	72
18	Cu(OTf) ₂ (20)	1,4-dioxane	70/24	58
19	Cu(OTf) ₂ (20)	1,4-dioxane	110/8	74

^a Reaction conditions: all reactions were carried out using **1a** (0.5 mmol), solvent (0.15 M) and catalyst (x mol%) in a sealed vial at indicated temperature and time unless otherwise noted. ^b Isolated yield. ^c See the SI (Tables S1–S4) for the detailed optimization studies.




Table 2 Scope and limitations of *N*-acyl- α -aminonitriles to 2,4,5-trisubstituted imidazoles^{a,b}



^a Reaction conditions: Compound **1a** (0.5 mmol), Cu(OTf)₂ (20 mol %), and 1,4-dioxane (0.15 M) were stirred at 90 °C for 7–20 h. ^b Isolated yields. ^c The presence of N1–N3 tautomeric forms was confirmed by NMR analysis. ^d Detected in trace amounts by TLC; not isolated due to multiple spots. ^e **2s** was detected in trace amounts and not isolated; the corresponding amide was obtained. ^f The mass was confirmed by HRMS analysis; however, isolation for pure NMR characterization was unsuccessful.

aromatics, including 3,4,5-tri-OMe (**1o**) and 3,4,5-trifluoro (**1p**), were compatible, producing **2o** and **2p** in 71% and 76% yield, respectively, with the structure of **2p** confirmed by single-crystal X-ray analysis.¹⁴ Fused aromatics and heteroaryl substrates, such as naphthyl (**1q**) and 2-thienyl (**1r**), furnished **2q** and **2r** in 62–68% yields. Substrates with aryl groups attached to the α -carbon, including *o*-Br (**1s**), *p*-OMe (**1t**), and 2-thienyl (**1u**), gave **2t** in 54% yield. For **1s**, only traces of **2s** were observed by TLC, and the major isolated product was the corresponding hydrolysed amide **2s''** in 52% yield. Similarly, **2u** was confirmed by HRMS analysis, while its isolation was unsuccessful.



Scheme 3 Synthetic applications and control studies.


Overall, *para*-substituted substrates generally gave higher yields and shorter reaction times, *meta*-substituents gave moderate results, and *ortho*-substitution lowered yields and required longer reactions, likely due to steric effects.

We further evaluated the scalability of the methodology on a gram-scale using the standard optimized conditions. Representative substrates bearing an electron-donating substituent (**1d**) and an electron-withdrawing substituent (**1h**) furnished the corresponding products **2d** and **2h** in moderate yields of 61% and 65%, respectively (Scheme 3a & b). In addition, reduction of carbonyl functional group in **2a** with NaBH₄ proceeded smoothly to afford the corresponding secondary alcohol **3a** in 84% yield (Scheme 3c).

A plausible mechanism, supported by control experiments and literature precedent,^{11,12,15} is outlined in Scheme 4. Coordination of Cu(OTf)₂ to the nitrile group of α -aminonitrile **1** generates intermediate **A**, increasing the electrophilicity of the nitrile carbon and enabling intramolecular 5-*exo*-dig cyclization to form imino-oxazolone **B**. DFT calculations using UB3LYP/def2-TZVP with a 1,4-dioxane solvent model (see SI for computational details) support preferential nitrogen coordination ($\Delta G = -8.04$ kcal mol⁻¹) over oxygen ($\Delta G = -3.35$ kcal mol⁻¹), with HOMO–LUMO overlap and a smaller band gap (2.13 eV) favouring the nitrile binding pathway (SI, Fig. S13). Further, to assess the ring-closure mechanism and stability of cyclized intermediate **B**, transition state free energy calculations were performed (see SI, Fig. S15). The process proceeds *via* transition state **A** with an activation free energy of $\Delta G^\ddagger = 23.4$ kcal mol⁻¹, leading to intermediate **B**, which is thermodynamically stabilized by -4.46 kcal mol⁻¹ relative to the initial reactant **1**.¹⁶ Next, nucleophilic addition of a second molecule of **1** to **B**, followed by ring opening, affords intermediate **C**. Owing to the lability of Cu(II) coordination, intramolecular exchange between imine and amide carbonyl binding facilitates cyclization to imidazoline **D**. Dehydration of **D** regenerates Cu(OTf)₂ and furnishes intermediate **E**, which upon isomerization and HCN elimination gives **F**. Proton transfer affords aromatic intermediate **G**, and final hydrolysis yields the 2,4,5-trisubstituted imidazole **2** along with amide **4**, confirmed by ¹H NMR spectroscopy (SI, Fig. S57).

In conclusion, a copper-catalyzed synthesis of *C*-acylimidazoles from open-chain analogues of Reissert compound is disclosed. The reaction is proposed to proceed through an imino-oxazolone





Scheme 4 Plausible reaction mechanism.

pathway, enabling access to 5-acylimidazole scaffolds. Broad aryl-substrate tolerance, gram-scale feasibility, and reduction of the carbonyl to the corresponding alcohol demonstrate the practicality of the method. A plausible mechanism was proposed based on control studies and DFT calculations. This work expands the synthetic toolbox for *N*-acyl- α -aminonitriles and provides a useful entry to acylimidazoles.

Conflicts of interest

There are no conflicts to declare.

Data availability

The supporting data has been provided as part of the supplementary information (SI). Supplementary information: Tables S1–S4 and Fig. S1 and S2. NMR spectra and further experimental details. See DOI: <https://doi.org/10.1039/d6cc00157b>.

CCDC 2518062 contains the supplementary crystallographic data for this paper.¹⁴

Acknowledgements

S. S., V. G. M., S. N and A. M. J., thank the SRM Institute of Science and Technology for the PhD fellowship. Dr G.C.S acknowledges Anusandhan National Research Foundation (ANRF) for Core Research Grant (File No: CRG/2022/006963). The authors wish to thank department of chemistry and Interdisciplinary Institute of Indian System of Medicine (IIISM), SRM Institute of Science and Technology for providing HRMS, NMR

and computational HPCC facilities. The authors also thank Prof. Jeh-Jeng Wang, Kaohsiung Medical University, Kaohsiung, Taiwan for providing support for SCXRD and HRMS analysis.

References

- (a) G. Yadav and R. Jain, *Eur. J. Med. Chem.*, 2025, **290**, 117524; (b) L. Zhang, X.-M. Peng, G. L. Damu, R.-X. Geng and C.-H. Zhou, *Med. Res. Rev.*, 2014, **34**, 340–437; (c) S. S. Alghamdi, R. S. Suliman, K. Almutairi, K. Kahtani and D. Aljatl, *Drug Des., Dev. Ther.*, 2021, **15**, 3289–3312; (d) N. Rani, A. Sharma and R. Singh, *Mini Rev. Med. Chem.*, 2013, **13**, 1812–1835.
- (a) Z. Jin, *Nat. Prod. Rep.*, 2016, **33**, 1268–1317; (b) M. R. U. Karim, E. Harunari, N. Oku, K. Akasaka and Y. Igarashi, *J. Nat. Prod.*, 2020, **83**, 1295–1299.
- (a) S. Gupta, M. A. Babu, R. Kumar, T. G. Singh, A. Goel, S. Rastogi, P. Sharma, Y. Tyagi, K. K. Goel and B. Kumar, *Chem. Biodivers.*, 2025, **22**, e202403020; (b) M. Gaba and C. Mohan, *Med. Chem. Res.*, 2016, **25**, 173–210.
- (a) Y. Yao, Z. Li, Y. Zhang, K. Xie, Y. Song and Y. Tang, *ACS Sens.*, 2025, **10**, 8489–8497; (b) J. Wang, X. Ding, Z. Lan, G. Liu, S. Hou and S. Hou, *Crit. Rev. Anal. Chem.*, 2024, **54**, 897–922; (c) S. Bureekaew, S. Horike, M. Higuchi, M. Mizuno, T. Kawamura, D. Tanaka, N. Yanai and S. Kitagawa, *Nat. Mat.*, 2009, **8**, 831–836; (d) C. Fodor, J. Bozi, M. Blazso and B. Ivan, *Macromolecules*, 2012, **45**, 8953–8960; (e) S.-S. Chen, *CrysEngComm*, 2016, **18**, 6543–6565; (f) X.-W. Zhu, D. Luo, X.-P. Zhou and D. Li, *Coord. Chem. Rev.*, 2022, **455**, 214354; (g) B. Radziszewski, *Ber. Dtsch. Chem. Ges.*, 1882, **15**, 1493–1496; (h) W. Rudolf and H. Roland, *Ber. Dtsch. Chem. Ges. (A and B Series)*, 1935, **68**, 1953–1961.
- (a) D. A. Shabalin and J. E. Camp, *Org. Biomol. Chem.*, 2020, **18**, 3950–3964; (b) G. Patel, D. K. Dewangan, N. Bhakat and S. Banerjee, *Curr. Res. Green Sustainable Chem.*, 2021, **4**, 100175.
- (a) B. Hu, Z. Wang, N. Ai, J. Zheng, X.-H. Liu, S. Shan and Z. Wang, *Org. Lett.*, 2011, **13**, 6362–6365; (b) H. Ma, X. Zhang, L. Chen and W. Yu, *J. Org. Chem.*, 2017, **82**, 11841–11847; (c) A. Anghyal, A. Demjén, J. Wöfling, L. G. Puskás and I. Kanizsai, *J. Org. Chem.*, 2020, **85**, 3587–3595; (d) S. M. Patel, P. E. Prasad, M. Bakthadoss and D. S. Sharada, *Org. Lett.*, 2021, **23**, 257–261.
- (a) K. Rajaguru, R. Suresh, A. Mariappan, S. Muthusubramanian and N. Bhuvanesh, *Org. Lett.*, 2014, **16**, 744–747; (b) Y. Zhu, C. Li, J. Zhang, M. She, W. Sun, K. Wan, Y. Wang, B. Yin, P. Liu and J. Li, *Org. Lett.*, 2015, **17**, 3872–3875; (c) S. Yugandar, S. Konda, G. Parameshwarappa and H. Ila, *J. Org. Chem.*, 2016, **81**, 5606–5622; (d) P. R. Adiyappa, S. Borra, A. Kamal and R. A. Maurya, *Eur. J. Org. Chem.*, 2016, 1269–1273; (e) S. K. Guchhait, N. Hura and A. P. Shah, *J. Org. Chem.*, 2017, **82**, 2745–2752; (f) M. B. Harisha, P. Dhanalakshmi, R. Suresh, R. R. Kumar, S. Muthusubramanian and N. Bhuvanesh, *ChemistrySelect*, 2019, **4**, 2954–2958; (g) P. Wu, X. Zhang and B. Chen, *Tetrahedron Lett.*, 2019, **60**, 1103–1107; (h) A. Takeda, H. Okai, K. Watabe and H. Iida, *J. Org. Chem.*, 2022, **87**, 10372–10376.
- A. Strecker, *Liebigs Ann. Chem.*, 1850, **75**, 27–45.
- (a) C. Grundke, N. Vierengel and T. Opatz, *Chem. Rec.*, 2020, **20**, 989–1016; (b) V. V. Kouznetsov and C. E. P. Galvis, *Tetrahedron*, 2018, **74**, 773–810; (c) N. Otto and T. Opatz, *Chem. – Eur. J.*, 2014, **20**, 13064–13077; (d) T. Opatz, *Synthesis*, 2009, 1941–1959; (e) H. Gröger, *Chem. Rev.*, 2003, **103**, 2795–2828; (f) D. Enders and J. P. Shilcock, *Chem. Soc. Rev.*, 2000, **29**, 359–373.
- (a) S. Swetha and G. C. Senadi, *Adv. Synth. Catal.*, 2022, **364**, 2872–2882; (b) S. Sathyendran, K. Muthu, K. Govindan, N.-Q. Chen, W.-Y. Lin and G. C. Senadi, *Org. Lett.*, 2023, **25**, 4086–4091; (c) S. Sathyendran and G. C. Senadi, *Asian J. Org. Chem.*, 2023, **12**, e202300433; (d) V. G. Moorthi, V. T. Seenivasan, S. Nokku, A. M. Joel, W.-Y. Lin and G. C. Senadi, *Chem. Commun.*, 2026, **62**, 5477–5481.
- (a) V. Boekelheide and J. C. Godfrey, *J. Am. Chem. Soc.*, 1953, **75**, 3679–3685; (b) F. D. Popp, L. E. Katz, C. W. Klinowski and J. M. Wefer, *J. Org. Chem.*, 1968, **33**, 4447–4450; (c) B. C. Uff and R. S. Budhram, *Synthesis*, 1978, 206; (d) F. D. Popp, *Adv. Heterocycl. Chem.*, 1979, **24**, 187–214; (e) J. V. Cooney and W. E. McEwen, *J. Org. Chem.*, 1981, **46**, 2570–2573; (f) J. V. Cooney, B. D. Beaver and W. E. McEwen, *J. Heterocycl. Chem.*, 1985, **22**, 635–642; (g) J.-P. Leblanc and H. W. Gibson, *J. Org. Chem.*, 1994, **59**, 1072–1077.
- (a) M. Uchibayashi, *Yakugaku Zasshi*, 1958, **78**, 845–849; (b) E. A. Younes, A. Q. Hussein, M. A. May and F. R. Fronczek,



- ARKIVOC, 2011, 2, 322–330; (c) M. Sedlák, R. Keder, J. Hanusek and A. Růžička, *J. Heterocycl. Chem.*, 2009, 42, 899–906; (d) G. Arnott, J. Clayden and S. D. Hamilton, *Org. Lett.*, 2006, 8, 5325–5328; (e) Y.-L. Zhong, J. Lee, R. A. Reamer and D. Askin, *Org. Lett.*, 2004, 6, 929–931.
- 13 For detailed isotopic studies with H₂¹⁸O refer supporting information Fig. S10 to Fig. S12.
- 14 CCDC 2518062: Experimental Crystal Structure Determination, 2026, DOI: [10.5517/ccdc.csd.cc2qj7tb](https://doi.org/10.5517/ccdc.csd.cc2qj7tb).
- 15 (a) W. E. McEwen, A. V. Grossi, R. J. MacDonald and A. P. Stamegna, *J. Org. Chem.*, 1980, 45, 1301–1308; (b) J. S. Fisk, R. A. Mosey and J. J. Tepe, *Chem. Soc. Rev.*, 2007, 36, 1432–1440.
- 16 N. Dhama, M. K. Tiwari, V. K. Vishvakarma, R. Yadav, M. Kumar and D. T. Masram, *Inorg. Chim. Acta*, 2026, 589, 122928.

