



Cite this: *Chem. Commun.*, 2026, 62, 6183

Received 12th January 2026,
Accepted 25th February 2026

DOI: 10.1039/d6cc00213g

rsc.li/chemcomm

Light-driven silane functionalization using disulfide and dichloromethane

Jiin Lee,^{†ab} Hyunji Kim^{†a} and Sunggi Lee^{ib}*^{ab}


Organosilicon compounds are of great importance in material and synthetic chemistry. Herein, we disclose a practical and general silane functionalization protocol that enables direct access to silanols and silyl ethers. The process features mild, environmentally benign conditions and enables the transformation of diverse silanes into valuable organosilicon compounds.

Organosilicon compounds have been broadly utilized as valuable building blocks in material science¹ and organic synthesis (Scheme 1).² Among them, silanols and silyl ethers find wide applications in silicon-based materials, metallasiloxane synthesis,³ C–H activation,⁴ organocatalysis,⁵ and medicinal chemistry.⁶ Motivated by the applications of organosilicon compounds, research has been actively conducted in the field of organic synthesis to develop methods for synthesizing these compounds. These are typically achieved through the hydrolysis of silyl chlorides⁷ or the oxidation of silanes with stoichiometric oxidants such as permanganate, osmium tetroxide, dioxirane, and ozone.⁸ However, the hydrolysis of silyl chlorides often suffers from low efficiency owing to the condensation of silanols to siloxanes, while oxidation of silanes with these reagents is limited by narrow substrate scope, poor functional group tolerance, and the generation of substantial waste. To address issues, transition-metal-catalyzed⁹ or photocatalytic¹⁰ methods have emerged as an attractive approach. However, most reported methods rely on noble metal catalysts and often suffer from the undesired formation of disiloxanes, arising from competitive reactions of silanols with hydrosilanes (Fig. 1A). In 2019, Wu and co-workers disclosed a photocatalytic method employing eosin Y, where hydrosilanes undergo chlorination and subsequent reaction with alcohols to afford the corresponding silyl ethers (Fig. 1B).¹¹ Although significant progress has been made, practical and robust methods for

accessing silanols and silyl ethers still limited. We hypothesized that the *in situ* generation of silyl chlorides from silanes under mild conditions would provide a versatile precursor, enabling their subsequent conversion into silanols and silyl ethers through reaction with oxygen nucleophiles.

Herein we report a practical, photocatalyst-free method for silane functionalization using disulfides and dichloromethanes under purple light irradiation (Fig. 1C). In our design, thiyl radicals were generated *in situ* from disulfides without the need for additional catalysts and efficiently engaged in hydrogen atom transfer (HAT) to form silyl radicals.¹² The resulting silyl radicals then underwent halogen atom transfer (XAT) with dichloromethane, thereby furnishing silyl chlorides under mild conditions. Furthermore, the *in situ* generated silyl chlorides can be selectively converted into either silanols (with water) or silyl ethers (with alcohols), offering modular access to multiple classes of valuable organosilicon compounds from the same hydrosilane precursor. This method provides an environmentally benign alternative to existing photocatalytic approaches, relying exclusively on inexpensive commercial reagents and expanding the synthetic utility of hydrosilanes.

We started our investigation of silanol synthesis using diphenylmethyl silane (**1a**) as a model substrate. After the



Scheme 1 Useful organosilicon compounds.


^a Department of Physics and Chemistry, DGIST, Daegu 42988, Republic of Korea.
E-mail: sunggi.lee@dgist.ac.kr

^b Center for Basic Science, DGIST, Daegu 42988, Republic of Korea


[†] These authors contributed equally to this work.



A Transition-metal catalyzed oxidation of silanes



B Wu's Eosin Y-catalyzed access to silyl ether



C This work




Fig. 1 Previous works and this work. (A) Transition-metal catalyzed oxidation of silanes. (B) Wu's Eosin Y-catalyzed access to silyl ether. (C) This work.

systemic screening, the optimal conditions were obtained as follows (see SI for further information): **1a** (1.0 equiv.), dicyclohexyl disulfide (**D1**, 20 mol%), H₂O (40 equiv.) in a solution of CH₂Cl₂ at room temperature under 390 nm irradiation for 24 hours (Table 1, entry 1). Under these conditions, the desired silanol **1b** was obtained in 72% isolated yield. Based on this result, several control experiments were conducted. Reducing the disulfide loading from 5 mol% to 1 mol%, the reaction efficiency was significantly dropped (entry 2). Additionally, diphenyl disulfide (**D2**) was not efficient for product formation due to the low BDE of aryl thiols (*ca.* 79 kcal mol⁻¹) (entry 3).¹³ When CH₂Cl₂ was replaced with other chlorine sources, including *n*-BuCl or 1,2-dichloroethane (DCE) or CH₃Cl, the yield decreased significantly (entry 4–6). Replacing 390 nm with 440 nm did not afford the desired product (entry 7). Without water, only trace amounts of silanol were observed, while silyl chloride was obtained in 82% yield (entry 8).

Table 1 Control experiments

Entry	Deviation from standard conditions ^a	Yield ^b [%]
1	None	97 (72 ^c)
2	1 mol% of D1	3
3	D2 instead of D1	8
4	<i>n</i> -BuCl instead of CH ₂ Cl ₂	10
5	DCE instead of CH ₂ Cl ₂	38
6	CHCl ₃ instead of CH ₂ Cl ₂	43
7	440 nm instead of 390 nm	Trace
8	Without H ₂ O	Trace ^d

^a Reaction conditions: **1a** (0.2 mmol), **D1** (5 mol%), H₂O (40 equiv.), CH₂Cl₂ (0.2 M), 390 nm irradiation at r.t. for 24 h. ^b Yield was determined by ¹H NMR spectroscopy using CH₂Br₂ as an internal standard. ^c Isolated yield. ^d Silyl chloride was obtained.

Table 2 Substrate scope for synthesis of silanols^a

2a , 95% (72%)	2b , 78% (74%) ^b	2c , 95% (89%)	2d , 97% (76%)	2e , 72%
2f , 97%	2g , 96%	2h , 90% (64%) ^c	2i , 96% (71%)	

^a Reaction conditions: **1** (0.2 mmol), **D1** (5 mol%), H₂O (40 equiv.) in DCM (0.2 M) at r.t. under 390 nm irradiation. ^b 48 h. ^c Addition of 1.5 equiv. of imidazole for 72 h. Yield determined by ¹H NMR spectroscopy using CH₂Br₂ as an internal standard; isolated yields are shown in parentheses.

With the optimized conditions in hand, various silanes were investigated to access corresponding silanols (Table 2). In general, a broad range of silanes bearing aryl, alkyl, and silyl substituents were smoothly converted into the corresponding silanols. Mono-, di- and triaryl substituted silanes afforded the desired product (**2a–2d**) with high yield. In addition, trialkylsilanes including *t*BuMe₂SiH, Et₃SiH, and *n*Bu₃SiH, were well tolerated, giving the excellent yield (**2e–2g**). Plus, sterically demanding *i*Pr₃SiH furnished silanol **2h** in 87% yield. Furthermore, tris(trimethylsilyl)silane (supersilane) also underwent smooth conversion to the silanol in 96% yield (**2i**), highlighting generality of this protocol.

Encouraged by the success of silanol formation, we sought to extend the protocol to the synthesis of silyl ethers using alcohols as nucleophiles. Reaction optimization revealed that the presence of alcohols together with silanes (1.5 equiv.) and with 1.5 equiv. of imidazole afforded corresponding silyl ethers (see SI for further optimization, Table S5). As shown in Table 3,

Table 3 Substrate scope for synthesis of silyl ethers^a

3a , 95% (76%)	3b , 98% (93%)	3c , 93% (87%)	3d , 99% (83%) ^b
3e , 92% (76%)	3f , 93% (90%)	3g , 99% (83%) ^c	3h , 43% ^b

^a Reaction conditions: 2-phenylethanol (1.0 equiv.), **1** (1.5 equiv.), imidazole (1.5 equiv.) **D1** (5 mol%) in CH₂Cl₂ (0.2 M) at r.t. under 390 nm irradiation. ^b 48 h. ^c 72 h. Yield determined by ¹H NMR spectroscopy using CH₂Br₂ as an internal standard; isolated yields are shown in parentheses.






Fig. 2 (A) Radical trapping experiments. ^aTris(trimethylsilyl)silane (1.0 equiv.), **D1** (5 mol%), ethyl acrylate (1.0 equiv.) in MeCN (0.2 M) at r.t. under 390 nm irradiation. (B) Plausible mechanism.

a wide range of silanes were compatible with this transformation, providing the desired silyl ethers. Several aryl-substituted silanes gave silyl ether with excellent yield (**3a–3c**). Also, trialkyl substituted silyl ether (**3d–3g**) were obtained in high yields. Additionally, supersilane underwent smooth transform to provide **3h** in a moderate yield.

Mechanistic studies were carried out to elucidate reaction mechanism (Fig. 2). Radical trapping experiments using ethyl acrylate resulted in the formation of the corresponding hydro-silylation product (**5a**), providing direct evidence for the generation of a silyl radical intermediate under the photochemical conditions (Fig. 2A). A plausible mechanism for this reaction is proposed in Fig. 2B. Under 390 nm purple light irradiation, the S–S bond in dicyclohexyl disulfide undergoes homolytic cleavage,¹³ generating thiyl radicals to produce a silyl radical through HAT. The silyl radical can undergo XAT with dichloromethane to generate silyl chloride. In the presence of H₂O or alcohol, the reaction furnishes the corresponding silanol or silyl ether. The resulting chloromethyl radical can abstract a hydrogen atom from thiol, thereby generating a thiyl radical and propagating the radical chain.

In conclusion, we have developed a simple and general method for functionalization of silanes using commercially available and environmentally benign reagents. This transformation proceeds under mild reaction conditions to produce various organosilicon compounds including silanols, silyl ether, and silyl chlorides. We believe this transformation offers a valuable platform for future applications in organosilicon chemistry and related fields.

H. K. and J. L. designed and carried out the experiments. J. L. and S. L. wrote the manuscript. S. L. supervised the project.

Conflicts of interest

There are no conflicts to declare.

Data availability

The data supporting this article have been included as part of the Supplementary Information (SI). Supplementary information includes experimental procedures, characterization data, and NMR spectra. See DOI: <https://doi.org/10.1039/d6cc00213g>.

Acknowledgements

This study was supported by the National Research Foundation (NRF) of Korea and funded by the Ministry of Science and ICT(RS-2024-00454580).

References

- (a) Y. Liu, I. Imae, A. Makishima and Y. Kawakami, Synthesis and characterization of poly(silphenylenesiloxane)s containing functional side groups, a study to high-temperature elastomer, *RSC Appl. Interfaces*, 2003, **4**, 27–34; (b) R. Murugavel, M. G. Walawalkar, M. Dan, H. W. Roesky and C. N. R. Rao, Transformations of Molecules and Secondary Building Units to Materials: A Bottom-Up Approach, *Acc. Chem. Res.*, 2004, **37**, 763–774.
- (a) E. W. Colvin, Silicon in organic synthesis, *Chem. Soc. Rev.*, 1978, **7**, 15–64; (b) M. Lalonde and T. Chan, Use of organosilicon reagents as protective groups in organic synthesis, *Synthesis*, 1985, 817–845.
- (a) S. E. Denmark and C. S. Regens, Palladium-Catalyzed Cross-Coupling Reactions of Organosilanols and Their Salts: Practical Alternatives to Boron- and Tin-Based Methods, *Acc. Chem. Res.*, 2008, **41**, 1486–1499; (b) R. Murugavel, A. Voigt, M. G. Walawalkar and H. W. Roesky, Hetero- and Metallasiloxanes Derived from Silanediols, Disilanols, Silanetriols, and Trisilanols, *Chem. Rev.*, 1996, **96**, 2205–2236.
- (a) Q.-L. Hu, K.-Q. Hou, J. Li, Y. Ge, Z.-D. Song, A. S. C. Chan and X.-F. Xiong, Silanol: a bifunctional group for peptide synthesis and late-stage functionalization, *Chem. Sci.*, 2020, **11**, 6070–6074; (b) Y. Lin, K.-Z. Jiang, J. Cao, Z.-J. Zheng, Z. Xu, Y.-M. Cui and L.-W. Xu, Iridium-Catalyzed Intramolecular C–H Silylation of Siloxane-Tethered Arene and Hydrosilane: Facile and Catalytic Synthesis of Cyclic Siloxanes, *Adv. Synth. Catal.*, 2017, **359**, 2247–2252; (c) M. Parasram and V. Gevorgyan, Silicon-Tethered Strategies for C–H Functionalization Reactions, *Acc. Chem. Res.*, 2017, **50**, 2038–2053.
- (a) J. W. Attard, K. Osawa, Y. Guan, J. Hatt, S.-I. Kondo and A. Mattson, Silanediol Anion Binding and Enantioselective Catalysis, *Synthesis*, 2019, 2107–2115; (b) K. M. Diemoz, J. E. Hein, S. O. Wilson, J. C. Fettinger and A. K. Franz, Reaction Progress Kinetics Analysis of 1,3-Disiloxanediols as Hydrogen-Bonding Catalysts, *J. Org. Chem.*, 2017, **82**, 6738–6747; (c) T. Min, J. C. Fettinger and A. K. Franz, Enantiocontrol with a Hydrogen-bond Directing Pyrrolidinylsilanol Catalyst, *ACS Catal.*, 2012, **2**, 1661–1666; (d) A. G. Schafer, J. M. Wieting, T. J. Fisher and A. E. Mattson, Chiral silanediols in anion-binding catalysis, *Angew. Chem.*, 2013, **52**, 11321–11324.
- (a) A. K. Franz and S. O. Wilson, Organosilicon Molecules with Medicinal Applications, *J. Med. Chem.*, 2013, **56**, 388–405; (b) G. K. Min, D. Hernández and T. Skrydstrup, Efficient Routes to Carbon-Silicon Bond Formation for the Synthesis of Silicon-Containing Peptides and Azasilaheterocycles, *Acc. Chem. Res.*, 2013, **46**, 457–470; (c) J.-L. Panayides, D. L. Riley, F. Hasenmaile and W. A. L. van Otterlo, The role of silicon in drug discovery: a review, *RSC Med. Chem.*, 2024, **15**, 3286–3344; (d) M. C. Parrott, M. Finnis, J. C. Luft, A. Pandya, A. Gullapalli, M. E. Napier and J. M. DeSimone, Incorporation and Controlled Release of Silyl Ether Prodrugs from PRINT Nanoparticles, *J. Am. Chem. Soc.*, 2012, **134**, 7978–7982; (e) R. Ramesh and D. S. Reddy, Quest for Novel Chemical Entities through Incorporation of Silicon in Drug Scaffolds, *J. Med. Chem.*, 2018, **61**, 3779–3798.
- (a) J. A. Cella and J. C. Carpenter, Procedures for the preparation of silanols, *J. Organomet. Chem.*, 1994, **480**, 23–26; (b) E. J. Corey and A. Venkateswarlu, Protection of hydroxyl groups as tert-butyl dimethylsilyl derivatives, *J. Am. Chem. Soc.*, 1972, **94**,



- 6190–6191; (c) L. J. Tyler, Phenylsilanetriol, *J. Am. Chem. Soc.*, 1955, 77, 770–771.
- 8 (a) W. Adam, R. Mello and R. Curci, O-Atom Insertion into Si–H Bonds by Dioxiranes: A Stereospecific and Direct Conversion of Silanes into Silanols, *Angew. Chem., Int. Ed. Engl.*, 1990, 29, 890–891; (b) M. Cavicchioli, V. Montanari and G. Resnati, Oxyfunctionalization reactions by perfluoro *cis*-2, 3-dialkyloxaziridines. Enantioselective conversion of silanes into silanols, *Tetrahedron Lett.*, 1994, 35, 6329–6330; (c) P. D. Lickiss and R. Lucas, Oxidation of sterically hindered organosilicon hydrides using potassium permanganate, *J. Organomet. Chem.*, 1996, 521, 229–234; (d) L. Spialter, L. Pazdernik, S. Bernstein, W. A. Swansiger, G. R. Buell and M. E. Freeburger, Mechanism of the reaction of ozone with the silicon-hydrogen bond, *J. Am. Chem. Soc.*, 1971, 93, 5682–5686; (e) K. Valliant-Saunders, E. Gunn, G. R. Shelton, D. A. Hrovat, W. T. Borden and J. M. Mayer, Oxidation of tertiary silanes by osmium tetroxide, *Inorg. Chem.*, 2007, 46, 5212–5219.
- 9 (a) S. Bähr, S. Brinkmann-Chen, M. Garcia-Borràs, J. M. Roberts, D. E. Katsoulis, K. N. Houk and F. H. Arnold, Selective enzymatic oxidation of silanes to silanols, *Angew. Chem., Int. Ed.*, 2020, 59, 15507–15511; (b) J. Y. Corey and J. Braddock-Wilking, Reactions of hydrosilanes with transition-metal complexes: formation of stable transition-metal silyl compounds, *Chem. Rev.*, 1999, 99, 175–292; (c) W.-S. Huang, H. Xu, H. Yang and L.-W. Xu, Catalytic Synthesis of Silanols by Hydroxylation of Hydrosilanes: From Chemoselectivity to Enantioselectivity, *Chem. – Eur. J.*, 2024, 30, e202302458; (d) M. Jeon, J. Han and J. Park, Catalytic Synthesis of Silanols from Hydrosilanes and Applications, *ACS Catal.*, 2012, 2, 1539–1549; (e) Y. Kikukawa, Y. Kuroda, K. Yamaguchi and N. Mizuno, Diamond-Shaped [Ag₄]⁴⁺ Cluster Encapsulated by Silicotungstate Ligands: Synthesis and Catalysis of Hydrolytic Oxidation of Silanes, *Angew. Chem., Int. Ed.*, 2012, 51, 2434–2437; (f) S. Li, H. Li, C.-H. Tung and L. Liu, Practical and selective bio-inspired iron-catalyzed oxidation of Si–H bonds to diversely functionalized organosilanols, *ACS Catal.*, 2022, 12, 9143–9152; (g) U. Prieto-Pascual, A. Rodríguez-Diéguez, Z. Freixa and M. A. Huertos, Tailor-made synthesis of hydrosilanols, hydrosiloxanes, and silanediols catalyzed by di-silyl rhodium(III) and iridium(III) complexes, *Inorg. Chem.*, 2023, 62, 3095–3105.
- 10 (a) J. Li, D. Xu, G. Shi, X. Liu, J. Zhang and B. Fan, Oxidation of Silanes to Silanols with Oxygen via Photoredox Catalysis, *ChemistrySelect*, 2021, 6, 8345–8348; (b) Y. Okada, M. Oba, A. Arai, K. Tanaka, K. Nishiyama and W. Ando, Diorganotelluride-Catalyzed Oxidation of Silanes to Silanols under Atmospheric Oxygen, *Inorg. Chem.*, 2010, 49, 383–385.
- 11 (a) X. Fan, P. Xiao, Z. Jiao, T. Yang, X. Dai, W. Xu, J. D. Tan, G. Cui, H. Su, W. Fang and J. Wu, Neutral-Eosin-Y-Photocatalyzed Silane Chlorination Using Dichloromethane, *Angew. Chem., Int. Ed.*, 2019, 58, 12580–12584; (b) X. Fan, M. Zhang, Y. Gao, Q. Zhou, Y. Zhang, J. Yu, W. Xu, J. Yan, H. Liu, Z. Lei, Y. C. Ter, S. Chanmungkalakul, Y. Lum, X. Liu, G. Cui and J. Wu, Stepwise on-demand functionalization of multihydrosilanes enabled by a hydrogen-atom-transfer photocatalyst based on eosin Y, *Nat. Chem.*, 2023, 15, 666–676.
- 12 J. Lee and S. Lee, Dehalogenative deuteration of alkyl and aryl bromides by thiyl radical catalysis under visible-light irradiation, *Chem. Commun.*, 2024, 60, 5526–5529.
- 13 F. Dénès, M. Pichowicz, G. Povie and P. Renaud, Thiyl Radicals in Organic Synthesis, *Chem. Rev.*, 2014, 114, 2587–2693.

