



Palladium-catalyzed denitrative C–N coupling of amides with nitroarenes enabled by Al(OTf)₃

 Linjie Yang,^{*abc} Xuejie Wang^c and Wanzhi Chen^{id}^c

Cite this: DOI: 10.1039/d6cc00697c

 Received 2nd February 2026,
 Accepted 18th March 2026

DOI: 10.1039/d6cc00697c


rsc.li/chemcomm

We report a Pd-catalyzed denitrative C–N coupling of amide nucleophiles with nitroarenes, providing direct and atom-economical access to *N*-aryl amides from readily available nitroarene feedstocks. Al(OTf)₃ is essential, both promoting the reaction and suppressing BrettPhos oxidation to maintain the active Pd–phosphine species. The method shows broad amide scope and delivers products in up to 95% yield.


The *N*-aryl amide scaffold represents a valuable structural motif widely found in biologically active molecules and functional materials.¹ Consequently, the development of efficient and versatile methods for synthesizing such compounds continues to attract significant interest. In efforts to expand the synthetic toolbox beyond classical amide-bond-forming protocols,² approaches involving transition-metal catalysis—particularly Pd-catalyzed C–N cross-coupling reactions—have drawn increasing attention (Scheme 1a). This appeal stems from the ready availability of amides as nucleophiles and the broad range of (hetero)aryl electrophiles available for coupling.³ Despite challenges associated with their inherently low nucleophilicity⁴ and potential catalytic inhibition through κ^2 -N,O coordination to the metal center,⁵ Pd-catalyzed C–N cross-coupling has nevertheless emerged as an efficient and reliable strategy for constructing *N*-aryl amides.⁶

Significant progress in amide *N*-arylation has been enabled by the development of phosphine⁷ and *N*-heterocyclic carbene (NHC) ligands⁸ with distinctive structural and electronic properties. Although ligand design has played a crucial role in advancing these transformations, persistent reactivity challenges continue to motivate exploration of alternative catalytic strategies and refinement of existing methodologies. One particularly effective approach


(a) Transition-Metal-Catalyzed Amidation of Aryl Electrophiles:



(b) Doberiner group, 2017:



(c) Organ group, 2017:



Scheme 1 (a) Transition-metal-catalyzed amidation of aryl electrophiles, (b) Doberiner group's work and (c) Organ group's work.

for accelerating the turnover-limiting transmetalation step in Pd-catalyzed amide coupling involves the use of Lewis acid additives, as independently demonstrated by the research groups of Doberiner (Scheme 1b)⁹ and Organ (Scheme 1c)⁸ in 2017.


In modern synthetic chemistry, achieving high atom economy and step efficiency is central to sustainable methodology design.¹⁰ In this context, employing nitroarenes as aryl donors in amide *N*-arylation offers a conceptually simple yet atom-economical strategy.¹¹ By enabling direct amide bond formation from readily available nitro compounds, such reactions streamline synthetic routes, minimize waste generation, and provide a more environmentally responsible alternative to traditional multistep protocols.

^a College of Chemistry and Environmental Engineering, Shenzhen University, Shenzhen 518060, China

^b Guangdong Key Laboratory for Biomedical Measurements and Ultrasound Imaging, National-Regional Key Technology Engineering Laboratory for Medical Ultrasound, School of Biomedical Engineering, Shenzhen University Medical School, Shenzhen 518060, China

^c Department of Chemistry, Zhejiang University, Hangzhou 310058, China. E-mail: 21937063@zju.edu.cn, chenwzz@zju.edu.cn






Scheme 2 (a) This work and (b) the proposed mechanism.

Building on prior studies by our group,¹² we hypothesized that relatively inert nitroarenes could serve as electrophilic partners in the Pd-catalyzed C–N coupling of amides. This approach would further broaden the scope of denitrative cross-coupling reactions (Scheme 2a). Guided by literature precedents and preliminary experiments, we speculated that under palladium catalysis, the use of electron-rich and sterically demanding phosphine or NHC ligands could facilitate cleavage of the C(sp²)-NO₂ bond in nitroarenes,¹³ thereby enabling amides to participate as nucleophiles. The proposed mechanism is illustrated in Scheme 2b. To ensure efficient reaction progress, suppression of the formation of the catalytically inactive κ^2 -amidate complex E was pursued in our current work by carefully optimizing the reaction conditions.

At the outset of this study, *p*-toluamide (**1a**) and α -nitronaphthalene (**2a**) were selected as model substrates to test the reaction parameters. The initial denitrative amidation conducted in toluene (150 °C) with Pd(acac)₂ (5 mol%), BrettPhos (10 mol%), and K₃PO₄ (3.0 equiv.) afforded no detectable product (Table 1, entry 1). Inspired by the work of Dobereiner⁹ and Stradiotto,¹⁴ Al(OTf)₃ (10 mol%) was introduced as a Lewis acid additive, and it furnished the coupling product **3a** in 37% isolated yield (entry 2). Subsequent optimization revealed K₃PO₄ to be the optimal base: K₂CO₃ or Cs₂CO₃ led to slightly lower yields, while strong bases, namely *t*-BuOK or DBU, completely suppressed the reaction (entries 3–6). Solvent screening identified PhCF₃ as superior to 1,4-dioxane, anisole, and *p*-xylene (entries 7–10). Ligand evaluation showed that BrettPhos outperformed other biarylphosphines (RuPhos, XPhos, *t*BuBrettPhos), while imidazo[1,5-*a*]pyridine-derived NHC ligands (NHC-HCl 1, NHC-HCl 2) also promoted the reaction effectively (entries 11–15). Ultimately, BrettPhos was identified as the optimal ligand. Increasing the loadings of both the catalyst and ligand improved the yield of **3a** to 85%, whereas deviations in temperature reduced efficiency (entries 16–18). Under optimized conditions—Pd(acac)₂ (10 mol%), BrettPhos (20 mol%),

Table 1 Optimization of the denitrative amidation reaction

Entry	Catalyst	Base	Solvent	Yield ^a (%)
1 ^b	BrettPhos	K ₃ PO ₄	Toluene	N.D.
2	BrettPhos	K ₃ PO ₄	Toluene	37
3	BrettPhos	K ₂ CO ₃	Toluene	20
4	BrettPhos	Cs ₂ CO ₃	Toluene	34
5	BrettPhos	<i>t</i> -BuOK	Toluene	N.D.
6	BrettPhos	DBU	Toluene	N.D.
7	BrettPhos	K ₃ PO ₄	Dioxane	39
8	BrettPhos	K ₃ PO ₄	Anisole	32
9	BrettPhos	K ₃ PO ₄	<i>p</i> -Xylene	30
10	BrettPhos	K ₃ PO ₄	PhCF ₃	40
11	RuPhos	K ₃ PO ₄	PhCF ₃	28
12	XPhos	K ₃ PO ₄	PhCF ₃	24
13	<i>t</i> BuBrettPhos	K ₃ PO ₄	PhCF ₃	31
14	NHC-HCl 1	K ₃ PO ₄	PhCF ₃	35
15	NHC-HCl 2	K ₃ PO ₄	PhCF ₃	36
16 ^c	BrettPhos	K ₃ PO ₄	PhCF ₃	85
17 ^{cd}	BrettPhos	K ₃ PO ₄	PhCF ₃	82
18 ^{ce}	BrettPhos	K ₃ PO ₄	PhCF ₃	79




Reaction conditions: *p*-toluamide **1a** (0.4 mmol), α -nitronaphthalene **2a** (0.6 mmol), Pd(acac)₂ (5 mol%), ligand (10 mol%), base (3.0 equiv.), Al(OTf)₃ (10 mol%) and solvent (2.0 mL, 0.2 M) at 150 °C under N₂ for 12 h. ^a Isolated yields. ^b No additive. ^c Pd(acac)₂ (10 mol%), ligand (20 mol%). ^d 140 °C. ^e 160 °C. N.D. = not detected.

Al(OTf)₃ (10 mol%), and K₃PO₄ (3.0 equiv.) in PhCF₃ at 150 °C and a reaction time of 12 h—the coupling product **3a** was obtained in 85% isolated yield (entry 16).


With the optimized conditions in hand, the nitroarene substrate scope was next examined (Scheme 3). Both 1-nitronaphthalene and nitrobenzene underwent smooth coupling to afford **3a** and **3b** in 85% and 86% yields, respectively. Scaling up the reaction of **2a** to 5.0 mmol afforded **3a** in 76% yield. The polycyclic aromatic substrate 1-nitropyrene exhibited moderate reactivity, yielding **3r** (52%). Alkyl-substituted nitrobenzenes afforded **3d–f** in 88–92% yields, irrespective of substitution pattern. Electron-donating substituents, namely methoxy (**3c**, **3j**) and *tert*-butyl (**3p**), were well tolerated, as were ester-functionalized substrates (**3n**). In contrast, nitroarenes bearing strongly electron-withdrawing groups (*e.g.*, cyano) failed to undergo coupling. A substrate containing a base-sensitive functionality, namely benzoyl (**3h**), gave a significantly reduced yield (15%). Fluorinated substrates, including those bearing trifluoromethoxy (**3g**), fluoro (**3i**), and trifluoromethyl (**3k**) groups, proved compatible. Furthermore, heteroaromatic nitro compounds, namely oxygen-containing heterocycles (**3l–m**), carbazole (**3o**), and dibenzothio-*phene* (**3q**), afforded the corresponding products in 30–91% yields.

Investigation of the amide substrate scope (Scheme 4) revealed similarly broad applicability. Benzamide afforded **4a**





Scheme 3 N-arylation of *p*-toluamide **1a** with nitroarenes under palladium catalysis.




Scheme 4 Scope of amide substrates.

in 78% yield, while *meta*- and *ortho*-methyl benzamides (**4b**, **4c**) were obtained in 88% and 84% yields, respectively, indicating negligible steric effects. The electron-donating groups methoxy (**4d**) and *tert*-butyl (**4e**) were well tolerated; however, the strongly electron-withdrawing substituents cyano and trifluoromethyl completely suppressed the reaction. The heteroaromatic amide furan-2-carboxamide delivered **4f** in 77% yield, and the fluorinated amide afforded **4g** in 53% yield. Additional aryl amides, including 1-naphthamide (**4h**) and 2-naphthamide (**4i**), provided excellent yields (90–93%). Aliphatic amides were also tested. For example, pivalamide (lacking any α -C–H bond) afforded **4j** in 95% yield. In contrast, cyclohexanecarboxamide gave **4k** in 61% yield—with this efficiency being only moderate likely due to deprotonation at the acidic α -C–H site, leading to competitive deprotonation/enolization and reduced chemo-selectivity under the reaction conditions, which can interfere with the coupling process.¹⁵ Secondary amides and lactams were not compatible with this protocol. Overall, taking our results together, the protocol provides a new and relatively general approach for accessing *N*-aryl amide substrates.

We also examined the influence of various Lewis acid additives on the reaction outcome (Scheme 5a). A series of air-stable metal triflates was tested, and the identity of the metal cation significantly affected the reactivity. Notably, the Group 13 metal triflates indium(III) triflate [In(OTf)₃] and Al(OTf)₃ exhibited the most pronounced promoting effects. Increasing the loading of Al(OTf)₃ to 20 mol% did not lead to any appreciable improvement in yield. In contrast, the borane-type Lewis acid B(C₆F₅)₃, previously employed by Organ and co-workers,⁸ proved ineffective in promoting the transformation in the current work.

We further investigated the C–N coupling reactions of amide nucleophiles with other electrophilic coupling partners, exemplified by dimethyl(phenyl)sulfonium triflate (**6a**) (Scheme 5b). In the absence of Al(OTf)₃, the desired *N*-phenyl amide product



Scheme 5 Effects of Lewis acid additives. (a) ^a Reaction conditions: **1a** (0.4 mmol), **2b** (0.6 mmol), Pd(acac)₂ (10 mol%), BrettPhos (20 mol%), K₃PO₄ (3.0 equiv.), Lewis acid (10 mol%) in PhCF₃ (2.0 mL, 0.2 M) at 150 °C for 12 h under an N₂ atmosphere. ^b Isolated yields. (b) ^a Reaction conditions: **1a** (0.4 mmol), **6a** (0.5 mmol), [(*t*BuBrettPhos)Pd(allyl)]OTf (2 mol%), K₃PO₄ (1.5 equiv.), Al(OTf)₃ (10 mol%) in 1,4-dioxane (2.0 mL, 0.2 M) at 110 °C for 12 h under an N₂ atmosphere. ^b Isolated yields.



was not detected; instead, thioanisole (*i.e.*, the demethylated byproduct) was formed as the major product. Upon addition of 10 mol% Al(OTf)₃, the desired coupling product **3b** was isolated in 85% yield. Moreover, Al(OTf)₃ effectively inhibited the oxidation of BrettPhos (Section II.5 of the Supporting Information), thereby maintaining a higher proportion of the active Pd-phosphine complex in the catalytic system and consequently enhancing the overall efficiency of the cross-coupling reaction.

In conclusion, we have developed a Pd-catalyzed denitrative C–N coupling between amide nucleophiles and nitroarenes using a catalytic amount of Lewis acid additives. This transformation employs nitroarenes as electrophilic partners. The Pd–Al cooperative catalytic system effectively inhibits the oxidation of the phosphine ligand. Importantly, the Lewis acid not only functions as an anion abstractor but also likely interacts with the amide substrate to form a catalytically active complex, accelerating the transmetalation step and preventing the formation of the inactive κ^2 -amidate species. Moreover, metal triflate additives may exert additional beneficial effects; further studies are required to elucidate their mechanistic role in Pd-catalyzed denitrative cross-coupling reactions.

Linjie Yang: conceived and performed the experiments, analyzed the data, and wrote the manuscript. Xuejie Wang: assisted with data analysis. Wanzhi Chen: administered the project, acquired funding, supervised the work, analyzed the data, and wrote the manuscript.

Conflicts of interest

There are no conflicts to declare.

Data availability

The data supporting this article have been included as part of the supplementary information (SI). Supplementary information: methods, experimental procedures, product characterization data, and NMR spectra. See DOI: <https://doi.org/10.1039/d6cc00697c>.

Acknowledgements

We thank the National Natural Science Foundation of China (No. 22071214) for financial support.

References

- (a) A. L. Simplicio, J. M. Clancy and J. F. Gilmer, *Molecules*, 2008, **13**, 519–547; (b) V. R. Pattabiraman and J. W. Bode, *Nature*, 2011, **480**, 471–479; (c) R. M. de Figueiredo, J. S. Suppo and J. M. Campagne, *Chem. Rev.*, 2016, **116**, 12029–12122; (d) A. R. Abdulkarim, H. S. Anbar, S. O. Zareei, A. A. Alfar, O. S. Al-Zoubi, E. G. Abdelkarim and M. I. El-Gamal, *Eur. J. Med. Chem.*, 2020, **188**, 112029.
- (a) J. R. Dunetz, J. Magano and G. A. Weisenburger, *Org. Process Res. Dev.*, 2016, **20**, 140–177; (b) E. Massolo, M. Pirola and M. Benaglia, *Eur. J. Org. Chem.*, 2020, 4641–4651; (c) P. Acosta-Guzmán, A. Ojedaporrás and D. Gamba-Sánchez, *Adv. Synth. Catal.*, 2023, **365**, 4359–4391.
- (a) C. L. Allen and J. M. J. Williams, *Chem. Soc. Rev.*, 2011, **40**, 3405; (b) S. Roy, S. Roy and G. W. Gribble, *Tetrahedron*, 2012, **68**, 9867–9923.
- (a) H. Sigel and R. B. Martin, *Chem. Rev.*, 1982, **82**, 385–426; (b) F. Milletti, L. Storchi, L. Goracci, S. Bendels, B. Wagner, M. Kansy and G. Cruciani, *Eur. J. Med. Chem.*, 2010, **45**, 4270–4279.
- K. I. Fujita, M. Yamashita, F. Puschmann, M. M. Alvarez-Falcon, C. D. Incarvito and J. F. Hartwig, *J. Am. Chem. Soc.*, 2006, **128**, 9044–9045.
- P. Ruiz-Castillo and S. L. Buchwald, *Chem. Rev.*, 2016, **116**, 12564–12649.
- (a) T. Ikawa, T. E. Barder, M. R. Biscoe and S. L. Buchwald, *J. Am. Chem. Soc.*, 2007, **129**, 13001–13007; (b) J. D. Hicks, A. M. Hyde, A. M. Cuezva and S. L. Buchwald, *J. Am. Chem. Soc.*, 2009, **131**, 16720–16734.
- S. Sharif, J. Day, H. N. Hunter, Y. Lu, D. Mitchell, M. J. Rodriguez and M. G. Organ, *J. Am. Chem. Soc.*, 2017, **139**, 18436–18439.
- J. Becica and G. E. Dobereiner, *ACS Catal.*, 2017, **7**, 5862–5870.
- P. Anastas and N. Eghbali, *Chem. Soc. Rev.*, 2010, **39**, 301–312.
- (a) K. Muto, T. Okita and J. Yamaguchi, *ACS Catal.*, 2020, **10**, 9856–9871; (b) M. Kashihara and Y. Nakao, *Acc. Chem. Res.*, 2021, **54**, 2928–2935; (c) K. Iizumi and J. Yamaguchi, *Org. Biomol. Chem.*, 2025, **23**, 1746–1772.
- (a) K. Chen, W. Chen, X. Yi, W. Chen, M. Liu and H. Wu, *Chem. Commun.*, 2019, **55**, 9287–9290; (b) W. Chen, K. Chen, W. Chen, M. Liu and H. Wu, *ACS Catal.*, 2019, **9**, 8110–8115; (c) W. Chen, W. Chen, M. Liu and H. Wu, *Org. Lett.*, 2022, **24**, 6983–6987; (d) W. Chen, K. Chen, X. Wang, L. Yang and W. Chen, *RSC Adv.*, 2024, **14**, 16624–16628; (e) L. Yang and W. Chen, *Org. Lett.*, 2025, **27**, 9721–9726.
- (a) M. R. Yadav, M. Nagaoka, M. Kashihara, R. L. Zhong, T. Miyazaki, S. Sakaki and Y. Nakao, *J. Am. Chem. Soc.*, 2017, **139**, 9423–9426; (b) F. Inoue, M. Kashihara, M. R. Yadav and Y. Nakao, *Angew. Chem., Int. Ed.*, 2017, **56**, 13307–13309; (c) J. Schleinitz, C. Chinchilla-Garzon, A. Perfetto, E. Escoudé, A. Makhloutah, G. Gontard, M. R. Vitale, A. Goujon, F. X. Felpin, L. Binet, I. Ciofini, P. Hudhomme and L. Grimaud, *Organometallics*, 2024, **44**, 29–35.
- M. J. Cotnam, N. Martinek and M. Stradiotto, *Eur. J. Org. Chem.*, 2023, e202301004.
- Z. Li, Y. Peng and T. Wu, *Org. Lett.*, 2021, **23**, 881–885.

