

# ChemComm

Chemical Communications

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: T. Wayama, R. Horinouchi, M. Arai, H. Ohashi, K. Watashi, S. Yoshida, S. Sato, S. Tsukamoto and H. Oguri, *Chem. Commun.*, 2026, DOI: 10.1039/D6CC01052K.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

## COMMUNICATION

## Design and Concise Synthesis of Halicyclamine-Inspired Bis-macrocycles Targeting DedA Membrane Proteins


Toshiaki Wayama,<sup>a</sup> Riku Horinouchi,<sup>b</sup> Masayoshi Arai,<sup>c\*</sup> Hirofumi Ohashi,<sup>d</sup> Koichi Watashi,<sup>d\*</sup> Satoshi Yoshida,<sup>e</sup> Sota Sato,<sup>ef†</sup> Sachiko Tsukamoto,<sup>g</sup> Hiroki Oguri<sup>a\*</sup>Received 00th January 20xx,  
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

**DedA membrane proteins promote the formation of double-membrane vesicles serving as replication platforms for RNA viruses, including SARS-CoV-2. Inspired by the only known inhibitor, halicyclamine A, we developed a concise biomimetic synthesis of halicyclamine-inspired macrocycles, identifying bis-macrocycles that suppress SARS-CoV-2 replication, while exhibiting DedA-associated activity in a bacterial overexpression assay.**

Positive-sense single-stranded RNA viruses, including severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), replicate in unique membrane compartments such as double-membrane vesicles (DMVs) derived from the endoplasmic reticulum and formed specifically in infected cells (Figure 1).<sup>1</sup> Biogenesis of these membranous replication organelles involves both viral and host factors. The host DedA superfamily, such as TMEM41B<sup>2</sup> and VMP1<sup>3</sup>, which function as lipid scramblases to remodel the lipid bilayer, has been reported to drive the maturation of the membrane structures into DMVs.<sup>4,5</sup> These DMVs are formed in the cells infected with positive-sense single-stranded RNA viruses, including not only SARS-CoV-2 but also hepatitis C virus, enteroviruses, and noroviruses.<sup>6</sup> While viral proteases and polymerases have been primary targets of antiviral therapeutics, formation of DMVs is expected to be an alternative target of drugs, especially those effective against multiple viruses. However, chemical modulation of host DedA

membrane proteins as an antiviral strategy remains largely unexplored.<sup>7</sup>



**Figure 1.** The estimated function of DedA membrane proteins in the replication cycle of SARS-CoV-2 and halicyclamine A (1), a reported inhibitor of bacterial DedA.

To date, the sponge-derived manzamine alkaloid halicyclamine A (1) has been reported as the only molecule that targets DedA membrane proteins (Figure 1).<sup>8</sup> Manzamine alkaloids,<sup>9,10</sup> exemplified by halicyclamine A (1), possess

<sup>a</sup> Department of Chemistry, Graduate School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

<sup>b</sup> School of Pharmaceutical Sciences, The University of Osaka, 1-6 Yamadaoka, Suita, Osaka 565-0871, Japan

<sup>c</sup> Graduate School and School of Pharmaceutical Sciences, The University of Osaka, 1-6 Yamadaoka, Suita, Osaka 565-0871, Japan

<sup>d</sup> Department of Drug Development, National Institute of Infectious Diseases, Japan Institute for Health Security, 1-23-1 Toyama, Shinjuku-ku, Tokyo 162-8640, Japan

<sup>e</sup> Department of Applied Chemistry, School of Engineering, The University of Tokyo, FS CREATION, Mitsui LINK Lab Kashiwanoha 1, 6-6-2 Kashiwanoha, Kashiwa, Chiba 277-0882, Japan

<sup>f</sup> Institute for Molecular Science, 5-1 Higashiyama, Myodaiji, Okazaki, Aichi 444-8787, Japan

<sup>g</sup> Department of Natural Medicines, Graduate School of Pharmaceutical Sciences, Kumamoto University, 5-1 Oehonmachi, Chuo-ku, Kumamoto 862-0973, Japan

\*Corresponding Authors


† Present address: JEOL Ltd., 3-1-2 Musashino, Akishima, Tokyo 196-8558, Japan



characteristic architectures comprising multicyclic bis-nitrogenated cores and macrocyclic alkyl loops, and display diverse biological activities such as antimalarial<sup>11</sup> and antituberculosis<sup>12</sup> effects. Their  $sp^3$ -rich macrocyclic scaffolds occupy an underexplored chemical space for the development of next-generation medium-sized molecule therapeutics.<sup>13</sup>

In this study, we report the concise synthesis of halicyclamine-inspired macrocyclic mimics and demonstrate activity consistent with DedA-associated modulation, thereby supporting their relevance for host-targeted antiviral strategies.

**Scheme 1.** Synthesis of halicyclamine-inspired macrocycles via biomimetic transannular cyclization (A), olefin metathesis (B), and macrolactonization (C). Compound **14** exhibited whole-molecule disorder in the crystal structure.



It has been proposed that halicyclamines are biosynthesized via an intramolecular transannular cyclization of a  $C_2$ -symmetric macrocyclic hypothetical intermediate **2** bearing a pair of 1,6-dihydropyridine (DHP) units.<sup>14</sup> Baldwin and co-workers chemically synthesized this highly unstable intermediate **2** and attempted transannular cyclizations; however, disproportionation of the 1,6-DHP units predominated, resulting in rapid decomposition within 1 h (Scheme 1A).<sup>15</sup> To overcome this problem, we strategically stabilized the 1,6-DHP moiety while preserving its reactivity toward formation of the bis-nitrogenated bicyclic scaffold. Specifically, 1,6-DHP **3** was designed to incorporate a geminal dimethyl group at C6 to completely suppress hydride transfer from the 1,6-DHP unit, together with an electron-withdrawing ester substituent at C3 to enhance oxidative stability.<sup>16</sup> Upon treatment with a Brønsted acid catalyst, macrocyclic  $C_2$ -symmetric intermediate **3** underwent a biomimetic, regio-controlled transannular cyclization between C4 and C5' of the two DHP units via an iminium intermediate **4**, affording the halicyclamine-type macrocycle **5**.<sup>17</sup> This compound **5** features a macrocyclic ring of the same size as halicyclamine B (**6**).<sup>18</sup> Despite this success, the previously reported route required 12 synthetic steps in total.


In this study, we report a redesigned synthetic approach that enables rapid and customizable access to bis-macrocylic halicyclamine analogues exhibiting activity consistent with DedA-associated modulation. To this end, we examined two alternative macrocyclization strategies, beginning with a ring-closing olefin metathesis (RCM)-based approach (Scheme 1B). When 1,6-DHP **7**, readily synthesized in four steps, was

subjected to a Brønsted acid-catalyzed regio-controlled dimerization between the C4 and C5' positions, the corresponding halicyclamine-type framework **9** was obtained in quantitative yield. Subsequent RCM using first-generation Grubbs catalyst **10** furnished halicyclamine-type bis-macrocycle **11** in six steps. The modest yield of this transformation is likely attributable to competing metathesis pathways during the two sequential RCM events, leading to undesired macrocycles and related byproducts.<sup>15c</sup> Single-crystal X-ray diffraction analysis confirmed that **11** adopts a three-dimensional architecture closely resembling that of halicyclamine A (**1**) and B (**6**)<sup>18</sup> (see Fig. S3–S4 for overlays of these structures). Overall, this approach reduced the total number of steps by approximately half compared with our previous biomimetic route<sup>17</sup> (Scheme 1A).

As a complementary strategy to olefin metathesis, we next explored a macrolactonization-based approach for the construction of **14** (Scheme 1C). Brønsted acid-catalyzed dimerization of 1,6-DHP **12**, which was likewise readily synthesized in four steps, afforded halicyclamine-type dimer **13** in quantitative yield. Subsequent site-selective hydrolysis of the two methyl ester groups in **13**, followed by macrolactonization under 2-methyl-6-nitrobenzoic anhydride (MNBA) and 4-dimethylaminopyridine *N*-oxide (DMAPO) conditions,<sup>19</sup> provided halicyclamine-type bis-macrocycle **14** in seven steps. Single-crystal X-ray diffraction analysis confirmed that **14** similarly reproduces the three-dimensional architecture of halicyclamine A (**1**) and B (**6**)<sup>18</sup> (Fig. S3–S4). Notably, both alternative synthetic routes (Schemes 1B and C) strategically




integrate our biomimetic dimerization platform with site-selective and operationally straightforward macrocyclization methods. These routes are substantially shorter than our previously reported synthesis of halicyclamine analogue **5** and provide rapid access to bis-macrocylic alkaloidal scaffolds that faithfully recapitulate the natural product architecture while enabling controlled structural diversification.<sup>20</sup>



**Figure 2.** Evaluation of inhibitory activity toward wild-type and DedA-overexpressed *Mycobacterium smegmatis*.

With a concise and modular synthetic route established, we next examined whether the synthesized halicyclamine analogues functionally engage DedA membrane proteins in a cellular context. Accordingly, their inhibitory activities were evaluated using an established DedA overexpression assay in *Mycobacterium smegmatis* and benchmarked against representative members of manzamine alkaloid family (Figure 2). Following the reported protocol,<sup>8</sup> a DedA-overexpressing *M. smegmatis* strain was prepared, and its responses to each compound were compared with those of the corresponding wild-type strain using an MTT-based viability assay.<sup>21</sup> In this assay format, compounds that exhibit reduced inhibitory effects against the DedA-overexpressing strain relative to the wild-type strain are interpreted as functionally engaging DedA membrane proteins, consistent with prior genetic and chemical validation

studies.<sup>8</sup> Halicyclamine A (**1**) was not available in sufficient quantity from natural sources; therefore, halicyclamine B (**6**), which is closely related in structure to the synthesized analogues, was selected as the reference compound.<sup>22</sup> The natural product halicyclamine B (**6**) displayed weak growth inhibition against the wild-type strain, with an MIC value of 100  $\mu\text{M}$ , whereas no inhibition was observed against the DedA-overexpressing strain at concentrations up to 100  $\mu\text{M}$ . Although the absolute potency was modest, this differential response indicates a discernible resistance shift associated with DedA overexpression. The concisely synthesized halicyclamine-type macrocycle **11** showed a closely comparable profile, exhibiting an MIC of 100  $\mu\text{M}$  against the wild-type strain and no detectable inhibition of the DedA-overexpressing strain up to 100  $\mu\text{M}$ , resulting in a resistance shift similar in magnitude to that of halicyclamine B (**6**). In contrast, the halicyclamine analogue **14**, assembled via macrolactonization, displayed enhanced inhibitory activity, with MIC values of 30  $\mu\text{M}$  against the wild-type strain and 60  $\mu\text{M}$  against the DedA-overexpressing strain. The larger separation between these values corresponds to a more pronounced resistance shift, exceeding that observed for the natural product **6**, and is consistent with enhanced DedA-related activity under these assay conditions. By comparison, macrocycle **15** lacking the central transannular bond<sup>17</sup> and the macrocyclization precursors **9** and **13** showed no detectable inhibitory activity against either strain. For reference, manzamine A (**16**), a representative macrocyclic marine alkaloid,<sup>23</sup> was also evaluated and exhibited no detectable inhibitory activity under the same assay conditions. Collectively, these results suggest that DedA-associated growth inhibition in *M. smegmatis* is preferentially linked to halicyclamine-type bis-macrocylic scaffolds and appears to depend on both the bis-nitrogenated core framework and the bis-macrocylic architecture.



**Figure 3.** Evaluation of inhibitory activity toward SARS-CoV-2 replication.

To gain initial insight into whether the designed halicyclamine-type analogues translate into functional antiviral activity, the anti-SARS-CoV-2 replication activities of the



synthetic analogues were evaluated in a cell-based SARS-CoV-2 infection assay (Figure 3). The synthetic bis-macrocyclic analogues **11** and **14** showed antiviral potency, with  $IC_{50}$  values of 1.7 and 1.4  $\mu$ M, respectively. In contrast, both macrocycle **15**, which lacks the central transannular bond,<sup>17</sup> and macrocyclization precursor **13** displayed only weak replication-inhibitory activity ( $IC_{50} > 10 \mu$ M) under the same conditions. This preliminary structure–activity relationship suggests that the bis-macrocyclic architecture, particularly the presence of the central transannular bond, substantially enhances antiviral potency. Collectively, these findings highlight halicyclamine-inspired bis-macrocyclic scaffolds as promising lead structures for antiviral drug discovery.

In summary, bis-macrocyclic analogues **11** and **14** were newly designed on the basis of halicyclamine A (**1**) and rapidly synthesized via a biomimetic dimerization strategy. DedA-associated activity of these analogues was supported by the differential responses observed between wild-type and DedA-overexpressing *Mycobacterium smegmatis* strains. In line with this DedA-associated activity, these halicyclamine-inspired analogues **11** and **14** potently suppressed SARS-CoV-2 replication in a cell-based infection assay.

Host DedA membrane proteins remain largely unexplored as targets of antiviral drug development. The present findings therefore suggest that these natural product-inspired analogues represent promising starting points for further investigation. Although DedA-associated activity was evaluated in this study using an established *M. smegmatis*-based assay, DedA homologues in humans and mycobacteria are expected to differ in sequence and structure. Moreover, SARS-CoV-2 replication is a complex process involving multiple host DedA-related homologues and viral proteins,<sup>5</sup> and the activity responses observed in the *M. smegmatis*-based assay may therefore not directly correspond to antiviral activity against SARS-CoV-2. Future work will focus on assessing inhibitory activity against human DedA homologues to clarify the translational potential of this strategy for the development of new host-directed antiviral agents.

## Data availability statement

The data underlying this study are available in the published article and its Supporting Information. Crystallographic data for compounds **11** and **14** have been deposited at the CCDC under 2519925 and 2519926, respectively.

## Author contributions

T. W. and H. Oguri designed the project. T. W. performed the organic synthesis. H. Ohashi and K. W. assessed antiviral activity against SARS-CoV-2, while T. W., R. H., and M. A. conducted assays using *Mycobacterium smegmatis*. T. W., S. Y., and S. S. performed X-ray crystallographic analysis. S. T. provided the authentic samples of natural products. T. W. and H. Oguri wrote the manuscript with input from all authors.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

This work was partially supported by JSPS KAKENHI (JP22H00346 and JP22H05127), the Astellas Foundation for Research on Metabolic Disorders, the Naito Foundation, and the Asahi Glass Foundation (H. Oguri). Additional support was provided by JSPS KAKENHI (JP25K21782) and JST MIRAI program (JPMJMI22G1) (K.W.) as well as JSPS KAKENHI (JP24K18457) (H. Ohashi). T.W. is a Research Fellow of JSPS DC2 and gratefully acknowledges financial support from JSPS KAKENHI (JP24KJ0618) to complete this research work. T. W. also acknowledges the Program for Leading Graduate Schools, World-leading Innovative Graduate Study Program for Materials Research, Industry and Technology (MERIT-WINGS). We were granted access to the X-ray diffraction instruments at KEK Photon Factory BL17A beamline (2023G604). This work was further inspired by the international and interdisciplinary environments of the JSPS Asian CORE Program, "Asian Chemical Biology Initiative" as well as the JSPS A3 Foresight Program. Figures were created using BioRender. We are grateful to Dr. Nobuto Kaneko (UTokyo) for his advice on the DFT calculations.

## Notes and references

- For reviews, see: (a) Fehr, A. R.; Perlman, S. *Methods Mol. Biol.*, **2015**, *1282*, 1. (b) V'kovski, P.; Kratzel, A.; Steiner, S.; Stalder, H.; Thiel, V. *Nat. Rev. Microbiol.*, **2021**, *19*, 155; (c) Yang, H.; Rao, Z. *Nat. Rev. Microbiol.*, **2021**, *19*, 685. (d) Soares, V. C.; Moreira, I. B. G.; Dias, S. S. G. S. *Viruses*, **2025**, *17*, 1064.
- (a) F. Moretti, P. Bergman, S. Dodgson, D. Marcellin, I. Claerr, J. M. Goodwin, R. DeJesus, Z. Kang, C. Antczak, D. Begue, D. Bonenfant, A. Graff, C. Genoud, J. S. Reece-Hoyes, C. Russ, Z. Yang, G. R. Hoffman, M. Mueller, L. O. Murphy, R. J. Xavier and B. Nyfeler, *EMBO Rep.*, **2018**, *19*, e45889; (b) K. Morita, Y. Hama, T. Izume, N. Tamura, T. Ueno, Y. Yamashita, Y. Sakamaki, K. Mimura, H. Morishita, W. Shihoya, O. Nureki, H. Mano and N. Mizushima, *J. Cell Biol.*, **2018**, *217*, 3817; (c) C. J. Shoemaker, T. Q. Huang, N. R. Weir, N. J. Polyakov, S. W. Schultz and V. Denic, *PLoS Biol.*, **2019**, *17*, e2007044; (d) J. D. Trimarco, B. E. Heaton, R. R. Chaparian, R. A. Binder, G. C. Gray, V. D. Menachery and N. S. Heaton, *PLoS Pathog.*, **2021**, *17*, e1009599.
- (a) N. J. Dusetti, Y. Jiang, M. I. Vaccaro, R. Tomasini, S. A. Azizi, E. L. Calvo, A. Ropolo, F. Fiedler, G. V. Mallo, J.-C. Dagorn and J. L. Iovanna, *Biochem. Biophys. Res. Commun.*, **2002**, *290*, 641; (b) A. Ropolo, D. Grasso, R. Pardo, M. L. Sacchetti, C. Archange, A. L. Re, M. Seux, J. Nowak, C. D. Gonzalez, J. L. Iovanna and M. I. Vaccaro, *J. Biol. Chem.*, **2007**, *282*, 37124; (c) Y. Tian, Z. Li, W. Hu, H. Ren, E. Tian, Y. U. Zhao, Q. Lu, X. Huang, P. Yang, X. Li, X. Wang, A. L. Kovács, L. Yu and H. Zhang, *Cell*, **2010**, *141*, 1042; (d) E. Itakura and N. Mizushima, *J. Cell Biol.*, **2011**, *192*, 17; (e) C. Kishi-Itakura, I. Koyama-Honda, E. Itakura and N. Mizushima, *J. Cell Sci.*, **2014**, *127*, 4089.
- (a) A. Ghanbarpour, D. P. Valverde, T. J. Melia and K. M. Reinisch, *Proc. Natl. Acad. Sci. U. S. A.*, **2021**, *118*, e2101562118; (b) D. Huang, B. Xu, L. U. Liu, L. Wu, Y. Zhu, A. Ghanbarpour, Y. Wang, F.-J. Chen, J. Lyu, Y. Hu, Y. Kang, W. Zhou, Y. Wang, X. Ding, W. Li, Z. Jiang, Z. Chen, X. Zhang, H. Zhou, J. Z. Li, C. Guo, W. Zheng, X. Zhang, P. Li, T. Melia, K. Reinisch and X.-W. Chen, *Cell Metab.*, **2021**, *33*, 1655; (c) Y. E.



- Li, Y. Wang, X. Du, T. Zhang, H. Y. Mak, S. E. Hancock, H. McEwen, E. Pandzic, R. M. Whan, Y. C. Aw, I. E. Lukmantara, Y. Yuan, Z. Dong, A. Don, N. Turner, S. Qi and H. Yang, *J. Cell Biol.*, 2021, **220**, e202103105; (d) K. M. Reinisch, X.-W. Chen and T. J. Melia, *Contact*, 2021, **4**, 25152564211024494; (e) T. Zhang, Y. E. Li, Y. Yuan, X. Du, Y. Wang, X. Dong, H. Yang and S. Qi, *Autophagy*, 2021, **17**, 2048.
- 5 For review, see: Y. Hama, H. Morishita and N. Mizushima, *EMBO Rep.*, 2022, **23**, e53894.
- 6 (a) J. Baggen, L. Persoons, E. Vanstreels, S. Jansen, D. V. Looveren, B. Boeckx, V. Geudens, J. D. Man, D. Jochmans, J. Wauters, E. Wauters, B. M. Vanaudenaerde, D. Lambrechts, J. Neyts, K. Dallmeier, H. J. Thibaut, M. Jacquemyn, P. Maes and D. Daelemans, *Nat. Genet.*, 2021, **53**, 435. (b) H. H. Hoffmann, W. M. Schneider, K. Rozen-Gagnon, L. A. Miles, F. Schuster, B. Razoogy, E. Jacobson, X. Wu, S. Yi, C. M. Rudin, R. M. MacDonald, L. K. McMullan, J. T. Poirier and C. M. Rice, *Cell*, 2021, **184**, 133.
- 7 For reviews, see: (a) W. Cui, Y. Duan, Y. Gao, W. Wang and H. Yang, *Structure*, 2024, **32**, 1301-1320; (b) T. Kronenberger, S. A. Laufer and T. Pillaiyar, *Drug Discov. Today*, 2023, **28**, 103579.
- 8 M. Arai, L. Liu, T. Fujimoto, A. Setiawan and M. Kobayashi, *Mar. Drugs*, 2011, **9**, 984.
- 9 (a) R. Sakai, T. Higa, C. W. Jefford and G. Bernardinelli, *J. Am. Chem. Soc.*, 1986, **108**, 6404; (b) J. I. Kobayashi, M. Tsuda, N. Kawasaki, K. Matsumoto and T. Adachi, *Tetrahedron Lett.*, 1994, **35**, 4383; (c) M. Jaspars, V. Pasupathy and P. Crews, *J. Org. Chem.*, 1994, **59**, 3253.
- 10 For reviews, see: (a) E. Magnier and Y. Langlois, *Tetrahedron*, 1998, **54**, 6201; (b) J.-F. Hu, M. T. Hamann, R. Hill and M. Kelly, *Alkaloids*, 2003, **60**, 207; (c) R. Duval and E. Poupon, *Biomimetic Organic Synthesis*, Wiley-VCH, 2011; (d) T. Kubota, S. Kurimoto and J. Kobayashi, *Alkaloids*, 2020, **84**, 1.
- 11 (a) K. K. H. Ang, M. J. Holmes, T. Higa, M. T. Hamann and U. A. K. Kara, *Antimicrob. Agents Chemother.*, 2000, **44**, 1645; (b) K. V. Rao, B. D. Santarsiero, A. D. Mesecar, R. F. Schinazi, B. L. Tekwani and M. T. Hamann, *J. Nat. Prod.*, 2003, **66**, 823; (c) J. D. Winkler, A. T. Londregan and M. T. Hamann, *Org. Lett.*, 2006, **8**, 2591.
- 12 M. Arai, M. Sobou, C. Vilchéze, A. Baughn, H. Hashizume, P. Pruksakorn, S. Ishida, M. Matsumoto, W. R. Jacobs and M. Kobayashi, *Bioorg. Med. Chem.*, 2008, **16**, 6732.
- 13 (a) K. A. El Sayed, M. Kelly, U. A. K. Kara, K. K. H. Ang, I. Katsuyama, D. C. Dunbar, A. A. Khan and M. T. Hamann, *J. Am. Chem. Soc.*, 2001, **123**, 1804; (b) S. Matsunaga, Y. Miyata, R. W. M. Van Soest and N. Fusetani, *J. Nat. Prod.*, 2004, **67**, 1758; (c) D. G. Jimenez, V. Poongavanam and J. Kihlberg, *J. Med. Chem.*, 2023, **66**, 5377; (d) F. Lovering, J. Bikker and C. Humblet, *J. Med. Chem.*, 2009, **52**, 6752.
- 14 (a) A. Kaiser, X. Billot, A. Gateau-Olesker, C. Marazano and B. C. Das, *J. Am. Chem. Soc.*, 1998, **120**, 8026; (b) J.-C. Wypych, T. M. Nguyen, P. Nuhant, M. Bénèche and C. Marazano, *Angew. Chem., Int. Ed.*, 2008, **47**, 5418; (c) I. Sinigaglia, T. M. Nguyen, J.-C. Wypych, B. Delpech and C. Marazano, *Chem. Eur. J.*, 2010, **16**, 3594. (e) Ye Tang, Lili Zhu and Ran Hong, *Tetrahedron Chem*, 2022, **3**, 100025.
- 15 (a) J. E. Baldwin and R. C. Whitehead, *Tetrahedron Lett.*, 1992, **33**, 2059; (b) J. E. Baldwin, T. D. W. Claridge, A. J. Culshaw, F. A. Heupel, V. Lee, D. R. Spring, R. C. Whitehead, R. J. Boughtflower, I. M. Mutton and R. J. Upton, *Angew. Chem., Int. Ed.*, 1998, **37**, 2661; (c) J. E. Baldwin, T. D. W. Claridge, A. J. Culshaw, F. A. Heupel, V. Lee, D. R. Spring and R. C. Whitehead, *Chem. Eur. J.*, 1999, **5**, 3154; (d) A. T. Piwko, B. G. Miller and J. M. Smith, *Nat. Prod. Rep.*, 2023, **40**, 964. (e) A. Leblond, A. Nguyen, C. Alcover, K. Leblanc, J.-F. Gallard, D. Joseph, E. Poupon and M. A. Beniddir, *Org. Lett.*, 2024, **26**, 2163. (f) S. I. Gómez, E. I. O. Nava, A. Dadawalla, D. Ashong, G. Chen and Q.-H. Chen, *Molecules*, 2025, **30**, 2907. [DOI: 10.1039/D6CC01052K](https://doi.org/10.1039/D6CC01052K)
- 16 (a) T. Wayama, Y. Arai and H. Oguri, *J. Org. Chem.*, 2022, **87**, 5938. (b) T. Wayama and H. Oguri, *Org. Lett.*, 2026, **28**, 982.
- 17 T. Wayama and H. Oguri, *Org. Lett.*, 2023, **25**, 3596.
- 18 B. Harrison, S. Talapatra, E. Lobkovsky, J. Clardy and P. Crews, *Tetrahedron Lett.*, 1996, **37**, 9151.
- 19 I. Shiina, H. Fukui and A. Sasaki, *Nat. Protoc.*, 2007, **2**, 2312.
- 20 Selected examples for synthetic studies, see: (a) G. A. Molander and F. Cadoret, *Tetrahedron Lett.*, 2011, **52**, 2199; (b) A. I. Lansakara, S. V. S. Mariappan and F. C. Pigge, *J. Org. Chem.*, 2016, **81**, 10266; (c) M. G. Banwell, M. J. Coster, N. L. Hungerford, M. J. Garson, S. Su, A. C. Kotzed and M. H. G. Munro, *Org. Biomol. Chem.*, 2012, **10**, 154; (d) A. G. Dalling, G. Späth and A. Fürstner, *Angew. Chem., Int. Ed.*, 2022, **61**, e202209651; (e) B. J. Smith and G. A. Sulikowski, *Angew. Chem., Int. Ed.*, 2010, **49**, 1599; (f) B. J. Smith, T. Qu, M. Mulder, M. J. Noetzel, C. W. Lindsley and G. A. Sulikowski, *Tetrahedron*, 2010, **66**, 4805; (g) Z. D. Firestone, T. A. Grigolo, F. G. Pernichelle and J. M. Smith, *J. Am. Chem. Soc.*, 2026, **148**, 6814.
- 21 (a) T. Mosmann, *J. Immunol. Methods*, 1983, **65**, 55; (b) G. Abate, A. Aseffa, A. Selassie, S. Goshu, B. Fekade, D. WoldeMeskal and H. Miörner, *J. Clin. Microbiol.*, 2004, **42**, 871.
- 22 H. Kato, A. H. El-Desoky, Y. Takeishi, T. Nehira, E. D. Angkouw, R. E. P. Mangindaan, N. J. de Voogd and S. Tsukamoto, *Bioorg. Med. Chem. Lett.*, 2019, **29**, 8.
- 23 (a) K. Eguchi, Y. Fujiwara, A. Hayashida, H. Horlad, H. Kato, H. Rotinsulu, F. Losung, R. E. P. Mangindaan, N. J. de Voogd, M. Takeya and S. Tsukamoto, *Bioorg. Med. Chem.*, 2013, **21**, 3831; (b) J. R. Palem, G. R. Bedadala, K. A. El Sayed and S.-c. V. Hsia, *Planta Med.*, 2011, **77**, 46; (c) M. Yousaf, N. L. Hammond, J. Peng, S. Wahyuono, K. A. McIntosh, W. N. Charman, A. M. S. Mayer and M. T. Hamann, *J. Med. Chem.*, 2004, **47**, 3512.



## Data Availability Statement

The data underlying this study are available in the published article and its Supporting Information. Crystallographic data for compounds **11** and **14** have been deposited at the CCDC under 2519925 and 2519926, respectively.

