










Cite this: DOI: 10.1039/d5ea00179j

Yields of perfluorocarboxylic acids from the atmospheric oxidation of Montreal Protocol related gases

M. P. Sulbaek Andersen,  ^{*ab} T. J. Wallington,  ^c J. B. Burkholder,  ^d
S. Madronich,  ^e M. L. Hanson,  ^f D. Van Hoomissen  ^d and K. R. Solomon  ^g

We present here a systematic evaluation of the molar yields of trifluoroacetic acid (TFA), perfluoropropanoic acid (PFPrA), and perfluorobutanoic acid (PFBA) from the atmospheric degradation of gases relevant to the Montreal Protocol and its Amendments. The yields are dependent on the molecular structure of the parent compound and the primary degradation products and radical intermediates formed. We incorporate new data into the Tropospheric Ultraviolet Visible (TUV) model and discuss how recent studies improve our understanding of the relative importance of the photochemical pathways for perfluoroaldehydes, $C_xF_{2x+1}C(O)H$, which are key degradation products from some chlorofluorocarbon (CFC) replacement compounds. We identify areas for further research that could advance our understanding of the environmental fate of precursors to short-chain length perfluorocarboxylic acids.

Received 31st December 2025
Accepted 9th February 2026

DOI: 10.1039/d5ea00179j

rsc.li/esatmospheres

Environmental significance

There is substantial scientific, technical, and policy interest in the sources and environmental impacts of trifluoroacetic acid (TFA) and other short chain perfluorocarboxylic acids (PFCAs), which belong to the broader group of chemicals known as per- and polyfluoroalkyl substances (PFAS). Quantifying the anthropogenic contribution to the past, present, and future PFCA loading in the environment requires robust estimates of yields from the atmospheric degradation of volatile fluorinated organic compounds. In this work, we present an updated and expanded evaluation of the molar yields of TFA, perfluoropropionic acid (PFPrA), and perfluorobutanoic acid (PFBA) in the atmospheric degradation of gases under the purview of the Montreal Protocol and other related gases.

1 Introduction

Perfluorocarboxylic acids (PFCAs) are a group of chemicals with the general formula $C_xF_{2x+1}C(O)OH$. PFCAs are part of the larger group of chemicals known as per- and polyfluoroalkyl substances (PFAS).^{1,2} The ionized form of PFCAs (salts) are ubiquitous contaminants present in precipitation and in river, lake, and ocean water. Concerns have been raised that PFCAs are toxic and threaten global ecosystems and human health. It is well established that long-chain members of the group such as perfluorooctanoic acid ($C_7F_{15}C(O)OH$, PFOA) bioaccumulate

and are hazardous. PFOA is regulated as part of national and international agreements.^{3,4} However, short-chain members of the group such as trifluoroacetic acid ($CF_3C(O)OH$, TFA) do not biomagnify and have been shown to have low toxicity in experimental studies.^{5,6} There is current research and regulatory interest in the concentrations of PFAS in the environment, anthropogenic sources, and effects on human and ecosystem health. Robust estimates of yields of PFCAs from the atmospheric degradation of volatile fluorinated organic compounds are essential for quantifying the anthropogenic contribution to the PFCA loading in environment in the past, present, and future.

Chlorofluorocarbons (CFCs), hydrochlorofluorocarbons (HCFCs), hydrofluorocarbons (HFCs), halons, perfluorocarbons (PFCs), hydrofluoroolefins (HFOs), hydrochlorofluoroolefins (HCFOs), hydrobromofluoroolefins (HFBOs), hydrofluoroethers and hydrochlorofluoroethers (HFES/HCFEs) are classes of commercial fluorinated organic compounds. Most of these gases are either controlled, or are substitutes for compounds controlled, by the Montreal Protocol on Substances that Deplete the Ozone Layer and its amendments. These compounds have been released into the atmosphere in significant amounts.⁷

^aDepartment of Chemistry and Biochemistry, California State University Northridge, Northridge, California, USA. E-mail: mads@sulbaek.dk

^bDepartment of Chemistry, University of Copenhagen, DK-2100 Copenhagen, Denmark

^cCenter for Sustainable Systems, School for Environment and Sustainability, University of Michigan, Ann Arbor, Michigan, USA

^dChemical Sciences Laboratory, National Oceanic and Atmospheric Administration, Boulder, Colorado, USA

^eNational Center for Atmospheric Research, Boulder, Colorado, USA

^fDepartment of Environment and Geography, University of Manitoba, Winnipeg, Manitoba, Canada

^gSchool of Environmental Sciences, University of Guelph, Guelph, Ontario, Canada



Some of these compounds can degrade to yield PFCAs.⁸ To assess the anthropogenic contribution to the historical, current, and future PFCA loading in the environment, the yields from the atmospheric degradation of volatile fluorinated organic compounds need to be established.⁹ Molar yields of TFA from 21 Montreal Protocol-related gases were evaluated by Madronich *et al.*¹⁰ The yields of two other short-chain acids, pentafluoropropionic acid (perfluoropropanoic acid, PFPrA), and heptafluorobutanoic acid (perfluorobutanoic acid, PFBA) were briefly discussed by Madronich *et al.*¹¹ These are the only PFCAs that can be formed from gases under the purview of the Montreal Protocol and its amendments or from related gases that are in use, but not currently controlled under the Montreal Protocol.⁷

The atmospheric fate of perfluorinated aldehydes is a critical factor in assessments of the yield of PFCAs in the degradation of Montreal Protocol related gases. Since the publication of the previous two assessments,^{10,11} new research has been published on the photolysis^{12–14} and reaction with OH radicals¹⁵ of $\text{CF}_3\text{C}(\text{O})\text{H}$. We incorporate the new data into the Tropospheric Ultraviolet Visible (TUV) model to provide estimates for the importance of photolysis in the atmospheric fate of $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{H}$ ($x = 1–3$). We discuss how the new studies improve our understanding of the relative importance of the photochemical pathways for perfluorinated aldehydes. We present an updated and expanded assessment of the molar yields of TFA, PFPrA, and PFBA from the atmospheric degradation of Montreal Protocol related gases, including CFCs, HCFCs, HFCs, PFCs, HFOs, HCFOs, HBFOs, HFES, halons, alcohols, and ketones.

2 Mechanisms of formation of PFCAs

Fig. 1 shows several possible routes to the formation of PFCAs. The photochemical pathways depend on the molecular structure of the parent compound and the primary degradation products and radical intermediates formed. Compounds with a $\text{C}_x\text{F}_{2x+1}\text{CF}$ -moiety (where for the purpose of this discussion of gases related to the Montreal Protocol $x \leq 3$) can lead to the formation of $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{F}$ as a degradation product.¹⁶ The fate of $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{F}$ depends on whether it is formed in the troposphere or the stratosphere. In the troposphere, the sole fate of $\text{CF}_3\text{C}(\text{O})\text{F}$ is incorporation into water droplets followed by hydrolysis to give TFA (on a time scale of 5–15 days¹⁷). In the stratosphere, the main fate ($\approx 90\%$) of $\text{CF}_3\text{C}(\text{O})\text{F}$ is photolysis¹⁸ with the remaining 10% transported to the troposphere where it is removed by hydrolysis.

Compounds with a CF_3CCl -moiety can degrade to give $\text{CF}_3\text{C}(\text{O})\text{Cl}$, which undergoes photolysis in competition with incorporation into water droplets. The tropospheric photolytic lifetime for $\text{CF}_3\text{C}(\text{O})\text{Cl}$ for an overhead sun is estimated to be 23 days.¹⁹ The atmospheric lifetime of $\text{CF}_3\text{C}(\text{O})\text{Cl}$ with respect to uptake and hydrolysis in cloud water is 5–30 days.¹⁷ On average, an estimated 60% of $\text{CF}_3\text{C}(\text{O})\text{Cl}$ in the troposphere is converted into TFA.²⁰ If $\text{CF}_3\text{C}(\text{O})\text{Cl}$ is formed as a degradation product in the stratosphere, its photolytic lifetime is approximately 16 days and conversion to TFA will be of little importance.⁸

Compounds with a perfluoroalkyl group can degrade to give perfluoroalkyl radicals. The sole atmospheric fate of alkyl radicals is addition of O_2 to give alkylperoxy radicals. Alkylperoxy radicals undergo self- and cross-reactions *via* the Russell mechanism.²¹ One channel of the Russell mechanism involves transfer of a hydrogen atom between peroxy radicals which gives alcohol and acyl products. For perfluoroalkylperoxy radicals this channel sets in motion a sequence of reactions leading to PFCAs. In sequence, $\text{C}_x\text{F}_{2x+1}\text{O}_2$ radicals react with α -hydrogen containing peroxy radicals (*e.g.*, CH_3O_2) to give perfluoroalcohols ($\text{C}_x\text{F}_{2x+1}\text{OH}$) that eliminate HF to give perfluoroacylfluorides ($\text{C}_{x-1}\text{F}_{2x-1}\text{C}(\text{O})\text{F}$) which then undergo hydrolysis to give PFCAs.²² Another channel of the Russell mechanism leads to the formation of alkoxy radicals. Linear perfluoroalkoxy radicals decompose rapidly by eliminating COF_2 resulting in a perfluoroalkyl radical with one less carbon atom than the parent, *i.e.*, the perfluorinated chain “unzips”. The result in the atmosphere is the formation of a series of PFCAs in small yields (1–10%).²³

HFES give fluorinated esters as intermediate degradation products. These are removed from the atmosphere by reaction with OH radicals and by dissolution in clouds and seawater. Hydrolysis of fluorinated esters can result in the formation of PFCAs.^{24,25}

Finally, compounds with a $\text{C}_x\text{F}_{2x+1}\text{CH}$ -moiety ($x \leq 3$) can give perfluoroaldehydes, $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{H}$, as a degradation product. These aldehydes are primarily formed from parent compounds whose atmospheric lifetimes are too short to allow transport to the stratosphere to any significant degree. Hence, the degradation of $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{H}$ generated from Montreal Protocol related gases occurs mainly under tropospheric conditions. Recent studies have improved our understanding of the relative importance of the photochemical pathways leading to formation of PFCAs from $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{H}$. These are discussed in the following section.

Table 1 summarizes the estimated molar yields of PFCAs (TFA, PFPrA, and PFBA) from the primary degradation products and radical intermediates in the processes described above.

3 Fate of $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{H}$, $x = 1–3$ and the impact on formation of PFCAs

The formation of PFCAs from $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{H}$ can occur through several different pathways. Reaction with OH radicals gives $\text{C}_x\text{F}_{2x+1}\text{CO}$ radicals. This reaction is thought to be a minor sink for $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{H}$, with an estimated lifetime of approximately 23 days.¹⁵ At 298 K and one atmosphere of air 1%, 50%, and 79% ($x = 1, 2$, and 3) of $\text{C}_x\text{F}_{2x+1}\text{CO}$ radicals decompose to give perfluoroalkyl radicals and CO. The remaining $\text{C}_x\text{F}_{2x+1}\text{CO}$ radicals add O_2 to form acylperoxy radicals, $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{O}_2$.²⁶ These acylperoxy radicals then react with NO, NO_2 , and HO_2 , radicals. Reaction of $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{O}_2$ with NO gives perfluorinated acyloxy radicals which decompose by eliminating CO_2 to give $\text{C}_x\text{F}_{2x+1}$ radicals. Reaction of $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{O}_2$ with NO_2 gives perfluorinated peroxyacyl nitrates. Reaction of $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{O}_2$ with HO_2 radicals leads to the formation of








Fig. 1 Key intermediates and degradation pathways leading to formation of TFA during atmospheric degradation of HCFCs, HFCs, HFOs, and HCFOs. Hydrolysis of trifluoroacetyl fluoride ($\text{CF}_3\text{C}(\text{O})\text{F}$) and trifluoroacetyl chloride ($\text{CF}_3\text{C}(\text{O})\text{Cl}$) leads to TFA in unity molar yield. Hydration of trifluoroacetaldehyde ($\text{CF}_3\text{C}(\text{O})\text{H}$) and reactions of the trifluoroacylperoxy radical ($\text{CF}_3\text{C}(\text{O})\text{O}_2$) can lead to formation of TFA. Approximate atmospheric lifetimes are indicated in parenthesis. In addition (not shown in figure), dissolution and hydrolysis of esters generated in the atmospheric oxidation of HFEs can lead to the formation of TFA. Similar mechanistic pathways lead to the formation of PFPrA and PFBA.

$\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{OH}$, e.g., ($39 \pm 4\%$ TFA from $\text{CF}_3\text{C}(\text{O})\text{O}_2$, at room temperature).^{26,32} This reaction is likely temperature dependent. In the analogous reaction of $\text{CH}_3\text{C}(\text{O})\text{O}_2$ radicals with HO_2 , the channel giving $\text{CH}_3\text{C}(\text{O})\text{OH}$ increases in importance as the temperature decreases.³³ Hence, the reaction of $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{O}_2$ with HO_2 radicals may produce increased yields of PFCA at lower temperatures, but until experimental studies have confirmed this, we chose to use the reported room temperature data.

Perfluorinated peroxyacyl nitrates are relatively stable and can undergo long range transport, e.g., trifluoro peroxyacetyl nitrate ($\text{CF}_3\text{C}(\text{O})\text{O}_2\text{NO}_2$, FPAN) has a lifetime >1 month above altitudes of about 3 km.³⁴ Two recent studies have investigated the reaction of FPAN with water vapor to establish the importance of this reaction and the potential molar yield of TFA and/or FPAN hydrate.^{35,36}



The formation of TFA was observed in both studies. An upper limit for the rate of the gas-phase reaction of FPAN with H_2O was established and it was concluded that this reaction is unlikely to be an important source of TFA ($<1\%$).³⁶ Still, the importance of heterogeneous reactions of FPAN with water to

the formation of TFA in the atmosphere are unclear. Similar uncertainty applies to analogous reactions of the larger perfluoro acyl peroxy nitrates $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{O}_2\text{NO}_2$, $x = 2$ and 3. Thermal decomposition of FPAN would reform the $\text{CF}_3\text{C}(\text{O})\text{O}_2$ radical. Thus, FPAN could represent a temporary reservoir for $\text{CF}_3\text{C}(\text{O})\text{O}_2$ and could impact the distribution of $\text{CF}_3\text{C}(\text{O})\text{O}_2$ in the atmosphere as a precursor for the formation of TFA (see further discussion below). Until further studies become available, we do not include hydrolysis of acyl peroxy nitrates in our evaluation of the yield of PFCAs from $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{O}_2$.

The formation of PFCAs from $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{O}_2$ depends on the local environment and the atmospheric abundance ratio of NO to HO_2 . The effective yield of TFA from $\text{CF}_3\text{C}(\text{O})$ in the troposphere was indirectly accessed by Sulbaek Andersen *et al.*²⁸ in a global modelling study of emissions of HCFO-1233(zd). In their work, the atmospheric lifetimes for $\text{CF}_3\text{C}(\text{O})\text{H}$ were 2 days and 20 days with respect to photolysis and reaction with OH radicals, respectively. With an overall TFA yield of 2%, the model results of Sulbaek Andersen *et al.* suggest that the molar yield of TFA in the reaction of OH radicals with $\text{CF}_3\text{C}(\text{O})\text{H}$ (equivalent to considering the yield of TFA from $\text{CF}_3\text{C}(\text{O})$), is on the order of $\sim 20\%$. For $\text{CF}_3\text{CF}_2\text{C}(\text{O})$, Sulbaek Andersen *et al.*²⁷ evaluated the importance of the NO/ HO_2 reactions and estimated a $\sim 10\%$ yield of PFPrA from the reaction of HO_2 with $\text{CF}_3\text{CF}_2\text{C}(\text{O})$. Using the approach of Sulbaek Andersen *et al.*²⁷ the





Table 1 Molar yields of PFCAs (TFA, PFPrA, and PFBA) from intermediates and radicals in the degradation of volatile fluorinated organic compounds relevant to the Montreal Protocol. An em dash (—) indicates that no formation (0% molar yield) of the acid is predicted

	Atmospheric fate	Lifetime ^e	Yield estimates (%)			Effective PFCA yield ^f (%)	Notes/references
			TFA	PFPrA	PFBA		
Primary degradation products							
CF ₃ C(O)F	UV photolysis (strat. only) Deposition/hydrolysis	≈100 days 5–15 days	— 100	— —	— —	TFA: 10 (strat.)–100 (trop.) TFA: 0 (trop.)–<10 (strat.)	Jubb <i>et al.</i> (2015) ¹⁸ Wallington <i>et al.</i> (1994) ¹⁷ Jubb <i>et al.</i> (2015) ¹⁸
CF ₃ CF ₂ C(O)F	UV photolysis (strat. only) ^c Deposition/hydrolysis	≈100 days 5–15 days	1–10 —	100	—	PFPrA: 10 (strat.)–100 (trop.) TFA: 0 (trop.)–<10 (strat.)	Wallington <i>et al.</i> (1994) ¹⁷ Jubb <i>et al.</i> (2015) ¹⁸
CF ₃ CF ₂ CF ₂ C(O)F	UV photolysis (strat. only) ^c Deposition/hydrolysis	≈100 days 5–15 days	1–10 —	—	100	PFPrA: 0 (trop.)–<10 (strat.) PFBA: 10 (strat.)–100 (trop.)	Wallington <i>et al.</i> (1994) ¹⁷
CF ₃ C(O)Cl	UV photolysis ^c	Strat: ~16 days	≤10	—	—	TFA: <10 (strat.)–60 (trop.)	Photolysis branching: CF ₃ C(O) + Cl/CF ₃ + C(O)Cl Tropospheric fate: Hayman <i>et al.</i> (1994) ²⁰
CF ₃ C(O)H	Deposition/hydrolysis OH reaction ^c UV photolysis Deposition/hydrolysis	Trop: 23 days 5–30 days 23 days 3 days ≥2 days	≤10 100 20 — ≤100	— — — — —	— — — — —	TFA: <58	OH reaction: → CF ₃ C(O) ~2% CF ₃ C(O) decomp Hurley <i>et al.</i> (2006) ²⁶ Photolysis: >98% CF ₃ + HCO OH reaction: → CF ₃ CF ₂ C(O) ~52% CF ₃ CF ₂ C(O) decomp Hurley <i>et al.</i> (2006) ²⁶ OH reaction: → CF ₃ CF ₂ C(O) decomp ~81% CF ₃ CF ₂ C(O) decomp Hurley <i>et al.</i> (2006) ²⁶
C ₂ F ₅ C(O)H	OH reaction ^c UV photolysis ^c Deposition/hydrolysis	23 days 0.8 days ≥2 days	1–10 1–10 —	10 — ≤100	— — —	TFA: <7, PFPrA: <28	
<i>n</i> -C ₃ F ₇ C(O)H	OH reaction ^c UV photolysis ^c Deposition/hydrolysis	23 days 0.5 days ≥2 days	1–10 1–10 —	1–10 1–10 —	5 — ≤100	TFA: <8 PFPrA: <8, PFBA: <20	
CF ₃ C(O)CF ₃	Deposition/hydrolysis	≥2 days	—	—	—	TFA: 20 ± 10	Hurley <i>et al.</i> (2006) ²⁶
CHF ₂ C(O)CF ₃	UV photolysis ^c	27 days	20 ± 10	—	—	TFA: 20 ± 10	(→ CF ₃ C(O) + CF ₃) Calvert <i>et al.</i> (2011) ¹⁹
(CF ₃) ₂ HCOC(O)F	UV photolysis ^c	n/a	20 ± 10	—	—	TFA: <100	(→ CF ₃ C(O) + CF ₃) Calvert <i>et al.</i> (2011) ¹⁹
C ₄ F ₉ OC(O)H	Deposition/hydrolysis OH reaction ^c Deposition/hydrolysis	2 years n/a	1–10 —	1–10 —	1–10 100	TFA: <5 PFPrA: <5, PFBA: <55	Kutsuna <i>et al.</i> (2004) ²⁵ Kutsuna <i>et al.</i> (2005) ²⁴ OH reaction and hydrolysis assumed equally important
C ₂ F ₅ CH ₂ OC(O)H	OH reaction ^c Deposition/hydrolysis	n/a n/a	<7 —	<28 100	— —	TFA: 100	Kutsuna <i>et al.</i> (2005) ²⁴ OH reaction and hydrolysis assumed equally important
CF ₃ CHF ₂ OC(O)H	OH reaction ^c Deposition/hydrolysis	n/a n/a	— 100	— —	— —	TFA: <5	Kutsuna <i>et al.</i> (2005) ²⁴ OH reaction and hydrolysis assumed equally important
<i>n</i> -C ₃ F ₇ OC(O)CH ₃	Deposition/hydrolysis OH reaction ^c Deposition/hydrolysis	n/a n/a n/a	100 1–10 —	— 1–10 —	— — —	TFA: <5, PFPrA: <55	Kutsuna <i>et al.</i> (2005) ²⁴ OH reaction and hydrolysis assumed equally important
C ₄ F ₉ OC(O)CH ₃	Deposition/hydrolysis ^c OH reaction ^c	n/a n/a	1–10 —	100 —	1–10 —	TFA: <5	Kutsuna <i>et al.</i> (2005) ²⁴ OH reaction and hydrolysis assumed equally important
CF ₃ C(O)OCHE ₂	Deposition/hydrolysis OH reaction ^c	n/a >1 year	1–10 —	— —	100 —	PFPrA: <5, PFBA: <55	Kutsuna <i>et al.</i> (2005) ²⁴ OH reaction and hydrolysis assumed equally important
C ₃ F ₅ C(O)OCHF ₂	Deposition/hydrolysis ^c OH reaction ^c	n/a >1 year	<20 100	— —	— —	TFA: 100	Kutsuna <i>et al.</i> (2004, 2005) ^{24,25} Hydrolysis assumed dominant
C ₃ F ₅ C(O)OCHF ₂	OH reaction ^c Deposition/hydrolysis	>1 year ~1 year	<10 —	≈10 —	— —	TFA: <10, PFPrA: <100	Sulbaek Andersen <i>et al.</i> (2024) ²⁷ Kutsuna <i>et al.</i> (2004, 2005) ^{24,25}

Table 1 (Contd.)

	Atmospheric fate	Lifetime ^e	Yield estimates (%)			Effective PFCA yield ^b (%)	Notes/references
			TFA	PFPrA	PFBA		
CF ₃ CHFCF ₂ OC(O)	OH reaction ^c	n/a	<110	—	—	TFA: 53, PFPrA: 50	Kutsuna <i>et al.</i> (2005), ²⁴ OH reaction and hydrolysis assumed equally important
CF ₂ CF ₃	Deposition/hydrolysis	n/a	<100	100	—	—	Kutsuna <i>et al.</i> (2005) ²⁴
C ₃ F ₇ CF(OC(O)CH ₃)	OH reaction ^c	n/a	100	—	100	TFA: <110	Hydrolysis assumed dominant
CF(CF ₂) ₂	Deposition/hydrolysis ^c	n/a	<110	<10	<5	PFPrA: <10, PFBA: <5	Kutsuna <i>et al.</i> (2005), ²⁴ OH reaction and hydrolysis assumed equally important
C ₂ F ₅ CF(OC(O)H)	OH reaction ^c	>2 years	100	100	—	TFA: 100, PFPrA: 50	Kutsuna <i>et al.</i> (2005), ²⁴ OH reaction and hydrolysis assumed equally important
CF(CF ₂) ₂	Deposition/hydrolysis	n/a	100	—	—	TFA: <110	Kutsuna <i>et al.</i> (2005) ²⁴
C ₃ F ₇ CF(OC(O)H)	OH reaction ^c	n/a	100	—	100	PFPrA: <10, PFBA: <5	Hydrolysis assumed dominant
CF(CF ₂) ₂	Deposition/hydrolysis ^c	n/a	<110	<10	<5	—	Ellis <i>et al.</i> (2004) ²²
Radical intermediates							
CF ₃ CF ₂	O ₂ /CH ₃ O ₂ → TFA	μs	1–10	—	—	TFA: <10	—
	O ₂ /NO → CF ₃ + CF ₂ O + NO ₂	μs	0	—	—	—	—
	O ₂ /CH ₃ O ₂ → PFPrA	μs	1–10	1–10	—	TFA: <10, PFPrA: <10	Ellis <i>et al.</i> (2004) ²²
	TFA						
	O ₂ /NO → CF ₃ CF ₂ + CF ₂ O + NO ₂	μs	—	—	—	—	—
	O ₂ /CH ₃ O ₂ → PFBA	μs	1–10	1–10	1–10	TFA: <10	Ellis <i>et al.</i> (2004) ²²
	PFPrA, TFA						
	O ₂ /NO → CF ₃ CF ₂ CF ₂ + CF ₂ O + NO ₂	mins	—	—	—	PFPrA: <10, PFBA: <10	—
	O ₂ /HO ₂ → TFA	mins	39	—	—	TFA: 20 ± 10	~2% CF ₃ C(O) undergoes prompt decomp. Hurley <i>et al.</i> (2006) ²⁶
	O ₂ /NO → CF ₃ + CO ₂ + NO ₂	mins	—	—	—	—	Sulbaek Andersen <i>et al.</i> (2018) ²⁸
	O ₂ /HO ₂ → PFPrA	mins	—	50	—	PFPrA: ≈ 10	~52% CF ₃ CF ₂ C(O) undergoes prompt decomp. Hurley <i>et al.</i> (2006) ²⁶
	O ₂ /NO → CF ₃ CF ₂ + CO ₂ + NO ₂ ^c	mins	1–10	—	—	—	—
	O ₂ /HO ₂ → PFBA	mins	—	—	53	PFBA: ≈ 5	~81% CF ₃ CF ₂ C(O) undergoes prompt decomp. Hurley <i>et al.</i> (2006) ²⁶
	O ₂ /NO → CF ₃ CF ₂ CF ₂ + CO ₂ + NO ₂ ^c	sec-mins	1–10	1–10	—	—	—
	+ RO ₂ → CF ₃ CHFO + RO + O ₂ ^c		25–81	—	—	TFA: 7–20	Wallington <i>et al.</i> (1996) ²⁹
	+ NO → CF ₃ CFHO + NO ₂ ^c		25–81	—	—	—	Wallington <i>et al.</i> (2017) ³⁰
	→ CF ₃ + HC(O)F + NO ₂		0	—	—	—	- See also reaction below
	+ O ₂ → CF ₃ C(O)F + HO ₂ ^c	μs-ms	25–81	—	—	TFA: 25–81	Pressure/temperature dep
	+ Δ → CF ₃ + HC(O)F	μs-ms	0	—	—	—	Wallington <i>et al.</i> (2017) ³⁰
	+ O ₂ → CF ₃ CF ₂ C(O)F + HO ₂ ^c	μs-ms	—	100	—	TFA: <10, PFPrA: <1	Mogelberg <i>et al.</i> (1997) ³¹
	+ Δ → CF ₃ CF ₂ + HC(O)F ^c		1–10	—	—	—	—

^a Stratospheric lifetime is a local not a global lifetime. ^b Effective PFCA yields include competitive direct (e.g. wet deposition/hydrolysis) and indirect sources (e.g. further reactions of radical intermediates) in the atmospheric oxidation of the individual species. ^c Loss process forms a product or radical intermediate that leads to the formation of TFA, PFPrA or PFBA.



yield of PFBA from atmospheric processing of $C_3F_7C(O)$ can be estimated as approximately 5%.

Long *et al.*³⁷ recently conducted a computational study and suggested that reaction with HO_2 radicals is an important sink for aldehydes in the atmosphere, incl. for $CF_3C(O)H$. For $CF_3C(O)H$, this reaction would produce a α -hydroxy substituted peroxy radical, which may undergo further reactions to yield TFA. Presently, there are no experimental data for this reaction, and its atmospheric importance remains unclear. Until experimental data becomes available, we have not included the reaction of HO_2 reaction with $C_xF_{2x+1}C(O)H$ in our evaluation of the yield of PFCAs from $C_xF_{2x+1}C(O)H$.

Photolysis is a major tropospheric sink for $C_xF_{2x+1}C(O)H$ and dominates in the stratosphere. While there have been few studies of the quantum yields and photolysis lifetimes of $C_2F_5C(O)H$ and $n-C_3F_7C(O)H$, the photolysis of $CF_3C(O)H$ has been extensively studied recently, largely due to concerns about photochemical production of HFC-23 (CHF_3).³⁸ The photolysis of $CF_3C(O)H$ in the troposphere proceeds mainly by two channels:






Fig. 2 Estimated average photolytic lifetime using TUV model of $CF_3C(O)H$ ($x = 1$, blue trace/circles), $C_2F_5C(O)H$ ($x = 2$, green trace/squares) and $n-C_3F_7C(O)H$ ($x = 3$, red trace/triangles) from 0 to 20 km at equinox. The hatched horizontal area denotes the density-weighted average altitude of the troposphere of 4.5 km. The insert shows the quantum yields used (at one atmosphere of pressure). For $CF_3C(O)H$, linear interpolation is used between the IUPAC recommended pressure dependent quantum yields at 248, 266, and 281 nm (blue solid trace), with an exponential decrease to a long-wavelength limit set to 360 nm. The quantum yields used for $C_2F_5C(O)H$ and $C_3F_7C(O)H$ follow those of $CF_3C(O)H$, with a shift of 10 nm towards the red (red/green dashed trace). A Stern–Volmer pressure dependency was applied in the TUV for all three compounds. The insert also shows the UV absorbance spectra of $CF_3C(O)H$, $C_2F_5C(O)H$ and $C_3F_7C(O)H$. See text for details.

Photolysis of $CF_3C(O)H$ is not a source of TFA. Since the evaluation of TFA yields by Madronich *et al.*¹⁰ three papers have been published that provide new measurements of the quantum yields for $CF_3C(O)H$.^{12–14} The results of these studies have been evaluated by the International Union of Pure and Applied Chemistry (IUPAC) Task Group on Atmospheric Chemical Kinetic Data Evaluation.³⁹ Estimates in the literature for the atmospheric lifetime for photolysis are in the range ~ 2.5 –13 days.^{12,40} Using the updated photochemical parameters, we have reevaluated the photolysis lifetime of $CF_3C(O)H$. Atmospheric photolysis coefficients were computed with the Tropospheric Ultraviolet Visible (TUV) model version 5.4, using the pseudo-spherical 4-stream discrete ordinates radiative solver option. Wavelength-dependent data were gridded according to the WMO2 scheme,⁴¹ spanning the range 121–750 nm with enhanced resolution (1 nm) over 220–420 nm to resolve for the strong spectral gradient in tropospheric actinic flux due to stratospheric ozone absorption. The pressure dependence was estimated from the total quantum yields at atmospheric pressure (taken as NTP, normal temperature and pressure), with air number density $M = 2.45 \times 10^{19}$ molecule cm^{-3} and the assumption that the reciprocal of the yield decreases linearly to unity at zero pressure (Stern–Volmer) at all wavelengths for which the yield is non-zero. Calculations were made for the Equator and $40^\circ N$, 21st March, in 15 min steps and averaged over 24 hours. Atmospheric conditions included cloud-free, aerosol-free skies, 0.1 ground albedo, with an ozone column of 250 Dobson Units at the Equator and 350 Dobson Units for $40^\circ N$. For input in the TUV model (see insert in Fig. 2 and SI 1.1), two new data sets were compiled:

(A) A linear interpolation in wavelength between the quantum yield datapoints provided by IUPAC (248, 266, 281 and 308 nm) with a Stern–Volmer pressure dependence.³⁹ Following the approach of Sulbaek Andersen *et al.* (2023), in this dataset we assume 335 nm for the long-wavelength limit,¹² consistent with the long-wavelength zero point for the quantum yield of $CH_3C(O)H$, which has a similar absorption spectrum as $CF_3C(O)H$.⁴²

(B) A linear interpolation between the quantum yield datapoints provided by IUPAC (248, 266 nm, QY = 1) to 281 nm, with an exponential decrease to a long-wavelength limit set to 360 nm. This approach assigns a more realistic wavelength dependence to the quantum yields at the longer wavelength region, which is particularly important for species that primarily undergo decomposition in the troposphere. The long-wavelength limit reflects the measured $CF_3C(O)H$ UV spectrum (see insert in Fig. 2).

For models A and B, inclusion of a Stern–Volmer pressure dependence increases the modeled photolysis frequencies resulting in a 20–40% decrease in estimated photolytic lifetime, on average. The actinic flux in the troposphere limits the importance of photolysis at wavelengths below 300 nm. However, $CF_3C(O)H$ has an absorption maximum at 300 nm, with significant absorption between 300 and 360 nm and modeled atmospheric lifetimes are highly sensitive to the assumed quantum yield fall-off and long-wavelength limit in the region between 308 and 360 nm.



Table 2 Molar yields of PFCAs (TFA, PFPrA, and PFBA) from volatile fluorinated organic compounds relevant to the Montreal Protocol, see SI for details^a

Compound	Formula	PFCAs molar yields (%)		
		TFA	PFPrA	PFBA
CFCs^b				
CFC-113a	CCl ₃ CF ₃	<10	—	—
CFC-114a	CCl ₂ FCF ₃	≈ 10	—	—
CFC-216ba	CClF ₂ CClFCF ₃	≈ 10	—	—
HCFCs/halons^b				
HCFC-124	CHClFCF ₃	100	—	—
HCFC-133a	CH ₂ ClCF ₃	<59	—	—
HCFC-225ca	CF ₃ CF ₂ CHCl ₂	<10	—	—
HCFC-233fb	CCl ₂ FCH ₂ CF ₃	<58	—	—
HCFC-243fa	CHCl ₂ CH ₂ CF ₃	<58	—	—
HCFC-244fa	CHFClCH ₂ CF ₃	<58	—	—
HCFC-253 fb	CH ₂ ClCH ₂ CF ₃	<58	—	—
Halon-2311 (Halothane) ^b	CHBrClCF ₃	60 ± 10	—	—
HFCs^b				
HFC-125	CHF ₂ CF ₃	<10	—	—
HFC-134a	CH ₂ FCF ₃	7–20	—	—
HFC-143a	CH ₃ CF ₃	<58	—	—
HFC-227ea	CF ₃ CHFCF ₃	100	—	—
HFC-236cb	CF ₃ CF ₂ CH ₂ F	<10	<1	—
HFC-236ea	CHF ₂ CHFCF ₃	≈ 100	—	—
HFC-236fa	CF ₃ CH ₂ CF ₃	20 ± 10	—	—
HFC-245fa	CHF ₂ CH ₂ CF ₃	<33	—	—
HFC-329p	CF ₃ CF ₂ CF ₂ CHF ₂	<10	<10	<10
HFC-365mfc	CH ₃ CF ₂ CH ₂ CF ₃	<53	—	—
HFC-43-10mee	CF ₃ CHFCHFCF ₂ CF ₃	54–60	54–60	—
HFEs/HCFEs				
HFE-236ea2 (Desflurane)	CHF ₂ OCHFCF ₃	<20	—	—
HFE-347mcf	C ₂ F ₅ CH ₂ OCHF ₂	<10	≈ 10	—
HFE-347mmz1 (Sevoflurane)	(CF ₃) ₂ C(O)HCH ₂ F	<95	—	—
HFE-365mcf3	C ₂ F ₅ CH ₂ OCH ₃	<6	<36	—
HFE-54-11meef	CF ₃ CHFCF ₂ OCH ₂ C ₂ F ₅	<103	25	—
HFE-7100	C ₄ F ₉ OCH ₃	<5	<5	<55
HFE-7200	C ₄ F ₉ OC ₂ H ₅	<5	<5	<55
HFE-7300	C ₂ F ₅ CF(OCH ₃)CF(CF ₃) ₂	100	50	—
HFE-7500	C ₃ F ₇ CF(OC ₂ H ₅)CF(CF ₃) ₂	<110	<10	<5
1-Ethoxy-1,1,2,2,2,3,3,3-hepta-fluoro-propane	C ₃ F ₇ OCH ₂ CH ₃	<5	<55	—
HCFE-235da2 (Isoflurane)	CHF ₂ OCHClCF ₃	≈ 98	—	—
HFOs/HCFOs/HBFOs				
HFO-1225zc	CF ₂ =CHCF ₃	<58	—	—
HFO-1234yf	CF ₃ CF=CH ₂	100	—	—
HFO-1234ze(<i>Z/E</i>)	(<i>E</i>)-CHF=CHCF ₃	<58	—	—
HFO-1243zf	CH ₂ =CHCF ₃	<58	—	—
HFO-1336mzz(<i>E/Z</i>)	(<i>E</i>)-CF ₃ CH=CHCF ₃	<116	—	—
HFO-1345zfc	CH ₂ =CHCF ₂ CF ₃	<7	<28	—
HFO-1438mzz(<i>E</i>)	(<i>E</i>)-CF ₃ CH=CHCF ₂ CF ₃	<65	<28	—
HFO-1447fz	CH ₂ =CHCF ₂ CF ₂ CF ₃	<8	<8	<20
HCFO-1233zd(<i>E</i>)	(<i>E</i>)-CF ₃ CH=CHCl	<58	—	—
HCFO-1233zd(<i>Z</i>)	(<i>Z</i>)-CF ₃ CH=CHCl	<58	—	—
2-Bromo-3,3,3-trifluoro-propene (2-BTP)	CF ₃ CBr=CH ₂	<58	—	—
Alcohols/ketones/nitriles				
2,2,2-Trifluoroethanol (TFE)	CF ₃ CH ₂ OH	<58	—	—
Perfluoro(2-methyl)-3-pentanone	CF ₃ CF ₂ C(O)CF(CF ₃) ₂	101–110	—	—
Heptafluorobutyro-nitrile	(CF ₃) ₂ CFCN	≈ 100	—	—

^a (—) indicates that no formation (0% molar yield) of the acid is predicted. ^b Compounds regulated under the Montreal Protocol and its amendments. Note that Halothane (a Halon) is exempt from regulation under the Montreal Protocol, but a switch by the healthcare industry to non-ozone-depleting alternatives means that emissions are declining.⁵²

Fig. 2 includes the trace of the resulting atmospheric photolysis lifetime for CF₃C(O)H. The troposphere extends to approximately 10 km altitude with a density weighted average

altitude of approximately 4.5 km.⁴³ At this altitude the average lifetime due to photolysis of CF₃C(O)H in the lower atmosphere is approximately 3 days. The modeled photolysis frequencies



using either model A or B produce near identical lifetime estimates (see SI Fig. 1.2). This can be explained by the fact that the derivative for the photolysis frequency with respect to wavelength ($dJ/d\lambda$, $s^{-1} \text{ nm}^{-1}$) is significantly larger for model A than model B in the range 308–330 nm, while an approximate equal, but opposite difference is observed for the range 335–360 nm, *i.e.*, the long-wavelength limit region (see SI Fig. 1.3). The region below ~ 300 nm, contributes very little to the photolytic lifetime of $\text{CF}_3\text{C}(\text{O})\text{H}$. Notably, the slightly lower 308 nm quantum yield value used in the quantum yield fit by Sulbaek Andersen *et al.*¹² ($QY = 0.16$ vs. 0.19 (IUPAC³⁹)) has an amplified impact on the predicted atmospheric lifetime, as this is the regional peak for $dJ/d\lambda$.

Relatively few studies have been conducted on the atmospheric lifetime of the longer chain-length perfluoro aldehydes, $\text{C}_2\text{F}_5\text{C}(\text{O})\text{H}$ and $n\text{-C}_3\text{F}_7\text{C}(\text{O})\text{H}$.^{44,45} For $\text{C}_2\text{F}_5\text{C}(\text{O})\text{H}$, Chiappero *et al.*⁴⁰ give annual averages for photolysis lifetimes of approximately 0.9 days at 11 km altitude and 2.5 days at 0 km, while Antinolo *et al.*⁴⁶ estimate a photolysis lifetime of 3.5 hours at 3.5 km altitude and solar zenith angle of 16° (summertime, noon, Spain). For $n\text{-C}_3\text{F}_7\text{C}(\text{O})\text{H}$, Chiappero *et al.*⁴⁰ averaged their measurements for $\text{CF}_3\text{C}(\text{O})\text{H}$ and $n\text{-C}_4\text{F}_9\text{C}(\text{O})\text{H}$ and used the resulting wavelength independent photolysis quantum yield (0.11) to estimate annual averages for the atmospheric lifetimes of approximately 0.75 days at 11 km altitude and 2 days at 0 km. Solignac *et al.*⁴⁷ studied the photolysis of $n\text{-C}_3\text{F}_7\text{C}(\text{O})\text{H}$ in the EUPHORE chamber in Valencia, Spain and estimated a photolytic lifetime of close to one day.

It is likely that the photolysis mechanism for $\text{C}_2\text{F}_5\text{C}(\text{O})\text{H}$ and $n\text{-C}_3\text{F}_7\text{C}(\text{O})\text{H}$ will be similar to that of $\text{CF}_3\text{C}(\text{O})\text{H}$. The peaks of the UV absorbance for $\text{C}_2\text{F}_5\text{C}(\text{O})\text{H}$ and $n\text{-C}_3\text{F}_7\text{C}(\text{O})\text{H}$ are nearly identical, and both shifted approximately 10 nm to the red in comparison to that of $\text{CF}_3\text{C}(\text{O})\text{H}$ (see insert in Fig. 2). Here, we use a self-consistent approach to model the photolytic lifetimes $\text{C}_2\text{F}_5\text{C}(\text{O})\text{H}$ and $n\text{-C}_3\text{F}_7\text{C}(\text{O})\text{H}$, assuming that the wavelength-dependent quantum yields of $\text{C}_2\text{F}_5\text{C}(\text{O})\text{H}$ and $n\text{-C}_3\text{F}_7\text{C}(\text{O})\text{H}$ are equal to that of $\text{CF}_3\text{C}(\text{O})\text{H}$, red-shifted by 10 nm, and following a Stern–Volmer pressure dependency as discussed above. As seen from Fig. 2, the atmospheric lifetimes with respect to photolysis for $\text{C}_2\text{F}_5\text{C}(\text{O})\text{H}$ and $n\text{-C}_3\text{F}_7\text{C}(\text{O})\text{H}$ are shorter than for $\text{CF}_3\text{C}(\text{O})\text{H}$, and both less than 1 day (0.8 and 0.5 days at 4.5 km altitude, respectively).

Finally, in addition to photolysis (major sink) and reaction with OH radicals (minor sink), $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{H}$ can undergo dry and wet deposition. On contact with liquid water, $\text{CF}_3\text{C}(\text{O})\text{H}$ reacts to give an aldehyde hydrate (gem-diol, $\text{CF}_3\text{CH}(\text{OH})_2$).⁴⁸ The gem-diol, at least in the gas-phase, reacts with OH radicals (lifetime of approximately 90 days) and generates TFA.⁴⁸ Unfortunately, key physical parameters needed to estimate dry and wet deposition velocities for $\text{CF}_3\text{C}(\text{O})\text{H}$ are presently unavailable. Measurements have not been reported for the hydration equilibrium constant, the Henry's Law Constant (H), or the effective HLC (H^*) for $\text{CF}_3\text{C}(\text{O})\text{H}$. Nielsen *et al.*⁴⁹ empirically estimated $H^* > 10^4 \text{ M atm}^{-1}$, which suggests a wet-deposition lifetime of a few days (4–8 days). Dry deposition may be competitive with wet deposition but is highly dependent on season and geographical location. This, and the absence of

any experimental data for $\text{CF}_3\text{C}(\text{O})\text{H}$, renders estimation of dry deposition velocities for $\text{CF}_3\text{C}(\text{O})\text{H}$ at least as uncertain as those estimated for wet deposition. The atmospheric lifetime of species such as HNO_3 , which deposit without surface resistance, provides an upper limit for the importance of dry and wet deposition for $\text{CF}_3\text{C}(\text{O})\text{H}$.⁵⁰ Dry deposition of HNO_3 occurs on a timescale of 2–3 days, wet deposition in precipitation is more rapid but episodic.⁵¹ We calculate the overall yield of TFA from $\text{CF}_3\text{C}(\text{O})\text{H}$ by assuming a 2-day lower limit for the deposition lifetime (*i.e.*, upper limit for the importance of dry and wet deposition) and that hydration is efficient, converting $\text{CF}_3\text{C}(\text{O})\text{H}$ into TFA in a yield of unity (100%). We make the same assumption for the longer chain-length aldehydes, $\text{C}_2\text{F}_5\text{C}(\text{O})\text{H}$ and $n\text{-C}_3\text{F}_7\text{C}(\text{O})\text{H}$.

Table 1 includes the estimated molar yields of PFCAs (TFA, PFPrA, and PFBA) from $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{H}$ based on the assumptions and caveats for the evaluated atmospheric sinks described above. For $\text{CF}_3\text{C}(\text{O})\text{H}$, based on the preceding analysis, we estimate an upper limit of $<58\%$ molar yield of TFA. The overall yield of TFA would be reduced to 27%, if the upper limit for the estimated wet-deposition lifetime (8 days) was used instead of a 2-day deposition lifetime. The parameterization of wet/dry deposition for $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{H}$ remains a significant source of uncertainty in quantifying an upper limit for PFCAs yields.

4 Conclusion

A systematic evaluation of the yields of TFA, PFPrA, and PFBA from the atmospheric degradation of gases relevant to the Montreal Protocol and its Amendments is presented. Table 2 lists the estimated overall yields of PFCAs from these compounds. These overall yields are obtained as the sum of the individual contributions from the yields of primary degradation products multiplied by their evaluated yields of PFCAs (see Table 1). Details for the yields of PFCAs for each compound are given in the SI Section 2. A reevaluation of the atmospheric photolytic lifetime for $\text{CF}_3\text{C}(\text{O})\text{H}$, $\text{C}_2\text{F}_5\text{C}(\text{O})\text{H}$, and $n\text{-C}_3\text{F}_7\text{C}(\text{O})\text{H}$ leads to revisions in the PFCAs yields of many of the species listed in Table 2 from those reported previously.^{10,11}

As seen from inspection of the values in Tables 1 and 2, there are substantial uncertainties in many of the estimated yields. For many compounds, it is only possible to provide upper limits for PFCAs yields. The uncertainties associated with the yield estimates have two origins. First, inherent uncertainties in the measurements of physical parameters *e.g.* rate coefficients, absorption cross sections, and quantum yields, which are typically 5–20% of the measured values and are relatively straightforward to quantify. Second, for PFCAs formation pathways with few, or no, direct measurements available, *e.g.*, wet deposition of $\text{CF}_3\text{C}(\text{O})\text{H}$, the yield estimates are based on semi-empirical extrapolations. The uncertainties associated with such extrapolations are significant, difficult to estimate and, in many cases, we can only provide upper limits for PFCAs yields.

Without knowledge of the fate of liquid phase $\text{CF}_3\text{C}(\text{O})\text{H}$, or its hydrate, after uptake into liquid water and deposition to surface waters, it is difficult to properly evaluate this contribution to the yields of TFA. However, the likely importance of wet



and dry deposition of $\text{CF}_3\text{C}(\text{O})\text{H}$, and the possibility that aqueous chemistry of $\text{CF}_3\text{C}(\text{O})\text{H}$ /hydrates would be an effective source of TFA require that this sink be assessed. Further work to better understand the aqueous chemistry of $\text{CF}_3\text{C}(\text{O})\text{H}$ /hydrates; hydrolysis of FPAN; quantum yields for photolysis of $\text{C}_2\text{F}_5\text{C}(\text{O})\text{H}$, and $n\text{-C}_3\text{F}_7\text{C}(\text{O})\text{H}$ at wavelengths and pressures relevant to the troposphere; and the atmospheric fate of fluorinated esters is also needed. Closing these knowledge gaps will significantly advance our understanding of the environmental fate of precursors to PFCAs such as TFA, PrPFA, and PFBA.

Author contributions

M. P. Sulbaek Andersen: conceptualization, formal analysis, investigation, methodology visualization, writing – original draft; T. J. Wallington: investigation, writing – review & editing; J. B. Burkholder: investigation, writing – review & editing; S. Madronich: investigation, writing – review & editing; M. L. Hanson: writing – review & editing; D. Van Hooymissen: writing – review & editing; K. R. Solomon: writing – review & editing.

Conflicts of interest

There are no conflicts to declare.

Data availability

Data supporting this article have been included as part of the supplementary information (SI). Supplementary information: SI 1 $\text{CF}_3\text{C}(\text{O})\text{H}$, $\text{C}_2\text{F}_5\text{C}(\text{O})\text{H}$ and $n\text{-C}_3\text{F}_7\text{C}(\text{O})\text{H}$ quantum yields, UV spectra and lifetimes with respect to photolysis. SI 1.1 $\text{CF}_3\text{C}(\text{O})\text{H}$ quantum yields used for TUV models. SI 1.2 $\text{CF}_3\text{C}(\text{O})\text{H}$ lifetimes. SI 1.3 Spectral contributions to the photolysis coefficient of $\text{CF}_3\text{C}(\text{O})\text{H}$ using the two different quantum yield models (Models A and B). SI 1.4 TUV model data input (tabulated). SI 2 Estimated perfluorocarboxylic acid yields from compounds listed in Table 2. See DOI: <https://doi.org/10.1039/d5ea00179j>.

Acknowledgements

We thank Stefan Reimann for helpful discussions.

References

- 1 R. C. Buck, J. Franklin, U. Berger, J. M. Conder, I. T. Cousins, P. de Voogt, A. A. Jensen, K. Kannan, S. A. Mabury and S. P. van Leeuwen, Perfluoroalkyl and polyfluoroalkyl substances in the environment: terminology, classification, and origins, *Integr. Environ. Assess.*, 2011, 7, 513.
- 2 Organisation for Economic Co-operation and Development (OECD), *OECD Environment, Health and Safety Publications Series on Risk Management No. 39. Toward a new comprehensive global database of per- and polyfluoroalkyl substances (PFASs): Summary Report on Updating the OECD 2007 List of Per- and Polyfluoroalkyl Substances (PFASs)*, 2018, [https://one.oecd.org/document/ENV/JM/MONO\(2018\)7/en/pdf](https://one.oecd.org/document/ENV/JM/MONO(2018)7/en/pdf), accessed December 31, 2025.
- 3 Stockholm Convention, *Stockholm Convention on Persistent Organic Pollutants (POPs)*, <https://www.pops.int/TheConvention/ThePOPs/TheNewPOPs/tabid/2511/Default.aspx>, accessed December 31, 2025.
- 4 Environmental Protection Agency, Human health toxicity assessment for perfluorooctanoic acid (PFOA) and related salts, *EPA Document 815R24006*, 2024. <https://www.epa.gov/system/files/documents/2024-05/final-human-health-toxicity-assessment-pfoa.pdf>, accessed December 31, 2025.
- 5 J. C. Boutonnet, P. Bingham, D. Calamari, C. d. Rooij, J. Franklin, T. Kawano, J.-M. Libre, A. McCulloch, G. Malinverno, J. M. Odom, G. M. Rusch, K. Smythe, I. Sobolev, R. Thompson and J. M. Tiedje, Environmental risk assessment of trifluoroacetic acid, *Human Ecol. Risk Assess.*, 1999, 5, 59.
- 6 K. R. Solomon, G. J. M. Velders, S. R. Wilson, S. Madronich, J. Longstreth, P. J. Aucamp and J. F. Bornman, Sources, fates, toxicity, and risks of trifluoroacetic acid and its salts: Relevance to substances regulated under the Montreal and Kyoto Protocols, *J. Toxicol. Environ. Health, Part B*, 2016, 19, 289.
- 7 World Meteorological Organization (WMO), *Scientific Assessment of Ozone Depletion: 2022, GAW Report No. 278*, WMO, Geneva, 2022.
- 8 J. B. Burkholder, R. A. Cox and A. R. Ravishankara, Atmospheric degradation of ozone depleting substances, their substitutes, and related species, *Chem. Rev.*, 2015, 115, 3704.
- 9 M. A. H. Khan, D. C. Mendes, R. E. T. Holland, M. de los Angeles Garavagno, A. J. Orr-Ewing, K. M. Stanley, S. J. O'Doherty, D. Young, M. K. Vollmer, A. J. Antony, F. Karamshahi, T. J. Wallington, C. J. Percival, A. Bacak, R. G. Derwent and D. E. Shallcross, Global modeling of trifluoroacetic acid surface concentration and deposition from the gas-phase oxidation of a wide range of precursor hydrofluoroolefins, *Environ. Sci.: Atmos.*, 2026, DOI: [10.1039/d5ea00108k](https://doi.org/10.1039/d5ea00108k), advance article.
- 10 S. Madronich, B. Sulzberger, J. D. Longstreth, T. Schikowski, M. P. Sulbaek Andersen, K. R. Solomon and S. R. Wilson, Changes in tropospheric air quality related to the protection of stratospheric ozone in a changing climate, *Photochem. Photobiol. Sci.*, 2023, 22, 1129.
- 11 S. Madronich, G. H. Bernhard, P. J. Neale, A. Heikkilä, M. P. Sulbaek Andersen, A. L. Andrady, P. J. Aucamp, A. F. Bais, A. T. Banaszak, P. J. Barnes, J. F. Bornman, L. S. Bruckman, R. Busquets, G. Chiodo, D.-P. Häder, M. L. Hanson, S. Hylander, M. A. K. Jansen, G. Lingham, R. M. Lucas, R. Mackenzie Calderon, C. Olsen, R. Ossola, K. K. Pandey, I. Petropavlovskikh, L. E. Revell, L. E. Rhodes, S. A. Robinson, T. M. Robson, K. C. Rose, T. Schikowski, K. R. Solomon, B. Sulzberger, T. J. Wallington, Q.-W. Wang, S.-Å. Wängberg, C. C. White, S. R. Wilson, L. Zhu and R. E. Neale, Continuing benefits of the Montreal Protocol and protection of the stratospheric ozone layer for human health and the environment, *Photochem. Photobiol. Sci.*, 2024, 23, 1087.



- 12 M. P. Sulbaek Andersen, S. Madronich, J. M. Ohide, M. Frausig and O. J. Nielsen, Photolysis of CF_3CHO at 254 nm and potential contribution to the atmospheric abundance of HFC-23, *Atmos. Environ.*, 2023, **314**, 120087.
- 13 J. D. Thomson, J. S. Campbell, E. B. Edwards, C. Medcraft, K. Nauta, K. M. Perez-Pena, J. A. Fisher, D. L. Osborn, S. H. Kable and C. S. Hansen, Fluoroform (CHF_3) production from CF_3CHO photolysis and implications for the decomposition of hydrofluoroolefins and hydrochlorofluoroolefins in the atmosphere, *J. Am. Chem. Soc.*, 2025, **147**, 33.
- 14 D. Van Hoomissen, A. Chattopadhyay, S. A. Montzka and J. B. Burkholder, CHF_3 (HFC-23) and CF_3CHO quantum yields in the pulsed laser photolysis of CF_3CHO at 248, 266, 281, and 308 nm, *ACS Earth Space Chem.*, 2025, **9**, 589.
- 15 F. Baumann, C. Fernholz, J. Lelieveld and J. N. Crowley, Kinetics of the reaction of CF_3CHO with OH between 204 K and 361 K, *Phys. Chem. Chem. Phys.*, 2025, **27**, 18907.
- 16 E. C. Tuazon and R. Atkinson, Tropospheric transformation products of a series of hydrofluorocarbons and hydrochlorofluorocarbons, *J. Atmos. Chem.*, 1993, **17**, 179.
- 17 T. J. Wallington, W. F. Schneider, D. R. Worsnop, O. J. Nielsen, J. Sehested, W. J. Debruyne and J. A. Shorter, The environmental impact of CFC replacements HFCs and HCFCs, *Environ. Sci. Technol.*, 1994, **28**, 320.
- 18 A. M. Jubbe, M. R. McGillen, R. W. Portmann, J. S. Daniel and J. B. Burkholder, An atmospheric photochemical source of the persistent greenhouse gas CF_4 , *Geophys. Res. Lett.*, 2015, **42**, 9505.
- 19 J. G. Calvert, A. Mellouki, J. J. Orlando, M. J. Pilling, and T. J. Wallington, *Mechanisms of Atmospheric Oxidation of the Oxygenates*, Oxford University Press 2011.
- 20 G. D. Hayman, M. E. Jenkin, T. P. Murrells and C. E. Johnson, Tropospheric degradation chemistry of HCFC-123 (CF_3CHCl_2): A proposed replacement chlorofluorocarbon, *Atmos. Environ.*, 1994, **28**, 421.
- 21 G. A. Russell, Deuterium-isotope effects in the autoxidation of aralkyl hydrocarbons. Mechanism of the interaction of peroxy radicals, *J. Am. Chem. Soc.*, 1957, **79**, 3871.
- 22 D. A. Ellis, J. W. Martin, A. O. De Silva, S. A. Mabury, M. D. Hurley, M. P. Sulbaek Andersen and T. J. Wallington, Degradation of Fluorotelomer Alcohols: A likely atmospheric source of perfluorinated carboxylic acids, *Environ. Sci. Technol.*, 2004, **38**, 3316.
- 23 T. J. Wallington, M. D. Hurley, J. Xia, D. J. Wuebbles, S. Sillman, A. Ito, J. E. Penner, D. A. Ellis, J. Martin, S. A. Mabury, O. J. Nielsen and M. P. Sulbaek Andersen, Degradation of fluorotelomer alcohols: A likely atmospheric source of perfluorinated carboxylic acids, *Environ. Sci. Technol.*, 2006, **40**, 924.
- 24 S. Kutsuna, L. Chen, T. Abe, J. Mizukado, T. Uchimaru, K. Tokuhashi and A. Sekiya, Henry's law constants of 2,2,2-trifluoroethyl formate, ethyl trifluoroacetate, and non-fluorinated analogous esters, *Atmos. Environ.*, 2005, **39**(32), 5884–5892.
- 25 S. Kutsuna, L. Chen, K. Ohno, K. Tokuhashi and A. Sekiya, Henry's law constants and hydrolysis rate constants of 2,2,2-trifluoroethyl acetate and methyl trifluoroacetate, *Atmos. Environ.*, 2004, **38**(5), 725–732.
- 26 M. D. Hurley, J. C. Ball, T. J. Wallington, M. P. Sulbaek Andersen, O. J. Nielsen, D. A. Ellis, J. W. Martin and S. A. Mabury, Atmospheric chemistry of $n\text{-C}_x\text{F}_{2x+1}\text{CHO}$ ($x = 1, 2, 3, 4$): Fate of $n\text{-C}_x\text{F}_{2x+1}\text{C}(\text{O})$ radicals, *J. Phys. Chem. A*, 2006, **110**, 12443.
- 27 M. P. Sulbaek Andersen, J. E. Borchert, C. Blair, M. L. Robin and O. J. Nielsen, Atmospheric chemistry of (E)- and (Z)- $\text{CF}_3\text{CF}_2\text{CH}=\text{CHCF}_2\text{CF}_3$ (HFO-153-10mczz): Kinetics and mechanisms of the reactions with Cl atoms, OH radicals, and O_3 , *J. Phys. Chem. A*, 2024, **128**(47), 10167–10180.
- 28 M. P. Sulbaek Andersen, J. A. Schmidt, A. Volkova and D. J. Wuebbles, A three-dimensional model of the atmospheric chemistry of E and Z- $\text{CF}_3\text{CH}=\text{CHCl}$ (HCFO-1233(zd) (E/Z)), *Atmos. Environ.*, 2018, **179**, 250–259.
- 29 T. J. Wallington, M. D. Hurley, J. M. Fracheboud, J. J. Orlando, G. S. Tyndall, J. Sehested, T. E. Møgelberg and O. J. Nielsen, Role of excited CF_3CFHO Radicals in the atmospheric chemistry of HFC-134a, *J. Phys. Chem.*, 1996, **100**(46), 18116–18122.
- 30 T. J. Wallington, M. P. Sulbaek Andersen and O. J. Nielsen, Atmospheric Chemistry of Halogenated Organic Compounds, in *Advances in Atmospheric Chemistry*, J. J. R. Barker, A. L. Steiner, and T. J. Wallington, World Scientific Publishing, 2017, vol. 1, ch. 5, pp. 305–402.
- 31 T. E. Møgelberg, J. Sehested, G. S. Tyndall, J. J. Orlando, J.-M. Fracheboud and T. J. Wallington, Atmospheric chemistry of HFC-236cb: Fate of the alkoxy radical $\text{CF}_3\text{CF}_2\text{CFHO}$, *J. Phys. Chem. A*, 1997, **101**(15), 2828–2832.
- 32 M. P. Sulbaek Andersen, C. Stensby, O. J. Nielsen, M. D. Hurley, J. C. Ball, T. J. Wallington, J. W. Martin, D. A. Ellis and S. A. Mabury, Atmospheric Chemistry of $n\text{-C}_x\text{F}_{2x+1}\text{CHO}$ ($x = 1, 3, 4$): Mechanism of the $\text{C}_x\text{F}_{2x+1}\text{C}(\text{O})\text{O}_2 + \text{HO}_2$ Reaction, *J. Phys. Chem. A*, 2004, **108**, 6325.
- 33 G. S. Tyndall, R. A. Cox, C. Granier, R. Lesclaux, G. K. Moortgat, M. J. Pilling, A. R. Ravishankara and T. J. Wallington, Atmospheric chemistry of small organic peroxy radicals, *J. Geophys. Res.*, 2001, **106**, 12157–12182.
- 34 T. J. Wallington, J. Sehested and O. J. Nielsen, Atmospheric chemistry of $\text{CF}_3\text{C}(\text{O})\text{O}_2$ radicals. Kinetics of their reaction with NO_2 and kinetics of the thermal decomposition of the product $\text{CF}_3\text{C}(\text{O})\text{O}_2\text{NO}_2$, *Chem. Phys. Lett.*, 1994, **226**, 563.
- 35 J. Salas, A. L. Cardona, M. A. B. Paci and F. E. Malanca, Trifluoroacetyl peroxyxynitrate ($\text{CF}_3\text{C}(\text{O})\text{O}_2\text{NO}_2$, FPAN) degradation in wet atmosphere, *Atmos. Environ.*, 2024, **327**, 120780.
- 36 O. J. Nielsen and M. P. Sulbaek Andersen, Trifluoroacetic acid production from trifluoroacetyl peroxyxynitrate degradation in the presence of water, *Atmos. Pollut. Res.*, 2025, DOI: [10.1016/j.apr.2025.102828](https://doi.org/10.1016/j.apr.2025.102828).
- 37 B. Long, Y. Xia and D. G. Truhlar, Quantitative Kinetics of HO_2 Reactions with Aldehydes in the Atmosphere: High-Order Dynamic Correlation, Anharmonicity, and Falloff Effects Are All Important, *J. Am. Chem. Soc.*, 2022, **114**, 19910–19920.



- 38 UNEP, *Report of the Scientific Assessment Panel in response to Decision XXXV/7: Emissions of HFC-23 (Lead authors: S. A. Montzka, J. B. Burkholder)*, UNEP, 2024, https://ozone.unep.org/system/files/documents/SAP_Report_on_HFC23_September2024.pdf.
- 39 IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, *Datasheet PF3*, <https://iupac-aeris.ipsl.fr/datasheets/pdf/PF3.pdf>, accessed December 31, 2025.
- 40 M. S. Chiappero, F. E. Malanca, G. A. Argüello, S. T. Wooldridge, M. D. Hurley, J. C. Ball, T. J. Wallington, R. L. Waterland and R. C. Buck, Atmospheric chemistry of perfluoroaldehydes ($C_xF_{2x+1}CHO$) and fluorotelomer aldehydes ($C_xF_{2x+1}CH_2CHO$): Quantification of the important role of photolysis, *J. Phys. Chem. A*, 2006, **110**(43), 11944–11953.
- 41 S. Madronich and G. Weller, Numerical integration errors in calculated tropospheric photodissociation rate coefficients, *J. Atmos. Chem.*, 1990, **10**, 289–300.
- 42 J. B. Burkholder, S. P. Sander, J. Abbatt, J. R. Barker, C. Cappa, J. D. Crouse, T. S. Dibble, R. E. Huie, C. E. Kolb, M. J. Kurylo, V. L. Orkin, C. J. Percival, D. M. Wilmouth and P. H. Wine, *Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation No. 19*, JPL Publication 19-5, Jet Propulsion Laboratory, Pasadena, 2019, <http://jpldataeval.jpl.nasa.gov>.
- 43 National Oceanic and Atmospheric Administration, *National Aeronautics and Space Administration and United States Air Force, U.S. Standard Atmosphere, 1976*, U.S. Government Printing Office, Washington, D.C., 1976, p. 227.
- 44 IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, *Datasheet PF6*, <https://iupac-aeris.ipsl.fr/datasheets/pdf/PF6.pdf>, accessed December 31, 2025.
- 45 IUPAC Task Group on Atmospheric Chemical Kinetic Data Evaluation, *Datasheet PF7*, <https://iupac-aeris.ipsl.fr/datasheets/pdf/PF7.pdf>, accessed December 31, 2025.
- 46 M. Antinolo, E. Jimenez, S. Gonzalez and J. Albaladejo, Atmospheric chemistry of CF_3CF_2CHO : Absorption cross sections in the UV and IR regions, photolysis at 308 nm, and gas-phase reaction with OH radicals ($T = 263\text{--}358\text{ K}$), *J. Phys. Chem. A*, 2014, **118**, 178.
- 47 G. Solignac, A. Mellouki, G. Le Bras, M. Yuqing and H. Sidebottom, The gas phase tropospheric removal of fluoroaldehydes ($C_xF_{2x+1}CHO$, $x = 3, 4, 6$), *Phys. Chem. Chem. Phys.*, 2007, **9**, 4200.
- 48 M. P. Sulbaek Andersen, A. Toft, O. J. Nielsen, M. D. Hurley, T. J. Wallington, H. Chishima, K. Tonokura, S. A. Mabury, J. W. Martin and D. A. Ellis, Atmospheric chemistry of perfluorinated aldehyde hydrates ($n\text{-}C_xF_{2x+1}CH(OH)_2$, $x = 1, 3, 4$): hydration, dehydration, and kinetics and mechanism of Cl atom and OH radical initiated oxidation, *J. Phys. Chem. A*, 2006, **110**, 9854.
- 49 O. J. Nielsen, M. P. Sulbaek Andersen and J. Frankin, Comment on “Assessing the atmospheric fate of trifluoroacetaldehyde (CF_3CHO) and its potential as a new source of fluoroform (HFC-23) using the AtChem2 box model” by Pérez-Peña et al. *Environ. Sci.: Atmos.*, 2023, **3**, 1767–1777, DOI: 10.1039/D3EA00120B,, *Environ. Sci.: Atmos.*, 2025, **5**, 530–534, DOI: [10.1039/D4EA00123K](https://doi.org/10.1039/D4EA00123K).
- 50 J. H. Seinfeld and S. N. Pandis, *Atmospheric Chemistry and Physics: from Air Pollution to Climate Change*. 2nd edn, John Wiley & Sons, New York, 2006.
- 51 CARB, *Toxic Air Contaminant Factsheet*, 1997, <https://ww2.arb.ca.gov/sites/default/files/classic/toxics/tac/factshts1997/nitricac.pdf>, accessed February 1, 2026.
- 52 M. K. Vollmer, T. S. Rhee, M. Rigby, D. Hofstetter, M. Hill, F. Schoenenberger and S. Reimann, Modern inhalation anesthetics: Potent greenhouse gases in the global atmosphere, *Geophys. Res. Lett.*, 2015, **42**, 1606–1611.

